

# 4 QF IV: Definiteness of Quadratic Forms

Prof.Dr.P.M.Bajracharya

School of Mathematical Sciences

T.U., Kirtipur

April 28, 2024

# Summary

- ① Classification of Quadratic Forms
- ② Constrained optimization

# Classification of Quadratic Forms

- The taxonomy of conic sections into ellipses, hyperbolas, and parabolas, as well as lines and points, is very useful for quadratic forms in 2 variables.

- 4 / 49

- 4 / 49

Let  $A$  be an  $n \times n$  symmetric matrix, and recall that  $Q(x) = x^T A x$  is the corresponding quadratic form. Then  $A$  (as well as  $Q$ ) is

- (a) **positive definite** if  $x^T Ax > 0$ , for all  $x \neq 0$
- (b) **negative definite** if  $x^T Ax < 0$ , for all  $x \neq 0$
- (c) **indefinite** if  $x^T Ax > 0$  for some  $x$ 's and  $x^T Ax < 0$  for others.
- (d) **positive semidefinite** if  $x^T Ax \geq 0$ , for all  $x \neq 0$
- (e) **negative semidefinite** if  $x^T Ax \leq 0$ , for all  $x \neq 0$ .








**Example 3.** The following QF is *indefinite*:


$$f(x, y) = x^2 + xy - y^2$$

because  $f(1, 1) = 1 > 0$ , but  $f(-1, 1) = -1 < 0$ .



$$f(x, y) = x^2 + 2xy + y^2$$

because  $f(x, y) = (x + y)^2 \geq 0$  for all  $x, y \in \mathbb{R}$ . Observe that  $f(x, y) = 0$  whenever  $x = -y$ .





Let  $A$  be a  $2 \times 2$  symmetric matrix. Then a quadratic form  $x^T A x$  is

- (a) positive definite if and only if the eigenvalues of  $A$  are both positive,
- (b) negative definite if and only if the eigenvalues of  $A$  are both negative,
- (c) indefinite if and only if  $A$  has both positive and negative eigenvalues,

- (d) positive semidefinite if and only if the eigenvalues of  $A$  are nonnegative,
- (e) negative semidefinite if and only if the eigenvalues of  $A$  are nonpositive.

Consequently, if  $|A| = 0$ , then the quadratic form is neither positive definite nor negative definite.

## Note that

This theorem can be generalized for an  $n \times n$  matrix.

## Example.

Determine the definiteness of the quadratic form  $Q(x) = x_1^2 + 2x_1x_2 + x_2^2$ .



**Solution.** This form can be written as

$$Q(x) = x^T \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} x.$$

**Solution.** This form can be written as

$$Q(x) = x^T \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} x.$$

Then we have

$$|A - \lambda I| = \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix}$$

**Solution.** This form can be written as

$$Q(x) = x^T \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} x.$$

Then we have

$$\begin{aligned} |A - \lambda I| &= \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix} \\ &= (1 - \lambda)^2 - 1 = \lambda(\lambda - 2). \end{aligned}$$

**Solution.** This form can be written as

$$Q(x) = x^T \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} x.$$

Then we have

$$\begin{aligned} |A - \lambda I| &= \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix} \\ &= (1 - \lambda)^2 - 1 = \lambda(\lambda - 2). \end{aligned}$$

Clearly, the eigenvalues of  $A$  are  $\lambda_1 = 2$  and  $\lambda_2 = 0$ . Hence the quadratic form is positive semi-definite.

## Example.

For which real numbers  $k$  is the quadratic form  $Q(x) = kx_1^2 - 6x_1x_2 + kx_2^2$  positive semi-definite?

**Example.**

For which real numbers  $k$  is the quadratic form  $Q(x) = kx_1^2 - 6x_1x_2 + kx_2^2$  positive semi-definite?

**Solution.** To determine the definiteness of this form we'll need to consider the matrix

$$\begin{pmatrix} k & -3 \\ -3 & k \end{pmatrix},$$

## Solution...

Its characterstic polynomial is

$$|A - \lambda I| = \begin{pmatrix} k - \lambda & -3 \\ -3 & k - \lambda \end{pmatrix}$$

## Solution...

Its characterstic polynomial is

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} k - \lambda & -3 \\ -3 & k - \lambda \end{vmatrix} \\ &= (k - \lambda)^2 - 9\end{aligned}$$



## Solution...

Its characterstic polynomial is

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} k - \lambda & -3 \\ -3 & k - \lambda \end{vmatrix} \\ &= (k - \lambda)^2 - 9 \\ &= (k - \lambda + 3)(k - \lambda - 3).\end{aligned}$$

## Solution...

Its characterstic polynomial is

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} k - \lambda & -3 \\ -3 & k - \lambda \end{vmatrix} \\ &= (k - \lambda)^2 - 9 \\ &= (k - \lambda + 3)(k - \lambda - 3).\end{aligned}$$

Clearly, the eigenvalues of  $A$  are  $\lambda_1 = k + 3$  and  $\lambda_2 = k - 3$ . In order for  $Q$  to be positive definite, both of these eigenvalues must be positive.

## Solution...

Its characterstic polynomial is

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} k - \lambda & -3 \\ -3 & k - \lambda \end{vmatrix} \\ &= (k - \lambda)^2 - 9 \\ &= (k - \lambda + 3)(k - \lambda - 3).\end{aligned}$$

Clearly, the eigenvalues of  $A$  are  $\lambda_1 = k + 3$  and  $\lambda_2 = k - 3$ . In order for  $Q$  to be positive definite, both of these eigenvalues must be positive. So  $k > 3$  is a necessary and sufficient condition for  $Q$  to be a positive definite quadratic form.

# Constrained optimization

Let  $x \in \mathbb{R}^n$  and  $A$  be an  $n \times n$  matrix. A quadratic form  $Q(x) = x^T A x$  may not have a maximum: given any vector  $x$ , we can increase the value of  $Q(x)$  by considering the vector  $x' = 2x$ .

Let  $x \in \mathbb{R}^n$  and  $A$  be an  $n \times n$  matrix. A quadratic form  $Q(x) = x^T A x$  may not have a maximum: given any vector  $x$ , we can increase the value of  $Q(x)$  by considering the vector  $x' = 2x$ .

The way to avoid this issue is to “constrain”  $x \in \mathbb{R}^n$ .

Let  $x \in \mathbb{R}^n$  and  $A$  be an  $n \times n$  matrix. A quadratic form  $Q(x) = x^T A x$  may not have a maximum: given any vector  $x$ , we can increase the value of  $Q(x)$  by considering the vector  $x' = 2x$ .

The way to avoid this issue is to “constrain”  $x \in \mathbb{R}^n$ . There are many ways to constrain  $x$ , but the one that leads to the spectral decomposition is to constrain  $x$  to the unit ball in  $\mathbb{R}^n$ .

Let  $x \in \mathbb{R}^n$  and  $A$  be an  $n \times n$  matrix. A quadratic form  $Q(x) = x^T A x$  may not have a maximum: given any vector  $x$ , we can increase the value of  $Q(x)$  by considering the vector  $x' = 2x$ .

The way to avoid this issue is to “constrain”  $x \in \mathbb{R}^n$ . There are many ways to constrain  $x$ , but the one that leads to the spectral decomposition is to constrain  $x$  to the unit ball in  $\mathbb{R}^n$ . That is, we try to solve the problem

## Example

Find  $x \in \mathbb{R}^n$  to attain  $\max x^T A x$  subject to  $x^T x = 1$ .



The requirement that a vector  $x$  in  $\mathbb{R}^n$  be a unit vector can be stated in several equivalent ways:


$$\|x\| = 1, \quad \|x\|^2 = 1, \quad x^T x = 1$$

and

$$x_1^2 + x_2^2 + \cdots + x_n^2 = 1. \quad (1)$$

The expanded version (1) of  $x^T x = 1$  is commonly used in applications.

Geometrically, the problem of finding the maximum and minimum values of  $x^T x$  subject to the requirement  $x^T x = 1$  amounts to finding the highest and lowest points on the intersection of the surface with the right circular cylinder determined by the unit circle. See the figure below.



When a quadratic form  $Q$  has no cross-product terms, it is easy to find the maximum and minimum of  $Q(x)$  for  $x^T x = 1$ .

When a quadratic form  $Q$  has no cross-product terms, it is easy to find the maximum and minimum of  $Q(x)$  for  $x^T x = 1$ .

### Example.

Find the maximum and minimum values of

$$Q(x) = 9x_1^2 + 4x_2^2 + 3x_3^2$$

subject to the constraint  $x^T x = 1$ .

**Solution.** Since  $x_2^2$  and  $x_3^2$  are nonnegative, observe that

$$4x_2^2 \leq 9x_2^2 \quad 3x_3^2 \leq 9x_3^2$$

**Solution.** Since  $x_2^2$  and  $x_3^2$  are nonnegative, observe that

$$4x_2^2 \leq 9x_2^2 \quad 3x_3^2 \leq 9x_3^2$$

and hence

$$Q(x) = 9x_1^2 + 4x_2^2 + 3x_3^2$$

**Solution.** Since  $x_2^2$  and  $x_3^2$  are nonnegative, observe that

$$4x_2^2 \leq 9x_2^2 \quad 3x_3^2 \leq 9x_3^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 9x_1^2 + 9x_2^2 + 9x_3^2 \end{aligned}$$



**Solution.** Since  $x_2^2$  and  $x_3^2$  are nonnegative, observe that

$$4x_2^2 \leq 9x_2^2 \quad 3x_3^2 \leq 9x_3^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 9x_1^2 + 9x_2^2 + 9x_3^2 \\ &= 9(x_1^2 + x_2^2 + x_3^2) \end{aligned}$$

**Solution.** Since  $x_2^2$  and  $x_3^2$  are nonnegative, observe that

$$4x_2^2 \leq 9x_2^2 \quad 3x_3^2 \leq 9x_3^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 9x_1^2 + 9x_2^2 + 9x_3^2 \\ &= 9(x_1^2 + x_2^2 + x_3^2) \\ &= 9x^T x = 9. \end{aligned}$$

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector.

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

To find the minimum value of  $Q(x)$ , we observe that

$$9x_1^2 \geq 3x_1^2 \quad 4x_2^2 \geq 3x_2^2$$

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

To find the minimum value of  $Q(x)$ , we observe that

$$9x_1^2 \geq 3x_1^2 \quad 4x_2^2 \geq 3x_2^2$$

and hence

$$Q(x) = 9x_1^2 + 4x_2^2 + 3x_3^2$$

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

To find the minimum value of  $Q(x)$ , we observe that

$$9x_1^2 \geq 3x_1^2 \quad 4x_2^2 \geq 3x_2^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 3x_1^2 + 3x_2^2 + 3x_3^2 \end{aligned}$$

So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

To find the minimum value of  $Q(x)$ , we observe that

$$9x_1^2 \geq 3x_1^2 \quad 4x_2^2 \geq 3x_2^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 3x_1^2 + 3x_2^2 + 3x_3^2 \\ &= 3(x_1^2 + x_2^2 + x_3^2) \end{aligned}$$



So the maximum value of  $Q(x)$  cannot exceed 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 9$  when  $x = (1, 0, 0)$ . Thus, 9 is the maximum value of  $Q(x)$  for  $x^T x = 1$ .

To find the minimum value of  $Q(x)$ , we observe that

$$9x_1^2 \geq 3x_1^2 \quad 4x_2^2 \geq 3x_2^2$$

and hence

$$\begin{aligned} Q(x) &= 9x_1^2 + 4x_2^2 + 3x_3^2 \\ &\leq 3x_1^2 + 3x_2^2 + 3x_3^2 \\ &= 3(x_1^2 + x_2^2 + x_3^2) \\ &= 3x^T x = 3. \end{aligned}$$

So the minimum value of  $Q(x)$  cannot be less than 9 when  $x$  is a unit vector.

So the minimum value of  $Q(x)$  cannot be less than 9 when  $x$  is a unit vector. Furthermore,  $Q(x) = 3$  when  $x = (1, 0, 0)$ . Thus, 3 is the minimum value of  $Q(x)$  for  $x^T x = 1$ . □

It is easy to see in the above example that the matrix of the quadratic form  $Q$  has eigenvalues 9, 4, and 3 and that the greatest and least eigenvalues equal, respectively, the (constrained) maximum and minimum of  $Q(x)$ .

It is easy to see in the above example that the matrix of the quadratic form  $Q$  has eigenvalues 9, 4, and 3 and that the greatest and least eigenvalues equal, respectively, the (constrained) maximum and minimum of  $Q(x)$ . The same holds true for any quadratic form, as well. This is what the next theorem claims.

## Theorem

Let  $A$  be a symmetric matrix, and

$$m = \min\{x^T Ax : \|x\| = 1\}, \quad M = \max\{x^T Ax : \|x\| = 1\}$$

Then  $M$  is the greatest eigenvalue  $\lambda_1$  of  $A$  and  $m$  is the least eigenvalue of  $A$ . The value of  $x^T Ax$  is  $M$  when  $x$  is a unit eigenvector  $u_1$  corresponding to  $M$ . The value of  $x^T Ax$  is  $m$  when  $x$  is a unit eigenvector corresponding to  $m$ .

**Proof.** Orthogonally diagonalize  $A$  as  $V\Lambda V^T$ .  
We observe that if  $x = Vy$ , then

$$x^T Ax = y^T \Lambda y$$

**Proof.** Orthogonally diagonalize  $A$  as  $V\Lambda V^T$ .  
We observe that if  $x = Vy$ , then

$$x^T Ax = y^T \Lambda y$$

and for all  $y$ ,

$$\|x\| = \|Vy\| = \|y\|,$$



**Proof.** Orthogonally diagonalize  $A$  as  $V\Lambda V^T$ .  
We observe that if  $x = Vy$ , then

$$x^T Ax = y^T \Lambda y$$

and for all  $y$ ,

$$\|x\| = \|Vy\| = \|y\|,$$

because  $V^T V = I$  and

$$\|Vy\| = (Vy)^T (Vy) = y^T V^T V y = y^T y = \|y\|^2.$$

**Proof.** Orthogonally diagonalize  $A$  as  $V\Lambda V^T$ .  
We observe that if  $x = Vy$ , then

$$x^T Ax = y^T \Lambda y$$

and for all  $y$ ,

$$\|x\| = \|Vy\| = \|y\|,$$

because  $V^T V = I$  and

$$\|Vy\|^2 = (Vy)^T (Vy) = y^T V^T V y = y^T y = \|y\|^2.$$

In particular,  $\|y\| = 1$  if and only if  $\|x\| = 1$ .

**Proof.** Orthogonally diagonalize  $A$  as  $V\Lambda V^T$ .  
We observe that if  $x = Vy$ , then

$$x^T Ax = y^T \Lambda y$$

and for all  $y$ ,

$$\|x\| = \|Vy\| = \|y\|,$$

because  $V^T V = I$  and

$$\|Vy\|^2 = (Vy)^T (Vy) = y^T V^T V y = y^T y = \|y\|^2.$$

In particular,  $\|y\| = 1$  if and only if  $\|x\| = 1$ .  
Thus  $x^T Ax$  and  $y^T \Lambda y$  assume the same set of values as  $x$  and  $y$  range over the set of all unit vectors.

**Proof...** To simplify notation, suppose that  $A$  is a  $3 \times 3$  matrix with eigenvalues  $a \geq b \geq c$ .

**Proof...** To simplify notation, suppose that  $A$  is a  $3 \times 3$  matrix with eigenvalues  $a \geq b \geq c$ . Arrange the (eigenvector) columns of  $V$  so that  $P = (u_1 \ u_2 \ u_3)$  and

$$\Lambda = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}.$$

Proof... We have

$$y^T \Lambda y = ay_1^2 + by_2^2 + cy_3^2$$

Proof... We have

$$\begin{aligned}y^T \Lambda y &= ay_1^2 + by_2^2 + cy_3^2 \\ &\leq ay_1^2 + ay_2^2 + ay_3^2\end{aligned}$$

**Proof...** We have

$$\begin{aligned}y^T \Lambda y &= ay_1^2 + by_2^2 + cy_3^2 \\&\leq ay_1^2 + ay_2^2 + ay_3^2 \\&= a(y_1^2 + y_2^2 + y_3^2)\end{aligned}$$



**Proof...** We have

$$\begin{aligned}y^T \Lambda y &= ay_1^2 + by_2^2 + cy_3^2 \\&\leq ay_1^2 + ay_2^2 + ay_3^2 \\&= a(y_1^2 + y_2^2 + y_3^2) \\&= \|y\|^2 = a.\end{aligned}$$

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .

However,  $y^T \Lambda y = a$  when  $y = e_1 = (1, 0, 0)$ , so in fact  $M = a$ .

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .  
 However,  $y^T \Lambda y = a$  when  $y = e_1 = (1, 0, 0)$ , so in fact  $M = a$ . Also, the  $x$  that corresponds to  $y = e_1$  is the eigenvector  $u_1$  of  $A$ ,

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .

However,  $y^T \Lambda y = a$  when  $y = e_1 = (1, 0, 0)$ , so in fact  $M = a$ . Also, the  $x$  that corresponds to  $y = e_1$  is the eigenvector  $u_1$  of  $A$ , because

$$x = V e_1 = (u_1 \ u_2 \ u_3) \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = u_1$$

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .  
However,  $y^T \Lambda y = a$  when  $y = e_1 = (1, 0, 0)$ , so in fact  $M = a$ . Also, the  $x$  that corresponds to  $y = e_1$  is the eigenvector  $u_1$  of  $A$ , because

$$x = V e_1 = (u_1 \ u_2 \ u_3) \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = u_1$$

Thus

$$M = a = e_1^T \Lambda e_1 = u_1^T A u_1,$$

which proves the statement about  $M$ .

**Proof...** Thus  $M \leq a$ , by definition of  $M$ .  
However,  $y^T \Lambda y = a$  when  $y = e_1 = (1, 0, 0)$ , so in fact  $M = a$ . Also, the  $x$  that corresponds to  $y = e_1$  is the eigenvector  $u_1$  of  $A$ , because

$$x = Ve_1 = (u_1 \ u_2 \ u_3) \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = u_1$$

Thus

$$M = a = e_1^T \Lambda e_1 = u_1^T A u_1,$$

which proves the statement about  $M$ . A similar argument shows that  $m$  is the least eigenvalue and this value of  $x^T A x$  is attained when

$$x = Ve_3 = u_3.$$

## Example

Find the maximum and minimum values of

$$Q(x) = 5x_1^2 + 5x_2^2 + 4x_1x_2$$

subject to the constraint  $x_1^2 + x_2^2 = 1$ .

**Solution.** The given quadratic form can be expressed in matrix notation as

$$Q(x) = (x_1 \ x_2) \begin{pmatrix} 5 & 2 \\ 2 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x^T A x.$$



**Solution...** To find the eigenvalues of  $A$ , we have

$$|A - \lambda I| = \begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix}$$

**Solution...** To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix} \\ &= (5 - \lambda)^2 - 4\end{aligned}$$

**Solution...** To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix} \\ &= (5 - \lambda)^2 - 4 \\ &= (7 - \lambda)(3 - \lambda).\end{aligned}$$

**Solution...** To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix} \\ &= (5 - \lambda)^2 - 4 \\ &= (7 - \lambda)(3 - \lambda).\end{aligned}$$

Therefore, the eigenvalues of  $A$  are

$$\lambda_1 = 7, \lambda_2 = 3.$$

**Solution...** To find the eigenvector associated with  $\lambda_1 = 7$ , we have

$$\begin{pmatrix} -2 & 2 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

**Solution...** To find the eigenvector associated with  $\lambda_1 = 7$ , we have

$$\begin{pmatrix} -2 & 2 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

This gives the equation

$$x_1 - x_2 = 0.$$

**Solution...** To find the eigenvector associated with  $\lambda_1 = 7$ , we have

$$\begin{pmatrix} -2 & 2 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

This gives the equation

$$x_1 - x_2 = 0.$$

So, the eigenvector associated with  $\lambda_1 = 7$  is

$$v_1 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

**Solution...** To find the eigenvector associated with  $\lambda_2 = 3$ , we have

$$\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$



**Solution...** To find the eigenvector associated with  $\lambda_2 = 3$ , we have

$$\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

This gives the equation

$$x_1 + x_2 = 0.$$

**Solution...** So, the eigenvector associated with  $\lambda_2 = 3$  is

$$v_2 = \begin{pmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

**Solution...** So, the eigenvector associated with  $\lambda_2 = 3$  is

$$v_2 = \begin{pmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

Thus, the constrained extrema are  
Constrained maximum:

$$Q(x) = 7 \text{ at } (x_1, x_2) = (1/\sqrt{2}, 1/\sqrt{2})$$

Constrained minimum:

$$Q(x) = 3 \text{ at } (x_1, x_2) = (-1/\sqrt{2}, 1/\sqrt{2}).$$



## Example

A rectangle is to be inscribed in the ellipse

$$4x^2 + 9y^2 = 36.$$

Use eigenvalue methods to find nonnegative values of  $x$  and  $y$  that produce the inscribed rectangle with maximum area.

**Solution.** The area  $z$  of the inscribed rectangle is given by

$$z = 4xy,$$

**Solution.** The area  $z$  of the inscribed rectangle is given by

$$z = 4xy,$$

so the problem is to maximize the quadratic form  $z = 4xy$  subject to the constraint  $4x^2 + 9y^2 = 36$ .

**Solution.** The area  $z$  of the inscribed rectangle is given by

$$z = 4xy,$$

so the problem is to maximize the quadratic form  $z = 4xy$  subject to the constraint  $4x^2 + 9y^2 = 36$ . In this problem, the graph of the constraint equation is an ellipse rather than the unit circle as required in the theorem,

**Solution.** The area  $z$  of the inscribed rectangle is given by

$$z = 4xy,$$

so the problem is to maximize the quadratic form  $z = 4xy$  subject to the constraint  $4x^2 + 9y^2 = 36$ . In this problem, the graph of the constraint equation is an ellipse rather than the unit circle as required in the theorem, but we can remedy this problem by rewriting the constraint as

$$(x/3)^2 + (y/2)^2 = 1$$



**Solution.** The area  $z$  of the inscribed rectangle is given by

$$z = 4xy,$$

so the problem is to maximize the quadratic form  $z = 4xy$  subject to the constraint  $4x^2 + 9y^2 = 36$ . In this problem, the graph of the constraint equation is an ellipse rather than the unit circle as required in the theorem, but we can remedy this problem by rewriting the constraint as


$$(x/3)^2 + (y/2)^2 = 1$$

and defining new variables,  $x_1$  and  $y_1$ , by the equations

$$x = 3x_1 \text{ and } y = 2y_1.$$

This enables us to reformulate the problem as follows:

Maximize  $z = 4xy = 24x_1y_1$   
subject to the constraint  $x_1^2 + y_1^2$ .



To solve this problem, we will write the quadratic form  $z = 24x_1y_1$  as

$$z = x^T A x = [x_1 \ y_1] \begin{pmatrix} 0 & 12 \\ 12 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

To find the eigenvalues of  $A$ , we have

$$|A - \lambda I| = \begin{vmatrix} -\lambda & 12 \\ 12 & -\lambda \end{vmatrix}$$

To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} -\lambda & 12 \\ 12 & -\lambda \end{vmatrix} \\ &= 12 - \lambda\end{aligned}$$

To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} -\lambda & 12 \\ 12 & -\lambda \end{vmatrix} \\ &= 12 - \lambda\end{aligned}$$

This shows that the eigenvalues of  $A$  is  $\lambda = 12$ .

To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} -\lambda & 12 \\ 12 & -\lambda \end{vmatrix} \\ &= 12 - \lambda\end{aligned}$$

This shows that the eigenvalues of  $A$  is  $\lambda = 12$ .  
To find the eigenvector associated with  $\lambda = 12$ ,  
we have

$$\begin{pmatrix} -12 & 12 \\ 12 & -12 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

To find the eigenvalues of  $A$ , we have

$$\begin{aligned}|A - \lambda I| &= \begin{vmatrix} -\lambda & 12 \\ 12 & -\lambda \end{vmatrix} \\ &= 12 - \lambda\end{aligned}$$

This shows that the eigenvalues of  $A$  is  $\lambda = 12$ .  
To find the eigenvector associated with  $\lambda = 12$ ,  
we have

$$\begin{pmatrix} -12 & 12 \\ 12 & -12 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0.$$

This gives the equation

$$x_1 - x_2 = 0.$$



So, the eigenvector associated with  $\lambda = 12$  is

$$v = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

So, the eigenvector associated with  $\lambda = 12$  is

$$v = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

Thus, the maximum area is  $z = 12$ , and this occurs at  $(x_1, y_1) = (1/\sqrt{2}, 1/\sqrt{2})$ . □

## Theorem

Let  $A$ ,  $\lambda_1$ , and  $u_1$  be as in the earlier theorem. Then the maximum value of  $x^T A x$  subject to the constraints

$$x^T x = 1, \quad x^T u_1 = 0$$

is the second greatest eigenvalue,  $\lambda_2$ , and this maximum is attained when  $x$  is an eigenvector  $u_2$  corresponding to  $\lambda_2$ .

This theorem can be proved by an argument similar to the one above in which the theorem is reduced to the case where the matrix of the quadratic form is diagonal.

This theorem can be proved by an argument similar to the one above in which the theorem is reduced to the case where the matrix of the quadratic form is diagonal. The next example gives an idea of the proof for the case of a diagonal matrix.

## Problem

Find the maximum value of

$$Q(x) = 9x_1^2 + 4x_2^2 + 3x_3^2$$

subject to the constraint  $x^T x = 1$  and  $x^T u_1 = 0$ ,  
where  $u_1 = (1, 0, 0)$ .

Here  $u_1$  is a unit eigenvector corresponding to the  
greatest eigenvalue  $\lambda = 9$  of the matrix of the  
quadratic form.

$$x_1 = 0.$$

**Solution.** If the coordinates of  $x$  are  $x_1, x_2, x_3$ , then the constraint  $x^T u_1 = 0$  means simply that  $x_1 = 0$ . For such a unit vector,  $x_2^2 + x_3^2 = 1$ , and

$$9x_1^2 + 4x_2^2 + 3x_3^2 = 4x_2^2 + 3x_3^2$$



**Solution.** If the coordinates of  $x$  are  $x_1, x_2, x_3$ , then the constraint  $x^T u_1 = 0$  means simply that  $x_1 = 0$ . For such a unit vector,  $x_2^2 + x_3^2 = 1$ , and

$$\begin{aligned} 9x_1^2 + 4x_2^2 + 3x_3^2 &= 4x_2^2 + 3x_3^2 \\ &\leq 4(x_2^2 + x_3^2) \end{aligned}$$

**Solution.** If the coordinates of  $x$  are  $x_1, x_2, x_3$ , then the constraint  $x^T u_1 = 0$  means simply that  $x_1 = 0$ . For such a unit vector,  $x_2^2 + x_3^2 = 1$ , and

$$\begin{aligned} 9x_1^2 + 4x_2^2 + 3x_3^2 &= 4x_2^2 + 3x_3^2 \\ &\leq 4(x_2^2 + x_3^2) \\ &= 4. \end{aligned}$$

**Solution.** If the coordinates of  $x$  are  $x_1, x_2, x_3$ , then the constraint  $x^T u_1 = 0$  means simply that  $x_1 = 0$ . For such a unit vector,  $x_2^2 + x_3^2 = 1$ , and

$$\begin{aligned} 9x_1^2 + 4x_2^2 + 3x_3^2 &= 4x_2^2 + 3x_3^2 \\ &\leq 4(x_2^2 + x_3^2) \\ &= 4. \end{aligned}$$

Thus the constrained maximum of the quadratic form does not exceed 4.

**Solution.** If the coordinates of  $x$  are  $x_1, x_2, x_3$ , then the constraint  $x^T u_1 = 0$  means simply that  $x_1 = 0$ . For such a unit vector,  $x_2^2 + x_3^2 = 1$ , and

$$\begin{aligned} 9x_1^2 + 4x_2^2 + 3x_3^2 &= 4x_2^2 + 3x_3^2 \\ &\leq 4(x_2^2 + x_3^2) \\ &= 4. \end{aligned}$$

Thus the constrained maximum of the quadratic form does not exceed 4. And this value is attained for  $x_1 = (0, 1, 0)$ , which is an eigenvector for the second greatest eigenvalue of the matrix of the quadratic form. □

The next theorem generalizes the previous theorem and gives a useful characterization of all the eigenvalues of  $A$ . The proof is omitted.

## Theorem

Let  $A$  be a symmetric  $n \times n$  matrix with an orthogonal diagonalization  $A = P\Lambda P^{-1}$ , where the entries on the diagonal of  $\Lambda$  are arranged so that  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$  and where the columns of  $P$  are corresponding unit eigenvectors  $u_1, u_2, \dots, u_n$ . Then for  $k = 2, \dots, n$ , the maximum value of  $x^T A x$  subject to the constraints

$$x^T x = 1, \quad x^T u_1 = 0, \dots, x^T u_{k-1} = 0$$

is the eigenvalue  $\lambda_k$ , and this maximum is attained at  $x = u_k$ .