

Unit 3: Spectral theory III

Prof.Dr.P.M.Bajracharya

School of Mathematical Sciences

T.U., Kirtipur

April 15, 2024

Summary

① Spectral decompositions

Spectral decompositions

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$.

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$.

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 3 - \lambda & 2 \\ 4 & 1 - \lambda \end{pmatrix}.$$

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 3 - \lambda & 2 \\ 4 & 1 - \lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = (3 - \lambda)(1 - \lambda) - 8$$

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 3 - \lambda & 2 \\ 4 & 1 - \lambda \end{pmatrix}.$$

It follows that

$$\begin{aligned} |A - \lambda I| &= (3 - \lambda)(1 - \lambda) - 8 \\ &= \lambda^2 - 4\lambda - 5 \end{aligned}$$

Let $A = \begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 3 - \lambda & 2 \\ 4 & 1 - \lambda \end{pmatrix}.$$

$$\begin{aligned} |A - \lambda I| &= (3 - \lambda)(1 - \lambda) - 8 \\ &= \lambda^2 - 4\lambda - 5 \\ &= (\lambda - 5)(\lambda + 1). \end{aligned}$$

Clearly, if $\lambda = 5$ or -1 , then

$$|A - \lambda I| = 0.$$

Clearly, if $\lambda = 5$ or -1 , then

$$|A - \lambda I| = 0.$$

So, the eigenvalues are $\lambda = 5, -1$.

$$|A - \lambda I| = 0.$$

$$|A - \lambda I| = 0.$$

So, the eigenvalues are $\lambda = 5, -1$. Let's compute the eigenvectors.

$$\underline{\lambda = 5}$$

$$\underline{\lambda = 5} \quad \begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\underline{\lambda = 5}$$

$$\begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow v_{\lambda=5} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

$$\underline{\lambda = -1} \quad \begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{array}{c} \underline{\lambda = 5} \\ \begin{pmatrix} -2 & 2 \\ 4 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow v_{\lambda=5} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} . \end{array}$$

$$\underline{\lambda = -1}$$

$$\begin{pmatrix} 4 & 2 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow v_{\lambda=-1} = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 \\ -2 \end{pmatrix}.$$

Observe that $v_{\lambda=5}$ and $v_{\lambda=-1}$ are linearly independent,

Observe that $v_{\lambda=5}$ and $v_{\lambda=-1}$ are linearly independent, but they are not orthogonal to each other. Thus, this matrix is an example of a matrix with two distinct eigenvalues, each of which has an associated eigenvector,

Observe that $v_{\lambda=5}$ and $v_{\lambda=-1}$ are linearly independent, but they are not orthogonal to each other. Thus, this matrix is an example of a matrix with two distinct eigenvalues, each of which has an associated eigenvector, where the two eigenvectors are linearly independent but not orthogonal.

Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix}.$$

Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = \lambda^2 + 1.$$

Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = \lambda^2 + 1.$$

Clearly, if $\lambda = \pm i$, then

$$|A - \lambda I| = 0.$$

Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = \lambda^2 + 1.$$

Clearly, if $\lambda = \pm i$, then

$$|A - \lambda I| = 0.$$

So, the eigenvalues are $\lambda = \pm i$. Let's compute the eigenvectors.

$$\underline{\lambda = i}$$

$$\frac{\lambda = i}{\begin{pmatrix} -i & 1 \\ -1 & -i \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}}$$

$$\begin{array}{c} \underline{\lambda = i} \\ \begin{pmatrix} -i & 1 \\ -1 & -i \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow v_{\lambda=i} = \begin{pmatrix} -i \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ i \end{pmatrix}. \end{array}$$

$$\lambda = -i$$

$$\frac{\lambda = -i}{\begin{pmatrix} -i & 1 \\ -1 & i \end{pmatrix}} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\frac{\lambda = -i}{\begin{pmatrix} -i & 1 \\ -1 & i \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}} \Rightarrow v_{\lambda=-i} = \begin{pmatrix} 1 \\ -i \end{pmatrix} = \begin{pmatrix} i \\ 1 \end{pmatrix}.$$

Example (An example with fewer than two eigenvectors). Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.

eigenvectors). Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ 0 & -\lambda \end{pmatrix}.$$

eigenvectors). Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ 0 & -\lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = \lambda^2.$$

Example (An example with fewer than two eigenvectors). Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} -\lambda & 1 \\ 0 & -\lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = \lambda^2.$$

Clearly, if $\lambda = 0$, then

$$|A - \lambda I| = 0.$$

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2.

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2. Let's compute the eigenvectors.

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2. Let's compute the eigenvectors. Since

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ 0 \end{pmatrix},$$

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2. Let's compute the eigenvectors. Since

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ 0 \end{pmatrix},$$

there is only one corresponding eigenvector,

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2. Let's compute the eigenvectors. Since

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ 0 \end{pmatrix},$$

there is only one corresponding eigenvector, which is

$$v_{\lambda=0} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Thus, $\lambda = 0$ is a degenerate eigenvalue with degeneracy equal to 2. Let's compute the eigenvectors. Since

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ 0 \end{pmatrix},$$

there is only one corresponding eigenvector, which is

$$v_{\lambda=0} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Thus, the given matrix is an example of a real-valued $n \times n$ matrix that has fewer than n eigenvectors.

The matrices considered above are “not nice” in the sense that they do not have a full set of n orthogonal real-valued eigenvectors and associated real-valued eigenvalues.

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$.

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{pmatrix}.$$

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{pmatrix}.$$

It follows that

$$|A - \lambda I| = (1 - \lambda)(4 - \lambda) - 4$$

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{pmatrix}.$$

It follows that

$$\begin{aligned} |A - \lambda I| &= (1 - \lambda)(4 - \lambda) - 4 \\ &= \lambda^2 - 5\lambda. \end{aligned}$$

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{pmatrix}.$$

It follows that

$$\begin{aligned} |A - \lambda I| &= (1 - \lambda)(4 - \lambda) - 4 \\ &= \lambda^2 - 5\lambda. \end{aligned}$$

Clearly, if $\lambda = 0$ or 5 , then

$$|A - \lambda I| = 0.$$

Example. Let $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$. Then

$$A - \lambda I = \begin{pmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{pmatrix}.$$

It follows that

$$\begin{aligned} |A - \lambda I| &= (1 - \lambda)(4 - \lambda) - 4 \\ &= \lambda^2 - 5\lambda. \end{aligned}$$

Clearly, if $\lambda = 0$ or 5 , then

$$|A - \lambda I| = 0.$$

So, the eigenvalues are $\lambda = 0, 5$. Let's compute the eigenvectors.

$$\underline{\lambda = 0}$$

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$.

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Then

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Then

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 + 4x_2 = 0. \end{cases}$$

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Then

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 + 4x_2 = 0. \end{cases}$$

Thus, $u_{\lambda=0} = \begin{pmatrix} -2x_2 \\ x_2 \end{pmatrix}$.

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Then

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 + 4x_2 = 0. \end{cases}$$

Thus, $u_{\lambda=0} = \begin{pmatrix} -2x_2 \\ x_2 \end{pmatrix}$. For example,

$$u_{\lambda=0} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}.$$

$\lambda = 0$ Let $u_{\lambda=0} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Then

$$\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x_1 + 2x_2 = 0 \\ 2x_1 + 4x_2 = 0. \end{cases}$$

Thus, $u_{\lambda=0} = \begin{pmatrix} -2x_2 \\ x_2 \end{pmatrix}$. For example,

$u_{\lambda=0} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$. Then the corresponding unit vector is given by

$$v_{\lambda=0} = \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix}.$$

$$\underline{\lambda = 5}$$

$$\underline{\lambda = 5}$$
$$\begin{pmatrix} -4 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\underline{\lambda = 5}$$
$$\begin{pmatrix} -4 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow u_{\lambda=5} = \begin{pmatrix} x_1 \\ 2x_1 \end{pmatrix}$$

$$\underline{\lambda = 5}$$
$$\begin{pmatrix} -4 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow u_{\lambda=5} = \begin{pmatrix} x_1 \\ 2x_1 \end{pmatrix}$$

For example, a corresponding unit vector is given by

$$v_{\lambda=5} = \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix}.$$

Note that

- The matrix A has two distinct eigenvalues.

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.
- The eigenvectors corresponding to two distinct eigenvalues are orthogonal.

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.
- The eigenvectors corresponding to two distinct eigenvalues are orthogonal. This is because

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.
- The eigenvectors corresponding to two distinct eigenvalues are orthogonal. This is because

$$u_{\lambda=0} \cdot u_{\lambda=5} = u_{\lambda=0}^T u_{\lambda=5}$$

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.
- The eigenvectors corresponding to two distinct eigenvalues are orthogonal. This is because

$$u_{\lambda=0} \cdot u_{\lambda=5} = u_{\lambda=0}^T u_{\lambda=5} = (-x_2 \ x_2) \begin{pmatrix} x_1 \\ 2x_1 \end{pmatrix} = 0.$$

Note that

- The matrix A has two distinct eigenvalues.
- One of the eigenvalues is 0, and this is a consequence of the fact that the columns of A are not linearly independent.
- The eigenvectors corresponding to two distinct eigenvalues are orthogonal. This is because

$$u_{\lambda=0} \cdot u_{\lambda=5} = u_{\lambda=0}^T u_{\lambda=5} = (-x_2 \ x_2) \begin{pmatrix} x_1 \\ 2x_1 \end{pmatrix} = 0.$$

Put

$$\lambda_1 = 5, \quad \lambda_2 = 0,$$

Put

$$\begin{aligned}\lambda_1 &= 5, & \lambda_2 &= 0, \\ v_1 &= v_{\lambda=5}, & v_2 &= v_{\lambda=0}.\end{aligned}$$

Put

$$\begin{aligned}\lambda_1 &= 5, & \lambda_2 &= 0, \\ v_1 &= v_{\lambda=5}, & v_2 &= v_{\lambda=0}.\end{aligned}$$

Further, put

$$V = (v_1 \quad v_2) = \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix},$$

Put

$$\begin{aligned}\lambda_1 &= 5, & \lambda_2 &= 0, \\ v_1 &= v_{\lambda=5}, & v_2 &= v_{\lambda=0}.\end{aligned}$$

Further, put

$$V = (v_1 \quad v_2) = \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix},$$

$$\Lambda = \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix}.$$

Put

$$\begin{aligned}\lambda_1 &= 5, & \lambda_2 &= 0, \\ v_1 &= v_{\lambda=5}, & v_2 &= v_{\lambda=0}.\end{aligned}$$

Further, put

$$V = (v_1 \quad v_2) = \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix},$$

$$\Lambda = \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix}.$$

Let us compute

$V\Lambda V^T.$

We have

$$V\Lambda V^T = \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix}$$

We have

$$\begin{aligned}
 V\Lambda V^T &= \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \\
 &= \begin{pmatrix} \sqrt{5} & 0 \\ 2\sqrt{5} & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix}
 \end{aligned}$$

We have

$$\begin{aligned}
 V\Lambda V^T &= \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \\
 &= \begin{pmatrix} \sqrt{5} & 0 \\ 2\sqrt{5} & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \\
 &= \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}.
 \end{aligned}$$

We now compute

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$

We now compute

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$

We have

$$\sum_{i=1}^2 \lambda_i v_i v_i^T = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T$$

We now compute

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$

We have

$$\begin{aligned} \sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} (1/\sqrt{5} \quad 2/\sqrt{5}) \end{aligned}$$

We now compute

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$

We have

$$\begin{aligned} \sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \end{pmatrix} \\ &\quad + 0 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \end{aligned}$$

We now compute

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$

We have

$$\begin{aligned} \sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \end{pmatrix} \\ &\quad + 0 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \\ &= \begin{pmatrix} 1 \\ 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} = A. \end{aligned}$$

Therefore,

$$A = 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \end{pmatrix} \\ + 0 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix}.$$

Therefore,

$$A = 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \end{pmatrix} \\ + 0 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix}.$$

Outer product

The product

$$\lambda_i v_i v_i^T$$

is called an **outer product**.

Expressing A as a product of three matrices and as a sum of outer products are called the **spectral decomposition** (or **eigendecomposition**) of A .

Expressing A as a product of three matrices and as a sum of outer products are called the **spectral decomposition** (or **eigendecomposition**) of A .

Basically, this will correspond to expressing A in terms of linear combinations of these special vectors and numbers.

Spectral decompositions of symmetric matrices are particularly well-behaved and have particularly nice properties, while the spectral decompositions of general matrices may or may not have these nice properties.

Spectral decompositions of symmetric matrices are particularly well-behaved and have particularly nice properties, while the spectral decompositions of general matrices may or may not have these nice properties.

By nice properties, we mean that for a given $n \times n$ matrix A we can find a full set of eigenvectors that form an orthonormal basis of \mathbb{R}^n .

Problem

$$\text{Let } A = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}.$$

Problem

Let $A = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}$. Express A as a product of three matrices and as a sum of outer products.

Solution: For this matrix, we have computed
Eigenvalues: $\lambda_1 = 3$, $\lambda_2 = 1$,

Solution: For this matrix, we have computed
Eigenvalues: $\lambda_1 = 3, \lambda_2 = 1$,
Eigenvectors: $v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Expressing A as a product of 3 matrices.

We have

$$V\Lambda V^T = (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix}$$

Expressing A as a product of 3 matrices.

We have

$$\begin{aligned} V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

Expressing A as a product of 3 matrices.

We have

$$\begin{aligned} V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} = A. \end{aligned}$$

Expressing A as a product of 3 matrices.

We have

$$\begin{aligned} V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} = A. \end{aligned}$$

Therefore,

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Expressing A as a sum of 2 outer products. We have

$$\sum_{i=1}^2 \lambda_i v_i v_i^T = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T$$

1. 2. 3.

$$\begin{aligned}\sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} + 1 \begin{pmatrix} 0 \\ 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}\end{aligned}$$

$$\begin{aligned}\sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} (1 \quad 0) + 1 \begin{pmatrix} 0 \\ 1 \end{pmatrix} (0 \quad 1) \\ &= \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} = A.\end{aligned}$$

Therefore,

$$A = 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} (1 \quad 0) + 1 \begin{pmatrix} 0 \\ 1 \end{pmatrix} (0 \quad 1).$$

For the time being, we will describe it in the context of the simple 2×2 matrices.

For the time being, we will describe it in the context of the simple 2×2 matrices.

To do this, let's enumerate the eigenvectors and eigenvalues in a way that will make it easier to generalize beyond 2×2 matrices.

To do this, let's enumerate the eigenvectors and eigenvalues in a way that will make it easier to generalize beyond 2×2 matrices.

- We enumerate eigenvalues in decreasing order.

For the time being, we will describe it in the context of the simple 2×2 matrices.

To do this, let's enumerate the eigenvectors and eigenvalues in a way that will make it easier to generalize beyond 2×2 matrices.

- We enumerate eigenvalues in decreasing order.
- If a matrix has two (or more) identical eigenvalues, then we repeat i in λ_i according to the multiplicity, and we can enumerate their associated orthogonal eigenvectors arbitrarily.

Given this numbering convention, for a 2×2 symmetric matrices, we have seen that there are two eigenvalue-eigenvector pairs (λ_i, v_i) that satisfy:

Given this numbering convention, for a 2×2 symmetric matrices, we have seen that there are two eigenvalue-eigenvector pairs (λ_i, v_i) that satisfy:

$$Av_i = \lambda_i v_i, \quad (1)$$

where $i \in \{1, 2\}$.

Given this numbering convention, for a 2×2 symmetric matrices, we have seen that there are two eigenvalue-eigenvector pairs (λ_i, v_i) that satisfy:

$$Av_i = \lambda_i v_i, \quad (1)$$

where $i \in \{1, 2\}$. The LHS and RHS of this equation are both vectors, i.e., 2×1 matrices.

Given this numbering convention, for a 2×2 symmetric matrices, we have seen that there are two eigenvalue-eigenvector pairs (λ_i, v_i) that satisfy:

$$Av_i = \lambda_i v_i, \quad (1)$$

where $i \in \{1, 2\}$. The LHS and RHS of this equation are both vectors, i.e., 2×1 matrices. Let's write the two equations for $i \in \{1, 2\}$ as a single matrix equation.

Given this numbering convention, for a 2×2 symmetric matrices, we have seen that there are two eigenvalue-eigenvector pairs (λ_i, v_i) that satisfy:

$$Av_i = \lambda_i v_i, \quad (1)$$

where $i \in \{1, 2\}$. The LHS and RHS of this equation are both vectors, i.e., 2×1 matrices. Let's write the two equations for $i \in \{1, 2\}$ as a single matrix equation. To do so, put

$$\Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad V = (v_1 \quad v_2).$$

Here, both are 2×2 matrices. If the i th eigenvalue is Λ_{ii} , then the corresponding i th eigenvector is the i th column of V .

Here, both are 2×2 matrices. If the i th eigenvalue is Λ_{ii} , then the corresponding i th eigenvector is the i th column of V .

Note that

Since the eigenvectors are unit-length and pair-wise orthogonal, V is an orthogonal matrix.

Here, both are 2×2 matrices. If the i th eigenvalue is Λ_{ii} , then the corresponding i th eigenvector is the i th column of V .

Note that

Since the eigenvectors are unit-length and pair-wise orthogonal, V is an orthogonal matrix.

Hence

$$VV^T = V^T V = I \in \mathbb{R}^{2 \times 2}$$

with $V^T = \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix}$.

Spectral decomposition: Expressing A as a product of three matrices.

Theorem (Spectral decomposing I)

Let v_1, v_2 be the eigenvectors associated with the eigenvalues λ_1, λ_2 of a 2×2 symmetric matrix A respectively. If

$$\Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad V = (v_1 \quad v_2),$$

then

$$A = V\Lambda V^T. \quad (2)$$

Proof.

We have

$$\begin{aligned} AV &= (Av_1 \quad Av_2) \\ &= (\lambda_1 v_1 \quad \lambda_2 v_2) \end{aligned}$$

Proof.

We have

$$\begin{aligned} AV &= (Av_1 \quad Av_2) \\ &= (\lambda_1 v_1 \quad \lambda_2 v_2) \\ &= \begin{pmatrix} \lambda_1 v_{11} & \lambda_2 v_{12} \\ \lambda_1 v_{21} & \lambda_2 v_{22} \end{pmatrix} \end{aligned}$$

Proof.

We have

$$\begin{aligned} AV &= (Av_1 \quad Av_2) \\ &= (\lambda_1 v_1 \quad \lambda_2 v_2) \\ &= \begin{pmatrix} \lambda_1 v_{11} & \lambda_2 v_{12} \\ \lambda_1 v_{21} & \lambda_2 v_{22} \end{pmatrix} \\ &= \begin{pmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \end{aligned}$$

Proof.

We have

$$\begin{aligned}
 AV &= (Av_1 \quad Av_2) \\
 &= (\lambda_1 v_1 \quad \lambda_2 v_2) \\
 &= \begin{pmatrix} \lambda_1 v_{11} & \lambda_2 v_{12} \\ \lambda_1 v_{21} & \lambda_2 v_{22} \end{pmatrix} \\
 &= \begin{pmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \\
 &= V\Lambda.
 \end{aligned}$$

Proof.

We have

$$\begin{aligned}
 AV &= (Av_1 \quad Av_2) \\
 &= (\lambda_1 v_1 \quad \lambda_2 v_2) \\
 &= \begin{pmatrix} \lambda_1 v_{11} & \lambda_2 v_{12} \\ \lambda_1 v_{21} & \lambda_2 v_{22} \end{pmatrix} \\
 &= \begin{pmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \\
 &= V\Lambda.
 \end{aligned}$$

Therefore, from this in view of $VV^T = I$, we obtain

$$A = V\Lambda V^T.$$



This theorem provides a decomposition of the matrix A into the product of three matrices:

This theorem provides a decomposition of the matrix A into the product of three matrices:

- an orthogonal matrix V (consisting of the eigenvectors of A),

This theorem provides a decomposition of the matrix A into the product of three matrices:

- an orthogonal matrix V (consisting of the eigenvectors of A),
- a diagonal matrix Λ (consisting of the eigenvalues of A), and

This theorem provides a decomposition of the matrix A into the product of three matrices:

- an orthogonal matrix V (consisting of the eigenvectors of A),
- a diagonal matrix Λ (consisting of the eigenvalues of A), and
- the transpose of that orthogonal matrix, V^T .

We can write Equation (2) in terms of individual elements as follows:

We can write Equation (2) in terms of individual elements as follows:

$$A = V\Lambda V^T$$

We can write Equation (2) in terms of individual elements as follows:

$$A = V\Lambda V^T = \begin{pmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_{11} & v_{21} \\ v_{12} & v_{22} \end{pmatrix}.$$

We can write Equation (2) in terms of individual elements as follows:

$$A = V\Lambda V^T = \begin{pmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_{11} & v_{21} \\ v_{12} & v_{22} \end{pmatrix}.$$

This simple eigendecomposition is the first example of the *spectral theorem*.

This decomposition of the form $A = V\Lambda V^T$ holds for general $n \times n$ symmetric matrices, and generalizations of it hold more generally.

This decomposition of the form $A = V\Lambda V^T$ holds for general $n \times n$ symmetric matrices, and generalizations of it hold more generally. In particular, if we consider computing $y = Ax$, this is the same as

$$y = V\Lambda V^T x = V(\Lambda(V^T x))$$

Spectral decomposition: expressing A as a sum of outer products.

We can write Equation (2) in terms of the columns of V (which recall are the eigenvectors, which are also the rows of V^T) and elements of Λ (which are the eigenvalues).

Spectral decomposition: expressing A as a sum of outer products.

We can write Equation (2) in terms of the columns of V (which recall are the eigenvectors, which are also the rows of V^T) and elements of Λ (which are the eigenvalues). This gives the following theorem:

Theorem (Spectral decomposition II)

Let v_1, v_2 be the eigenvectors associated with the eigenvalues λ_1, λ_2 of a 2×2 symmetric matrix A respectively. Then

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T. \quad (3)$$

Proof.

Let

$$\Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad V = (v_1 \quad v_2),$$

Proof.

Let

$$\Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad V = (v_1 \quad v_2),$$

Then by Spectral theorem I, we have

$$\begin{aligned} A &= V\Lambda V^T \\ &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 v_1^T \\ \lambda_2 v_2^T \end{pmatrix}. \end{aligned}$$

Proof.

Let

$$\Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad V = (v_1 \quad v_2),$$

Then by Spectral theorem I, we have

$$\begin{aligned} A &= V\Lambda V^T \\ &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 v_1^T \\ \lambda_2 v_2^T \end{pmatrix}. \end{aligned}$$

This gives

$$A = \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T.$$



Equation (3) says that if we do this for every i and sum up the corresponding matrices, then we get the original matrix A .

Equation (3) says that if we do this for every i and sum up the corresponding matrices, then we get the original matrix A . Thus, the decomposition given by Equation (3) expresses the matrix A in terms of the sum of 2 terms, each of which is the outer product of an eigenvector with its transpose, multiplied/scaled by the corresponding eigenvalue.

This decomposition holds for general $n \times n$ symmetric matrices, in which case

$$A = \sum_{i=1}^n \lambda_i v_i v_i^T,$$

and generalizations of it hold more generally.

$$A = \sum_{i=1}^n \lambda_i v_i v_i^T,$$

$$A = \sum_{i=1}^n \lambda_i v_i v_i^T,$$

In particular, by the spectral theorem II, the equation $y = Ax$ is equivalent to the equation

$$y = \lambda_1 v_1 v_1^T x + \lambda_2 v_2 v_2^T x.$$

$$y = \lambda_1 v_1 v_1^T x + \lambda_2 v_2 v_2^T x. \quad (4)$$

How do we express the matrix $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ in two standard forms?

How do we express the matrix $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ in two standard forms?

Eigenvectors: $v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

$v_2 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}$, then we obtain

$$\begin{aligned}\sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 1 \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} (1/\sqrt{2} \quad 1/\sqrt{2}) \\ &\quad + 1 \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix} (1/\sqrt{2} \quad -1/\sqrt{2}) \\ &= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} + \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = A.\end{aligned}$$

Therefore,

$$A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} + \frac{1}{\sqrt{2}} \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}.$$

$v_2 = \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}$, then we obtain

$$\begin{aligned}\sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\ &= 1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} + 1 \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix} \begin{pmatrix} 1/2 & 1/2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix} \\ &= \begin{pmatrix} 3/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix} \neq A.\end{aligned}$$

we choose $v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, and $v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, then we obtain

$$\begin{aligned} V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = A. \end{aligned}$$

Therefore,

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

If we choose $v_1 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}$, and $v_2 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}$, then we obtain

$$\begin{aligned} V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \\ &= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = A. \end{aligned}$$

Therefore,

$$A = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}.$$

$$(\cdot, \cdot)_0, (\cdot, \cdot)_T$$

Reflection matrix. How do we express the matrix $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ in two standard forms?

Solution: For this matrix, we have computed
Eigenvalues: $\lambda_1 = 1$, $\lambda_2 = -1$,

Eigenvectors: $v_1 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}$, $v_2 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}$.

Therefore,

$$A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

1 0 1

$$\begin{aligned} V\Lambda V^T &= \begin{pmatrix} v_1 & v_2 \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\ &= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \\ &= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = A. \end{aligned}$$

Therefore,

$$A = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}$$

How do we express the matrix $A = \begin{pmatrix} 9 & -2 \\ -2 & 6 \end{pmatrix}$ in two standard forms?

Eigenvalues: $\lambda_1 = 5, \lambda_2 = 10,$

Eigenvectors: $v_1 = \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix}$, $v_2 = \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix}$.

Expressing A as a sum of 2 outer products. We have

$$\begin{aligned}
 \sum_{i=1}^2 \lambda_i v_i v_i^T &= \lambda_1 v_1 v_1^T + \lambda_2 v_2 v_2^T \\
 &= 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} (1/\sqrt{5} \quad 2/\sqrt{5}) \\
 &\quad + 10 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} (-2/\sqrt{5} \quad 1/\sqrt{5}) \\
 &= \begin{pmatrix} 1 \\ 2 \end{pmatrix} (1 \quad 2) + \begin{pmatrix} -2 \\ 1 \end{pmatrix} (-2 \quad 1) \\
 &= \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} + \begin{pmatrix} 8 & -4 \\ -4 & 2 \end{pmatrix} \\
 &= \begin{pmatrix} 9 & -2 \\ -2 & 6 \end{pmatrix} = A.
 \end{aligned}$$

Therefore,

$$A = 5 \begin{pmatrix} 1/\sqrt{5} \\ 2/\sqrt{5} \end{pmatrix} (1/\sqrt{5} \quad 2/\sqrt{5}) \\ + 10 \begin{pmatrix} -2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix} (-2/\sqrt{5} \quad 1/\sqrt{5}).$$

Expressing A as a product of 3 matrices. We have

$$\begin{aligned}
 V\Lambda V^T &= (v_1 \quad v_2) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} v_1^T \\ v_2^T \end{pmatrix} \\
 &= \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 5 & 10 \\ -20 & 10 \end{pmatrix} \\
 &= \frac{1}{5} \begin{pmatrix} 45 & -10 \\ -10 & 30 \end{pmatrix} \\
 &= \begin{pmatrix} 9 & 2 \\ -2 & 6 \end{pmatrix} = A
 \end{aligned}$$

Therefore,

$$A = \begin{pmatrix} 1/\sqrt{5} & -2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix} \begin{pmatrix} 5 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ -2/\sqrt{5} & 1/\sqrt{5} \end{pmatrix}.$$