



Introduction to General Relativity

AMATH 475



Prof. Eduardo Martin-martinez

Preface

Disclaimer Much of the information on this set of notes is transcribed directly/indirectly from the lectures of AMATH 475 during Winter 2020 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

Some of the notes (especially special relativity part) are projected to the screen instead of using blackboards. They can be found on [professor's course page](#).

For any questions, send me an email via <https://notes.sibeliusp.com/contact/>.

You can find my notes for other courses on <https://notes.sibeliusp.com/>.

Sibeliuss Peng

Contents

Preface	1
0 Pre-Math	3
0.1 Index notation	3
0.2 Vectors and one-forms	4
0.3 Tensor	5
0.4 Levi-Civita symbol	6

Pre-Math

0.1 Index notation

$$A = \begin{pmatrix} A^1_1 & A^1_2 \\ A^2_1 & A^2_2 \end{pmatrix} \quad B = \begin{pmatrix} B^1_1 & B^1_2 \\ B^2_1 & B^2_2 \end{pmatrix}$$

$$(A \cdot B)^a_b = A^a_c B^c_b = B^c_b A^a_c \quad \text{sum over all possible } c$$

Identify followings:

$$\begin{aligned} B_\kappa^\nu A_\mu^\kappa &= A_\mu^\kappa B_\kappa^\nu = C_\mu^\nu = (A \cdot B)_\mu^\nu \\ A^\kappa_\mu B_\kappa^\nu &= D_\mu^\nu = (A^T)_\mu^\kappa B_\kappa^\nu = (A^T \cdot B)_\mu^\kappa \\ A_\kappa^\nu B_\mu^\kappa &= E_\mu^\nu = (B \cdot A)_\mu^\nu \\ A^\kappa_\mu B^\nu_\kappa &= (A^T)_\mu^\kappa (B^T)_\kappa^\nu = \left((B \cdot A)^T \right)_\mu^\nu \end{aligned}$$

$$\mathbf{v} = v^1 \mathbf{e}_1 + v^2 \mathbf{e}_2 \quad \{\mathbf{e}_1, \mathbf{e}_2\} \text{ Basis 1.}$$

$$\mathbf{v} = v'^a \mathbf{e}_a = v'^a \mathbf{e}'_a \quad \{\mathbf{e}'_1, \mathbf{e}'_2\} \text{ Basis 2.}$$

Change of basis matrix Λ

$$\begin{aligned} \mathbf{e}'_a &= \Lambda_a^b \mathbf{e}_b \\ v'^a &= \tilde{\Lambda}^a_b v^b \end{aligned}$$

$$\begin{aligned}
v^a \mathbf{e}_a &= v'^a \mathbf{e}'_a \\
&= \tilde{\Lambda}^a_b v^b \Lambda_a^c \mathbf{e}_c \\
&= \tilde{\Lambda}^a_b \Lambda_a^c v^b \mathbf{e}_c \\
&= \underbrace{\left(\tilde{\Lambda}^T \right)_b^a}_{\delta_b^c} \Lambda_a^c v^b \mathbf{e}_c \\
&= v^b \mathbf{e}_b \\
\\
\implies \left(\tilde{\Lambda}^T \right)_b^a \Lambda_a^c &= \delta_b^c \\
\tilde{\Lambda}^T \cdot \Lambda &= \mathbb{1}
\end{aligned}$$

$\tilde{\Lambda}^T$ is the inverse transpose of Λ

covariant and contravariant object

A covariant object is an object that under change of basis transforms like the elements of a basis. Λ . (sub-indices)

A contravariant object transforms like components of vectors. $(\tilde{\Lambda} = (\Lambda^T)^{-1})$. (super-indices)

0.2 Vectors and one-forms

one-form

Let V be a vector space. A one-form is a linear map $\omega : V \rightarrow \mathbb{R}$.

or we write: $(\omega, \cdot) : V \rightarrow \mathbb{R}$ and $(\omega, \mathbf{v}) \in \mathbb{R}$.

dual vector space

The set of all one-forms on V (call V^*) is a vector space as well called the dual vector space to V .

dual basis

Let $\{\Upsilon_1, \Upsilon_2, \dots\}$ (or $\{\Upsilon_i\}$) be a basis of V so that any $\mathbf{v} \in V$ can be written as $\mathbf{v} = v^i \Upsilon_i$.

We define the dual basis (of V^*) to $\{\Upsilon_i\}$ as $\{\omega^i\}$ such that $\omega^i(\Upsilon_j) = \delta_j^i$.

For a one form ω we denote its “components of the basis Υ ” as $(\omega, \Upsilon_m) = \omega_m$

Proposition 0.1

The dual basis of V^* is actually a basis of V^* .

The action of $\omega \in V^*$ on a vector $\mathbf{v} = v^\mu \Upsilon_\mu \in V$ is

$$(\omega, \mathbf{v}) = (\omega, v^\mu \Upsilon_\mu) = v^\mu \omega_\mu$$

Let’s prove $\{\Upsilon^a\}$ is linear independent.

Proof:

A linear comb. $c_a \Upsilon^a$ acts on a vector $\mathbf{v} = v^a \Upsilon_a$

$$\begin{aligned} (c_a \Upsilon^a, \mathbf{v}) &= c_a (\Upsilon^a, \mathbf{v}) \\ &= c_a (\Upsilon^a, v^b \Upsilon_b) \\ &= c_a v^b \underbrace{(\Upsilon^a, \Upsilon_b)}_{\delta_b^a} \\ &= c_a v^b \delta_b^a = c_a v^a \end{aligned}$$

For LI,

$$\begin{aligned} c_a \Upsilon^a = 0 &\iff c_a = 0 \quad \forall a \\ c_a v^a = 0 \quad \forall \mathbf{v} &\iff c_a = 0 \end{aligned}$$

□

vectors: take one-forms $\rightarrow \mathbb{R}$ one-forms: take vectors $\rightarrow \mathbb{R}$

0.3 Tensor

type (m, n) tensor

A type (m, n) tensor is a multilinear map that

$$\mathbf{T} : V^n \otimes (V^*)^m \rightarrow \mathbb{R}$$

Components of \mathbf{T} :

$$\mathbf{T}(\Upsilon_{a_1}, \dots, \Upsilon_{a_n}, \Upsilon^{b_1}, \dots, \Upsilon^{b_m}) = T_{a_1 \dots a_n}{}^{b_1 \dots b_m}$$

1. Tensor product takes $\binom{m}{n}$ and $\binom{m'}{n'} \rightarrow \binom{m+m'}{n+n'}$ tensor
2. Contraction takes $\binom{m}{n} \rightarrow \binom{m-1}{n-1}$

Example:

$$1. T_a^b, S_c^d.$$

$$(\mathbf{T} \otimes \mathbf{S})_a^b{}_c^d = T_a^d S_c^d = P_a^b{}_c^d$$

$$2. T_a^{bc} \rightarrow c^b T_a^{ba}$$

$$v^a, w_b \begin{cases} v^a \omega_b \\ v^a \omega_a \end{cases}$$

If you have a favorite type $(2, 0)$ symmetric tensor \mathbf{g}

$$v_\mu = g_{\mu\nu} v^\nu$$

$g^{\mu\nu} :=$ components of the inverse of $\mathbf{g}_{\mu\nu}$

$$v^\nu = g^{\mu\nu}$$

then

$$g^{\mu\nu} g_{\nu\sigma} = \delta_\sigma^\mu$$

$$g_{\mu\nu} v^\mu w^\nu = v_\mu w^\mu = \mathbf{v} \cdot \mathbf{w}$$

$$||\mathbf{v}||^2 = g_{\nu\mu} v^\mu v^\nu$$

Then we can define the angle

$$\frac{\mathbf{v} \cdot \mathbf{w}}{||\mathbf{w}|| ||\mathbf{v}||} := \cos \theta$$

$$T_\mu{}^\nu = g^{\nu\sigma} T_{\mu\sigma}$$

$$T^{\mu\nu} = g^{\nu\sigma} g^{\mu\rho} T_{\sigma\rho}$$

$$g_\mu^\nu = g^{\nu\sigma} g_{\sigma\mu} = \sigma_\mu^\nu$$

0.4 Levi-Civita symbol

Levi-Civita symbol $\epsilon^{abc\dots}, \epsilon_{abc\dots}$

- is antisymmetric

- $\epsilon^{1234\dots} = 1, \epsilon_{1234} = 1$

$$\epsilon^{123} = 1, \quad \epsilon^{213} = -1, \quad \epsilon^{312} = 1, \quad \epsilon^{113} = 0$$


$$\epsilon^{123456} = 1, \quad \epsilon^{612453} = -1$$

Idea just see the permutations

Levi-Civita symbol

$$\varepsilon_{a_1 a_2 a_3 \dots a_n} = \begin{cases} +1 & \text{if } (a_1, a_2, a_3, \dots, a_n) \text{ is an even permutation of } (1, 2, 3, \dots, n) \\ -1 & \text{if } (a_1, a_2, a_3, \dots, a_n) \text{ is an odd permutation of } (1, 2, 3, \dots, n) \\ 0 & \text{otherwise} \end{cases}$$

Here is a short-cut:



odd number crossings, so odd permutation.

Note that $\det(M) := \epsilon_{ijk\dots} M^i_1 M^j_2 M^k_3 \dots$

Exercise.

prove $\epsilon^{i_1 i_2 \dots i_n} \epsilon_{j_1 j_2 \dots j_n} = n! \delta_{i_j}^{j_j} = 1, \dots, n$

$$\begin{aligned} \epsilon^{ijk} \epsilon_{ilm} &= \delta_l^j \delta_m^k - \delta_m^j \delta_l^k \\ \epsilon^{ijmn} \epsilon_{klmn} &= 2(\delta_k^i \delta_l^j - \delta_l^i \delta_k^j) \end{aligned}$$

Prove $\vec{A} \times (\vec{B} \times \vec{C}) = (\vec{A} \cdot \vec{C})\vec{B} - (\vec{A} \cdot \vec{B})\vec{C}$

Proof.

Let $\vec{F} = \vec{A} \times (\vec{B} \times \vec{C})$ $\vec{D} = \vec{B} \times \vec{C}$

Then

$$\begin{aligned} D^k &= \epsilon^k_{ij} B^i C^j \\ F^l &= \epsilon^l_{mk} A^m D^k \implies F^l = \epsilon^l_{mk} \epsilon^k_{ij} A^m B^i C^j \end{aligned}$$

Then

$$\begin{aligned} F^l &= (\delta_i^l \delta_{mj} - \delta_j^l \delta_{mi}) A^m B^i C^j \\ &= \delta_i^l \delta_{mj} A^m B^i C^j - \delta_j^l \delta_{mi} A^m B^i C^j \\ &= B^l (A_j C^j) - C^l (A_i B^i) \end{aligned}$$

where we use

$$\vec{A} \cdot \vec{B} = A^i B_i$$

□

Index

C

covariant and contravariant object . 4

D

dual basis..... 4

dual vector space 4

L

Levi-Civita symbol..... 7

O

one-form 4

T

type (m,n) tensor 5