



Groups and Rings

PMATH 347



William Slofstra

Preface

Disclaimer Much of the information on this set of notes is transcribed directly/indirectly from the lectures of PMATH 347 during Spring 2020 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

Spring 2020 classes online only. So the grading scheme:

- Participation: 4%
- Quizzes: 32%
- Written homework: 32%
- Final takehome exam: 32%

For any questions, send me an email via <https://notes.sibeliusp.com/contact/>.


You can find my notes for other courses on <https://notes.sibeliusp.com/>.

Sibeliusp Peng

Contents

Preface	1
I Group Theory	4
1 Introduction to Groups	5
1.1 Binary Operations	5
1.2 Associativity and commutativity	7
1.3 Identities and inverses	9
1.4 Groups	11
1.4.1 Terminology	12
1.4.2 Additive notation	13
1.4.3 Multiplicative table	14
1.4.4 Order of elements	14
1.5 Dihedral groups	15
1.5.1 Special elements of D_{2n}	17
1.6 Permutation groups	18
1.6.1 Representations	19
1.6.2 Cycles	21
2 Subgroups	22
2.1 Subgroups	22
2.2 Subgroups generated by a set	25
2.2.1 Lattice of subgroups	27
2.3 Cyclic groups	27
2.3.1 $\mathbb{Z}/n\mathbb{Z}$	29
3 Homomorphisms	32
3.1 Homomorphisms	32
3.2 Homomorphisms and subgroups	34
3.2.1 Application: subgroups of cyclic groups	36
3.3 Isomorphisms	37
3.4 Cosets	40
3.5 The index and Lagrange's theorem	43
3.6 Proof of Lagrange's theorem	46
3.6.1 Equivalence relations	48
3.7 Normal subgroups	50

3.8	Normalizers and the center	52
4	Products	54
4.1	Product groups	54
4.2	Homomorphisms between products	55
4.3	Unique factorizations & internal direct products	57
5	Quotient groups and the isomorphism theorems	60
5.1	Quotient groups	60
5.2	The universal property of quotients	62
5.3	The first isomorphism theorem	65
5.4	The correspondence theorem	66
5.5	The third isomorphism theorem	70
5.6	The second isomorphism theorem	72
6	Group actions	75
6.1	Group actions	75



PART I:

GROUP THEORY

It is important to realize, with or without the historical context, that the reason the abstract definitions are made is because it is useful to isolate specific characteristics and consider what structure is imposed on an object having these characteristics.

Abstract Algebra, Third Edition

Introduction to Groups

1.1 Binary Operations

week 1

If we randomly ask someone on the street: *What's math about?* The answer we might get is **numbers**. It always comes with **operations**.

Objects	Operations
Natural numbers \mathbb{N}	addition $+$ subtraction $-$ multiplication \cdot division with remainders
Integers \mathbb{Z}	negation $x \mapsto -x$
Rational number \mathbb{Q}	multiplicative inversion $x \mapsto 1/x$
Real numbers \mathbb{R}	k th roots, etc
$\mathbb{Z}/n\mathbb{Z}$	modular arithmetic and operations

Then we realized that math is not just about numbers. We later have **elementary algebra**:

Objects	Operations
Expressions with variables	operations with variables
Functions	Pointwise operations $+$, $-$, \cdot and Composition \circ

Then ..., and (leaving lots of stuff out), we have **linear algebra**:

Objects	Operations
Vectors	Vector addition $+$, scalar multiplication \cdot
Matrices	$+$, $-$, scalar and matrix multiplication \cdot

Then *what's algebra about?*

Pre-university answer:

- manipulating expr involving indeterminates (variables):

If $a, b \in \mathbb{R}$, $ax = b$ and $a \neq 0$, then $x = \frac{b}{a}$.

- solving eqs by applying ops to both sides:
If A, B are matrices, $AX = B$ and A is invertible, then $X = A^{-1}B$.

Key idea: algebra is about operations

Then *what operations should we study?* Polynomials in several vars; functions, pointwise ops and function composition... *Are there other operations we should study?* Then we introduce **abstract algebra**: try to answer this question by studying operations abstractly, and seeing what the possibilities are.

binary operation

A binary operation on a set X is a function $b : X \times X \rightarrow X$.

Notation:

- Any letter (b, m) or symbol $(+, \cdot)$
- function notation

$$b : X \times X \rightarrow X : (x, y) \mapsto b(x, y)$$

or inline notation

$$+ : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} : (x, y) \mapsto x + y$$

Typically use inline notation with symbols and function notation with letters.

- There are lots of symbols to choose from: $a + b, a \times b, a \cdot b, a \circ b, a \oplus b, a \otimes b, a \odot b, a \diamond b, a \heartsuit b, a \spadesuit b, a * b, a \bullet b, a \boxplus b, a \boxtimes b, a \uplus b$
- If there's no chance of confusion, can even drop symbol completely:

$$X \times X \rightarrow X : (a, b) \mapsto ab$$

Example:

- Addition $+$ is a binary op on \mathbb{N} , but subtraction $-$ is not, since $a - b$ is not necessarily a natural number.
- Subtraction $=$ is a binary op on \mathbb{Z} .
- If $(V, +, \cdot)$ is a vector space over a field \mathbb{K} , then $+$ is a binary op on V , but \cdot is not, since \cdot is a function $\mathbb{K} \times V \rightarrow V$.^a

^aWe'll define fields later, now think of $\mathbb{K} = \mathbb{R}$ or \mathbb{C} .

k-ary operation

A k -ary operation on a set X is a function

$$\underbrace{X \times X \times \cdots \times X}_{k \text{ times}} \rightarrow X$$

A 1-ary operation is called a unary operation.

Example:

Negation $\mathbb{Z} \rightarrow \mathbb{Z} : x \mapsto -x$ is a unary operation.

Taking the multiplicative inverse $x \mapsto 1/x$ is not a unary operation on \mathbb{Q} , since $1/0$ is not defined, but it is a unary operation on

$$\mathbb{Q}^\times := \{a \in \mathbb{Q} : a \neq 0\}$$

Now let's discuss some properties that binary ops might satisfy.

1.2 Associativity and commutativity

associative

A binary operation $\boxtimes : X \times X \rightarrow X$ is associative if

$$a \boxtimes (b \boxtimes c) = (a \boxtimes b) \boxtimes c$$

for all $a, b, c \in X$.

Many operations we've mentioned so far are associative:

- Addition and multiplication for $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, polynomials, and functions
- Vector addition, matrix addition and multiplication
- Modular addition and multiplication on $\mathbb{Z}/n\mathbb{Z}$
- Function composition

Note that Subtraction and division are not associative. Subtraction is adding negative numbers, same for division. So we aren't that interested in subtraction and division, and focus on associative operations.

Here we introduce an informal definition: A **bracketing** of a sequence $a_1, \dots, a_n \in X$ is a way of inserting brackets into $a_1 \boxtimes \dots \boxtimes a_n$ so that the expression can be evaluated.

Example:

The bracketings of a_1, \dots, a_4 are

$$a_1 \boxtimes (a_2 \boxtimes (a_3 \boxtimes a_4))$$

$$a_1 \boxtimes ((a_2 \boxtimes a_3) \boxtimes a_4)$$

$$\begin{aligned}
& (a_1 \boxtimes a_2) \boxtimes (a_3 \boxtimes a_4) \\
& (a_1 \boxtimes (a_2 \boxtimes a_3)) \boxtimes a_4 \\
& ((a_1 \boxtimes a_2) \boxtimes a_3) \boxtimes a_4
\end{aligned}$$

Proposition 1.1

A binary operation $\boxtimes : X \times X \rightarrow X$ is associative if and only if for all finite sequences $a_1, \dots, a_n \in X, n \geq 1$, every bracketing of a_1, \dots, a_n evaluated to the same element of X .

Note

If \boxtimes is associative, can use notation $a_1 \boxtimes a_2 \boxtimes \dots \boxtimes a_n$ without choosing a bracketing.

Proof:

\Leftarrow The two bracketings $a \boxtimes (b \boxtimes c)$ and $(a \boxtimes b) \boxtimes c$ of a, b, c evaluate to the same element of X for all sequences of length 3.

\Rightarrow Proof is by induction. Base cases are $n = 1, 2, 3$.

For $n = 1, 2$, there's only one bracketing. For $n = 3$ follows from defn of associativity.

Suppose prop is true for all sequences of length $k, 1 \leq k < n$.

Let w be a bracketing of a_1, \dots, a_n .

$w = w_1 \boxtimes w_2$ where w_1 is a bracketing of a_1, \dots, a_k , w_2 is a bracketing of a_{k+1}, \dots, a_n , for some $k < n$.

By induction,

$$w_1 = (\dots((a_1 \boxtimes a_2) \boxtimes a_3) \dots \boxtimes a_k) \quad \text{and} \quad w_2 = (a_{k+1} \boxtimes \dots (a_{n-1} \boxtimes a_n) \dots)$$

Therefore

$$\begin{aligned}
w &= (\dots((a_1 \boxtimes a_2) \boxtimes a_3) \dots \boxtimes a_k) \boxtimes w_2 = (a_{k+1} \boxtimes \dots (a_{n-1} \boxtimes a_n) \dots) \\
&= (\dots(a_1 \boxtimes a_2) \dots \boxtimes a_{k-1}) \boxtimes (a_k \boxtimes (a_{k+1} \boxtimes \dots a_n) \dots) \\
&= \dots \\
&= (a_1 \boxtimes (a_2 \boxtimes \dots (a_n \boxtimes a_n) \dots))
\end{aligned}$$

□

commutative

A binary operation $\boxtimes : X \times X \rightarrow X$ is commutative (also known as abelian) if $a \boxtimes b = b \boxtimes a$ for all $a, b \in X$.

Fact The word “abelian” comes from the surname of Niels Henrik Abel (1802-1829).

Many familiar operations are commutative: addition and multiplication on $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$; vector and matrix addition; modular addition and multiplication on $\mathbb{Z}/n\mathbb{Z}$. The following operation are **not** commutative: subtraction and division; function composition; matrix multiplication.

Therefore, subtraction and division are not commutative or associative. Function composition and matrix multiplication are not commutative, but are associative. We are not going to worry about the first type of operation, but we are interested in operations of the second type.

First half of the course: group theory – single associative operation, not necessarily commutative.

Second half of the course: ring theory – two associative operations (like addition and multiplication on \mathbb{Z}), focus on commutative case.

1.3 Identities and inverses

Let \boxtimes be a binary operation on a set X .

identity

An element $e \in X$ is an identity for \boxtimes if

$$e \boxtimes x = x \boxtimes e = x$$

for all $x \in X$.

Example:

The zero element 0 of \mathbb{Z} is an identity for $+$. $1 \in \mathbb{Q}$ is identity for \cdot . $0 \in \mathbb{Q}$ is not identity for \cdot .

Lemma 1.2

If $e, e' \in X$ are both identities for \boxtimes , then $e = e'$.

Proof:

$$e = e \boxtimes e' = e'$$

□

inverse

Let \boxtimes be a binary operation on X with identity element e . An element y is a left inverse for x (w.r.t. \boxtimes) if $y \boxtimes x = e$, a right inverse if $x \boxtimes y = e$, and an inverse if $x \boxtimes y = y \boxtimes x = e$.

Example:

$-n$ is an inverse for $n \in \mathbb{Z}$ w.r.t. $+$.

$n \in \mathbb{Z}$ does not have an inverse w.r.t. \cdot unless $n = \pm 1$.

If $x \in \mathbb{Q}$ is non-zero, then $1/x$ is an inverse of x w.r.t. \cdot . The element 0 does not have an inverse.

Lemma 1.3

Let \boxtimes be an **associative** binary op with an identity e . If y_L and y_R are left and right inverse of x respectively, then $y_L = y_R$.

Proof:

$$y_L = y_L \boxtimes e = y_L \boxtimes (x \boxtimes y_R) = (y_L \boxtimes x) \boxtimes y_R = e \boxtimes y_R = y_R$$

□

Corollary 1.4

- If x has both a left and right inverse, then x has an inverse.
- Inverses are unique.

invertible

An element a is invertible if it has an inverse, in which case the inverse is denoted by a^{-1} .

Exercise

It's possible to have a left (resp. right inverse), but not be invertible. Also, left and right inverses don't have to be unique (unless an element has both).

Lemma 1.5

1. If \boxtimes has an identity e , then e is invertible, and $e^{-1} = e$.
2. If a is invertible, then so is a^{-1} , and $(a^{-1})^{-1} = a$.
3. If \boxtimes is associative, and a and b are invertible, then so is $a \boxtimes b$, and $(a \boxtimes b)^{-1} = b^{-1} \boxtimes a^{-1}$.

Proof:

1. $e \boxtimes e = e$
2. $a \boxtimes a^{-1} = a^{-1} \boxtimes a = e$, so a is clearly an inverse to a^{-1} .
3. $(a \boxtimes b) \boxtimes (b^{-1} \boxtimes a^{-1}) = a \boxtimes (b \boxtimes b^{-1}) \boxtimes a^{-1} = a \boxtimes e \boxtimes a^{-1} = a \boxtimes a^{-1} = e$, and similarly $(b^{-1} \boxtimes a^{-1}) \boxtimes (a \boxtimes b) = e$.

□

Proposition 1.6

Let \boxtimes be an associative binary operation on X with identity e , and let x and y be variables taking values in X .

An element $a \in X$ is invertible if and only if the equations

$$a \boxtimes x = b \text{ and } y \boxtimes a = b$$

have unique solutions for all $b \in X$.

Proof:

\Leftarrow A solution to $ax = e$ is a right inverse of a , and a solution to $ya = e$ is a left inverse. If a both have a left and right inverse, then it has an inverse.

\Rightarrow Suppose a is invertible. Then

$$a \boxtimes (a^{-1}b) = (a \boxtimes a^{-1}) \boxtimes b = e \boxtimes b = b$$

so $a^{-1} \boxtimes b$ is a solution to $a \boxtimes x = b$.

If x_0 is a solution to $a \boxtimes x = b$, then

$$a^{-1} \boxtimes b = a^{-1} \boxtimes (a \boxtimes x_0) = (a^{-1} \boxtimes a) \boxtimes x_0 = e \boxtimes x_0 = x_0$$

So $a^{-1} \boxtimes b$ is the unique solution to $a \boxtimes x = b$.

Similarly $b \boxtimes a^{-1}$ is the unique solution to $y \boxtimes a = b$. □

Proposition 1.7: Cancellation property

Let \boxtimes be an associative binary operation, and $a \in X$. Then

1. If a has a left inverse and $a \boxtimes u = a \boxtimes v$, then $u = v$.
2. If b has a right inverse and $u \boxtimes a = v \boxtimes a$, then $u = v$.

Proof:

1. $u = a^{-1} \boxtimes a \boxtimes u = a^{-1} \boxtimes a \boxtimes v = v$
2. similar. □

1 and 2 also hold for $n \in \mathbb{Z}$ w.r.t. \cdot if $n \geq 0$, even though n is not invertible for $n \neq \pm 1$.

1.4 Groups

group

A **group** is a pair (G, \boxtimes) , where

1. G is a set, and
2. \boxtimes is an associative binary operation on G such that

- (a) \boxtimes has an identity e , and
- (b) every element $g \in G$ is invertible with respect to \boxtimes .

abelian

A group is **abelian** (or commutative) if \boxtimes is abelian.

finite

A group is **finite** if G is a finite set.

order

The **order** of G the number of elements in G if G is finite, and $+\infty$ if G is infinite. The order of G is denoted by $|G|$.

1.4.1 Terminology

Usually we refer to (G, \boxtimes) simply as G , and just assume the operation is given. (Note: we still need to clearly specify the operation for each group we work with).

It's cumbersome to write \boxtimes all the time, so usually we use one of the following options:

- Use \cdot as the standard symbol, write $g \cdot h$ for the product of $g, h \in G$
- Drop the symbol entirely, write gh for the product of $g, h \in G$.

The identity of G is denoted by e (or e_G when we want to make the group clear). 1 and 1_G are also used.

Since every element of a group G is invertible, g^{-1} is defined for all $g \in G$. The function $G \rightarrow G : g \mapsto g^{-1}$ can be regarded as a unary operation on G .

Consider $\iota : G \rightarrow G : g \mapsto g^{-1}$. Since $(g^{-1})^{-1} = g$, $\iota \circ \iota = \text{Id}_G$, the identity map $G \rightarrow G$. In particular, ι is a bijection, both injective and surjective.

If $g \in G$, then

$$g^n := \underbrace{g \cdots g}_{n \text{ times}} \text{ and } g^{-n} := (g^{-1})^n = (g^n)^{-1}$$

Exercise

If $m, n \in \mathbb{Z}$, then $(g^n)^m = g^{mn}$.

If $g, h \in G$, then

$$(gh)^n = ghgh \cdots gh,$$

which is not necessarily the same as $g^n h^n$ if G is not abelian.

Example: Groups

$\mathbb{N}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ are all groups under operation $+$. The identity is 0 and the inverse of n is $-n$. These groups have infinite order. They are infinite abelian groups.

$\mathbb{Z}/n\mathbb{Z}$ is also a group under $+$. The identity is $0 = [0]$, and the inverse of $[m]$ is $-[m] = [-m]$. This group is finite, with order $|\mathbb{Z}/n\mathbb{Z}| = n$. It is a finite abelian group.

If $(V, +, \cdot)$ is a vector space, then $(V, +)$ is group. The identity element is 0, and the inverse of v is $-v$.

Example: Not a group?! & Trivial group

\mathbb{Z} is not a group with respect to \cdot , since most elements do not have an inverse.

\mathbb{Q} is also not a group with respect to \cdot , since 0 does not have an inverse.

\mathbb{Q}^\times is a group with respect to \cdot .

Every group has to contain at least one element, the identity. So the simplest possible group is $\{1\}$ with operation $1 \cdot 1 = 1$. This is called the **trivial group**.

A non-abelian example

All the examples previously are abelian.

Let $\text{GL}_n(\mathbb{K})$ denote the invertible $n \times n$ matrices with entries in a field \mathbb{K} .

Proposition 1.8

$\text{GL}_n(\mathbb{K})$ is a group under matrix multiplication (called the **general linear group**). For $n \geq 2$, $\text{GL}_n(\mathbb{K})$ is non-abelian.

Proof:

If A and B are invertible matrices, then AB is also invertible, so matrix multiplication is an associative binary operation $\text{GL}_n(\mathbb{K})$. The identity matrix is an identity, and every element has an inverse by definition, so $\text{GL}_n(\mathbb{K})$ is a group.

Exercise

Find matrices A, B such that $AB \neq BA$.

□

1.4.2 Additive notation

Standard notation for operation in a group is gh . This is called **multiplicative notation**. For groups like $(\mathbb{Z}, +)$, it is confusion to write mn instead of $m + n$, since mn already has another meaning. For abelian groups G , there is another convention called **additive notation**. In additive notation, we write the group operation as $g + h$. The identity is denoted by 0 or 0_G . Inverse are denoted by $-g$. Writing g^n in additive notation gives

$$\underbrace{g + g + \dots + g}_{n \text{ times}}$$

so rather than g^n we use ng . Similarly g^{-n} is $-ng$.

Multiplicative notation	Additive notation
$g \cdot h$ or gh	$g + h$
e_G or 1_G	0_G
g^{-1}	$-g$
g^n	ng

Table 1.1: Comparison between multiplicative and additive notation

For nonabelian groups we always use multiplicative notation. For abelian groups, we can choose either.

Note the potential for conflict between the two conventions. We must be clear about what convention we are using!

For groups like $(\mathbb{Z}, +)$, we could denote the operation by mn , but it's clearer to write $m + n$. For groups like (Q^\times, \cdot) , we could denote the operation by $x + y$, but it is clearer to write $x \cdot y$ or xy .

1.4.3 Multiplicative table

multiplicative table

The multiplicative table of a group G is a table with rows and columns indexed by the elements of G . The cell for row g and column h contains the product gh .

The multiplication table contains the complete info of the group G . It is defined for finite and infinite groups, but makes the most sense for finite groups.

Example: $\mathbb{Z}/2\mathbb{Z}$

The multiplication table for $\mathbb{Z}/2\mathbb{Z}$ is

	0	1
0	0	1
1	1	0

1.4.4 Order of elements

order

If G is a group, then the order $g \in G$ is

$$|g| := \min\{k \geq 1 : g^k = e_G\} \cup \{+\infty\}$$

Some easy properties:

- $|g| = 1$ if and only if $g = e_G$.
- If $g^n = 1$, then $g^{n-1}g = gg^{n-1} = g^n = 1$, so $g^{n-1} = g^{-1}$. In particular, if $|g| = n < +\infty$, then $g^{-1} = g^{n-1}$.

Example: $\mathbb{Z}/n\mathbb{Z}$

We use additive notation for $\mathbb{Z}/n\mathbb{Z}$, so g^n is written as ng , $e = 0$. For this group, $k1 = 0$ if and only if n divides k , so $|1| = n$.

Lemma 1.9

$g^n = e$ if and only if $g^{-n} = e$, so in particular $|g| = |g^{-1}|$.

Proof:

We have $g^{-n} = (g^n)^{-1}$. Since $g \mapsto g^{-1}$ is a bijection,

$$g^n = e \text{ if and only if } (g^n)^{-1} = e^{-1} = e.$$

But g^{-n} also equals $(g^{-1})^n$, so


$$\{k \geq 1 : g^k = e\} = \{k \geq 1 : (g^{-1})^k = e\}$$

and this implies $|g| = |g^{-1}|$. □

1.5 Dihedral groups

n-gon

A regular polygon P_n with n vertices, $n \geq 3$, is called an n -gon.



To be specific: set $v_k = (\cos 2\pi k/n, \sin 2\pi k/n) = e^{2\pi i k/n}$

Get n -gon by drawing line segment from v_k to v_{k+1} for all $0 \leq k \leq n$ (where $v_n := v_0$)

symmetry

A symmetry of the n -gon P_n is an invertible linear transformation $T \in \text{GL}_2(\mathbb{R})$ such that $T(P_n) = P_n$.

dihedral group

The set of symmetries of P_n is called the dihedral group, and is denoted by D_{2n} (or D_n).

In this course, we use D_{2n} .

Note

We think of matrices and invertible linear transformations interchangeably.

Matrix multiplication = composition of transformations.

Proposition 1.10

D_{2n} is a group under composition.

Proof:

Later. Key point: $S, T \in D_{2n} \implies ST \in D_{2n}$. □

v_i and v_j are **adjacent** in P_n if connected by line segment.

Lemma 1.11

1. If $T \in D_{2n}$ then $(T(v_0), T(v_1))$ are adjacent
2. If $S, T \in D_{2n}$ and $S(v_i) = T(v_i)$, $i = 0, 1$ then $S = T$.

Proof:

1. v_0, v_1 are adjacent, T is linear
 2. v_0 and v_1 are linearly independent.
-

Corollary 1.12

$$|D_{2n}| \leq 2n$$

Proof:

Let A be the set of adjacent pairs $(v_i, v_j)^a$, so $|A| = 2n$. By Lemma 1.11, $D_{2n} \rightarrow A : T \mapsto (T(v_0), T(v_1))$ is well-defined and injective. □


^aordered pairs

For every pair of adjacent vertices (v_i, v_j) , is there an element $T \in D_{2n}$ with $T(v_0) = v_i, T(v_1) = v_j$?


If the answer is yes, then $|D_{2n}| = 2n$.

1.5.1 Special elements of D_{2n}

Let $s \in D_{2n}$ be rotation by $2\pi/n$ radians, so $|s| = n$ (i.e., $s^n = 2, s^k \neq e$ for $1 \leq k < n$).



Let r be reflection through the x -axis:




$|r| = 2$, i.e. $r^2 = e, r \neq e$.

$r(v_0) = v_0$. $r(v_1)$ is now the vertex before v_0 , rather than the vertex after v_0 .

If we try to put these two elements together:

1. $s^i, 0 \leq i < n$: sends $v_0 \mapsto v_i, v_1 \mapsto v_{i+1}$ (notes: $v_n = v_0, s^0$ is the identity)
2. $s^i r, 0 \leq i < n$: sends $v_0 \mapsto v_i, v_1 \mapsto v_{i-1}$ (notes: $v_{-1} = v_{n-1}$)



Proposition 1.13

$D_{2n} = \{s^i r^j : 0 \leq i < n, 0 \leq j < 2\}$, so $|D_{2n}| = 2n$.

What is rs ?

$rs(v_0) = r(v_1) = v_{n-1}$ and $rs(v_1) = r(v_2) = v_{n-2}$. So

$$rs = s^{n-1}r = s^{-1}r$$

Corollary 1.14

D_{2n} is a finite nonabelian group.

Exercise

$$D_{2n} = \{s^i r^j : 0 \leq i < n, 0 \leq j < 2\}$$

$$|D_{2n}| = 2n$$

$$s^n = e, r^2 = e, rs = s^{-1}r$$

These relations are enough to completely determine D_{2n} .

What's group theory about?

Basic answer: study sets with one binary op. A better answer: group theory is study of symmetry. If we resize or rotate P_n , then symmetries are the same.

Kleinian view of geometry:

- D_{2n} captures what it means to be a regular n -gon
- More generally, geometry is about study of symmetries

1.6 Permutation groups

If X is a set, let $\text{Fun}(X, X)$ be set of functions $X \rightarrow X$. Then

$$\circ : \text{Fun}(X, X) \times \text{Fun}(X, X) \rightarrow \text{Fun}(X, X) : (f, g) \mapsto f \circ g$$

is an associative operation with an identity Id_X . Let $S_X = \{f \in \text{Fun}(X, X) : f \text{ is a bijection}\}$

Proposition 1.15

S_X is a group under \circ .

Proof:

See homework. □

symmetric/permutation group

Let $n \geq 1$. The symmetric group (or permutation group) S_n is the group S_X with $X = \{1, \dots, n\}$.

Elements of S_n are bijections $\pi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$

What makes a function $\pi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ a bijection?

Every element of $\{1, \dots, n\}$ must appear in the list $\pi(1), \dots, \pi(n)$, and no element can appear twice (\Leftarrow redundant by pig.-hole princ.)

How many elements in S_n ?

n choices for $\pi(1)$, $n - 1$ choices for $\pi(2)$, ..., 1 choice for $\pi(n)$. So $n(n - 1) \cdots 1 = n!$ choices $\implies |S_n| = n!$.

Note $|S_1| = 1! = 1$, so S_1 is the trivial group.

1.6.1 Representations

Elements of S_n are called **permutations**. There are a number of different ways to represent permutations:

1. **Two-line representation:**

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 5 & 1 & 4 & 2 & 3 \end{pmatrix}$$

2. **One-line representation:**

$$\pi = 651423$$

This representation saves space than the previous one, but it is hard to do operations in group theory. The one below seems counter-intuitive, but convenient for doing operations.

3. Note $\pi(1) = 6, \pi(6) = 3, \pi(3) = 1$. Say (163) is a **cycle** of π .

Disjoint cycle representation: write down cycles of π

$$\pi = (163)(25)(4) = (163)(25)$$

We typically drop cycles of length 1.

Identity is empty in disjoint cycle notation, so just use e .

The convention is that we start from the lowest item in the cycle, and sort the cycles by their lowest items.

Multiplication

Multiplication can be done in two-line or disjoint cycle notation

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 5 & 1 & 4 & 2 & 3 \end{pmatrix} = (163)(25)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 6 & 4 & 5 & 3 & 1 \end{pmatrix} = (126)(345)$$

$$\pi\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 3 & 4 & 2 & 1 & 6 \end{pmatrix} = (15)(234)$$

Note i comes from the right: $\pi(\sigma(i))$.

(It's a bit of a pain in one-line notation, so we don't use one-line notation often in group theory)

Inversion

We can also take inverse in two-line or disjoint cycle notation

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 5 & 1 & 4 & 2 & 3 \end{pmatrix} = (163)(25)$$

$$\pi^{-1} = \begin{pmatrix} 6 & 5 & 1 & 4 & 2 & 3 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix} \stackrel{*}{=} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 5 & 6 & 4 & 2 & 1 \end{pmatrix} = (136)(25)$$

*: swap two rows and sort the columns by the first row. Disjoint cycle notation is even easier.

If $\pi(i) = j$, then $\pi^{-1}(j) = i$, so cycles of π^{-1} are cycles of π in opposite order.

fixed points

The fixed points of a permutation $\pi \in S_n$ are the numbers $1 \leq i \leq n$ such that $\pi(i) = i$.

support set

The support set of $\pi \in S_n$ is

$$\text{supp}(\pi) = \{1 \leq i \leq n : \pi(i) \neq i\}$$

disjoint

π and σ are disjoint if $\text{supp}(\pi) \cap \text{supp}(\sigma) = \emptyset$

Example:

$$\text{supp}((163)(25)) = \{1, 2, 3, 5, 6\}$$

Remark:

In general, $\text{supp}(\pi)$ are numbers that appear in disjoint cycle representation of π (when cycles of length one are dropped).

$$\text{supp}(\pi) = \emptyset \text{ if and only if } \pi = e$$

$$\text{supp}(\pi^{-1}) = \text{supp}(\pi)$$

If $i \in \text{supp}(\pi)$, then $\pi(i) \in \text{supp}(\pi)$

commute

Two elements g, h in a group G commute if $gh = hg$.

Lemma 1.16

If $\pi, \sigma \in S_n$ are disjoint, then $\pi\sigma = \sigma\pi$.

Proof:

Suppose $1 \leq i \leq n$. If $i \in \text{supp}(i)$, then $\pi(i) \in \text{supp}(\pi)$. Since π, σ disjoint, $i, \pi(i) \notin \text{supp}(\sigma)$. So $\pi(\sigma(i)) = \pi(i) = \sigma(\pi(i))$.

By symmetry, $\pi(\sigma(i)) = \sigma(\pi(i))$ if $i \in \text{supp}(\sigma)$.

If $i \notin \text{supp}(\pi) \cup \text{supp}(\sigma)$, then $\pi(\sigma(i)) = i = \sigma(\pi(i))$.

So $\pi(\sigma(i)) = \sigma(\pi(i))$ for all $i \implies \pi\sigma = \sigma\pi$. □

1.6.2 Cycles

k-cycle

A k -cycle is an element of S_n with disjoint cycle notation $(i_1 i_2 \cdots i_k)$.

Suppose cycles of $\pi \in S_n$ are c_1, \dots, c_k . We can regard c_i as an element of S_n , $\pi = c_1 \cdot c_2 \cdots c_k$ as product in S_n . c_i and c_j are disjoint, so $c_i c_j = c_j c_i$. Note that order of cycles in disjoint cycle representation doesn't matter.

Example:

$$\pi = (163)(25) = (25) \cdot (163)$$

We can also get an interesting prospective on this formula for the inverse of π in the disjoint cycle notation. If c_1, \dots, c_k are cycles of π , then $\pi = c_1 c_2 \cdots c_k$ as product in S_n .

c_i and c_j are disjoint, so $c_i c_j = c_j c_i$.

$$\pi^{-1} = c_k^{-1} \cdots c_1^{-1} = c_1^{-1} \cdots c_k^{-1}$$

Example:

If c and c' are non-disjoint cycles, then they don't necessarily commute:

$$(12)(23) = (123) \text{ while } (23)(12) = (123)^{-1} = (132) \neq (12)(23).$$

If π is a permutation, then π commutes with π^i for all i since $\pi^{i+1} = \pi\pi^i = \pi^i\pi$, so π and π^i commute. However, note that they don't necessarily have disjoint support sets.

Subgroups

2.1 Subgroups

week 2

subgroup

Let (G, \cdot) be a group. A subset $H \subseteq G$ is a **subgroup** if

- (a) for all $g, h \in H$, $g \cdot h \in H$ (H is **closed under products**),
- (b) for all $g \in H$, $g^{-1} \in H$ (H is **closed under inverses**), and
- (c) $e_G \in H$.

Notation $H \leq G$.

Example:

$$\mathbb{Z} \leq \mathbb{Q}^+ := (\mathbb{Q}, +)$$

$$\mathbb{Q}_{>0} := \{x \in \mathbb{Q} : x > 0\} \leq \mathbb{Q}^\times.$$

To check this: if $x, y \in \mathbb{Q}$, $x, y > 0$, then $xy > 0 \implies xy \in \mathbb{Q}_{>0}$.

Also, if $x > 0$, then $1/x > 0 \implies 1/x \in \mathbb{Q}_{>0}$.

Example: More complicated

Let $G = D_{2n}$, s rotation.

$H = e = s^0, s, s^2, \dots, s^{n-1}$ is a subgroup of D_{2n} .

Proof:

Claim $s^i \in H$ for all $i \in \mathbb{Z}$.

Proof Let $i = nk + r, 0 \leq r < n$. Then $s^i = s^{nk+r} = (s^n)^k s^r = s^r$, since $s^n = e$. ■

Now check subgroup: if $s^i, s^j \in H$, then $s^{i+j} \in H$. If $s^i \in H$, then $s^{-i} \in H$. Finally, $e \in H$ by construction. □

H is the smallest subgroup containing s . The notation for H is $\langle s \rangle$.

Example: \mathbb{Z}

Let $G = \mathbb{Z} = (\mathbb{Z}, +)$.

If $m \in \mathbb{Z}$, then $m\mathbb{Z} := \{km : k \in \mathbb{Z}\} = \{n \in \mathbb{Z} : m|n\}$ is a subgroup of \mathbb{Z} .

In particular, if $m = 0$, then $0\mathbb{Z} = \{0\}$ is a subgroup of \mathbb{Z} , which is called the **trivial subgroup**.

trivial subgroup

If G is a group, $\{e\}$ is a subgroup called the **trivial subgroup**.

proper subgroup

Also, H is a subgroup of G . A subgroup H is **proper** if $H \neq G$. Notation: $H < G$.

H is proper nontrivial subgroup if $\{e\} \neq H < G$.

Example: Not subgroups

$\mathbb{Q}_{\geq 0} := \{x \in \mathbb{Q} : x \geq 0\}$ is not a subgroup of \mathbb{Q}^+ . We can verify as follows: If $x, y \in \mathbb{Q}_{\geq 0}$, then $x + y \in \mathbb{Q}_{\geq 0}$. Also $0 \in \mathbb{Q}_{\geq 0}$. But if $x \in \mathbb{Q}_{\geq 0}$, then $-x \notin \mathbb{Q}_{\geq 0}$ unless $x = 0$.

\mathbb{Q}^\times is not a subgroup of (\mathbb{Q}, \cdot) because (\mathbb{Q}, \cdot) is not a group.

Proposition 2.1

If H is a subgroup of (G, \boxtimes) , then $(H, \boxtimes|_{H \times H})$ is a group, such that

- (a) the identity of H is $e_H = e_G$, and
- (b) the inverse of $g \in H$ is the same as the inverse of g in G .

Proof:

First, why is $\boxtimes|_{H \times H}$ a binary operation on H ?

Recall \boxtimes is a function $G \boxtimes G \rightarrow G$ which implies $\boxtimes|_{H \times H}$ is a function $H \times H \rightarrow G$ if we restrict its domain. But if $g, h \in H$, then $g \boxtimes h \in H$. So we can think of $\boxtimes|_{H \times H}$ as function $H \times H \rightarrow H$. For the rest of this proof, we just denote this function by \boxtimes .

Since \boxtimes is associative, $\tilde{\boxtimes}$ is also associative.

$e_H = e_G$ is identity for $\tilde{\boxtimes}$.

If $g \in H$, then inverse g^{-1} with respect to \boxtimes is in H by the definition of subgroup.

Since $g\tilde{\boxtimes}g^{-1} = g\boxtimes g^{-1} = e_G = e_H$, and similarly $g^{-1}\boxtimes g = e_H$, g^{-1} is inverse of g with respect to $\tilde{\boxtimes}$.

So $(H, \tilde{\boxtimes})$ is a group. □

Call $\tilde{\boxtimes}$ the **operation induced by \boxtimes on H** . Usually just refer to $\tilde{\boxtimes}$ as \boxtimes .

Example:

\mathbb{Z} is subgroup \mathbb{Q} with operation $+$.

If H is group of (G, \cdot) , then H is group with operation \cdot .

Proposition 2.2

H is subgroup if and only if

- (a) H is non-empty, and
- (b) $gh^{-1} \in H$ for all $g, h \in H$.

Proof:

\Rightarrow If H is a subgroup of G , then $e_G \in H$, so $H \neq \emptyset$. Also if $g, h \in H$, then $h^{-1} \in H$, so $gh^{-1} \in H$.

\Leftarrow By (a), there is some element $x \in H$. In part (b), let $g = h := x$, then $xx^{-1} = e_G = e_H \in H$.

Also by (b), $e_G \cdot x^{-1} = x^{-1} \in H$ (closed under inverses).

If $x, y \in H$, then $y^{-1} \in H$, so $xy = x(y^{-1})^{-1} \in H$ (closed under inverses). □

Example:

Let $(V, +, \cdot)$ be a vector space.

If W is a subspace of V , then W is a subgroup of $(V, +)$.

Check:

- $0 \in W$ so W is non-empty.
- If $v, w \in W$, then $v - w \in W$.

Conclusion: W is subgroup.

Proposition 2.3

Suppose H is a finite subset of G . Then H is a subgroup of G if and only if

- (a) H is non-empty, and
- (b) $gh \in H$ for all $g, h \in H$.

Proof:

Since H is nonempty, suppose $g \in H$. By induction, we can show $g^n \in H$ for all $n \in \mathbb{N}$. Since H is finite, sequence $g, g^2, g^3, \dots \in H$ must eventually repeat. So $g^i = g^j$ for some $1 \leq i < j \implies g^n = e$ for $n = j - i$. Since $i < j$, then $n \geq 1$, therefore $g^n = e \in H$.

Now we need to show it is closed under inverses.

- $n = 1$, then $g = e = g^{-1}$.
- $n > 1$, then $g^{n-1} = g^{-1} \in H$.

□

2.2 Subgroups generated by a set

Proposition 2.4

Suppose \mathcal{F} is a non-empty set of subgroups of G . Then

$$L := \bigcap_{H \in \mathcal{F}} H$$

is a subgroup of G .

Proof:

First we check it is non-empty. Since $e_G \in H$ for all $H \in \mathcal{F}$, then $e_G \in K \implies K$ is non-empty.

Suppose $x, y \in K$, then

$$\begin{aligned} \implies x, y &\in H \quad \forall H \in \mathcal{F} \\ \implies y^{-1} &\in H \quad \forall H \in \mathcal{F} \\ \implies xy^{-1} &\in H \quad \forall H \in \mathcal{F} \\ \implies xy^{-1} &\in K \end{aligned}$$

By Proposition 2.3, K is a subgroup of G .

□

subgroup generated by S in G

Let S be a subset of group G . The **subgroup generated by S in G** is

$$\langle S \rangle := \bigcap_{S \subseteq H \leq G} H$$

Note

Intersection is non-empty because $S \subseteq G \leq G$.

If $S \subseteq K \leq G$, then $\langle S \rangle \subseteq K$. So say that $\langle S \rangle$ is smallest subgroup of G containing S .

To simplify the notation: If $S = \{s_1, s_2, \dots\}$, often write $\langle S \rangle = \langle s_1, s_2, \dots \rangle$.

We can write the trivial subgroup as $\langle \emptyset \rangle = \langle e \rangle = \{e\}$.

Example: D_{2n}

Let s be the rotation generator of D_{2n} . Let $K = \{s^0 = e, s^1, s^2, \dots, s^{n-1}\}$.

As previously checked, K is a subgroup of D_{2n} .

Since $s \in K, \langle s \rangle \subseteq K$.

On the other hand, can show by induction that $s^i \in \langle s \rangle$ for all $i \in \mathbb{Z}$.

So $K \subseteq \langle s \rangle \implies \langle s \rangle = K$.

$\langle s \rangle$ is constructed by taking all products of s with itself. Can we generalize this example?

Here we introduce a notation: If $S \subset G$, let $S^{-1} = \{s^{-1} : s \in S\}$.

Proposition 2.5

If $S \subset G$, let

$$K = \{e\} \cup \{s_1 \cdots s_k : k \geq 1, s_1, \dots, s_k \in S \cup S^{-1}\}$$

Then $\langle S \rangle = K$.

Proof:

Claim 1 $S \subseteq K \subseteq \langle S \rangle$

Proof It is easy to show that $S \subseteq K$. We simply let $k = 1$ and s_1 to be any element of S .

To show the second part, we know $e \in \langle S \rangle$. Then we can prove by induction that $s_1 \cdots s_k \in \langle S \rangle$ for all $k \geq 1, s_1, \dots, s_k \in S \cup S^{-1}$. ■

Claim 2 K is a subgroup of G .

Proof $e \in K$ by construction.

Suppose $x, y \in K$,

$$x = s_1 \cdots s_k, k \geq 0, s_1, \dots, s_k \in S \cup S^{-1}$$

$$y = t_1 \cdots t_\ell, \ell \geq 0, t_1, \dots, t_\ell \in S \cup S^{-1}$$

Then $xy = s_1 \cdots s_k t_1 \cdots t_\ell \in K$ by construction. Also, $x^{-1} = s_k^{-1} \cdots s_1^{-1} \in K$ since $s_k^{-1}, \dots, s_1^{-1} \in S \cup S^{-1}$. So K is a subgroup. ■


$S \subseteq K$, and $\langle S \rangle$ is smallest subgroup containing $S \implies \langle S \rangle \subseteq K$. Thus $\langle S \rangle = K$. □

2.2.1 Lattice of subgroups

Before concluding this section, it is interesting to mention one closed related subject which the lattice of subgroups of G .

Subgroups of G are ordered by set inclusion \subseteq . If $H_1, H_2 \leq G$, and $H_1 \subseteq H_2$, then $H_1 \leq H_2$, so we also write this order as \leq . Set of subgroups of G with order \leq is called the **lattice of subgroups of G** . We don't need to deal with formal definitions and properties here.

The picture below shows the subgroups of \mathbb{Z} , where $k\mathbb{Z}$ denotes the set containing all integers that are divisible by k .



Subgroup below $H_1, H_2 \leq G$ in the lattice is $H_1 \cap H_2$. In the picture above, it is $2\mathbb{Z} \cap 3\mathbb{Z} = 6\mathbb{Z}$. Intuitively, a number is divisible by 2 and 3, which is the same thing as being divisible by 6.

What about the subgroup above H_1 and H_2 ? The subgroup above H_1, H_2 is $\langle H_1 \cup H_2 \rangle$.

2.3 Cyclic groups

generate

A subset S of a group G **generates** G if $\langle S \rangle = G$.

cyclic

A group G is **cyclic** if $G = \langle a \rangle$ for some $a \in G$.

Example:

$\mathbb{Z} = \langle 1 \rangle = \langle -1 \rangle$ (generators are not unique)

$\mathbb{Z}/n\mathbb{Z} = \langle [1] \rangle = \langle -[1] \rangle$

\mathbb{Q}^+ is not cyclic

If G is a group, then $\langle a \rangle$ is a cyclic group for any $a \in G$ (called the **cyclic subgroup generated by a**).

Lemma 2.6

1. If $a \in G$, then $\langle a \rangle = \{a^i : i \in \mathbb{Z}\}$.
2. If $|a| = n$, then $\langle a \rangle = \{a^i : 0 \leq i < n\}$.

Proof:

1. Follows from Proposition 2.5 about $\langle S \rangle$.
2. See argument for $\langle s \rangle$ in D_{2n} .

□

Remark:

In the first part of Lemma 2.6, it does not mean each element in the subgroup can be uniquely represented in the form of a^i .

Then we have two questions:

- In (2), can $|\langle a \rangle|$ be smaller than n ?
- Does $|\langle a \rangle|$ determine $|a|$?

Proposition 2.7

If $G = \langle a \rangle$, then $|G| = |a|$.

This proposition also applies to infinite groups.

Proof:

From part 2 of Proposition 2.6, we know that there are at most n elements in $\langle a \rangle$, then $|G| \leq |a|$.

Suppose $|G| = n < +\infty$. Then the sequence $a^0, a^1, \dots, a^n \in G$ must have repetition. Thus there is $0 \leq i < j \leq n$ with $a^i = a^j$. Then with the similar argument before, $a^{j-i} = e$, which implies that $|a| \leq n$.

Thus $|a| \leq |G| \implies |a| = |G|$.

□

Remark:

It is worth thinking that what happens if $|G| = \infty$ and it seems the proof only works with finite order. If $|G| = \infty$, then $|G| \leq |a|$ will force $|a|$ to be infinite.

Example: \mathbb{Z}

$G = \mathbb{Z}$:

- Infinite cyclic group
- Generators are $+1$ and -1

- Order of $m \in \mathbb{Z}$ is

$$|m| = \begin{cases} +\infty & m \geq 0 \\ 1 & m = 0 \end{cases}$$

- Cyclic subgroups: $\langle m \rangle = m\mathbb{Z} = \{km : k \in \mathbb{Z}\}$. (Note difference in $\langle a \rangle$ in additive and multiplicative notation)

All subgroups of \mathbb{Z} are cyclic

2.3.1 $\mathbb{Z}/n\mathbb{Z}$

Can we analyze $\mathbb{Z}/n\mathbb{Z}$ in the same way? Recall $\mathbb{Z}/n\mathbb{Z}$ is the set of congruence classes mod n . We denote congruence class of $a \in \mathbb{Z}$ by $[a]$, or just a . For example, in $\mathbb{Z}/5\mathbb{Z}$, $3 = 8$.

Then we might wonder:

- What are the generators?
- What are the orders of elements?
- What are the subgroups?

Before we explore these questions, it is nice to have the following lemma which works for arbitrary group G .

Generators

Lemma 2.8

Suppose $G = \langle S \rangle$. Then $G = \langle T \rangle$ if and only if $S \subseteq \langle T \rangle$.

Proof:

It's relatively easy to prove.

\Rightarrow If $G = \langle T \rangle$, and we know $S \subseteq G$, then $S \subseteq \langle T \rangle$.

\Leftarrow If $S \subseteq \langle T \rangle$, and we know $\langle S \rangle$ is the smallest subgroup containing S , then $\langle T \rangle$ must contain the subgroup generated by S , which is $\langle S \rangle = G$, thus $G \subseteq \langle T \rangle$. And $\langle T \rangle$ is a subgroup as well, then $G = \langle T \rangle$. \square

What does this mean in our example? So $\mathbb{Z}/n = \langle [a] \rangle$ if and only if $[1] \in \langle [a] \rangle$.

$$\begin{aligned} [1] \in \langle [a] \rangle &\iff xa = 1 \pmod{n} \text{ for some } x \in \mathbb{Z} \\ &\iff xa - 1 = yn \text{ for some } x, y \in \mathbb{Z} \\ &\iff xa + yn = 1 \text{ for some } x, y \in \mathbb{Z} \\ &\iff \gcd(a, n) = 1 \end{aligned}$$

So $\langle [a] \rangle = \mathbb{Z}/n\mathbb{Z}$ if and only if $\gcd(a, n) = 1$.

Order of elements

Lemma 2.9

If G is a group, $g \in G$, $g^n = e$, then $|g| \mid n$.

Proof:

Exercise. □

If $a \in \mathbb{Z}$, then $n[a] = 0$, so $|[a]| \mid n$.

Lemma 2.10

Suppose $a \mid n$. Then $|[a]| = \frac{n}{a}$.

Proof:

If $n = ka$, then $\ell[a] \neq 0$ for $1 \leq \ell < k$ and $k[a] = [ka] = 0$, so $|[a]| = k$. □

Lemma 2.11

Suppose $a \in \mathbb{Z}$, and let $b = \gcd(a, n)$. Then $\langle [a] \rangle = \langle [b] \rangle$.

Proof:

Since $b \mid a$, there is k such that $a = kb$, then $[a] \in \langle [b] \rangle \implies \langle [a] \rangle \subseteq \langle [b] \rangle$.

By properties of \gcd , there is $x, y \in \mathbb{Z}$ such that $xa + yn = b$. So $[b] = x[a] \implies [b] \in \langle [a] \rangle \implies \langle [b] \rangle \subseteq \langle [a] \rangle$.

Therefore $\langle [a] \rangle = \langle [b] \rangle$. □

Using these lemmas, we can find order for a general element in $\mathbb{Z}/n\mathbb{Z}$.

Proposition 2.12

Suppose $a \in \mathbb{Z}$. Then

$$|[a]| = \frac{n}{\gcd(a, n)}$$

Proof:

Let $b = \gcd(a, n)$. Then $\langle [a] \rangle = \langle [b] \rangle$. So

$$|[a]| = |\langle [a] \rangle| = |\langle [b] \rangle| = |[b]|$$

Finally

$$|[b]| = \frac{n}{b}$$

□

Subgroups

Corollary 2.13

Let $n \geq 1$.

- The order d of any cyclic subgroup of $\mathbb{Z}/n\mathbb{Z}$ divides n .
- For every $d|n$, there is a unique subgroup of $\mathbb{Z}/n\mathbb{Z}$ of order d . It is generated by $[a]$, where $a = \frac{n}{d}$.

Proof:

If $|\langle [a] \rangle| = d$, then $d = |[a]| \mid n$ by Lemma 2.9. Also, $d = \frac{n}{\gcd(a, n)}$, and by Lemma 2.11, $\langle [a] \rangle = \langle [\frac{n}{d}] \rangle$.

Conversely, if $d|n$ and $a = \frac{n}{d}$, then $|\langle [a] \rangle| = d$. □

Example:

Cyclic subgroups of $\mathbb{Z}/6\mathbb{Z}$ are

- $\langle 6 \rangle = \{0\}$
- $\langle 2 \rangle = \{0, 2, 4\}$
- $\langle 3 \rangle = \{0, 3\}$
- $\langle 1 \rangle = \{0, 1, 2, 3, 4, 5\} = \mathbb{Z}/6\mathbb{Z}$.

Cyclic subgroups of $\mathbb{Z}/p\mathbb{Z}$, p prime

- $\langle p \rangle = \langle 0 \rangle$
- $\langle 1 \rangle = \mathbb{Z}/p\mathbb{Z}$

Every subgroup of a cyclic group is cyclic. So Corollary 2.13 is a complete list of subgroups of $\mathbb{Z}/n\mathbb{Z}$. Every cyclic group is isomorphic to one of $\mathbb{Z}/n\mathbb{Z}$, $n \geq 1$, or \mathbb{Z} .

Homomorphisms

3.1 Homomorphisms

homomorphism

Let G and H be groups. A function $\phi : G \rightarrow H$ is a **homomorphism** (or **morphism**) if

$$\phi(g \cdot h) = \phi(g) \cdot \phi(h)$$

for all $g, h \in G$.

Example:

\mathbb{K} field, $\mathbb{K}^\times = \{a \in \mathbb{K}, a \neq 0\}$ with operation \cdot .

$\text{GL}_n \mathbb{K} \rightarrow \mathbb{K}^\times : A \mapsto \det(A)$ is a homomorphism because $\det(AB) = \det(A) \det(B)$ for all invertible matrices A, B .

Let $\mathbb{R}_{>0} = \{x \in \mathbb{R} : x > 0\} \leq \mathbb{R}^\times$. $\mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0} : x \mapsto \sqrt{x}$ is a homomorphism since $\sqrt{xy} = \sqrt{x}\sqrt{y}$.

Additive notation: $\phi : (G, +) \rightarrow (H, +)$ is a homomorphism if $\phi(x + y) = \phi(x) + \phi(y)$ for all $x, y \in G$. For example, $\phi : \mathbb{Z} \rightarrow \mathbb{Z} : k \mapsto mk$ is a homomorphism for any $m \in \mathbb{Z}$, since

$$\phi(x + y) = m(x + y) = mx + my = \phi(x) + \phi(y) \quad \forall x, y \in \mathbb{Z}$$

If V, W are vector spaces, and $T : V \rightarrow W$ is a linear transformation, then T is a homomorphism from $(V, +)$ to $(W, +)$, since $T(v + w) = T(v) + T(w)$ for all $v, w \in V$.

Mixed notation: $\mathbb{R}^+ \rightarrow \mathbb{R}^\times : x \mapsto e^x$ is a homomorphism since $e^{x+y} = e^x \cdot e^y$ for all $x, y \in \mathbb{R}^+$.

$\mathbb{R}^+ \rightarrow \mathbb{R}^+ : x \mapsto e^x$ is not a homomorphism since $e^{x+y} \neq e^x + e^y$ for some $x, y \in \mathbb{R}^+$ (e.g. $x = y = 0$).

Lemma 3.1

Suppose $\phi : G \rightarrow H$ is a homomorphism. Then

- (a) $\phi(e_G) = e_H$
- (b) $\phi(g^{-1}) = \phi(g)^{-1}$
- (c) $\phi(g^n) = \phi(g)^n$ for all $n \in \mathbb{Z}$
- (d) $|\phi(g)| \mid |g|$ for all $g \in G$ ($n \mid \infty$ for all $n \in \mathbb{N}$)

Proof:

- (a) $\phi(e_G) = \phi(e_G^2) = \phi(e_G) \cdot \phi(e_G)$
so $e_H = \phi(e_G)^{-1} \cdot \phi(e_G) = \phi(e_G)^{-1} \cdot \phi(e_G) \cdot \phi(e_G) = \phi(e_G)$.
- (b) $e_H = \phi(e_G) = \phi(gg^{-1}) = \phi(g)\phi(g^{-1})$ and similarly $\phi(g^{-1})\phi(g) = e_H$, so $\phi(g^{-1})$ is the unique inverse of $\phi(g)$.
- (c) Use induction for $n \geq 0$, use part (b) for $n < 0$.
- (d) If $|g| = n < +\infty$, then $g^n = e_G$ so $\phi(g)^n = \phi(g^n) = \phi(e_G) = e_H$. This implies^a $|\phi(g)| \mid n$.

□

^aSee homework

Lemma 3.2

If $H \leq G$, and H is considered as a group with the induced operation from G , then $i : H \rightarrow G : x \mapsto x$ is a homomorphism.

Proof:

$$i(g \cdot h) = g \cdot h = i(g) \cdot i(h)$$

□

Lemma 3.3

If $\phi : G \rightarrow H$ and $\psi : H \rightarrow K$ are homomorphisms, then $\psi \circ \phi$ is a homomorphism.

Proof:

$$\psi \circ \phi(g \cdot h) = \psi(\phi(g) \cdot \phi(h)) = \psi(\phi(g)) \cdot \psi(\phi(h)).$$

□

Corollary 3.4

If $\phi : G \rightarrow H$ is a homomorphism, $K \leq G$, then the **restriction** $\phi|_K$ is a homomorphism.

Proof:

$$\phi_K = \phi \circ i, \text{ where } i : K \rightarrow G \text{ is the inclusion } x \mapsto x.$$

□

3.2 Homomorphisms and subgroups

If $f : X \rightarrow Y$ is a function, $S \subseteq X$, then $f(S) := \{f(x) : x \in S\}$

Proposition 3.5

If $\phi : G \rightarrow H$ is a homomorphism, and $K \leq G$, then $\phi(K) \leq H$.

Proof:

Since K is non-empty, $\phi(K)$ is non-empty.

If $x, y \in \phi(K)$, then $x = \phi(x_0), y = \phi(y_0)$ for $x_0, y_0 \in K$.

So $xy^{-1} = \phi(x_0)\phi(y_0)^{-1} = \phi(x_0)\phi(y_0^{-1}) = \phi(x_0y_0^{-1}) \in \phi(K)$, since $x_0y_0^{-1} \in K$. \square

image

If $\phi : G \rightarrow H$ is a homomorphism, the **image** of ϕ is the subgroup $\text{Im } \phi = \phi(G) \leq H$.

Example:

Let $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^\times : x \mapsto e^x$. $e^x > 0$ for all $x \in \mathbb{R}$, so $\text{Im } \phi \subseteq \mathbb{R}_{>0}$. If $y \in \mathbb{R}_{>0}$, then $y = \phi(\log y)$, so $\text{Im } \phi = \mathbb{R}_{>0}$.

If $K \leq G$ and $i : K \rightarrow G$ is inclusion, then $\text{Im } i = K$.

$\phi : \mathbb{Z} \rightarrow \mathbb{Z} : k \mapsto mk$ for some $m \in \mathbb{Z}$. $\phi(\mathbb{Z}) = m\mathbb{Z}$.

Lemma 3.6

If $\phi : G \rightarrow H$ is a homomorphism with $\text{Im } \phi \leq K \leq H$, then the function $\tilde{\phi} : G \rightarrow K : x \mapsto \phi(x)$ is also a homomorphism with $\text{Im } \tilde{\phi} = \text{Im } \phi \leq K$.

Proof:

$$\begin{aligned} \tilde{\phi}(x \cdot y) &= \phi(x \cdot y) \\ &= \phi(x) \cdot \phi(y) \text{ in } H \\ &= \tilde{\phi}(x) \cdot \tilde{\phi}(y) \text{ in } K \end{aligned}$$

Also $\tilde{\phi}(G) = \phi(G)$, regarded as a subset of K . \square

Usually just refer to $\tilde{\phi}$ as ϕ .

Lemma 3.7

A homomorphism $\phi : G \rightarrow H$ is surjective if and only if $\text{Im } \phi = H$.

Proof:

Obvious from definition. \square

Corollary 3.8

ϕ induces a surjective homomorphism $\tilde{\phi} : G \rightarrow K$, where $K = \text{Im } \phi$.

Remark:

From Lemma 3.7, if ϕ is not surjective, then $\text{Im } \phi < H$, then we can let $K = \text{Im } \phi$, and then construct a surjective homomorphism by Lemma 3.6.

Because this is a bit abstract, it is helpful to go through some examples.

Recall the previous example: Let $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^\times : x \mapsto e^x$. $e^x > 0$ for all $x \in \mathbb{R}$. This is not surjective, because $\text{Im } \phi = \mathbb{R}_{>0}$. If we restrict the codomain to be $\mathbb{R}_{>0}$, then it is surjective.

Similarly for $\phi : \mathbb{Z} \rightarrow \mathbb{Z} : k \mapsto mk$ for some $m \in \mathbb{Z}$, but it induced surjective homomorphism $\mathbb{Z} \rightarrow m\mathbb{Z}$.

Proposition 3.9

Let $\phi : G \rightarrow H$ be a homomorphism. If $S \subseteq G$, then $\phi(\langle S \rangle) = \langle \phi(S) \rangle$.

Proof:

$\phi(S^{-1}) = \{\phi(s^{-1}) : s \in S\} = \{\phi(s)^{-1} : s \in S\} = \phi(S)^{-1}$. So

$$\begin{aligned} \phi(\langle S \rangle) &= \phi(\{s_1 \cdots s_k : k \geq 0, s_1, \dots, s_k \in S \cup S^{-1}\}) \\ &= \{\phi(s_1) \cdots \phi(s_k) : k \geq 0, s_1, \dots, s_k \in S \cup S^{-1}\} \\ &= \{t_1 \cdots t_k : k \geq 0, t_1, \dots, t_k \in \phi(S) \cup \phi(S)^{-1}\} \\ &= \langle \phi(S) \rangle \end{aligned}$$

□

Remark:

We used the fact that $\phi(S \cup S^{-1}) = \phi(S) \cup \phi(S^{-1})$, but it doesn't work for intersection.

If $f : X \rightarrow Y$ is a function, and $S \subseteq Y$, then $f^{-1}(S) := \{x \in X : f(x) \in S\}$.

Proposition 3.10

If $\phi : G \rightarrow H$ is a homomorphism, $K \leq H$, then $\phi^{-1}(K) \leq G$.

Proof:

$\phi(e_G) = e_H \in K$, so $e_G \in \phi^{-1}(K)$.

If $x, y \in \phi^{-1}(K)$, then $\phi(x), \phi(y) \in K$. Thus $\phi(xy^{-1}) = \phi(x)\phi(y)^{-1} \in K$. Hence $xy^{-1} \in \phi^{-1}(K)$. Thus it is a subgroup of G . □

kernel

If $\phi : G \rightarrow H$ is a homomorphism, then the **kernel** of ϕ is the subgroup $\ker \phi := \phi^{-1}(e_H) = \{g \in G : \phi(g) = e_H\} \leq G$.

Example:

For $\det : \mathrm{GL}_n \mathbb{K} \rightarrow \mathbb{K}^\times$, $\ker \det = \{A \in \mathrm{GL}_n : \det(A) = 1\}$.

This subgroup of $\mathrm{GL}_n \mathbb{K}$ is called the **special linear group**, and is denoted by $\mathrm{SL}_n \mathbb{K}$.

If $\phi : \mathbb{Z} \rightarrow \mathbb{Z} : k \mapsto mk$, then $\phi(k) = 0$ if and only if $mk = 0$, so

$$\ker \phi = \begin{cases} \{0\} & m \neq 0 \\ \mathbb{Z} & m = 0 \end{cases}$$

If $\phi : \mathbb{R} \rightarrow \mathbb{R}^\times : x \mapsto e^x$, then $e^x = 1$ if and only if $x = 0$, so $\ker \phi = \{0\}$.

We can generalize the last example into the following proposition.

Proposition 3.11

A homomorphism $\phi : G \rightarrow H$ is injective if and only if $\ker \phi = \{e_G\}$.

Proof:

\Rightarrow If ϕ is injective, then $\phi(x) = \phi(e_H) = \phi(e_G)$ if and only if $x = e_G$, so $\ker \phi = \{e_G\}$.

\Leftarrow Suppose $\ker \phi = \{e_G\}$, and $\phi(x) = \phi(y)$. Then $\phi(xy^{-1}) = \phi(x)\phi(y)^{-1} = e_H$, so $xy^{-1} \in \ker \phi$.

But then $xy^{-1} = e_G$, so $x = y$ which implies that ϕ is injective. \square

3.2.1 Application: subgroups of cyclic groups**Proposition 3.12**

If H is a subgroup of a cyclic group G , then H is cyclic.

Proof:

We need following facts:

1. All subgroups of \mathbb{Z} are of the form $m\mathbb{Z} = \langle m \rangle$, hence cyclic.
2. G is cyclic if and only if there is surjective homomorphism $\mathbb{Z} \rightarrow G$.
3. If $f : X \rightarrow Y$ is a surjective function, and $S \subseteq Y$, then $f(f^{-1}(S)) = S$.

The first two are in the homework. The last one is not hard to see.

Since G is cyclic, there is a surjective homomorphism $\phi : \mathbb{Z} \rightarrow G$.

Since all subgroups of \mathbb{Z} are cyclic, there is $m \in \mathbb{Z}$ such that $\phi^{-1}(H) = \langle m \rangle$.

Let $\psi : \mathbb{Z} \rightarrow \mathbb{Z}$ be homomorphism with $\psi(k) = mk$.

Then $\phi \circ \psi : \mathbb{Z} \rightarrow G$ is homomorphism.

$$\phi \circ \psi(\mathbb{Z}) = \phi(m\mathbb{Z}) = \phi(\phi^{-1}(H)) = H.$$

Then we can restrict codomain of $\phi \circ \psi$ to get surjective homomorphism $\mathbb{Z} \rightarrow H$.

Hence H is cyclic. □

3.3 Isomorphisms

in/sur/bi-jective

Let $f : X \rightarrow Y$ be a function. Then f is:

1. **injective** if for all $x, y \in X$, $f(x) = f(y) \implies x = y$,
2. **surjective** if for all $y \in Y$, $\exists x \in X$ with $f(x) = y$, and
3. **bijective** if f is both injective and surjective.

Proposition 3.13

$f : X \rightarrow Y$ is a bijection if and only if there is a function $g : Y \rightarrow X$ such that $f \circ g = 1_Y$ and $g \circ f = 1_X$.

If g exists, then it is unique, and we denote it by f^{-1} .

isomorphism

A homomorphism $\phi : G \rightarrow H$ is an **isomorphism** if ϕ is a bijection.

Lemma 3.14

$\phi : G \rightarrow H$ is an isomorphism if and only if $\ker \phi = \{e_G\}$ and $\text{Im } \phi = H$.

Example:

$\mathbb{R}^+ \rightarrow \mathbb{R}_{>0} : x \mapsto e^x$ is an isomorphism.

If $\phi : G \rightarrow H$ is injective, then ϕ induces an isomorphism $G \rightarrow \text{Im } \phi$.

Proposition 3.15

Suppose $\phi : G \rightarrow H$ is an isomorphism. Then $\phi^{-1} : H \rightarrow G$ is also an isomorphism.

Proof:

ϕ^{-1} is also a bijection, so just need to show that it is a homomorphism.

If $g, h \in H$, then

$$\phi(\phi^{-1}(g) \cdot \phi^{-1}(h)) = \phi(\phi^{-1}(g))\phi(\phi^{-1}(h)) = g \cdot h$$

So ϕ^{-1} is a homomorphism, hence isomorphism. \square

Corollary 3.16

A homomorphism $\phi : G \rightarrow H$ is an isomorphism if and only if there is a homomorphism $\psi : H \rightarrow G$ such that

- $\psi \circ \phi = 1_G$, and
- $\phi \circ \psi = 1_H$.

Proof:

\Rightarrow If ϕ is an isomorphism, then can take $\psi = \phi^{-1}$.

\Leftarrow If ψ exists, then ϕ is a bijection.

 \square **isomorphic**

We say that G and H are **isomorphic** if there is an isomorphism $\phi : G \rightarrow H$.

Notation: $G \cong H$.

Key facts:

- If $G \cong H$ then $H \cong G$.

Proof:

If $\phi : G \rightarrow H$ is an isomorphism, then $\phi^{-1} : H \rightarrow G$ is an isomorphism. \square

- If $G \cong H$ and $H \cong K$ then $G \cong K$.

Proof:

If $\phi : G \rightarrow H$ is an isomorphism and $\psi : H \rightarrow K$ is an isomorphism, then $\psi \circ \phi$ is an isomorphism. \square

- $G \cong G$.

Proof:

$1_G : G \rightarrow G$ is an isomorphism. \square

Idea If $G \cong H$, then G and H are identical as groups.

If $\phi : G \rightarrow H$ is an isomorphism, then

- $|G| = |H|$

- G is abelian if and only if H is abelian
- $|g| = |\phi(g)|$ for all $g \in G$
- $K \subseteq G$ is a subgroup of G if and only if $\phi(K)$ is a subgroup of H

Proposition 3.17

If G and H are cyclic groups, then $G \cong H$ if and only if $|G| = |H|$.

Proof:

Suppose $|G| = \langle a \rangle$, $H = \langle b \rangle$.

\Leftarrow Assume that $|G| = |H|$.

Claim $a^i = a^j$ for $i < j$ if and only if $|a| \mid j - i$.

Proof

\Leftarrow If $a^i = a^j$ then $a^{j-i} = e$.

\Rightarrow If $|a| \mid j - i$, then $j - i = k|a|$. So $a^{j-i} = a^{k|a|} = e \implies a^j = a^i$. ■

Note: if $|a| = +\infty$, $a^i \neq a^j$ for all $i \neq j \in \mathbb{Z}$.

Then we define a function $\phi : G \rightarrow H : a^i \mapsto b^i$.

Well-defined? $|a| = |G| = |H| = |b|$.

$a^i = a^h \implies |a| \mid j - i \implies |b| \mid j - i \implies b^i = b^j$

Homomorphism? $\phi(a^i \cdot a^j) = \phi(a^{i+j}) = b^{i+j} = b^i \cdot b^j = \phi(a^i) \cdot \phi(a^j)$ for all $a^i, a^j \in G$.

Inverse? $\psi : H \rightarrow G : b^i \mapsto a^i$ is well-defined. Clearly ψ is inverse to ϕ .

Thus ϕ is isomorphism $\implies G \cong H$.

\Rightarrow If $G \cong H$, then $|G| = |H|$ which holds for all groups. Same cardinality thus same order.

□

Corollary 3.18

Suppose G is a cyclic group.

- If $|G| = +\infty$, then $G \cong \mathbb{Z}$.
- If $|G| = n < +\infty$, then $G \cong \mathbb{Z}/n\mathbb{Z}$.

Corollary 3.19

Cyclic groups are abelian.

multiplicative form of cyclic groups

Let a be formal indeterminate (can use any letter). Let

- $C_\infty = \{a^i : i \in \mathbb{Z}\}, a^i \cdot a^j = a^{i+j}$
- $C_n = \{a^i : i \in \mathbb{Z}/n\mathbb{Z}\}, a^i \cdot a^j = a^{i+j}$

Of course we have $C_\infty \cong \mathbb{Z}$ via $a^i \mapsto i$, and $C_n \cong \mathbb{Z}/n\mathbb{Z}$ via $a^i \mapsto i$.

3.4 Cosets

week 3

Recall linear subspaces are motivation for definition of subgroups. Let $T : V \rightarrow W$ be a linear transformation. (so T is also a group homomorphism $(V, +) \rightarrow (W, +)$). $\ker T = \{x \in V : T(x) = 0\} = \text{“solutions to } Tx = 0\text{”}$.

What are solutions to $Tx = b$?

They can be empty: $Tx = b$ has a solution if and only if $b \in \text{Im } T$. If $b \in \text{Im } T$, and $Tx = b$ has a solution x_0 , then all other solutions are of the form $x_0 + x_1$, for $x_1 \in \ker T$.

Conclusion: space of solutions has form $x_i + \ker T$. $x_0 + \ker T$ is called an **affine** subspace. (it's like a linear subspace, but doesn't have to contain 0). We can still talk about the dimension.

coset

If $S \subseteq G$, and $g \in G$, we let

$$gS = \{gh : h \in S\} \quad \text{and} \quad Sg = \{hg : h \in S\}$$

If $H \leq G$, gH is called a **left coset** of H in G and Hg is called a **right coset** of H in G .

Remark:

We also refer these sets: left/right translate of S by g .

For abelian groups, $gH = Hg$.

Additive notation: coset of H in $(G, +)$ is $g + H$.

Example:

U subspace of vector space $(V, +, \cdot)$, cosets of U are affine subspaces $v + U$ for $v \in V$.

Given $m \in \mathbb{Z}$, cosets of $m\mathbb{Z}$ are sets

$$a + m\mathbb{Z} = \{a + km : k \in \mathbb{Z}\} = \{x \in \mathbb{Z} : x \equiv a \pmod{m}\}$$

We can think of the cosets as the sets of solutions to system of equations.

Example: Dihedral group $\langle s \rangle$

Recall $D_{2n} = \{s^i r^j : 0 \leq i < n, j \in \{0, 1\}\}$.

Let $H = \langle s \rangle = \{e = s^0, s^1, \dots, s^{n-1}\}$

What are the right cosets of H ?

$$\begin{aligned} H &= He \\ Hr &= \{r, sr, \dots, s^{n-1}r\} \\ Hs^i &= \{s^i, s^{i+1}, \dots, s^{n-1}, e, s^1, \dots, s^{i-1}\} = H \\ Hs^i r &= \{s^i r, s^{i+1}r, \dots, s^{n-1}r, r, sr, \dots, s^{i-1}r\} = Hr \end{aligned}$$

Conclusion: right cosets are H and Hr .

Also $D_{2n} = H \sqcup Hr$, where \sqcup is disjoint union.

What about the left cosets of $H = \langle s \rangle$?

Exercise

- use $rs = s^{-1}r$ to show $s^i = rs^{-i}$ for all $i \in \mathbb{Z}$.
- if $S \subseteq G$, $g, h \in G$, then $ghS = g(hS)$. This follows from the associativity of the group.

With these facts,

$$\begin{aligned} s^i H &= H \\ s^i r H &= r s^{-i} H = r H \end{aligned}$$

Conclusion: left cosets of H are H, rH

$$\begin{aligned} rH &= \{r, rs, rs^2, \dots, rs^{n-1}\} \\ &= \{r, s^{-1}r, s^{-2}r, \dots, s^{1-n}r\} \\ &= \{r, s^{-1}r, s^{-2}r, \dots, sr\} \\ &= \{r, s^{n-1}r, s^{n-2}r, \dots, sr\} \\ &= Hr \end{aligned}$$

Example: Dihedral group $\langle r \rangle$

What about $H = \langle r \rangle = \{e, r\}$?

Left cosets: $rH = \{r, e\} = H$ and $s^i H = \{s^i, s^i r\} = s^i r H$.

Conclusion: Left cosets are $s^i H, 0 \leq i < n$, and

$$D_{2n} = \bigsqcup_{i=0}^{n-1} s^i H$$

Right cosets: $Hr = \{r, e\} = H$ and $HS^i = \{s^i, rs^i\} = \{s^i, s^{-i}r\}$
 $HS^i r = \{s^i r, s^{-i}\} = HS^{-i}$

Conclusion: Right cosets are $HS^i, 0 \leq i < n$, and $D_{2n} = \bigsqcup_{i=0}^{n-1} HS^i$.

In this case, left cosets and right cosets are different.

set of left/right cosets

If $H \leq G$, let

$$G/H = \{gH : g \in G\} = \{S \subseteq G : S = gH \text{ for some } g \in G\}$$

be the **set of left cosets** of H in G , and

$$H \backslash G = \{Hg : g \in G\} = \{S \subseteq G : S = Hg \text{ for some } g \in G\}$$

be the **set of right cosets** of H in G .

Remark:

It is read as $G \bmod H$. We count each subset once.

We are very interested in trying to understand G/H and $H \backslash G$.

Example: D_{2n}

$$D_{2n}/\langle s \rangle = \{\langle s \rangle, r\langle s \rangle\}$$

$$D_{2n}/\langle r \rangle = \{s^i \langle r \rangle, 0 \leq i < n\}$$

Example: $\mathbb{Z}/n\mathbb{Z}$

Consider $n\mathbb{Z} \leq \mathbb{Z}$.

$a + n\mathbb{Z} = \{x \in \mathbb{Z} : x \equiv a \pmod{n}\} =: [a]$. Thus

$$\begin{aligned} \mathbb{Z}/n\mathbb{Z} &= \{a + n\mathbb{Z} : a \in \mathbb{Z}\} \\ &= \{a + n\mathbb{Z} : 0 \leq a < n\} \\ &= \{[a] : 0 \leq a < n\} \end{aligned}$$

Big question for next week: for $H \leq G$, is G/H always a group? spoiler: no...

Suppose $\phi : G \rightarrow K$ is a homomorphism, let $H = \ker \phi$. Note that $\phi(x) = b$ has a solution x for $b \in K$ if and only if $b \in \text{Im } \phi$.

Lemma 3.20

Suppose $\phi(x_0) = b$. The set of solutions $\phi^{-1}(\{b\})$ to $\phi(x) = b$ is $x_0H = Hx_0$.

Proof:

Suppose $\phi(x_1) = b$. Then $\phi(x_0^{-1}x_1) = \phi(x_0)^{-1}\phi(x_1) = b^{-1}b = e$. Thus $x_0^{-1}x_1 \in H$. Therefore $x_1 = x_0(x_0^{-1}x_1) \in Hx_0$.

Conversely, if $x_1 = x_0h$ for $h \in H$, then $\phi(x_1) = \phi(x_0h) = \phi(x_0)\phi(h) = be = b$. Therefore, every element of x_0H is a solution.

Same argument for right cosets shows set of solutions is Hx_0 . \square

In this case, left cosets are right cosets.

Lemma 3.21

Suppose $\phi(x_0) = b$. Then set of solutions to $\phi(x) = b$ is $x_0 \cdot \ker \phi$.

Proposition 3.22

If $\phi : G \rightarrow K$ is a homomorphism, then there is a bijection between $G/\ker \phi$ and $\text{Im } \phi$.

Proof:

$g \cdot \ker \phi \in G/\ker \phi$ is the set of solutions to $\phi(x) = b$ where $b = \phi(g)$. As a result, $\phi(g \cdot \ker \phi) = \{b\}$, $b \in \text{Im } \phi$.

In the other direction, given $b \in \text{Im } \phi$, $g \ker \phi = \phi^{-1}(\{b\})$.

From Lemma 3.21, we see these two mappings are inverses of each other, thus bijection. \square

Example:

Suppose $G = \mathbb{Z}$, $K = \mathbb{Z}/n\mathbb{Z}$.

From tutorial: there is a homomorphism $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z} : a \mapsto [a]$.

$\ker \phi = n\mathbb{Z}$, $\text{Im } \phi = \mathbb{Z}/n\mathbb{Z}$.

Elements of $\mathbb{Z}/n\mathbb{Z} = \{[a] : 0 \leq a < n\} = \{a + n\mathbb{Z} : 0 \leq a < n\}$

$a + n\mathbb{Z}$ is the set of solutions of $[x] \equiv [a]$ in $\mathbb{Z}/n\mathbb{Z}$.

3.5 The index and Lagrange's theorem

Given $H \leq G$, how many left cosets does H have in G ?

index

The **index** of H in G is

$$[G : H] := \begin{cases} |G/H| & G/H \text{ is finite} \\ +\infty & G/H \text{ is infinite} \end{cases}$$

Theorem 3.23: Lagrange's theorem

If $H \leq G$, then

$$|G| = [G : H] \cdot |H|$$

Remark:

Why are we use left cosets here for index? Why not right cosets? Anything holds for left cosets should also be expected hold for right cosets with the order of product reversed. Lagrange's theorem didn't mention the order of product. Thus we should expect it holds for right cosets as well. Thus when G is finite, Lagrange's theorem should imply the number of left cosets is equal to the number of right cosets.

Proposition 3.24

The function $\phi : G/H \rightarrow H \setminus G : S \mapsto S^{-1}$ is a bijection.

Proof:

First we check ϕ is well defined: if we are given left coset S , then S^{-1} is a right coset.

Suppose $S \in G/H$, so $S = gH$ for some $g \in G$. Then

$$\begin{aligned} S^{-1} &= \{(gh)^{-1} : h \in H\} \\ &= \{h^{-1}g^{-1} : h \in H\} \\ &\stackrel{*}{=} \{hg^{-1} : h \in H\} \\ &= Hg^{-1} \end{aligned}$$

*: because $H \rightarrow H : h \mapsto h^{-1}$ is a bijection.

So ϕ is well-defined, and same argument shows $\psi : H \setminus G \rightarrow G/H : S \mapsto S^{-1}$ is well-defined.

Finally, ψ is an inverse to ϕ . □

Thus can use either left or right cosets to define index:

Corollary 3.25

If $H \leq G$ then

$$[G : H] = \begin{cases} |H \setminus G| & H \setminus G \text{ is finite} \\ +\infty & H \setminus G \text{ is infinite} \end{cases}$$

Theorem 3.26: Lagrange's theorem (detailed)

If $H \leq G$, then $|G| = [G : H] \cdot |H|$. (In particular, $|H|$ divides $|G|$.) Furthermore, if G is finite, then $[G : H] = \frac{|G|}{|H|}$.

Remark:

We don't want to use the second formula if $|G|$ and $|H|$ both are infinite. See proof in the next section.

Example:

$G = D_{2n}$, $H = \langle s \rangle$, $|D_{2n}| = 2n$, $|H| = n$, so $[G : H] = 2$.

$G = D_{2n}$, $H = \langle r \rangle$, $|D_{2n}| = 2n$, $|H| = 2$, so $[G : H] = n$.

$G = \mathbb{Z}$, $H = m\mathbb{Z}$. $|G| = |H| = +\infty$, $[G : H] = |\mathbb{Z}/m\mathbb{Z}| = m$. So $|G| = [G : H] \cdot |H|$, but we don't get any info about $[G : H]$ from Lagrange's theorem. However, it still gives us some info in many cases.

Corollary 3.27

If $x \in G$, then $|x|$ divides $|G|$.

Proof:

$|x| = |\langle x \rangle|$ and $|\langle x \rangle|$ divides $|G|$. □

Proposition 3.28

If $|G|$ is prime, then G is cyclic.

Proof:

Here we don't treat $+\infty$ as a prime number, and 1 is not a prime number.

Let $x \in G$, $x \neq e$. Then $|x| \neq 1$, and $|x| \mid |G|$, so $|x| = |G|$. Since $|\langle x \rangle| = |x| = |G|$, $G = \langle x \rangle$. □

Order	Known groups
1	Trivial group
2	$\mathbb{Z}/2\mathbb{Z}$
3	$\mathbb{Z}/3\mathbb{Z}$
4	$\mathbb{Z}/4\mathbb{Z}$, ??
5	$\mathbb{Z}/5\mathbb{Z}$
6	$\mathbb{Z}/6\mathbb{Z}$, $D_6 = S_3$, ??
7	$\mathbb{Z}/7\mathbb{Z}$
8	$\mathbb{Z}/8\mathbb{Z}$, D_8 , ??
9	$\mathbb{Z}/9\mathbb{Z}$, ??

Table 3.1: Groups of small order

?? = could be more groups.

Corollary 3.29

If $\phi : G \rightarrow K$ is a homomorphism, then $|\text{Im } \phi| = [G : \ker \phi]$, and hence divides $|G|$.

Proof:

There is a bijection $G/\ker \phi \rightarrow \text{Im } \phi$, so $|\text{Im } \phi| = [G : \ker \phi]$. Then Lagrange's theorem implies $[G : H]$ divides $|G|$ for any $H \leq G$. \square

Note

Lagrange's theorem also implies that $|\text{Im } \phi|$ divides $|K|$.

Exercise

If G, K are groups, then $\phi : G \rightarrow K : g \mapsto e_K$ is a homomorphism (called the **trivial homomorphism**).

$\phi : G \rightarrow K$ is the trivial homomorphism if and only if $\text{Im } \phi = \{e\}$, the trivial subgroup.

Corollary 3.30

If G and K have coprime order, then the only homomorphism $\phi : G \rightarrow K$ is the trivial homomorphism.

3.6 Proof of Lagrange's theorem

How to prove this theorem?

Recall

$$\begin{aligned} D_{2n} &= \{s^i r^j : 0 \leq i < n, j \in \{0, 1\}\} \\ &= \langle s \rangle \sqcup r \langle s \rangle \quad (|s| = n) \\ &= \bigsqcup_{i=0}^{n-1} s^i \langle r \rangle \quad (|r| = 2) \end{aligned}$$

In example, cosets of H are disjoint, we can divide G into $[G : H]$ sets of size $|H|$. Does this work in general? Need to better understand cosets.

Proposition 3.31

Let $H \leq G$, and suppose $g, k \in G$. Then the following are equivalent:

- (a) $g^{-1}k \in H$
- (b) $k \in gH$
- (c) $gH = kH$
- (d) $gH \cap kH \neq \emptyset$

Example:

$hH = H$ if and only if $h \in H$. (This is from (c) and (a))

Proof:

(a) \Rightarrow (b) If $g^{-1}k = h \in H$, then $k = gh \in gH$.

(b) \Rightarrow (c) Suppose $k = gh$ for $h \in H$. If $h' \in H$, then $kh' = g(hh') \in gH$, since

$hh' \in H$. So $kU \subseteq gH$.

For the reverse inclusion, notice that $g = kh^{-1} \in kH$. If $h' \in H$, then $gh' = k(h^{-1}h') \in kH$, so $gH \subseteq kH$.

(c) \Rightarrow (d) Since $e \in H$, then $g \in gH$, so $gH \neq \emptyset$. If $gH = kH$, then $gH \cap kH = gH \neq \emptyset$.

(d) \Rightarrow (a) Suppose $x \in gH \cap kH$. Then $x = gh_1 = kh_2$ for $h_1, h_2 \in H$. Multiply on the left by g^{-1} , right by h_2^{-1} . So $g^{-1}k = h_1h_2^{-1} \in H$.

□

partition

Let X be a set. A **partition** of X is a subset \mathcal{Q} of 2^X such that

- (a) $\bigcup_{S \in \mathcal{Q}} S = X$, and
- (b) $S \cap T = \emptyset$ for all $S \neq T \in \mathcal{Q}$.

Here 2^X denotes set of subsets of X .

Exercise

If $\mathcal{Q} \subseteq 2^X$, then the following are equivalent:

- \mathcal{Q} is a partition
- $X = \bigsqcup_{S \in \mathcal{Q}} S$
- Every element of X is contained in exactly one element of \mathcal{Q} .

Corollary 3.32

If $H \leq G$, then G/H is a partition of G .

Proof:

Let $g \in G$, then $g \in gH$, so every element of G belongs to some element of G/H . Consequently, $\bigcup_{S \in G/H} S = G$.

Suppose $S \neq T \in G/H$ (so $S = gH$, $T = kH$ for some $g, k \in G$). If $S \cap T \neq \emptyset$, then $S = T$ by parts (c) and (d) of Proposition 3.31. So $S \cap T = \emptyset$. □

Lemma 3.33

If $S \subseteq G$, $g \in G$, then $S \rightarrow gS : h \mapsto gh$ is a bijection.

Proof:

Inverse is $gS \rightarrow S : h \mapsto g^{-1}h$. □

Consequence: If H is finite, and $g \in G$, then $|gH| = |H|$.

Now we can prove the Lagrange's theorem.

Proof:

If $|H| = +\infty$ then $|G| = +\infty$. Since cosets are disjoint, if $[G : H] = +\infty$ then $|G| = +\infty$.

Suppose $|H|, [G : H]$ are finite.

By Lemma 3.33, $|gH| = |H|$ for all $g \in G$.

Since G/H is a partition of G , G is a disjoint union of $[G : H]$ subsets, all of size $|H|$.

Conclude that $|G| = [G : H] \cdot |H|$. \square

3.6.1 Equivalence relations

relation \sim

Let X be a set. A **relation** \sim on X is a subset of $X \times X$.

Notation: $a \sim b$ if $(a, b) \in \sim$.

Example:

$=$ on X . $\leq, <, >, \geq$ on \mathbb{N} (or any ordered set). \subseteq on 2^X .

equivalence relation

A relation \sim on X is an **equivalence relation** if

- $x \sim x$ for all $x \in X$ (reflexivity)
- $x \sim y \implies y \sim x$ for all $x, y \in X$ (symmetry), and
- $x \sim y$ and $y \sim z$ for all $x, y, z \in X$ (transitivity).

Example:

$=$ on X . \equiv_m , congruence mod m , is an equivalence relation on \mathbb{Z} .

$\leq, <$ on \mathbb{N}, \mathbb{R} , etc. are not equivalence relations.

Isomorphism \cong is an equivalence relation on the *proper class* of groups. Note that there is no set of all sets, or set of all groups.

equivalence class

If \sim is an equivalence relation on X , the **equivalence class** of $x \in X$ is $[x] = [x]_\sim := \{y \in X : x \sim y\}$.

Proposition 3.34

Let \sim be an equivalence relation on X . If $x, y \in X$ then the following are equivalent:

- (a) $x \sim y$
- (b) $y \in [x]$
- (c) $[x] = [y]$
- (d) $[x] \cap [y] \neq \emptyset$

Proof:

- (a) \Rightarrow (b) Follows immediately from definition of equivalent classes.
- (b) \Rightarrow (c) Assume $y \in [x]$. If $z \in [y]$, then $x \sim y \sim z$, and by transitivity, $z \in [x]$. Thus $[y] \subseteq [x]$. Also $x \sim y \implies y \sim x$, which implies $[x] \subseteq [y]$.
- (c) \Rightarrow (d) Assume $[x] = [y]$, $[x] \cap [y] = [x] \supset \{x\} \neq \emptyset$.
- (d) \Rightarrow (a) If $x \in [x] \cap [y]$, then $x \sim z \sim y \implies x \sim y$. □

Corollary 3.35

If \sim is an equivalence relation on X , then $\{[x]_\sim : x \in X\}$ is a partition of X .

Proof:

Since $x \sim x$, $x \in [x]$. Therefore, every element x belongs to some equivalent class. If two equivalent class intersect, they must be equal. Thus X is a disjoint union of its equivalent classes. □

Thus equivalence relation \implies partition. It turns out we can go the opposite direction:

Lemma 3.36

If \mathcal{Q} is a partition of X , then there is an equivalence relation \sim on X such that $\{[x]_\sim : x \in X\} = \mathcal{Q}$.

Proof:

Every element $x \in X$ is contained in a unique set $S_x \in \mathcal{Q}$. Define \sim by saying $x \sim y$ if and only if $S_x = S_y$. This defines an equivalence relation. □

Proposition 3.37

If $H \leq G$, define a relation \sim_H on G by $g \sim_H k$ if $g^{-1}k \in H$. Then \sim_H is an equivalence relation, and the equivalence class of $g \in G$ is $[g] = gH$.

Remark:

From the proposition, we would say $h \sim e$ if and only if $h \in H$.

Proposition 3.37 follows from that cosets partition G . Proposition 3.31 is a special

case of Proposition 3.34. Thus we can prove that \sim_H is equivalence class directly, and use Proposition 3.37 to prove Proposition 3.31.

3.7 Normal subgroups

Recall Proposition 3.31, by symmetry:

Proposition 3.38

Let $H \leq G$, and suppose $g, k \in G$. Then the following are equivalent:

- (a) $kg^{-1} \in H$
- (b) $k \in Hg$
- (c) $Hg = Hk$
- (d) $Hg \cap Hk \neq \emptyset$

Caution: $g^{-1}k \in H$ does not necessarily imply $kg^{-1} \in H$.

Lemma 3.39

If $H \leq G$ and $Hg = hH$ for $g, h \in G$, then $gH = Hg$.

Proof:

$g \in Hg = hH$, so $gH = hH$. □

normal subgroup

A subgroup $N \leq G$ is a **normal subgroup** if $gN = Ng$ for all $g \in G$.

Notation: $N \trianglelefteq G$.

conjugate of h by g

If $g, h \in G$, the **conjugate of h by g** is ghg^{-1} .

Conjugates come up in linear algebra in change of basis and diagonalization.

Recall: $gS = \{gh : h \in S\}$, $Sg = \{hg : h \in S\}$. So $gSg^{-1} = \{ghg^{-1} : h \in S\}$.

As previously mentioned, $g(hS) = (gh)S$, $(Sg)h = S(gh)$, $g(Sh) = (gS)h$, $eS = S = Se$.

So $gN = Ng$ if and only if $gNg^{-1} = N$. Here we

Also: $S \subseteq T$ if and only if $gS \subseteq gT$ if and only if $Sg \subseteq Tg$.

Proposition 3.40

Let $N \leq G$. Then the following are equivalent:

- (1) $N \trianglelefteq G$ ($gN = Ng \forall g \in G$)
- (2) $gNg^{-1} = N$ for all $g \in G$
- (3) $gNg^{-1} \subseteq N$ for all $g \in G$
- (4) $G/N = N \setminus G$
- (5) $G/N \subseteq N \setminus G$
- (6) $N \setminus G \subseteq G/N$

Proof:

We've already done $(1) \iff (2)$. Clearly $(2) \implies (3)$.

To see $(3) \implies (2)$, suppose $gNg^{-1} \subseteq N$ for all $g \in G$. Given $g \in G$, we know $g^{-1}Ng \subseteq N$ by apply assumption to g^{-1} . Thus $N \subseteq gNg^{-1}$. Hence $N = gNg^{-1}$, so (2) holds.

By definition, $(1) \implies (4) \implies (5), (6)$.

$(5) \implies (1)$: Suppose $G/N \subseteq N \setminus G$. If $g \in G$, then $gN = Nh$ for some $h \in G$. By Lemma 3.39, $gN = Ng$.

$(6) \implies (1)$: Similar. □

Example:

$\langle s \rangle \leq D_{2n}$: Already seen $G/\langle s \rangle = \langle s \rangle \setminus G$. So $\langle s \rangle \trianglelefteq D_{2n}$. Can also check $s^i \langle s \rangle s^{-i} = \langle s \rangle$, $r \langle s \rangle r^{-1} = \langle s \rangle$ (since $rs^i r^{-1} = s^{-i}$).

$\langle r \rangle \leq D_{2n}$: $G/\langle r \rangle \neq \langle r \rangle \setminus G$, so $\langle r \rangle$ is not normal. Indeed, $sr s^{-1} = s^2 r \notin \langle r \rangle$ for $n \geq 3$.

If G is abelian, then all subgroups are normal.

If $\phi : G \rightarrow K$ is a homomorphism, then $\ker \phi$ is normal. Previously, we have proved $G/\ker \phi \equiv$ solution sets to equations $\phi(x) = b, b \in \text{Im } \phi = \ker \phi \setminus G$. Alternatively, we can use statement (2): if $x \in \ker \phi, g \in G$, then $\phi(gxg^{-1}) = \phi(g)\phi(x)\phi(g)^{-1} = \phi(g)\phi(g)^{-1} = e$, so $gxg^{-1} \in \ker \phi \implies g(\ker \phi)g^{-1} \subseteq \ker \phi$.

The subgroup relation \leq is transitive: If $H \leq G$ (G considered as group) and $K \leq H$ (H considered as group) then $K \leq G$. Normally we just say $K \leq H \leq G \implies K \leq G$.

The normal subgroup relation \trianglelefteq is **not** transitive: Consider $H = \langle r, s^2 \rangle \leq D_8$. $rs^2 = s^{4-2}r = s^2r \implies rs^2r^{-1} = s^2$. We know $H \trianglelefteq D_8$, and H is abelian. Since H is abelian, then $\langle r \rangle \trianglelefteq H$. However, $\langle r \rangle \not\trianglelefteq D_8$.

3.8 Normalizers and the center

normalizer of S in G

Let $S \subseteq G$. Then $N_G(S) := \{g \in G : gSg^{-1} = S\}$ is called the **normalizer of S in G** .

Lemma 3.41

$N_G(S) \leq G$.

Proof:

$eSe = S$, so $e \in N_G(S)$.

If $g, h \in N_G(S)$, then $ghS(gh)^{-1} = g(hSh^{-1})g^{-1} = gSg^{-1} = S$, so $gh \in N_G(S)$.

If $g \in N_G(S)$, then $g^{-1}Sg = g^{-1}(gSg^{-1})g = eSe = S$. So $g^{-1} \in N_G(S)$. \square

Lemma 3.42

Suppose $H \leq G$. Then $H \trianglelefteq G$ if and only if $N_G(H) = G$.

Corollary 3.43

If $G = \langle S \rangle$, and $H \leq G$, then $H \trianglelefteq G$ if and only if $gHg^{-1} = H$ for all $g \in S$.

Proof:

$H \trianglelefteq G$ if and only if $N_G(H) = G$ if and only if $S \subseteq N_G(H)$. \square

Remark:

It will be helpful to check a subgroup is normal. Warning: it is possible to have $gHg^{-1} \subseteq H$ and $g \notin N_G(H)$.

Lemma 3.44

If $|g| < +\infty$, and $gHg^{-1} \subseteq H$, then $g \in N_G(H)$.

Proof:

Prove by induction. If $gHg^{-1} \subseteq H$, then $g^iHg^{-i} \subseteq H$ for all $i \geq 0$.
(Use $g(g^{i-1}Hg^{-(i-1)})g^{-1} \subseteq gHg^{-1}$).

If $|g| = n < +\infty$, then $g^{-1}Hg = g^{n-1}Hg^{-(n-1)} \subseteq H$. We multiply g on the left and g^{-1} on the right, then $H \subseteq gHg^{-1}$, conclude $gHg^{-1} = H$. \square

Corollary 3.45

Suppose $G = \langle S \rangle$ is finite, and $H \leq G$. If $gHg^{-1} \subseteq H$ for all $g \in S$, then $H \trianglelefteq G$.

Remark:

If G is a finite group, this process makes checking whether the group is normal even faster.

center of G

If G is a group, the **center of G** is $Z(G) = \{g \in G : gh = hg \text{ for all } h \in G\}$.

Example:

$$Z(\mathrm{GL}_n \mathbb{C}) = \{\lambda 1 : \lambda \neq 0\}$$

Proposition 3.46

$$Z(G) \trianglelefteq G.$$

Proof:

Exercise. □

Products

4.1 Product groups

Proposition 4.1

Suppose (G_1, \cdot_1) , (G_2, \cdot_2) are groups. Then $G_1 \times G_2$ is a group under operation

$$(g_1, g_2) \cdot (h_1, h_2) = (g_1 \cdot_1 h_1, g_2 \cdot_2 h_2)$$

for $g_i, h_i \in G_i$, $i = 1, 2$.

Proof:

Exercise. □

product of G_1 and G_2

If G_1, G_2 are groups, the group $G_1 \times G_2$ with operation from Proposition 4.1 is called the **product of G_1 and G_2** .

Example: the Klein 4-group

Obviously $|G_1 \times G_2| = |G_1| \cdot |G_2|$.

So the group $(\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$ has order 4. Called the **Klein 4-group**.

Multiplication table:

	(0,0)	(0,1)	(1,0)	(1,1)
(0,0)	(0,0)	(0,1)	(1,0)	(1,1)
(0,1)	(0,1)	(0,0)	(1,1)	(1,0)
(1,0)	(1,0)	(1,1)	(0,0)	(0,1)
(1,1)	(1,1)	(1,0)	(0,1)	(0,0)

All elements have order 2, so it's not cyclic. Identity is $(0, 0)$. In general, identity of $G_1 \times G_2$ is (e_{G_1}, e_{G_2}) .

Proposition 4.2

Suppose $G = H \times K$. Let $\tilde{H} = \{(h, e_k) : h \in H\}$, $\tilde{K} = \{(e_H, k) : k \in K\}$. Then

- (a) $\tilde{H}, \tilde{K} \leq G$.
- (b) $H \rightarrow \tilde{H} : h \mapsto (h, e)$ and $K \rightarrow \tilde{K} : k \mapsto (e, k)$ are isomorphisms.

Proof:

Exercise. □

Remark:

So we can think of H and K as subgroups of $H \times K$. $H \times K$ can have lots of other subgroups as well. Here we listed the two particularly important ones.

Let $G = H \times K$, $\tilde{H} = H \times \{e\}$, $\tilde{K} = \{e\} \times K \leq H \times K$.

Lemma 4.3

If $h \in \tilde{H}$, $k \in \tilde{K}$, then $hk = kh$.

Proof:

Exercise. □

4.2 Homomorphisms between products

Corollary 4.4

If $\phi : H \times K \rightarrow G$ is a homomorphism, then $\phi(h)\phi(k) = \phi(k)\phi(h)$ for all $h \in \tilde{H}$, $k \in \tilde{K}$.

Proof:

Immediate. □

Now consider the converse of this corollary.

Lemma 4.5

If $\alpha : H \rightarrow G$, $\beta : K \rightarrow G$ are homomorphisms, such that $\alpha(h)\beta(k) = \beta(k)\alpha(h)$ for all $h \in H$, $k \in K$, then $\gamma : H \times K \rightarrow G : (h, k) \mapsto \alpha(h)\beta(k)$ is a homomorphism.

Proof:

$$\begin{aligned} \gamma((x, y) \cdot (z, w)) &= \gamma((xz, yw)) \\ &= \alpha(xz)\beta(yw) \\ &= \alpha(x)\alpha(z)\beta(y)\beta(w) \\ &= \alpha(x)\beta(y)\alpha(z)\beta(w) \\ &= \gamma(x, y)\gamma(z, w) \end{aligned}$$

for all $x, z \in H$, $y, w \in K$. □

Notation: the homomorphism γ is called $\alpha \cdot \beta$. This is not entirely standard. You should mention this homomorphism if you use this notation.

Remark:

You might wonder why Lemma 4.5 is called the converse of corollary. In Corollary 4.4, given ϕ , we can get homomorphisms: $H \rightarrow G : h \mapsto (h, e)$ and apply ϕ to it, similar for K .

Corollary 4.6

If $\alpha : H \rightarrow H'$, $\beta : K \rightarrow K'$ are homomorphisms, then $\gamma : H \times K \rightarrow H' \times K' : (h, k) \mapsto (\alpha(h), \beta(k))$ is a homomorphism.

Proof:

Define $\tilde{\alpha} : H \rightarrow H' \times K' : h \mapsto (\alpha(h), e)$ and $\tilde{\beta} : K \rightarrow H' \times K' : k \mapsto (e, \beta(k))$. $\tilde{\alpha}, \tilde{\beta}$ are homomorphisms (exercise), and that $\tilde{\alpha}(x)\tilde{\beta}(y) = \tilde{\beta}(y)\tilde{\alpha}(x)$ for all $x \in H, y \in K$.

Then $\gamma((x, y)) = (\alpha(x), e) \cdot (e, \beta(y)) = \tilde{\alpha}(x) \cdot \tilde{\beta}(y)$ so $\gamma = \tilde{\alpha} \cdot \tilde{\beta}$. \square

Notation: the homomorphism γ is called $\alpha \times \beta$. This notation is quite standard, which is safer to use.

Corollary 4.7

If $\alpha : H \rightarrow H'$, $\beta : K \rightarrow K'$ are isomorphisms, then $\alpha \times \beta : H \times K \rightarrow H' \times K'$ is an isomorphism.

Proof:

$\alpha \times \beta$ has inverse $\alpha^{-1} \times \beta^{-1}$. \square

Proposition 4.8

$G \rightarrow G \times \{e\} : g \mapsto (g, e)$ is an isomorphism.

Proof:

Exercise. \square

Using products, can complete list of groups of order p^2 :

Proposition 4.9

Suppose p is prime, $|G| = p^2$. Then either G is cyclic, or $G \cong (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/p\mathbb{Z})$.

Proof:

Exercise. \square

Recall our table of small order:

Order	Known groups
1	Trivial group
2	$\mathbb{Z}/2\mathbb{Z}$
3	$\mathbb{Z}/3\mathbb{Z}$
4	$\mathbb{Z}/4\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$
5	$\mathbb{Z}/5\mathbb{Z}$
6	$\mathbb{Z}/6\mathbb{Z}, D_6 = S_3, ??$
7	$\mathbb{Z}/7\mathbb{Z}$
8	$\mathbb{Z}/8\mathbb{Z}, D_8, ??$
9	$\mathbb{Z}/9\mathbb{Z}, (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$

How do we know if a group is a product?

Recall Proposition 4.2. Corollary: $H \times K \rightarrow \tilde{H} \times \tilde{K} : (h, k) \rightarrow ((h, e), (e, k))$ is an isomorphism. So we are looking for two subgroups \tilde{H}, \tilde{K} which satisfy these properties:

- if $h \in \tilde{H}, k \in \tilde{K}$, then $hk = kh$.
- every element $g \in G$ can be written as $g = \tilde{h}\tilde{k}$ for unique $\tilde{h} \in \tilde{H}, \tilde{k} \in \tilde{K}$.

4.3 Unique factorizations & internal direct products

Given $S, T \subseteq G$, let $ST = \{gh : g \in S, h \in T\}$.

Lemma 4.10

$G = ST$ if and only if every element $g \in G$ can be written as $g = hk$ for some $h \in S, k \in T$.

Example:

$$D_{2n} = \{s^i r^j\} = \langle s \rangle \cdot \langle r \rangle.$$

Suppose $G = HK$ for $H, K \leq G$. When does $g = hk$ for unique $h \in H, k \in K$? Uniqueness means that if $g = hk = h'k'$ for $h, h' \in H, k, k' \in K$, then $h = h'$ and $k = k'$.

It is easy to find necessary condition: If $e \neq g \in H \cap K$, then $g = g \cdot e = e \cdot g$, then factorization is not unique. So if factorization is unique, $H \cap K = \{e\}$. It turns out this is also a sufficient condition.

Lemma 4.11

Suppose $G = HK$ for $H, K \leq G$. Then every element $g \in G$ can be written as $g = hk$ for unique $h \in H, k \in K$ if and only if $H \cap K = \{e\}$.

Proof:

We've proved it is necessary. Suppose $H \cap K = \{e\}$. If $g = hk = h'k'$, then $(h')^{-1}h = k'k^{-1} \in H \cap K$. Thus $(h')^{-1}h = k'k^{-1} = e$. This implies $h = h', k = k'$. \square

internal direct product

We say that G is the **internal direct product** of subgroups $H, K \leq G$ if

- (a) $HK = G$,
- (b) $H \cap K = \{e\}$, and
- (c) $hk = kh$ for all $h \in H, k \in K$.

Remark:

To make the condition (b) and (c) hold, we put the word “direct” here.

Example:

$H \times K$ is the internal direct product of $\tilde{H} = H \times \{e\}$ and $\tilde{K} = \{e\} \times K$.

D_{2n} is not the internal direct product of $\langle s \rangle$ and $\langle r \rangle$ because $sr \neq rs$.

Theorem 4.12

Suppose G is the internal direct product of H and K . Then $\phi : H \times K \rightarrow G : (h, k) \mapsto hk$ is an isomorphism.

Proof:

Let $i_H : H \rightarrow G : h \mapsto h$ and $i_K : K \rightarrow G : k \mapsto k$. By part (c) of definition, $i_H(h)i_K(k) = i_K(k)i_H(h)$ for all $h \in H, k \in K$. So $\phi = i_H \cdot i_K$ is a homomorphism.

By Lemma 4.11, every element $g \in G$ can be written as $g = hk$ for unique $h \in H, k \in K$. Thus ϕ is a bijection, then ϕ is an isomorphism. \square

Lemma 4.13

If G is internal direct product of H, K , then $H, K \trianglelefteq G$.

Proof:

Suppose $g \in G$, so $g = hk$, $h \in H, k \in K$. Then

$$kHk^{-1} = \{khk^{-1} : h \in H\} = \{kk^{-1}h : h \in H\} = H,$$

so $gHg^{-1} = hkHk^{-1}h^{-1} = hHh^{-1} \subseteq H$. So $H \trianglelefteq G$. Proof for K is similar. \square

Proposition 4.14

G is the internal direct product of $H, K \leq G$ if and only if

- (a) $G = HK$, and
- (b) $H \cap K = \{e\}$.
- (c) $H, K \trianglelefteq G$.

Before proving the proposition, we introduce a definition:

commutator

The **commutator** of $g, h \in G$ is $[g, h] := g \cdot h \cdot g^{-1} \cdot h^{-1}$.

Lemma 4.15

If $g, h \in G$, then $[g, h] = e$ if and only if $gh = hg$.

Proof:

This is the proof of Proposition 4.14.

We have proved \Rightarrow .

If $h \in H, k \in K$, then $[h, k] = (hkh^{-1})k^{-1} \in K$ since $K \trianglelefteq G$. But $[h, k] = h(kh^{-1}k^{-1}) \in H$ since $H \trianglelefteq G$. So $[h, k] \in H \cap K = \{e\} \implies [h, k] = e$. Therefore, $hk = kh$ for all $h \in H, k \in K$, thus G is indeed an internal direct product. \square

Quotient groups and the isomorphism theorems

week 4

5.1 Quotient groups

Recall an example: $\mathbb{Z}/n\mathbb{Z} = \{[a] : 0 \leq a < n\}$. In this example, $\mathbb{Z}/n\mathbb{Z}$ is a group, with operation $[a] + [b] = [a + b]$. *Can we generalize this example? Can we define a group structure on G/H by $[g] \cdot [h] = [gh]$? or $gH \cdot hH = ghH$? (Here we regard gH and hH as elements in G/H instead of sets.)* Big problem: this might not be **well-defined**.

relation

A **relation** R between two sets X and Y is a subset of $X \times Y$. Notation $a R b$ if $(a, b) \in R$.

A relation R is a **function** from $X \rightarrow Y$ if

- (a) for all $x \in X$, there is $y \in Y$ such that $x R y$, and
- (b) for all $x \in X$, $y, z \in Y$, if $x R y$ and $x R z$ then $y = z$.

Can define relation \rightarrow between $G/H \times G/H$ and G/H by $([g], [h]) \rightarrow [gh]$ for all $g, h \in G$? Yes. It is properly defined, we just need to find a subset of $X \times Y$ (in this case $(G/H \times G/H) \times G/H$).

Is this relation a function? For (a), if $x = ([g], [h])$, can take $y = [gh]$. What about (b)?

Lemma 5.1

The relation \rightarrow between $G/H \times G/H$ and G/H defined by $([g], [h]) \rightarrow [gh]$ is a function if and only if H is normal. Furthermore, if H is normal, then $ghH = gH \cdot hH$, the setwise product. (Recall $S \cdot T = \{xy : x \in S, y \in T\}$)

Proof:

\Rightarrow Suppose \rightarrow is a function. Suppose $g \in G, h \in H$. Then $([g], [g^{-1}]) \rightarrow [e]$. Since $g^{-1} \cdot gh = h \in H$ from Proposition 3.37, then $g \sim_H gh$, then $[g] = [gh]$, and $([gh], [g^{-1}]) \rightarrow [ghg^{-1}]$. Since \rightarrow is a function, $[ghg^{-1}] = [e]$. But this means $ghg^{-1} \sim_H e$, i.e., $ghg^{-1} \in H$. Since this holds for all $g \in G, h \in H$. Hence $H \trianglelefteq G$.

\Leftarrow First let's prove H normal $\implies ghH = gH \cdot hH$.

Note that $gH \cdot hH = gh(h^{-1}Hh) \cdot H$. If H is normal, then $h^{-1}Hh \subseteq H$, then $(h^{-1}Hh) \cdot H \subseteq H$. Since $e \in H^{-1}Hh$, $(h^{-1}Hh) \cdot H = H$ if we take e on the left and every element of H on the right. Thus if H is normal, then $gH \cdot hH = ghH$.

Suppose that $(S, T) \rightarrow R$ and $(S, T) \rightarrow R'$ for $S, T, R, R' \in G/H$. Then $R = S \cdot T = R'$ by the definition of equivalent class. So \rightarrow is a function. \square

G/N is called the **quotient of G by N** , or a **quotient group**.

Elements of G/N can be written as gN or $[g]$ or \bar{g} .

Group operation can be stated as $gN \cdot hN = ghN$ or $[g] \cdot [h] = [gh]$ or $\bar{g} \cdot \bar{h} = \overline{gh}$

q (defined in the following theorem) is called the **quotient map** or **quotient homomorphism**.

Theorem 5.2

Let $N \trianglelefteq G$. Then the setwise product $gN \cdot hN = ghN$ makes G/N into a group. Further more, the function $q : G \rightarrow G/N : g \mapsto gN$ is a surjective homomorphism with $\ker q = N$.

Proof:

$([g] \cdot [h]) \cdot [k] = [gh] \cdot [k] = [ghk] = [g] \cdot ([h] \cdot [k])$ for all $[g], [h], [k] \in G/N$, so \cdot is associative.

$[e] \cdot [g] = [e \cdot g] = [g] = [g \cdot e] = [g] \cdot [e]$ for all $[g] \in G/N$, so $[e] = N$ is an identity.

$[g] \cdot [g^{-1}] = [gg^{-1}] = [e] = [g^{-1}g] = [g^{-1}] \cdot [g]$ for all $[g] \in G/N$, so every element of G/N has an inverse.

q clearly surjective, and $q(gh) = [gh] = [g] \cdot [h] = q(g) \cdot q(h)$. $q(g) = [g] = [e]$ if and only if $g \in N$, so $\ker q = N$. \square

We previously proved that if $\phi : G \rightarrow K$ is a homomorphism then $\ker \phi \trianglelefteq G$.

Corollary 5.3

Let $N \trianglelefteq G$. Then there is a group K and homomorphism $\phi : G \rightarrow K$ such that $N = \ker \phi$.

Proof:

Take $K = G/N$, and $q : G \rightarrow G/N$ the quotient homomorphism. Then $\ker q = N$. \square

Example:

$\mathbb{Z}/n\mathbb{Z}$: can now define this using theorem, no need to rely on pre-existing definition.

$D_{2n}/\langle s \rangle$: Cosets are $\langle s \rangle = \{s^i : 0 \leq i < n\}$ and $\langle s \rangle r = \{s^i r : 0 \leq i < n\}$

Multiplication table:

	$\langle s \rangle$	$\langle s \rangle r$
$\langle s \rangle$	$\langle s \rangle$	$\langle s \rangle r$
$\langle s \rangle r$	$\langle s \rangle r$	$\langle s \rangle$

So $D_{2n}/\langle s \rangle \cong \mathbb{Z}/2\mathbb{Z}$.

If N not normal: $\langle r \rangle$ has left cosets $s^i \langle r \rangle = \{s^i, s^i r\}, 0 \leq i < n$. If we take two left cosets and do setwise product:

$$\langle r \rangle \cdot s \langle r \rangle = \{s, sr, s^{-1}r, s^{-1}\}$$

is not a left coset of $\langle r \rangle$. Also $e \sim_{\langle r \rangle} r$, $e \cdot s = s$ is in a different coset from $r \cdot s = s^{-1}r$ so $[g] \cdot [h] = [gh]$ is not a well-defined operation.

See $D_{2n}/Z(D_{2n})$ on homework.

Example: projective general linear group

$\mathrm{GL}_n(\mathbb{K})/Z(\mathrm{GL}_n \mathbb{K})$: Recall $Z(\mathrm{GL}_n \mathbb{K}) = \{\lambda I : \lambda \neq 0\}$.

If M is invertible, $[M] = \{\lambda M : \lambda \neq 0\}$.

$$[M] \cdot [N] = \{\lambda_1 \lambda_2 MN : \lambda_1, \lambda_2 \neq 0\} = [MN]$$

We can view $\mathrm{GL}_n(\mathbb{K})$ as group of invertible linear transformations of \mathbb{K}^n (acts on vectors).

$\mathrm{GL}_n(\mathbb{K})/Z(\mathrm{GL}_n \mathbb{K})$ is invertible transformations of lines through origin in \mathbb{K}^n .

$\mathrm{GL}_n(\mathbb{K})/Z(\mathrm{GL}_n \mathbb{K})$ is called the **projective general linear group**, and is denoted by $\mathrm{PGL}_n(\mathbb{K})$. It is a very important group in some areas of geometry.

In general, can look at:

- $G/Z(G)$, any group G
- $G/\ker \phi$, any homomorphism $\phi : G \rightarrow K$
- G/N , any group G and normal subgroup $N \trianglelefteq G$

How do we find the group structure on G/N ? It might be hard.

5.2 The universal property of quotients

Suppose $N \trianglelefteq G$. What are the homomorphisms $\psi : G/N \rightarrow K$?

$$\begin{array}{ccc}
 & \xrightarrow{\psi \circ q} & \\
 G & \searrow q \quad \nearrow \psi & K \\
 & G/N &
 \end{array}$$

Every such ψ gives a homomorphism $\psi \circ q : G \rightarrow K$ (this homomorphism is sometimes also called lift, pullback of ψ to G). Not every homomorphism from $G \rightarrow K$ is a lift of ψ . What homomorphisms $G \rightarrow K$ are lift of some homomorphism ψ ?

$$\begin{array}{ccc}
 & \xrightarrow{\phi} & \\
 G & \searrow q \quad \nearrow \psi? & K \\
 & G/N &
 \end{array}$$

If we start with $\phi : G \rightarrow K$, when does there exist ψ such that $\phi = \psi \circ q$? Given ϕ , when can fill in ψ so that diagram **commutes**? Here “commute” means if we start at any point in the diagram and go to any other point, it doesn’t matter what path we take to get there, we get the same function.

Theorem 5.4: Universal property of quotients

Suppose $\phi : G \rightarrow K$ is a homomorphism, and $N \trianglelefteq G$. Let $q : G \rightarrow G/N$ be the quotient homomorphism. Then there is a homomorphism $\psi : G/N \rightarrow K$ such that $\psi \circ q = \phi$ if and only if $N \subseteq \ker \phi$. Further more, if ψ exists, then it is unique.

In other words, can fill in dashed line so that diagram “commutes” if and only if $N \subseteq \ker \phi$.

$\text{Hom}(G, K)$

If G, K are groups, let $\text{Hom}(G, K)$ be the set of (homo)morphisms $G \rightarrow K$.


Corollary 5.5

For any groups G, K , and $N \trianglelefteq G$, the function $q^* : \text{Hom}(G/N, K) \rightarrow \{\phi \in \text{Hom}(G, K) : N \subseteq \ker \phi\} : \psi \mapsto \psi \circ q$ is a bijection.

Recall we previously proved:

Theorem 5.6: Universal property of products

Let $\alpha : H \rightarrow G$ and $\beta : K \rightarrow G$ be homomorphisms, and let $i_H : H \rightarrow H \times K$ and $i_K : K \rightarrow H \times K$ be the inclusions of H and K in the product of $H \times K$. Then there is a homomorphism $\phi : H \times K \rightarrow G$ such that $\phi \circ i_H = \alpha$ and $\phi \circ i_K = \beta$ if and only if $\alpha(h)\beta(k) = \beta(k)\alpha(h)$ for all $h \in H, k \in K$.



Corollary 5.7

There is a bijection between $\text{Hom}(H \times K, G)$ and

$$\{(\alpha, \beta) \in \text{Hom}(H, G) \times \text{Hom}(K, G) : \alpha(h)\beta(k) = \beta(k)\alpha(h) \text{ for all } h \in H, k \in K\}$$

Remark:

We won't formally define the term *universal property*. Often that's held off until grad level. But even if we want to define it now, we would need some category theory. Intuitively, we can think about it as a type of theorem setting up a bijection between some sets and set of homomorphisms.

We still need to prove Theorem 5.4. Before we get into the proof, let's prove the following lemma:

Lemma 5.8

If $\alpha : G \rightarrow H$ is surjective, $\psi_i : H \rightarrow K, i = 1, 2$ are such that $\psi_1 \circ \alpha = \psi_2 \circ \alpha$, then $\psi_1 = \psi_2$.

Proof:

If $h \in H$, then there is $g \in G$ with $\alpha(g) = h$. So $\psi_1(h) = \psi_2(\alpha(g)) = \psi_2(h)$. We conclude that $\psi_1 = \psi_2$. \square

With this lemma, we can dive into the proof of Theorem 5.4:

Proof:

\Rightarrow If ψ exists, and $n \in N$, then $\phi(n) = \psi(q(n)) = \psi(e) = e$ so $N \subseteq \ker \phi$.

\Leftarrow Suppose $N \subseteq \ker \phi$. Define $\psi : G/N \rightarrow K : [g] \mapsto \phi(g)$. To show ψ is well-defined, note that if $[g] = [h]$, then $g^{-1}h \in N \subseteq \ker \phi$, so $\phi(g)^{-1}\phi(h) = \phi(g^{-1}h) = e$, so $\phi(g) = \phi(h)$.

Clearly $\psi \circ q(g) = \psi([g]) = \phi(g)$ for all $g \in G$, so $\psi \circ q = \phi$.

If $[g], [h] \in G/N$, then

$$\psi([g] \cdot [h]) = \psi([gh]) = \phi(gh) = \phi(g)\phi(h) = \psi([g])\psi([h])$$

so ψ is a homomorphism.

If $\psi' : G/N \rightarrow K$ is another homomorphism with $\psi' \circ q = \phi$ then $\psi' \circ q = \psi \circ q$. Since q is surjective, by Lemma 5.8, $\psi' = \psi$. So uniqueness holds. \square

Remark:

Equivalent way to define ψ : $\phi(gN) = \phi(g)\phi(N) = \phi(g)\{e\} = \{\phi(g)\}$. So if $S \in G/N$, then $\phi(S) = \{b\}$, a singleton set. Can define $\psi(S) = b$ for $b \in K$ such that $\phi(S) = \{b\}$.

5.3 The first isomorphism theorem

Recall: If $\phi : G \rightarrow K$ is a homomorphism then $[G : \ker \phi] = |\operatorname{Im} \phi|$. We prove this by setting up a bijection $\psi : G/\ker \phi \rightarrow \operatorname{Im} \phi$ defined by $\psi(S) = b$, where $b \in K$ is such that $\phi(S) = \{b\}$. This looks like what we just did! Now we know $G/\ker \phi$ is a group, $|G/\ker \phi| = [G : \ker \phi] = |\operatorname{Im} \phi|$. Maybe this bijection is an isomorphism?

Theorem 5.9: First isomorphism theorem

Suppose that $\phi : G \rightarrow K$ is a homomorphism. Then there is an isomorphism $\psi : G/\ker \phi \rightarrow \operatorname{Im} \phi$ such that $\phi = \psi \circ q$, where $q : G \rightarrow G/\ker \phi$ is the quotient homomorphism.

Proof:

$\ker \phi \subseteq \ker \phi$, so by universal property there is a homomorphism $\psi : G/\ker \phi \rightarrow K$ with $\psi \circ q = \phi$.

For $g \in G$, $\psi([g]) = \phi(g)$, so plainly $\operatorname{Im} \psi = \operatorname{Im} \phi$. Thus we can regard ψ as surjective homomorphism $G/\ker \phi \rightarrow \operatorname{Im} \phi$.

ψ agrees with the function $G/\ker \phi \rightarrow \operatorname{Im} \phi$ defined previously, so ψ is a bijection. Therefore ψ is an isomorphism.

Alternatively, we can prove it from the scratch. If $\psi([g]) = e$, then $\phi(g) = e$, so $g \in \ker \phi$ which implies $[g] = [e]$. So ψ is injective. Thus it is isomorphism. \square

The first isomorphism theorem is the best way to determine G/N .

Example: $\operatorname{GL}_n \mathbb{K} / \operatorname{SL}_n \mathbb{K}$

Recall $\operatorname{SL}_n(\mathbb{K}) \trianglelefteq \operatorname{GL}_n(\mathbb{K})$ is defined as the kernel of homomorphism $\det : \operatorname{GL}_n \mathbb{K} \rightarrow \mathbb{K}^\times$.

The image of \det is $\operatorname{Im} \det = \mathbb{K}^\times$. By first isomorphism theorem, $\operatorname{GL}_n \mathbb{K} / \operatorname{SL}_n \mathbb{K} \cong \mathbb{K}^\times$.

Example: \mathbb{R}/\mathbb{Z}

Consider $\mathbb{Z} \trianglelefteq \mathbb{R}^+$. What is \mathbb{R}/\mathbb{Z} ?

Have homomorphism $\exp : \mathbb{R} \rightarrow \mathbb{C}^\times : x \mapsto e^{2\pi i x}$. Thus $e^{2\pi i x} = 1$ if and only if $x \in \mathbb{Z}$.

$\operatorname{Im} \exp = \{a \in \mathbb{C} : |a| = 1\} =: S^1$ (the **circle group**).

So $\mathbb{R}/\mathbb{Z} \cong S^1$

In general, to find G/N , we can find a group K and homomorphism $\phi : G \rightarrow K$ such that $\ker \phi = N$. Then we can conclude $G/N \cong \operatorname{Im} \phi$.

Sometimes we can also turn this around and use first isomorphic theorem to find $\operatorname{Im} \phi$.

5.4 The correspondence theorem

a.k.a. the fourth isomorphism theorem. We want to understand subgroups of G/N using $q : G \rightarrow G/N$. Recall: Suppose $f : X \rightarrow Y$ is a function, $S \subseteq X, T \subseteq Y$. Then

- $f(S) := \{f(x) : x \in S\}$, and
- $f^{-1}(T) := \{x \in X : f(x) \in T\}$

We previously proved:

Proposition 3.5

If $\phi : G \rightarrow H$ is a homomorphism, $K \leq G$, then $\phi(K) \leq H$. (a.k.a. pushforward, image of K)

Proposition 3.10

If $\phi : G \rightarrow H$ is a homomorphism, $K \leq H$, then $\phi^{-1}(K) \leq G$. (a.k.a. pullback of K)

If $f : X \rightarrow Y$ is a bijection, $f^{-1}(f(S)) = S$ and $f(f^{-1}(T)) = T$. Thus if $\phi : G \rightarrow H$ is an isomorphism, we get a bijection.

$$\begin{array}{ccc} & K \mapsto \phi(K) & \\ \text{Subgroups of } G & \xleftrightarrow{\hspace{1.5cm}} & \text{Subgroups of } H \\ & \phi^{-1}(K') \leftarrow K' & \end{array}$$

Furthermore:

- $K_1 \leq K_2 \iff \phi(K_1) \leq \phi(K_2)$
- $\phi(K_1 \cap K_2) = \phi(K_1) \cap \phi(K_2)$
- K is normal $\iff \phi(K)$ is normal
- $\phi(\langle S \rangle) = \langle \phi(S) \rangle$. This holds for any homomorphisms. $\phi^{-1}(\langle S \rangle) = \langle \phi^{-1}(S) \rangle$ doesn't have to hold if ϕ is not isomorphism.
- $[G : K] = [H : \phi(K)]$

Some identities for bijections don't hold for general functions:

Always holds	Don't always hold
$f(A) \subseteq f(B)$ if $A \subseteq B$	$f(A \cap B) = f(A) \cap f(B)$
$f^{-1}(A) \subseteq f^{-1}(B)$ if $A \subseteq B$	$f^{-1}(f(A)) = A$
$f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$	$f(f^{-1}(B)) = B$
$f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$	
$f(A \cup B) = f(A) \cup f(B)$	

Everything in the left column holds for any function f . Everything in the right column

holds when f is a bijection, but not for general functions f .

Order is preserved:

Lemma 5.10

If $\phi : G \rightarrow H$ is a homomorphism, then:

- (a) If $K_1 \leq K_2 \leq G$, then $f(K_1) \leq f(K_2)$
- (b) If $K_1 \leq K_2 \leq H$, then $f^{-1}(K_1) \leq f^{-1}(K_2)$

Note that we can't say $K_1 \leq K_2$ if and only if $\phi(K_1) \leq \phi(K_2)$ since $\phi^{-1}(\phi(K)) \neq K$ in general.

Also, pullback preserves intersection:

Lemma 5.11

If $\phi : G \rightarrow H$ is a homomorphism, and $K_1, K_2 \leq H$, then $\phi^{-1}(K_1 \cap K_2) = \phi^{-1}(K_1) \cap \phi^{-1}(K_2)$.

Suppose $f : X \rightarrow Y$ is a surjection, then we can move $f(f^{-1}(B)) = B$ from the right column to the left column:

Always holds	Don't always hold
$f(A) \subseteq f(B)$ if $A \subseteq B$	$f(A \cap B) = f(A) \cap f(B)$
$f^{-1}(A) \subseteq f^{-1}(B)$ if $A \subseteq B$	$f^{-1}(f(A)) = A$
$f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$	$f(f^{-1}(B)) = B$
$f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$	
$f(A \cup B) = f(A) \cup f(B)$	
$f(f^{-1}(B)) = B$	

Lemma 5.12

If $\phi : G \rightarrow H$ is a surjective homomorphism, and $K \leq H$, then $\phi(\phi^{-1}(K)) = K$.

Sub(G)

If G is a group, let $\text{Sub}(G)$ denote set of subgroups of G .

If $\phi : G \rightarrow H$ is a homomorphism, have induced functions $\phi : \text{Sub}(G) \rightarrow \text{Sub}(H)$ and $\phi^{-1} : \text{Sub}(H) \rightarrow \text{Sub}(G)$.

If ϕ is surjective, by Lemma 5.12, then ϕ is *left* inverse to ϕ^{-1} . (It might not have inverse. Sometimes we use ϕ^* .)

So $\phi^{-1} : \text{Sub}(H) \rightarrow \text{Sub}(G)$ is injective. Question: *What's the image of ϕ^{-1} in $\text{Sub}(G)$?*

Lemma 5.13

Let $\phi : G \rightarrow H$ be a homomorphism. Then

- (a) If $K \leq H$, then $\ker \phi \leq \phi^{-1}(K)$.
- (b) If $\ker \phi \leq K \leq G$, then $\phi^{-1}(\phi(K)) = K$.

Proof:

- (a) If $K \leq H$, then $\ker \phi \leq \phi^{-1}(K)$.
- (b) It's clear that $K \leq \phi^{-1}(\phi(K))$.

Suppose $y \in \phi^{-1}(\phi(K))$. Then $\phi(y) \in \phi(K)$, so $\phi(y) = \phi(k)$ for some $k \in K$. Since $\phi(k^{-1}y) = e$, $k^{-1}y \in \ker \phi \subseteq K$, thus $y \in K$. We conclude that $\phi^{-1}(\phi(K)) \subseteq K$. \square

From this lemma, we can conclude: $K = \phi^{-1}(K') \iff \ker \phi \leq K$.

When we combine Lemma 5.12 and Lemma 5.13, we get the following theorem:

Theorem 5.14: Correspondence theorem

Let $\phi : G \rightarrow H$ be a surjective homomorphism. Then there is bijection

$$\begin{array}{ccc}
 \begin{array}{l} \text{Subgroups} \\ K \text{ of } G \text{ s.t.} \\ \ker \phi \leq K \end{array} & \begin{array}{c} \xrightarrow{K \mapsto \phi(K)} \\ \xleftarrow{\phi^{-1}(K') \leftarrow K'} \end{array} & \begin{array}{l} \text{Subgroups} \\ K' \text{ of } H \end{array}
 \end{array}$$

Furthermore, if $\ker \phi \leq K, K_1, K_2 \leq G$ then

- (a) $K_1 \leq K_2 \iff \phi(K_1) \leq \phi(K_2)$
- (b) $\phi(K_1 \cap K_2) = \phi(K_1) \cap \phi(K_2)$
- (c) $K \text{ is normal} \iff \phi(K) \text{ is normal}$

Proof:

Since ϕ is surjective, $\phi(\phi^{-1}(K')) = K'$ for all $K' \leq H$. Conversely, if $\ker \phi \leq K \leq G$, then $\phi^{-1}(\phi(K)) = K$. So ϕ and ϕ^{-1} are inverses on the specified sets. So they are bijections.

- (a) follows from fact that ϕ and ϕ^{-1} are inverses and preserve \leq . For instance, if $\phi(K_1) \leq \phi(K_2)$ then $K_1 = \phi^{-1}(\phi(K_1)) \leq \phi^{-1}(\phi(K_2)) = K_2$
- (b) $\phi^{-1}(\phi(K_1) \cap \phi(K_2)) = \phi^{-1}(\phi(K_1)) \cap \phi^{-1}(\phi(K_2)) = K_1 \cap K_2$ since $\phi(\phi^{-1}(K)) = K$, $\phi(K_1 \cap K_2) = \phi(K_1) \cap \phi(K_2)$,
- (c) Exercise. \square

What about quotient groups?

If $N \trianglelefteq G$, then $q : G \rightarrow G/N$ is a surjection, so we have

Theorem 5.15: Correspondence theorem for quotient groups

Let $N \trianglelefteq G$. Then there is a bijection

$$\begin{array}{ccc} \text{Subgroups } N \leq K \leq G & \begin{array}{c} \xrightarrow{K \mapsto q(K)} \\ \xleftarrow{q^{-1}(K') \mapsto K'} \end{array} & \text{Subgroups } K' \text{ of } G/N \end{array}$$

Furthermore, if $N \leq K, K_1, K_2 \leq G$ then

- (a) $K_1 \leq K_2 \iff q(K_1) \leq q(K_2)$
- (b) $q(K_1 \cap K_2) = q(K_1) \cap q(K_2)$
- (c) $K \text{ is normal} \iff q(K) \text{ is normal}$

Recall from first isomorphism theorem: If $\phi : G \rightarrow H$ is a surjective homomorphism, then $G/\ker \phi \cong H$. So there is a bijection between $\text{Sub}(H)$ and $\text{Sub}(G/\ker \phi)$.

Exercise

Check that

$$\left. \begin{array}{l} \text{first isomorphism theorem} \\ \text{subgroup correspondence for isomorphisms} \\ \text{correspondence theorem for quotient groups} \end{array} \right\} \implies \begin{array}{l} \text{correspondence theorem} \\ \text{for surjective homomorphisms} \end{array}$$

Suppose $N \trianglelefteq G$ and $N \leq K \leq G$, we immediately see that $N \trianglelefteq K$ since $kNk^{-1} \subseteq N$ for all $k \in K \subseteq G$.

Let $q_G : G \rightarrow G/N$ be quotient map. Since $N \trianglelefteq K$, also have quotient map $q_K : K \rightarrow K/N$.

$$\begin{array}{ccc} K & \xrightarrow{i_K} & G \\ q_K \downarrow & \searrow q_G \circ i & \downarrow q_G \\ K/N & \xrightarrow{kN \mapsto kN} & G/N \end{array}$$

It's easy to see $\ker q_G \circ i = N$, and by first isomorphism theorem, we get an isomorphism $\psi : K/N \rightarrow \text{Im } q \circ i_K = q(K)$ such that $\psi \circ q_K = q_G \circ i$. In other words, if $k \in K$, then $\psi(kN) = q(k) = kN$. Let's summarize this into a proposition:

Proposition 5.16

Suppose $N \trianglelefteq G$ and $N \leq K \leq G$. Let $q : G \rightarrow G/N$ be the quotient map. Then the function $K/N \rightarrow q(K) \leq G/N : kN \mapsto kN$ is an isomorphism.

Because of this isomorphism, we use the following notation:

K/N

If $N \trianglelefteq G$ and $N \leq K \leq G$, then the subgroup $q(K)$ corresponding to K in G/N is denoted by K/N .

Example: D_{2n}

Let $G = D_{2n}$, $N = \langle s \rangle$, where s is rotation generator.

Subgroups of D_{2n} containing N correspond to subgroups of $D_{2n}/N = \mathbb{Z}_2$.

\mathbb{Z}_2 has two subgroups, \mathbb{Z}_2 and $\{e\}$.

So there are only two subgroups of D_{2n} containing N .

Example: $\mathrm{GL}_n \mathbb{K}$

$\mathrm{GL}_n \mathbb{K} / \mathrm{SL}_n \mathbb{K} \cong \mathbb{K}^\times$, so subgroups of $\mathrm{GL}_n \mathbb{K}$ containing $\mathrm{SL}_n \mathbb{K}$ correspond to subgroups of \mathbb{K}^\times (of which there can be lots: $\{1, -1\}$, $\{2^x | x \in \mathbb{Z}\}$)

5.5 The third isomorphism theorem

Suppose $N \trianglelefteq G$ and $N \leq K \leq G$. From correspondence theorem: $K \trianglelefteq G$ if and only if $K/N \trianglelefteq G/N$. Suppose $K/N \trianglelefteq G/N$. What's $(G/N)/(K/N)$?

Third isomorphism theorem, informal version

$$(G/N)/(K/N) \cong G/K.$$

Example:

Suppose $n|m$, so that $m\mathbb{Z} \leq n\mathbb{Z}$.

Then $(\mathbb{Z}/m\mathbb{Z})/(n\mathbb{Z}/m\mathbb{Z}) \cong \mathbb{Z}/n\mathbb{Z}$.

$$(\mathbb{Z}/20\mathbb{Z})/(5\mathbb{Z}/20\mathbb{Z}) \cong \mathbb{Z}/5\mathbb{Z}$$

Theorem 5.17: Third isomorphism theorem

Let $N \trianglelefteq G$ and $N \leq K \trianglelefteq G$. Let:

- q_1 be the quotient map $G \rightarrow G/N$,
- q_2 be the quotient map $G/N \rightarrow (G/N)/(K/N)$, and

- q_3 be the quotient map $G \rightarrow G/K$.

Then there is an isomorphism $\psi : G/K \rightarrow (G/N)/(K/N)$ such that $\psi \circ q_3 = q_2 \circ q_1$.

$$\begin{array}{ccc}
 G & \xrightarrow{q_1} & G/N \\
 q_3 \downarrow & & \downarrow q_2 \\
 G/K & \xrightarrow{\psi} & (G/N)/(K/N)
 \end{array}$$

Proof:

Note that $\ker q_1 \circ q_1 = (q_2 \circ q_1)^{-1}(\{e\}) = q_1^{-1}(q_2^{-1}(\{e\})) = q_1^{-1}(K/N) = K$

and q_1, q_2 surjective, $\text{Im } q_2 \circ q_1 = (G/N)/(K/N)$.

By first isomorphism theorem, there is an isomorphism $\psi : G/K \rightarrow (G/N)/(K/N)$ such that $\psi \circ q_3 = q_2 \circ q_1$. \square

What if K isn't normal? Then G/K isn't a group, and neither is $(G/N)/(K/N)$. However we can still talk about $[G : K]$ and $[G/N : K/N]$.

Proposition 5.18

If $N \trianglelefteq G$ and $N \leq K \leq G$, then $[G : K] = [G/N : K/N]$.

In fact, there's no reason to use quotient spaces. This holds for surjective homomorphisms.

Proposition 5.19

Let $\phi : G \rightarrow H$ be a surjective homomorphism, and suppose $\ker \phi \leq K \leq G$. Then $[G : K] = [H : \phi(K)]$.

These two propositions are actually equivalent by the first isomorphism theorem.

Proof:

Define a function $f : G/K \rightarrow H/\phi(K) : gK \mapsto \phi(g)\phi(K)$.

Well-defined: If $gK = hK$, then $h^{-1}g \in K \implies \phi(h)^{-1}\phi(g) = \phi(h^{-1}g) \in \phi(K)$. So $\phi(g)\phi(K) = \phi(h)\phi(K)$.

Since ϕ is surjective, f is onto.

Suppose $f(gK) = f(hK)$, so $\phi(g)\phi(K) = \phi(h)\phi(K)$. Then $\phi(h^{-1}g) = \phi(h)^{-1}\phi(g) \in \phi(K) \implies h^{-1}g \in \phi^{-1}(\phi(K)) = K$. So $gK = hK$, and f is injective.

We conclude that f is a bijection. \square

5.6 The second isomorphism theorem

Recall *internal direct product* and lemma from products:

Lemma 4.11

Suppose $G = HK$ for $H, K \leq G$. Then every element $g \in G$ can be written as $g = hk$ for unique $h \in H, k \in K$ if and only if $H \cap K = \{e\}$.

We didn't use the fact $G = HK$ in the proof. If $HK \subsetneq G$, then we won't be able to write $g = hk$, but for the g that we can write in the form of hk , the factorization is unique if and only if $H \cap K = \{e\}$. Then we can reword this lemma a little more generally.

Suppose $H, K \leq G$.

Lemma 5.20

Every element of HK can be written as hk for unique $h \in H, k \in K$, if and only if $H \cap K = \{e\}$.

If $H \cap K = \{e\}$, then $|HK| = |H| \cdot |K|$.

What is $|HK|$ if $H \cap K \neq \{e\}$?

$HK = \bigcup_{h \in H} hK$, a union of cosets of K .

Let $X = \{hK : h \in H\} \subseteq G/K$.

Then X is a partition of HK , so $|HK| = |X| \cdot |K|$.

How large is X ?

Lemma 5.21

Let $H, K \leq G$. If $h_1, h_2 \in H$, then $h_1K = h_2K$ if and only if $h_1(H \cap K) = h_2(H \cap K)$.

Proof:

$$h_1K = h_2K \iff h_1^{-1}h_2 \in K \iff h_1^{-1}h_2 \in H \cap K.$$

But $h_1^{-1}h_2 \in H \cap K$ if and only if $h_1H \cap K = h_2H \cap K$. This uses the fact that $H \cap K \leq H$. □

Rephrasing: Consider equivalence relations \sim_K on G , $\sim_{H \cap K}$ on H . If $h_1, h_2 \in H$, then $h_1 \sim_K h_2 \iff h_1 \sim_{H \cap K} h_2$.

Corollary 5.22

$H/H \cap K \rightarrow X : hH \cap K \rightarrow hK$ is a bijection.

Proof:

From Lemma 5.21, it is well-defined, injective. Surjective obvious. \square

Now going back to what we are trying to do: If $H, K \leq G$, $X = \{hK : h \in H\}$ partitions HK , so $|HK| = |X| \cdot |K|$.

$|X| = [H : H \cap K]$, so $|HK| = [H : H \cap K]|K|$. Using $[H : H \cap K] \cdot |H \cap K| = |H|$ (by Lagrange's theorem), we have

Proposition 5.23

If $H, K \leq G$, then $|HK| |H \cap K| = |H| |K|$.

Note that this proposition also holds when H and K are infinite. Another way to think about this formula if H, K finite:

$[H : H \cap K] = |X| = \frac{|HK|}{|K|}$ LHS is an index, RHS is a fraction. Is RHS an index as well? The problem: HK not necessarily a group. Thus RHS might not be an index. But when is HK a group?

Proposition 5.24

Let $H, K \leq G$. Then $HK \leq G \iff HK = KH \iff KH \subseteq HK$.

Proof:

(1) \Rightarrow (2) If $HK \leq G$, and $h \in H, k \in K$, then $h, k \in HK$, so $kh \in HK$. Also $k^{-1}h^{-1} \in HK$, so $k^{-1}h^{-1} = h_0k_0$. Hence $hk = (k^{-1}h^{-1})^{-1} = k_0^{-1}h_0^{-1} \in KH$. Sp $KH \subseteq HK$ and $HK \subseteq KH \implies HK = KH$

(3) \Rightarrow (1) Conversely, suppose $KH \subseteq HK$. We always have $e \in HK$. If $x, y \in HK$, then $x = h_0k_0, y = h_1k_1$ for $h_0, h_1 \in H, k_0, k_1 \in K$. Since $KH \subseteq HK$, $k_0^{-1}(h_0^{-1}h_1) = h_2k_2$ for $h_2 \in H, k_2 \in K$. So $x^{-1}y = k_0^{-1}h_0^{-1}h_1k_1 = h_2(k_2k_1) \in HK$. \square

Corollary 5.25

If $KH \subseteq HK$, then $[H : H \cap K] = [HK : K]$.

When is $KH \subseteq HK$?

Sufficient condition:

for all $h \in H$, there is $h' \in H$ such that $Kh = h'K$.

Recall: if $Kh = h'K$, then $h'K = hK$.

Rephrase condition: $hKh^{-1} = K$, for all $h \in H$, i.e., $H \subseteq N_G(K)$.

Corollary 5.26

If $H \subseteq N_G(K)$, then $HK \leq G$, and hence $[H : H \cap K] = [HK : K]$.

What else does the condition $H \subseteq N_G(K)$ imply?

We know $hKh^{-1} = K$, $kKk^{-1} = K$, so

$$H, K \subseteq N_{HK}(K) \implies N_{HK}(K) = HK \implies K \trianglelefteq HK.$$

If $k \in H \cap K, h \in H$, then $hkh^{-1} \in H \cap K$. So $H \cap K \trianglelefteq H$.

Theorem 5.27: Second isomorphism theorem

Suppose $H \subseteq N_G(K)$. Then $HK \leq G$, $K \trianglelefteq HK$, and $H \cap K \trianglelefteq H$. Further more, if $i_H : H \rightarrow HK$ is the inclusion, $q_1 : H \rightarrow H/H \cap K$ and $q_2 : HK \rightarrow HK/K$ are the quotient maps, then there is an isomorphism $\psi : H/H \cap K \rightarrow HK/K$ such that $\psi \circ q_1 = q_2 \circ i_H$.

$$\begin{array}{ccc} H & \xrightarrow{i_H} & HK \\ q_1 \downarrow & & \downarrow q_2 \\ H/H \cap K & \xrightarrow{\psi} & HK/K \end{array}$$

Proof:

Already shown $HK \leq G$, $K \trianglelefteq HK$, $H \cap K \trianglelefteq H$.

If $h \in H, k \in K$, then $hkhK = hK$. So $HK/K = \{gK : g \in HK\} = \{hK : h \in H\}$. Hence $\text{Im } q_2 \circ i_H = \{hK : h \in H\} = HK/K$.

$$\ker q_2 \circ i_H = i_H^{-1}(q_2^{-1}(\{e\})) = i_H^{-1}(K) = H \cap K$$

By the first isomorphism theorem, there is an isomorphism ψ as desired. \square

Example: $\text{PGL}_n \mathbb{C}$

Recall that $\text{PGL}_n \mathbb{C} = \text{GL}_n \mathbb{C} / Z(\text{GL}_n \mathbb{C})$.

Let $K = Z(\text{GL}_n \mathbb{C}) = \{\lambda 1 : \lambda \neq 0\}$. Since $K \trianglelefteq G$, $N_G(K) = G$. Let $H = \text{SL}_n \mathbb{C} = \{M \in G : \det M = 1\} \trianglelefteq G = N_G(K)$, so $HK \leq G$. Suppose $M \in \text{GL}_n \mathbb{C}$, let $\lambda = \det M$. Then $\det \lambda^{-1/n} M = \lambda^{-1} \det M = 1$, $\lambda^{-1/n} M \in \text{SL}_n \mathbb{C}$.

Conclusion: $G = HK$.

$C_n := H \cap K = \{\lambda 1 : \lambda^n = 1\} = \{e^{2\pi i k/n} 1 : k = 0, \dots, n-1\}$ (Note: $C_n \cong \mathbb{Z}/n\mathbb{Z}$). Second isomorphism $\implies \text{PGL}_n \mathbb{C} \cong \text{SL}_n \mathbb{C} / C_n$. In Lie theory, this kind of calculation can be quite useful.

Group actions

6.1 Group actions

Two group actions

Example: S_n

Permutation S_n of $\{1, \dots, n\}$ form a group.

This means that we can multiply permutations together. e.g.,

$$(12)(34)(24) = (1234)$$

However, that's not all there is to permutations: we can also plug in numbers from $1, \dots, n$

$$((12)(34))(3) = 4$$

We say that S_n **acts** on $\{1, \dots, n\}$

Example: $GL_n \mathbb{C}$

Similarly, for $GL_n \mathbb{C}$, we can do more than multiply matrices: We can also multiply matrices and vectors. Given $A \in GL_n \mathbb{C}$, $v \in \mathbb{C}^n$, can take $A \cdot v \in \mathbb{C}^n$. Say that $GL_n \mathbb{C}$ **acts** on \mathbb{C}^n .

Clearly the notion of actions is important to groups.

action

Let G be a group. A **(left) action** of G on a set X is a function $\cdot : G \times X \rightarrow X$ such that

- (a) $e \cdot x = x$ for all $x \in X$, and
- (b) $g \cdot (h \cdot x) = (gh) \cdot x$ for all $g, h \in G, x \in X$.

Example:

S_n acts on $\{1, \dots, n\}, n \geq 1$.

$\text{GL}_n \mathbb{K}$ acts on \mathbb{K}^n (prove by the associativity of matrix multiplication).

Let X be any set, G any group. We can define an action of G on X by $g \cdot x = x$ for all $g \in G, x \in X$. This is called the **trivial action** of G on X .

Proof:

(a) clear, $g \cdot (h \cdot x) = g \cdot x = x = (gh) \cdot x$. □

Proposition 6.1

Let X be a set. The group S_X (of invertible functions $X \rightarrow X$ under composition \circ) acts on X via $f \cdot x = f(x)$.

Proof:

The identity 1 in S_X is the identity function, so $1 \cdot x = 1(x) = x$. If $f, g \in S_X$, then $(f \circ g)(x) = f(g(x)) = g \cdot (g \cdot x)$. □

Note

We typically stick with the notation $f(x)$, rather than $f \cdot x$. Recall $S_n = S_{\{1, \dots, n\}}$.

Lemma 6.2

If G acts on X , and $H \leq G$, then H acts on X by the restricted action $X \times X \rightarrow X : (h, x) \mapsto h \cdot x$.

Proof:

Immediate. □

Alternative way to show $\text{GL}_n \mathbb{K}$ acts on \mathbb{K}^n : observe $\text{GL}_n \mathbb{K} \leq S_{\mathbb{K}^n}$.

However, note that groups aren't tied to a particular action:

Example:

D_{2n} was defined as subgroup of $\text{GL}_2 \mathbb{R}$, so it acts on \mathbb{R}^2 .

However, D_{2n} also acts on the vertices v_0, \dots, v_{n-1} of the n -gon.

In fact, this action determines elements of D_{2n} :

$$s^i \text{ sends } v_0 \text{ to } v_i, v_1 \text{ to } v_{i+1} \quad s^i r \text{ sends } v_0 \text{ to } v_i, v_1 \text{ to } v_{i-1}$$

This dihedral group action on the vertices of the n -gon is an instance of the following pattern:

invariant under the action of G

If G acts on X , a subset $Y \subseteq X$ is **invariant under the action of G** if $g \cdot y \in Y$ for all $g \in G, y \in Y$.

Lemma 6.3

If G acts on X and Y is an invariant subset, then G acts on Y via $G \times Y \rightarrow Y :$
 $(g, y) \mapsto g \cdot y$.

Example:

$\{0\}$ is an invariant subset of \mathbb{K}^n under the action of $\mathrm{GL}_n \mathbb{K}$. In this case, the action of $\mathrm{GL}_n \mathbb{K}$ on $\{0\}$ is trivial.

Index

A

abelian 12
action 75
associative 7

B

binary operation 6

C

center of G 53
commutative 8
commutator 59
commute 20
conjugate of h by g 50
coset 40
cyclic 27

D

dihedral group 16
disjoint 20

E

equivalence class 48
equivalence relation 48

F

finite 12

fixed points 20

G

generate 27
group 11

H

homomorphism 32

I

identity 9
image 34
in/sur/bi-jective 37
index 43
internal direct product 58
invariant under the action of G 76
inverse 9
invertible 10
isomorphic 38
isomorphism 37

K

k -ary operation 7
 k -cycle 21
kernel 36

M

multiplicative form of cyclic groups .. 40

multiplicative table..... 14

N

n-gon 15

normal subgroup 50

normalizer of S in G 52

O

order 12, 14

P

partition 47

product of G_1 and G_2 54

proper subgroup 23

R

relation 60

relation \sim 48

S

set of left/right cosets 42

subgroup 22

subgroup generated by S in G 25

support set 20

symmetric/permutation group 18

symmetry 15

T

trivial subgroup 23