



Introduction to Optimization

CO 255



Ricardo Fukasawa

Preface

Disclaimer Much of the information on this set of notes is transcribed directly/indirectly from the lectures of CO 255 during Winter 2020 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

For any questions, send me an email via <https://notes.sibeliusp.com/contact/>.

You can find my notes for other courses on <https://notes.sibeliusp.com/>.

Sibeliusp Peng

Contents

Preface	1
0 Info	3
1 Introduction	4
2 Linear Optimization	6
2.1 Determining Feasibility	7
2.2 Fourier-Motzkin Elimination	8
2.3 Certifying Optimality	12
2.4 Possible Outcomes	16
2.5 Duals of generic LPs	16
2.5.1 Cheat Sheet	17
2.6 Other interpretations of dual	19
2.7 Complementary Slackness	21
2.7.1 Geometric Interpretation of C.S.	23
2.8 Geometry of Polyhedra	24
2.9 Simplex Algorithm	29
2.9.1 CANONICAL FORM	31
2.9.2 Iteration of simplex	34
2.9.3 Mechanics of Simplex	35

Info

Ricardo: MC 5036. OH: M 1:30 - 3pm
TA: Adam Brown: MC 5462. OH: F 10-11am

Books (not required)

- Intro to Linear Opt. Bertsimas
- Int Programming. Conforti

Grading

- assns: 20% (≈ 5)
- mid: 30% (Feb 11 in class)
- final: 50%

Introduction

Given a set S , and a function $f : S \rightarrow \mathbb{R}$. An optimization problem is:

$$\begin{array}{ll} \max & f(x) \\ \underbrace{\text{s.t.}}_{\text{subject to}} & x \in S \end{array} \quad (\text{OPT})$$

- S **feasible region**
- A point $\bar{x} \in S$ is a **feasible solution**
- $f(x)$ is **objective function**

(OPT) means: “Find a feasible solution x^* such that $f(x) \leq f(x^*), \forall x \in S$ ”

- Such x^* is an **optimal solution**
- $f(x^*)$ is **optimal value**

Other ways to write (OPT):

$$\begin{aligned} \max \{ & f(x), x \in S \} \\ \max_{x \in S} & f(x) \end{aligned}$$

Analogous problem

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & x \in S \end{array}$$

Note

$$\begin{array}{ll} \max & f(x) \\ \text{s.t.} & x \in S \end{array} = -1 \left(\begin{array}{ll} \min & -f(x) \\ \text{s.t.} & x \in S \end{array} \right)$$

Problem x^* may not exist

a) Problem is unbounded:

$$\forall M \in \mathbb{R}, \exists \bar{x} \in S, \text{ s.t. } f(\bar{x}) > M$$

b) $S = \emptyset$, i.e. (OPT) is **INFEASIBLE**

c) There may not exist x^* achieving supremum.

Example:

$$\begin{array}{ll} \max & x \\ \text{s.t.} & x < 1 \end{array}$$

supremum

$$\sup\{f(x) : x \in S\} = \begin{cases} +\infty & \text{if OPT unbounded} \\ -\infty & \text{if } S = \emptyset \\ \min\{x : x \geq f(x), \forall x \in S\} & \text{otherwise} \end{cases}$$

always exist and are well-defined

infimum

$$\inf\{f(x) : x \in S\} = -1 \cdot \sup\{-f(x) : x \in S\}$$

From this point on, we will abuse notation and say $\max\{f(x) : x \in S\}$ is $\sup\{f(x) : x \in S\}$.

One way to specify that I want an opt. sol. (if exists) is

$$x^* \in \operatorname{argmax}\{f(x) : x \in S\}$$

2

Linear Optimization (Programming) (LP)

$$S = \{x \in \mathbb{R}^n : Ax \leq b\}$$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $f(x) = c^T x$, $c \in \mathbb{R}^n$.

↓

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \end{array} \quad (\text{LP})$$

Note

$$A = \begin{pmatrix} | & & | \\ A_1 & \cdots & A_n \\ | & & | \end{pmatrix} \quad A = \begin{pmatrix} - & a_1^T & - \\ & \vdots & \\ - & a_m^T & - \end{pmatrix}$$

Clarifying

$$u, v \in \mathbb{R}^n, \quad u \leq v \iff u_j \leq v_j, \forall j \in 1, \dots, n$$

Note

$u \not\leq v$ is not the same as $u > v$

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \not\leq \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Example:

$$\begin{array}{ll} \max & 2x_1 + 0.5x_2 \\ \text{s.t.} & x_1 \leq 2 \\ & x_1 + x_2 \leq 2 \\ & x \geq 0 \end{array}$$

- Strict ineq. not allowed

halfspace, hyperplane, polyhedron

Let $h \in \mathbb{R}^n, h_0 \in \mathbb{R}$.

$\{x \in \mathbb{R}^n : h^T x \leq h_0\}$ is a **halfspace**.

$\{x \in \mathbb{R}^n : h^T x = h_0\}$ is a **hyperplane**.

$Ax \leq b$ is a **polyhedron** (i.e. intersection of finitely many halfspaces).

Example:

n products, m resources. Producing $j \in \{1, \dots, n\}$ given c_j profit/unit and consumes a_{ij} units of resource i , $\forall i \in \{1, \dots, m\}$. There are b_i units available $\forall i \in \{1, \dots, m\}$.

$$\begin{aligned} \max \quad & \sum_{j=1}^n c_j x_j \\ \text{s.t.} \quad & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad \forall i = 1, \dots, m \\ & x \geq 0 \end{aligned}$$

which is an LP.

2.1 Determining Feasibility

Given a polyhedron

$$P = \{x \in \mathbb{R}^n : Ax \leq b\}$$


either find $\bar{x} \in P$ or show $P = \emptyset$.

Idea In 1-d, easy. \rightarrow Reduce problem in dimension n to one in dimension $n - 1$.

Notation Let $S = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^p : Ax + Gy \leq b\}$, then

$$\text{proj}_x S := \{x \in \mathbb{R}^n : \exists y \text{ so that } (x, y) \in S\}$$

is the (orthogonal) *projection* of S onto x .



We will find if $P = \emptyset$ by looking at $\text{proj}_{x_1, \dots, x_{n-1}}(P)$

2.2 Fourier-Motzkin Elimination

Call a_{ij} entries of A . Let

$$\begin{aligned} M &:= \{1, 2, \dots, m\} \\ M^+ &:= \{i \in M : a_{in} > 0\} \\ M^- &:= \{i \in M : a_{in} < 0\} \\ M^0 &:= \{i \in M : a_{in} = 0\} \end{aligned}$$

For $i \in M^+$:

$$a_i^T x \leq b_i \iff \sum_{j=1}^n a_{ij} x_j \leq b_i \iff \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} x_j + x_n \leq \frac{b_i}{a_{in}}, \quad \forall i \in M^+ \quad (1)$$

For $i \in M^-$

$$a_i^T x \leq b_i \iff \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} x_j - x_n \leq \frac{b_i}{-a_{in}}, \quad \forall i \in M^- \quad (2)$$

For $i \in M^0$

$$a_i^T x \leq b_i \iff \sum_{j=1}^{n-1} a_{ij} x_j \leq b_i, \quad \forall i \in M^0 \quad (3)$$

$$P = \{x \in \mathbb{R}^n : (1)(2)(3)\}$$

Define

$$\sum_{j=1}^{n-1} \left(\frac{a_{ij}}{a_{in}} - \frac{a_{kj}}{a_{kn}} \right) x_j \leq \frac{b_i}{a_{in}} - \frac{b_k}{a_{kn}}, \quad \forall i \in M^+, \forall k \in M^- \quad (4)$$

Theorem 2.1

$$(\bar{x}_1, \dots, \bar{x}_{n-1}) \text{ satisfies (3), (4)} \iff \exists \bar{x}_n : (\bar{x}_1, \dots, \bar{x}_n) \in P$$

Proof:

\Leftarrow If $(\bar{x}_1, \dots, \bar{x}_n)$ satisfies (1), (2), (3) then $(\bar{x}_1, \dots, \bar{x}_{n-1})$ satisfies (3) and adding (1), (2) \Rightarrow $(\bar{x}_1, \dots, \bar{x}_{n-1})$ satisfies (4)

\Rightarrow If $(\bar{x}_1, \dots, \bar{x}_{n-1})$ satisfies (4)

$$\sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \bar{x}_j - \frac{b_i}{a_{in}} \leq \sum_{j=1}^{n-1} \frac{a_{kj}}{a_{kn}} \bar{x}_j - \frac{b_k}{a_{kn}}, \quad \forall i \in M^+, k \in M^-$$

Let

$$\bar{x}_n := \max_{i \in M^+} \left\{ \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \bar{x}_j - \frac{b_i}{a_{in}} \right\}$$

$$\implies \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \bar{x}_j - \frac{b_i}{a_{in}} \leq -\bar{x}_n, \quad \forall i \in M^+$$

and

$$-\bar{x}_n \leq \sum_{j=1}^{n-1} \frac{a_{kj}}{a_{kn}} \bar{x}_j - \frac{b_k}{a_{kn}}, \quad \forall k \in M^-$$

$$\implies (\bar{x}_1, \dots, \bar{x}_n) \in P$$

□

Note

Proof assumes M^+, M^- are nonempty. But statement holds regardless.

(if M^+ or $M^- = \emptyset$ then (4) yields no constraints)

Algorithm 1: Fourier-Motzkin

- 1 $A^n = A, b^n = b$
- 2 given A^i, b^i obtain A^{i-1}, b^{i-1} (A^{i-1} has one less column than A^i column than A^i) by applying the steps described

$$P_i := \{x \in \mathbb{R}^i : A^i x \leq b^i\}$$

then

$$P_{i-1} = \text{proj}_{x_1, \dots, x_{i-1}} P_i$$

- 3 Keep applying projection until $i = 1$.

$$P_0 = \emptyset \iff P_n = P = \emptyset$$

Let

$$P_i^n = P_i \times \mathbb{R}^{n-i} = \{x \in \mathbb{R}^n (A^i, 0)x \leq b^i\}$$

not hard to see $P_i^n = \emptyset \iff P_i = \emptyset$

Notice that

$$P_0 = \emptyset \iff P_0^n = \emptyset, P_0^n = \{0 \leq b^0\}$$

Example:

$$P_2 = \left\{ x \in \mathbb{R}^2 : \begin{array}{rrcl} x_1 & +x_2 & \leq & 1 \\ -x_1 & & \leq & 0 \\ & -x_2 & \leq & -2 \\ -3x_1 & -3x_2 & \leq & -6 \end{array} \right\}$$

draw the graph, clearly empty

$$M^+: \frac{1}{2}x_1 + x_2 \leq \frac{1}{2}$$

$$M^-: -x_2 \leq -2 \quad -x_1 - x_2 \leq -2$$

$$M^0: -x_1 \leq 0$$

$$P_1 = \left\{ x_1 \in \mathbb{R} : \begin{array}{ll} -x_1 & \leq 0 \\ \frac{1}{2}x_1 & \leq -\frac{3}{2} \\ -\frac{1}{2}x_1 & \leq -\frac{3}{2} \end{array} \right\}$$

$$M^+: x_1 \leq -3$$

$$M^-: -x_1 \leq 0 \text{ and } -x_1 \leq -3$$

$$P_0^2 = \left\{ x \in \mathbb{R}^2 : \begin{array}{l} 0 \leq -3 \\ 0 \leq -6 \end{array} \right\} = \emptyset$$

$$\text{Here } b^0 = \begin{pmatrix} -3 \\ -6 \end{pmatrix}$$

Remark:

Inequality in P_i^n :

- All inequalities are obtained by a nonnegative combination of inequality in P_{i+1}^n
 \implies all nonnegative combination of inequalities in P .
- If all A, b are rational then so are all A^i, b^i
- If $b = 0, b_i = 0, \forall i$

Theorem 2.2: Farkas' Lemma

$$P = \{x \in \mathbb{R}^n : Ax \leq b\} = \emptyset \iff \begin{array}{l} u^T A = 0 \\ \exists u \in \mathbb{R}^m : u^T b < 0 \\ u \geq 0 \end{array}$$

Proof:

(\Leftarrow) Suppose \bar{x} satisfies $A\bar{x} \leq b$.

$$0 = u^T A\bar{x} \leq u^T b < 0$$

which is impossible.

(\Rightarrow) If $P = \emptyset$. Apply Fourier-Motzkin until we get

$$P_0^n = \emptyset = \{x \in \mathbb{R}^n : 0x \leq b^0\}$$

i.e. there exists j for which $b_j^0 < 0$.

If we look at corresponding constraint in P_0^n is

$$0^T x \leq b_j^0$$

which can be obtained by a vector u such that $u^T A = 0, u^T b = b_j^0, u \geq 0$.

□

Farkas' Lemma (alternate statement)

Exactly one of the following has a solution:

a) $Ax \leq b$

$$u^T A = 0$$

b) $u^T b < 0$

$$u \geq 0$$

Farkas' Lemma (Different Form)

Exactly one of the following has a solution:

a) $Ax = b$
 $x \geq 0$

b) $u^T A \geq 0$
 $u^T b < 0$

Proof:

(Sketch)

$$P = \left\{ x : \begin{array}{l} Ax = b \\ x \geq 0 \end{array} \right\} = \left\{ x : \underbrace{\begin{pmatrix} A \\ -A \\ -I \end{pmatrix}}_{A'} x \leq \underbrace{\begin{pmatrix} b \\ -b \\ -0 \end{pmatrix}}_{b'} \right\}$$

Apply original Farkas' Lemma to get $P = \emptyset \iff \exists u_1 \in \mathbb{R}^m, u_2 \in \mathbb{R}^m, v \in \mathbb{R}^n$:

$$u_1^T A - u_2^T A - v = 0$$

$$u_1^T b - u_2^T b < 0$$

$$u_1, u_2, v \geq 0$$

Let $u = (u_1 - u_2)$

$$u^T A - v = 0 \implies u^T A \geq 0, \quad u^T b < 0$$

□

Consider a linear programming (LP):

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \end{array} \quad (\text{LP})$$

Theorem 2.3: Fundamental Theorem of Linear Programming

(LP) has exactly one of 3 outcomes:

- a) Infeasible
- b) Unbounded
- c) There exists an optimal solution.

Proof:

Let's assume a), b) don't hold.

If $n = 1$, then (LP) has an optimal solution. (Why?)

Else, define

$$\begin{array}{ll} \max & z \\ \text{s.t.} & z - c^T x \leq 0 \\ & Ax \leq b \end{array} \quad (\text{LP}')$$

(LP') is also not in case a) or b). (Why?)

Also if (x^*, z^*) is an optimal solution to (LP'), then x^* is an optimal solution to (LP). (Why?)

Apply Fourier-Motzkin to

$$\left\{ (x, z) : \begin{array}{l} z - c^T x \leq 0 \\ Ax \leq b \end{array} \right\}$$

Until we are left with a polyhedron

$$\{z \in \mathbb{R} : A'z \leq b'\}$$

Now $\max_{\text{s.t. } A'z \leq b'} z$ is not cases a) or b). (Why?)

→ can get an optimal solution z^* to such problem. Apply Fourier-Motzkin back to get (x^*, z^*) optimal solution to (LP'). (Why?) \square

2.3 Certifying Optimality

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \end{array} \quad (\text{LP})$$

and let $\bar{x} \in P = \{x : Ax \leq b\}$

Question Can we certify that \bar{x} is optimal?

Example:

$$\begin{aligned} \max \quad & 2x_1 + x_2 \\ & x_1 + 2x_2 \leq 2 \\ \text{s.t.} \quad & x_1 + x_2 \leq 2 \\ & x_1 - x_2 \leq 0.5 \end{aligned}$$

Consider $\bar{x} = (0, 1)^T$ is clearly NOT optimal.

$x^* = (1, 0.5)^T$ and $c^T x^* = 2.5$. Any feasible solution satisfies

$$\begin{array}{rcl} x_1 + 2x_2 & \leq 2 & \times 1/3 \\ x_1 + x_2 & \leq 2 & \times 1 \\ + \quad x_1 - x_2 & \leq 0.5 & \times 2/3 \\ \hline 2x_1 + x_2 & \leq 3 & \end{array}$$

Instead do $1 \times 1st$ constraint + $1 \times 3rd$ constraint $\implies 2x_1 + x_2 \leq 2.5$

In general:

$$\begin{array}{rcl} x_1 + 2x_2 & \leq 2 & \times y_1 \\ x_1 + x_2 & \leq 2 & \times y_2 \\ + \quad x_1 - x_2 & \leq 0.5 & \times y_3 \\ \hline (y_1 + y_2 + y_3)x_1 + (2y_1 + y_2 - y_3)x_2 & \leq 2y_1 + 2y_2 + 0.5y_3 \end{array}$$

As long as $y_1, y_2, y_3 \geq 0$ and

$$\begin{aligned} y_1 + y_2 + y_3 &= 2 \\ 2y_1 + y_2 - y_3 &= 1 \end{aligned}$$

This leads to the following linear program:

$$\begin{aligned} \min \quad & 2y_1 + 2y_2 + 0.5y_3 \\ & y_1 + y_2 + y_3 = 2 \\ \text{s.t.} \quad & 2y_1 + y_2 - y_3 = 1 \\ & y_1, y_2, y_3 \geq 0 \end{aligned}$$

This is called the dual LP.

In general:

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax \leq b \end{aligned} \tag{P}$$

Dual of (P)

$$\begin{aligned} \min \quad & b^T y \\ \text{s.t.} \quad & y^T A = c^T \\ & y \geq 0 \end{aligned} \tag{D}$$

Remark:

We call (P) primal LP.

Theorem 2.4: Weak Duality

Let \bar{x} feasible for (P), \bar{y} feasible for (D). Then $c^T \bar{x} \leq b^T \bar{y}$.

Proof:

$$c^T \bar{x} = \bar{y}^T (A\bar{x}) \leq \bar{y}^T b$$

where we used $A\bar{x} \leq b$ and $\bar{y} \geq 0$. □

Corollary 2.5

Several results:

- If (P) is unbounded then (D) is infeasible.
- If (D) is unbounded then (P) is infeasible.

Note

(P) and (D) can both be infeasible.

- If \bar{x} is feasible for (P) \bar{y} feasible for (D) $c^T \bar{x} = b^T \bar{y}$, then \bar{x} optimal for (P), \bar{y} optimal for (D).

Theorem 2.6: Strong Duality

x^* is optimal for (P) $\iff \exists y^*$ feasible for (D) such that $c^T x^* = b^T y^*$.

Proof:

(\Leftarrow) ✓

(\Rightarrow) Is (D) infeasible?

$$\text{Suppose } \left\{ y \in \mathbb{R}^n : \begin{aligned} A^T y &= c \\ y &\geq 0 \end{aligned} \right\} = \emptyset$$

$$(\text{Alternate version of Farkas' Lemma}) \exists u : \begin{aligned} u^T A &\geq 0 \\ u^T c &< 0 \end{aligned} \iff \exists d : \begin{aligned} Ad &\leq 0 \\ c^T d &> 0 \end{aligned}$$

Take look at $x' = x^* + d$, then

$$\begin{aligned} Ax' &= Ax^* + Ad \leq b \\ c^T x' &= c^T x^* + c^T d > c^T x^* \end{aligned}$$

Contradiction. Thus (D) has an optimal solution y^* .

Now let $\gamma = b^T y^*$, and let $\theta := \left\{ x \in \mathbb{R}^n : \begin{array}{l} Ax \leq b \\ -c^T x \leq -\gamma \end{array} \right\}$.

If $\theta = \emptyset$, by Farkas'

$$\exists \left(\frac{\bar{y}}{\bar{\lambda}} \right) : \begin{cases} \left(\frac{\bar{y}}{\bar{\lambda}} \right)^T \begin{pmatrix} A \\ -c^T \end{pmatrix} = 0 \\ \left(\frac{\bar{y}}{\bar{\lambda}} \right)^T \begin{pmatrix} b \\ -\gamma \end{pmatrix} < 0 \\ \left(\frac{\bar{y}}{\bar{\lambda}} \right) \geq 0 \end{cases} \iff \begin{array}{l} A^T \bar{y} = c \bar{\lambda} \\ b^T \bar{y} < \gamma \bar{\lambda} \\ \bar{y} \geq 0 \\ \bar{\lambda} \geq 0 \end{array}$$

Case 1: $\bar{\lambda} > 0$.

Let $y' = \frac{\bar{y}}{\bar{\lambda}}$. Then we have

$$A^T y' = A^T \frac{\bar{y}}{\bar{\lambda}} = c \quad \text{and} \quad b^T y' = b^T \frac{\bar{y}}{\bar{\lambda}} < \gamma \quad \text{and} \quad y' = \frac{\bar{y}}{\bar{\lambda}} \geq 0$$

Contradicts optimality of y^* .

$$A^T y = 0$$

Case 2: $\bar{\lambda} = 0$. Then $b^T y < 0$

$$\bar{y} \geq 0$$

Now we can do the same thing previously. Let $y' = y^* + \bar{y}$, then

$$A^T y' = A^T y^* + A^T \bar{y} = c$$

and

$$y' = y^* + \bar{y} \geq 0$$

$$b^T y' = b^T y^* + b^T \bar{y} < b^T y^*$$

Contradicts optimality of y^* .

Thus $\theta \neq \emptyset$.

Let $\bar{x} \in \theta$,

$$c^T x^* \underbrace{\leq}_{\text{weak duality}} b^T y^* = \gamma \underbrace{\leq}_{\bar{x} \in \theta} c^T \bar{x} \leq c^T x^*$$

where the last inequality is because \bar{x} feasible for (P), x^* optimal for (P).

□

2.4 Possible Outcomes

See [here](#).

2.5 Duals of generic LPs

$$\begin{array}{rcll}
 \max & 2x_1 + 3x_2 - 4x_3 & & \\
 & x_1 & +7x_3 & \leq 5 \\
 & & 2x_2 - x_3 & \geq 3 \\
 \text{s.t.} & x_1 & +x_3 & = 8 \\
 & & x_2 & \leq 6 \\
 & x_1 & & \geq 0 \\
 & & x_2 & \leq 0
 \end{array}$$

$$\begin{array}{rcl}
 \max & (2, 3, -4)x & \\
 \text{s.t.} & \begin{pmatrix} 1 & 0 & 7 \\ 0 & -2 & 1 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} x \leq \begin{pmatrix} 5 \\ -3 \\ 8 \\ -8 \\ 6 \\ 0 \\ 0 \end{pmatrix} &
 \end{array}$$

and dual

$$\begin{array}{rcl}
 \min & (5, -3, 8, -8, 6, 0, 0)y & \\
 \text{s.t.} & \begin{pmatrix} 1 & 0 & 1 & -1 & 0 & -1 & 0 \\ 0 & -2 & 0 & 0 & 1 & 0 & 1 \\ 7 & 1 & 1 & -1 & 0 & 0 & 0 \end{pmatrix} y = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} \quad \text{and } y \geq 0 & (D_1)
 \end{array}$$

$$\begin{array}{rcl}
 \min & (5, -3, 8, -8, 6)y & \\
 \text{s.t.} & \begin{pmatrix} 1 & 0 & 1 & -1 & 0 \\ 0 & -2 & 0 & 0 & 1 \\ 7 & 1 & 1 & -1 & 0 \end{pmatrix} y \begin{matrix} \geq \\ \leq \\ = \end{matrix} \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} \quad \text{and } y \geq 0 & (D_2)
 \end{array}$$

Claim (y_1^*, \dots, y_5^*) is optimal for $(D_2) \iff (y_1^*, \dots, y_5^*, y_6^*, y_7^*)$ optimal for (D_1) with

$$\begin{aligned}
 y_6^* &= y_1^* + y_3^* - y_4^* - 2 \\
 y_7^* &= 3 - (-2y_2^* + y_5^*)
 \end{aligned}$$

$$\begin{array}{ll} \min & (5, 3, 8, 6)y \\ \text{s.t.} & \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ 7 & -1 & 1 & 0 \end{pmatrix} y \begin{array}{l} \geq \\ \leq \\ = \end{array} \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} \quad \text{and} \quad y_1 \geq 0, y_2 \leq 0 \quad y_4 \geq 0 \end{array} \quad (D_3)$$

Claim Opt value of (D_2) and (D_3) are same.

In general

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \\ & x \geq 0 \end{array} \quad (P) \quad \left| \quad \begin{array}{ll} \min & b^T y \\ \text{s.t.} & A^T y \leq c \\ & y \geq 0 \end{array} \quad (D)$$

2.5.1 Cheat Sheet

Here or

Primal (max)		Dual (min)	
Constraint	\leq	≥ 0	Variable
	\geq	≤ 0	
	$=$	free	
Variable	\geq	≥ 0	Constraint
	\leq	≤ 0	
	free	$=$	

Remark:

This is not symmetric... The way you can remember it is by thinking natural variables in real life, like you cannot have negative number of cars and so on...

Q What if you start with a minimization LP as primal?

Example:

$$\begin{array}{ll} \min & x_1 - x_2 \\ & 2x_1 + 3x_2 \leq 5 \\ \text{s.t.} & x_1 - x_2 \geq 3 \\ & x_1 + 5x_2 = 7 \\ & x_1 \geq 0, x_2 \leq 0 \end{array} \quad (P)$$

Rewrite as:

$$-1 \times \begin{pmatrix} \max & -x_1 + x_2 \\ \downarrow \\ \text{s.t.} & \dots \end{pmatrix}$$

Will lead to finding dual:

$$\begin{array}{ll} \max & 5y_1 + 3y_2 + 7y_3 \\ \downarrow & \\ & 2y_1 + y_2 \leq 1 \\ \text{s.t.} & 3y_1 - y_2 + 5y_3 \geq -1 \\ & y_1 \leq 0, y_2 \geq 0, y_3 \text{ free} \end{array}$$

Also

- Weak duality holds.

If \bar{x} feasible for (P), \bar{y} feasible for (D), then $c^T \bar{x} \geq b^T \bar{y}$.

- Strong duality holds

Note

The dual of the dual of (P) is (P).

Example:

Given a simple undirected graph $G = (V, E)$. $M \subseteq E$ is a *matching* if every vertex $v \in V$ is incident to ≤ 1 edge in M .

See examples of matching in [CO 342](#) or [MATH 249](#).

Max cardinality matching

Find matching M with largest $|M|$.

Define $x_e = \begin{cases} 1, & \text{if } e \in M \\ 0, & \text{otherwise} \end{cases}$.

$$\begin{array}{ll} \max & \sum_{e \in E} x_e \\ \downarrow & \\ & \sum_{e \in \delta(v)} x_e \leq 1, \quad \forall v \in V \\ \text{s.t.} & \\ & 0 \leq x_e, \quad \forall e \in E \end{array}$$

where $\delta(v)$ = set of edges in E incident to v .

$$\begin{array}{ll} \min & \sum_{v \in V} y_v \\ \downarrow & \\ \text{s.t.} & y_u + y_v \geq 1, \quad \forall e = uv \in E \\ & y \geq 0 \end{array}$$

2.6 Other interpretations of dual

Example:

			Resources	
		Per unit Profit	Per unit consumption	
			A	B
Product	1	5	2	3
	2	3	4	1
Available Resources			15	10

$$\begin{aligned}
 & \max \quad 5x_1 + 3x_2 \\
 & \downarrow \\
 & \quad 2x_1 + 4x_2 \leq 15 \\
 & \text{s.t.} \quad 3x_1 + x_2 \leq 10 \\
 & \quad x \geq 0
 \end{aligned}$$

Suppose somebody wants to buy A, B from me. What is the lowest price I should ask?

Let y_A, y_B be prices:

$$\begin{aligned}
 & \min \quad 15y_A + 10y_B \\
 & \downarrow \\
 & \quad 2y_A + 3y_B \geq 5 \\
 & \text{s.t.} \quad 4y_A + y_B \geq 3 \\
 & \quad y \geq 0
 \end{aligned}$$

Example: Zero-Sum

Alice, Bob play game. A: m choices. B: n choices. Alice play i , Bob plays j , Bob pays Alice M_{ij} dollars.

		Alice		
		R	P	S
Bob	R	0	1	-1
	P	-1	0	1
	S	1	-1	0

Zero-sum: Amount won by Alice - Amount won by Bob = 0

Let $y \in \mathbb{R}_+^m$, Alice's probability distribution.

Let $x \in \mathbb{R}_+^n$, Bob's probability distribution.

Expected Amount Bob pays Alice:

$$\sum_{i=1}^m \sum_{j=1}^n y_i M_{ij} x_j = y^T M x$$

$$P = \left\{ x \in \mathbb{R}^n : \sum x_j = 1, x \geq 0 \right\}$$

$$Q = \left\{ y \in \mathbb{R}^m : \begin{array}{l} \sum y_i = 1 \\ y \geq 0 \end{array} \right\}$$

Alice wants $\max_{y \in Q} \left\{ \min_{x \in P} y^T M_x \right\}$. Bob wants $\min_{x \in P} \left\{ \max_{y \in Q} y^T M_x \right\}$.

Suppose $\bar{y} \in Q$ is fixed. Bob's problem is

$$\begin{aligned} \min_{x \in P} \bar{y}^T M_x &= \min \sum_{j=1}^n \left(\sum_{i=1}^m M_{ij} \bar{y}_i \right) x_j \\ &\downarrow \\ \text{s.t.} \quad &\sum_{j=1}^n x_j = 1 \\ &x \geq 0 \end{aligned}$$

This is equivalent to picking smallest number in

$$\begin{aligned} &\left\{ \sum_{i=1}^m M_{ij} \bar{y}_i \right\}_{j=1}^n \\ \Rightarrow \max_{y \in Q} \min_{x \in P} y^T M_x &= \max_{y \in Q} \left\{ \begin{array}{l} \max u \\ \downarrow \\ \text{s.t.} \quad u \leq y^T M e_j, \quad \forall j = 1, \dots, n \end{array} \right\} \\ &= \begin{array}{l} \max u \\ \downarrow \\ \text{s.t.} \quad u \leq y^T M e_j, \quad \forall j = 1, \dots, n \\ y^T = 1 \\ y \geq 0 \end{array} \end{aligned}$$

Similarly Bob's problem:

$$\begin{aligned} \min v & \\ \downarrow & \\ \text{s.t.} \quad &v \geq e_i^T M x, \quad \forall i = 1, \dots, m \\ &x^T = 1 \\ &x \geq 0 \end{aligned}$$

There are x^*, y^* for which strategy values match \rightarrow Nash's Equilibrium.

Now get back to Farkas' Lemma Theorem 2.2. ¹

Proof:

$$\begin{aligned} \max \quad &0^T x \\ \downarrow & \\ \text{s.t.} \quad &Ax \leq b \end{aligned} \tag{P}$$

¹Rephrase it a little bit: Exactly one of the two has a solution (i) $Ax \leq b$ (ii) $u^T \dots$

$$\begin{array}{ll}
\min & b^T u \\
\downarrow & \\
\text{s.t.} & u^T A = 0 \\
& u \geq 0
\end{array} \tag{D}$$

(D) is always feasible ($u = 0$).

If $\exists \bar{x} : A\bar{x} \leq b$, \bar{x} optimal for (P) \implies optimal for (D) has value 0.
 $\implies \nexists u$ satisfying (i).

And the converse is also true. □

2.7 Complementary Slackness (C.S.)

Let x^*, y^* be feasible for primal and dual respectively.

Complementary Slackness

Abbreviated as C.S.

- i) Either $x_j^* = 0$ or corresponding dual constraint is tight at y^* , $\forall j = 1, \dots, n$.
- ii) Either $y_i^* = 0$ or corresponding primal constraint is tight at x^* , $\forall i = 1, \dots, m$.

Example:

$$\begin{array}{ll}
\min & x_1 - x_2 \\
\downarrow & \\
& 2x_1 + 3x_2 \leq 5 \\
\text{s.t.} & x_1 - x_2 \geq 3 \\
& x_1 + 5x_2 = 7 \\
& x_1 \geq 0, x_2 \leq 0
\end{array} \tag{P}$$

$$\begin{array}{ll}
\max & 5y_1 + 3y_2 + 7y_3 \\
\downarrow & \\
& 2y_1 + y_2 + y_3 \leq 1 \\
\text{s.t.} & 3y_1 - y_2 + 5y_3 \geq -1 \\
& y_1 \leq 0, y_2 \geq 0
\end{array} \tag{D}$$

- i) $x_1^* = 0$ OR $2y_1^* + y_2^* + y_3^* = 1$
 $x_2^* = 0$ OR $3y_1^* - y_2^* + 5y_3^* = -1$
- ii) $y_1^* = 0$ OR $2x_1^* + 3x_2^* = 5$
 $y_2^* = 0$ OR $x_1^* - x_2^* = 3$
 $y_3^* = 0$ OR $x_1^* + 5x_2^* = 7$

Theorem 2.7

Let x^*, y^* be feasible for primal/dual respectively. TFAE^a

- a) x^* opt for primal AND y^* opt. for dual
- b) Obj. value of $x^* =$ Obj. value of y^*
- c) x^*, y^* satisfy C.S.

^athe following are equivalent

Proof:

a) \iff b) done.

b) \iff c) Proof for

$$\begin{array}{ll} \max & c^T x \\ \downarrow & \\ \text{s.t.} & Ax \leq b \\ & x \geq 0 \end{array} \qquad \begin{array}{ll} \min & b^T y \\ \downarrow & \\ \text{s.t.} & A^T y \geq c \\ & y \geq 0 \end{array}$$

Note

$$A^T y \geq c \iff \sum_{i=1}^m a_{ij} y_i \geq c_j, \quad \forall j = 1, \dots, n$$

$$\begin{aligned} c^T x^* &= \sum_{j=1}^n c_j x_j^* \\ &\leq \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} y_i^* \right) x_j^* \\ &= \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} x_j^* \right) y_i^* \\ &\leq \sum_{i=1}^m b_i y_i^* = b^T y^* \end{aligned}$$

where first and second inequalities come from $x \geq 0, y \geq 0$ respectively.

(b) $c^T x^* = b^T y^* \iff$ C.S. holds. (Just play with some strict inequality conditions)

□

Example:

$$\begin{array}{ll} \max & x_1 + x_2 \\ \downarrow & \\ \text{s.t.} & x_1 + x_2 \leq 1 \end{array} \qquad \begin{array}{ll} \min & y \\ \downarrow & \\ & y = 1 \\ \text{s.t.} & y = 1 \\ & y \geq 0 \end{array}$$

Consider a pair $x^* = (0, 0), y^* = 1$ which violates CS.

2.7.1 Geometric Interpretation of C.S.

$$\begin{array}{ll} \max & c^T x \\ \downarrow & \\ \text{s.t.} & Ax \leq b \end{array} \qquad \begin{array}{ll} \min & c^T y \\ \downarrow & \\ \text{s.t.} & A^T y = c \\ & y \geq 0 \end{array}$$

$$A = \begin{pmatrix} - & a_1^T & - \\ & \vdots & \\ - & a_m^T & - \end{pmatrix}$$


C.S. says $a_i^T x^* = b_i$ or $y_i^* = 0$.

$$A^T y = c \implies \begin{pmatrix} | & | & \cdots & | \\ a_1 & a_2 & \cdots & a_m \\ | & | & & | \end{pmatrix} y = c \implies \sum_{i=1}^m a_i y_i = c$$

C.S. says c is a nonnegative combination of tight constraint at x^* .

Example:

$$\begin{array}{ll} \max & 2x_1 + 0.5x_2 \\ \downarrow & \\ & x_1 \leq 2 \\ & x_2 \leq 2 \\ \text{s.t.} & x_1 + x_2 \leq 3 \\ & x_1, x_2 \geq 0 \end{array}$$



Theorem 2.8

$$\begin{array}{ll} \max & c^T x \\ \downarrow & \\ \text{s.t.} & Ax \leq b \end{array} \quad (\text{P})$$

is unbounded iff (P) is feasible and $\exists d \in \mathbb{R}^n : \begin{array}{l} c^T d > 0 \\ Ad \leq 0 \end{array}$.

Proof:

\Rightarrow) Let \bar{x} feasible for (P), $\bar{x} + \lambda d$ is also feasible for (P) $\forall \lambda \geq 0$.

$c^T(\bar{x} + \lambda d)$ can be made arbitrary large.

\Leftarrow) Hard exercise but doable.

□

2.8 Geometry of Polyhedra

line segment




$\bar{x}, \bar{y} \in \mathbb{R}^n$ the line segment between \bar{x}, \bar{y} is

$$\left\{ x \in \mathbb{R}^n : \begin{array}{l} x = \lambda \bar{x} + (1 - \lambda) \bar{y} \\ \text{for some } \lambda \in [0, 1] \end{array} \right\}$$

convex set

S is a convex set if $\forall x, y \in S$, line segment between x, y is contained in S .

Example:



NOT a convex set

Polyhedra are convex sets. $P = \{x : Ax \leq b\}$. $\bar{x}, \bar{y} \in P$ then

$$A(\underbrace{\lambda}_{\geq 0} \bar{x} + \underbrace{(1 - \lambda)}_{\geq 0} \bar{y}) \leq \lambda b + (1 - \lambda)b = b$$

convex combination

Given $x^1, \dots, x^k \in \mathbb{R}^n$. We say \bar{x} is a convex combination of x^1, \dots, x^k if $\exists \lambda$:

$$\begin{aligned}\bar{x} &= \sum_{i=1}^k \lambda_i x^i \\ 1 &= \sum_{i=1}^k \lambda_i \\ \lambda &\geq 0\end{aligned}$$

Optimal solution seems to be happen at “corners”.

Let P be a polyhedron $P = \{x \in \mathbb{R}^n : Ax \leq b\}$.

vertex

\bar{x} is a vertex of P if $\exists c$: \bar{x} is unique optimal solution to

$$\begin{aligned}\max \quad & c^T x \\ \text{s.t.} \quad & Ax \leq b\end{aligned}$$

extreme point

\bar{x} is an extreme point of P if $\nexists u, v \in P \setminus \{\bar{x}\}$ such that \bar{x} is in line segment between u, v .

basic feasible solution

$\bar{x} \in P$ is a basic feasible solution of P if there are n linearly independent tight constraints at \bar{x} .

Note

Constraints

$$a_i^T x \leq b_i, \quad \forall i = 1, \dots, m$$

are linearly independent if $\{a_i\}_{i=1}^m$ are linearly independent.

Theorem 2.9

Let $\bar{x} \in P$. TFAE:

- a) \bar{x} is a vertex of P .
- b) \bar{x} is a basic feasible solution of P .
- c) \bar{x} is a extreme point of P .

Proof:a) \implies c) Suppose $\exists u, v \in P \setminus \{\bar{x}\}$ such that

$$\bar{x} = \lambda u + (1 - \lambda)v$$

for some $\lambda \in (0, 1)$. Consider c for which \bar{x} is an optimal solution to

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & x \in P \end{aligned}$$

$$\implies \begin{aligned} c^T \bar{x} &\geq c^T u \\ c^T \bar{x} &\geq c^T v \end{aligned}$$

and

$$c^T \bar{x} = \underbrace{\lambda}_{\geq 0} c^T u + \underbrace{(1 - \lambda)}_{\geq 0} c^T v \leq \lambda c^T \bar{x} + (1 - \lambda) c^T \bar{x} = c^T \bar{x}$$

$$\implies c^T u = c^T v = c^T \bar{x}$$

 $\implies \bar{x}$ NOT a vertex.c) \implies b) Suppose \bar{x} is not a BFS. Let $i \subseteq \{1, \dots, m\}$ be the index set of tight constraint at \bar{x} . Consider

$$a_i^T d = 0, \quad \forall i \in I \tag{*}$$

But since \bar{x} not BFS, $\exists \bar{d} \neq 0$ satisfying $(*)$.^a

$$x(\epsilon) = \bar{x} + \epsilon \bar{d}$$

$$a_i^T x(\epsilon) = a_i^T \bar{x} \leq b_i, \quad \forall i \in I$$

$$a_i^T x(\epsilon) = \underbrace{a_i^T \bar{x}}_{< b_i} + \epsilon a_i^T \bar{d} \leq b_i, \quad \forall i \notin I$$

which is satisfied if $|\epsilon|$ is small enough. $x(\epsilon) \in P$ if $|\epsilon|$ is small enough.

But then

$$\bar{x} = \frac{1}{2}x(\epsilon) + \frac{1}{2}x(-\epsilon)$$

b) \implies a) Let $I \subseteq \{1, \dots, m\}$ index set of tight constraint at \bar{x} .

Define

$$c := \sum_{i \in I} a_i$$

Then $\forall x \in P$

$$c^T x = \sum_{i \in I} a_i^T x \leq \sum_{i \in I} b_i$$

And

$$c^T \bar{x} = \sum_{i \in I} a_i^T \bar{x} = \sum_{i \in I} b_i$$

$\implies \bar{x}$ is optimal solution to

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & x \in P \end{array} \quad (**)$$

If $x' \in P$ is optimal solution to $(**)$, then

$$a_i^T x' = b_i, \quad \forall i \in I \quad (***)$$

But since there are n linear independent constraints in I , \bar{x} is unique solution to $(***)$. $\implies x' = \bar{x}$.

□

^aby Rank-Nullity Theorem.


Q When does P have extreme points?

line

Let $\bar{x}, \bar{d} \in \mathbb{R}^n$, $\bar{d} \neq 0$. The set


$$\{x \in \mathbb{R}^n : x = \bar{x} + \lambda \bar{d} \text{ for some } \lambda \in \mathbb{R}\}$$

is called a line.



We say a polyhedron P has a line if $\exists \bar{x}, \bar{d}$ has a line if $\exists \bar{x}, \bar{d}$ s.t. $\bar{x} \in P, \bar{d} \neq 0$ and

$$\{x \in \mathbb{R}^n : x = \bar{x} + \lambda \bar{d} \text{ for some } \lambda \in \mathbb{R}\} \subseteq P$$



Proposition 2.10

$P = \{x \in \mathbb{R}^n : Ax \leq b\}$ has a line iff $P \neq \emptyset$ and $\exists \bar{d} \neq 0$ such that $A\bar{d} = 0$

$\iff P \neq \emptyset$ and $\text{rank}(A) < n$

Proof:

Exercise.

□

Theorem 2.11

$P = \{x \in \mathbb{R}^n : Ax \leq b\}$ has an extreme point

$\iff P \neq \emptyset$ and P has no lines.

Proof:

Exercise. □

pointed polyhedron

A non-empty polyhedron is called pointed if it has no lines.

Note

not pointed does not imply bounded. For example, in \mathbb{R}^2 , $x \geq 0$ and $y \geq 0$.

Theorem 2.12

Let $P \neq \emptyset$ pointed polyhedron. If $\max_{x \in P} c^T x$ (LP) has an optimal solution, it has an optimal solution that is an extreme point.

Proof:

Let \bar{x} be an optimal solution to (LP) with largest number of linear independent tight constraints.

Suppose there are $\leq n - 1$ linear independent tight constraints at \bar{x} .

Pick $\bar{d} \neq 0$ such that $a_i^T \bar{d} = 0, \forall i \in I$, where I is the index set of tight constraints. By the exact same argument as before, $\bar{x} \pm \epsilon \bar{d} \in P$ for ϵ small enough. But

$$c^T(\bar{x} \pm \epsilon \bar{d}) = c^T \bar{x} \pm \epsilon c^T \bar{d}$$

$$\implies c^T \bar{d} = 0$$

$$\implies c^T d(\bar{x} \pm \epsilon d) = c^T \bar{x}$$



Since P is pointed, $\exists \bar{\epsilon}$ for which

$$\bar{x} \pm \bar{\epsilon} \in P$$

and one of them not in P if $|\epsilon| > \bar{\epsilon}$. That can only happen if

$$a_k^T(\bar{x} + \bar{\epsilon} \bar{d}) = b_k \quad \text{or} \quad a_k^T(\bar{x} - \bar{\epsilon} \bar{d}) = b_k$$

for some $k \notin I$.

$\implies a_k^T \bar{d} \neq 0, \implies a_k$ is linear independent from $\{a_i\}_{i \in I}$ since non-zero cannot be linear combination of zeros. Contradiction to choice of \bar{x} . \square

2.9 Simplex Algorithm

Standard Equality Form

A linear program is in Standard Equality Form (SEF) if it is of the form

$$\begin{array}{ll} \max & c^T x \\ \downarrow & \\ \text{s.t.} & Ax = b \\ & x \geq 0 \end{array}$$

Proposition 2.13

Given any linear program, there exists an equivalent LP in SEF.

Example:

$$\begin{array}{ll} \max & x_1 + 2x_2 + x_3 \\ \downarrow & \\ & 3x_1 + x_2 \leq 5 \\ \text{s.t.} & -x_1 + x_3 \geq 6 \\ & x_1 \leq 0, x_3 \geq 0 \end{array} \quad (\text{P1})$$

$$\begin{aligned} x'_1 &= -x_1 \geq 0 \text{ and} \\ x_2 &= x_2^+ - x_2^- \text{ where } x_2^+ \geq 0, x_2^- \geq 0 \end{aligned}$$

We introduce

$$s_1 = 5 - 3x_1 - x_2 \geq 0, \quad s_2 = -x_1 + x_3 - 6 \geq 0$$

Then

$$\begin{array}{ll} \max & -x'_1 + 2x_2^+ - 2x_2^- + x_3 \\ \downarrow & \\ & -3x'_1 + 2x_2^+ - x_2^- + s_1 = 5 \\ \text{s.t.} & x'_1 + x_3 - s_2 = 6 \\ & x'_1, x_2^+, x_2^-, x_3, s_1, s_2 \geq 0 \end{array} \quad (\text{P2})$$

x feasible for (P1) $\iff (x'_1, x_2^+, x_2^-, x_3, s_1, s_2)$ feasible for (P2) and they have same cost.

Assumption $A \in \mathbb{R}^{m \times n} \rightarrow \text{rank}(A) = m$. This is WLOG. Since if

$$a_i = \sum_{k \neq i} \lambda_k a_k$$

Either

$$b_i \neq \sum_{k \neq i} \lambda_k b_k$$

in which case (SEF) is infeasible. Or $a_i^T x = b_i$ is redundant. So it can be removed from (SEF).

Note

$\{x : Ax = b, x \geq 0\}$ is *pointed* polyhedron (if nonempty).

Structure of BFS Any feasible solution has m linear independent tight constraints ($n - m$) extra tight constraint must come from $x_j \geq 0$.

Let $B \subseteq \{1, \dots, n\}$ such that $|B| = m$ and A_B ² is invertible.

$N = \{1, \dots, n\} \setminus B$. $x_N = 0$, i.e. $x_j = 0, \forall j \in N$.

Feasible solutions obtained this way are precisely BFS.

Example:

$$\begin{array}{ll} \max & (3 \quad 2 \quad 1 \quad 4) x \\ \downarrow & \\ \text{s.t.} & \begin{pmatrix} 1 & 2 & -1 & 0 \\ 2 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 5 \\ 7 \end{pmatrix} \\ & x \geq 0 \end{array}$$

If we pick

$$\begin{array}{ll} B = \{1, 2\} & A_B = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \\ N = \{3, 4\} & A_N = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \\ C_B = (3 \quad 2)^T & C_N = (1 \quad 4)^T \end{array}$$

$$x_B = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad x_N = \begin{pmatrix} x_3 \\ x_4 \end{pmatrix}$$

$$B = \{1, 3\}, B = \{2, 4\}, A_B = \begin{pmatrix} 1 & -1 \\ 2 & 0 \end{pmatrix}, A_N = \begin{pmatrix} 2 & 0 \\ 1 & 1 \end{pmatrix}$$

$$C_B = \begin{pmatrix} 3 \\ 1 \end{pmatrix}, C_N = \begin{pmatrix} 2 \\ 4 \end{pmatrix}, x_B = \begin{pmatrix} x_1 \\ x_3 \end{pmatrix}, x_N = \begin{pmatrix} x_2 \\ x_4 \end{pmatrix}$$

If we set $x_N = 0$ (for $B = \{1, 3\}$) we are left with

$$\begin{pmatrix} 1 & -1 \\ 2 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_3 \end{pmatrix} = \begin{pmatrix} 5 \\ 7 \end{pmatrix}$$

This has a unique solution $x_1 = 3.5, x_3 = -1.5$, but not feasible.

² A_B is submatrix obtained by picking columns of A indexed by B . Such B is called a basis.

If we pick $B = \{1, 2\}$

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 5 \\ 7 \end{pmatrix}$$

$\underbrace{x_3 = x_4 = 0}_{x_N}, x_1 = 3, x_2 = 1$, which is feasible.

In general,

$$Ax = b \iff A_B x_B + \cancel{A_N x_N}^0 = b$$

has unique solution $x_b = A_B^{-1}b$.

For any basis B , the corresponding *basic solution* is

$$\begin{pmatrix} x_B \\ x_N \end{pmatrix} = \begin{pmatrix} A_B^{-1}b \\ 0 \end{pmatrix}$$

If $A_B^{-1}b \geq 0$, then it is a *BFS*.

2.9.1 Canonical Form

Let B be a feasible basis (i.e. corresponding basis solution is feasible).

$$\begin{aligned} Ax = b &\iff A_B x_B + A_N x_N = b \\ &\iff x_B + A_B^{-1} A_N x_N = A_B^{-1}b \end{aligned}$$

Now let's take a look at objective.

$$\begin{aligned} c^T x &= c_B^T x_B + c_N^T x_N - \cancel{c_B^T (x_B + A_B^{-1} A_N x_N - A_B^{-1}b)} \\ &= (c_N^T - c_B^T A_B^{-1} A_N) x_N + c_B^T A_B^{-1}b \end{aligned}$$

Thus (SEF) is said to be in canonical form for B if it is written as

$$\begin{aligned} \max \quad & \overbrace{(c_N^T - c_B^T A_B^{-1} A_N)}^{\text{Reduced costs}} x_N + c_B^T A_B^{-1}b \\ \downarrow \\ \text{s.t.} \quad & x_B + A_B^{-1} A_N x_N = A_B^{-1}b \\ & x_B, x_N \geq 0 \end{aligned}$$

Example:

Back to our previous example...

$B = \{1, 2\}$. Rewriting in canonical form for B :

$$\begin{aligned} A_B^{-1} &= \begin{pmatrix} -1/3 & 2/3 \\ 2/3 & -1/3 \end{pmatrix} \\ A_B A &= \begin{pmatrix} 1 & 0 & 1/3 & -2/3 \\ 0 & 1 & 2/3 & -1/3 \end{pmatrix} \end{aligned}$$

$$c_B^T A_B^{-1} A_N = (3 \quad 2) \begin{pmatrix} 1/3 & -2/3 \\ 2/3 & -1/3 \end{pmatrix} = (7/3 \quad -8/3)$$

$$c_N^T - c_B^T A_B^{-1} A_N = (-4/3 \quad 4/3)$$

Then

$$\begin{aligned} \max \quad & (0 \quad 0 \quad -4/3 \quad 4/3)x + 11 \\ \downarrow \\ \text{s.t.} \quad & \begin{pmatrix} 1 & 0 & 1/3 & -2/3 \\ 0 & 1 & 2/3 & -1/3 \end{pmatrix} x = \begin{pmatrix} 3 \\ 1 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

is in canonical form for $B = \{1, 2\}$.

Example:

$$\begin{aligned} \max \quad & (1 \quad 3 \quad -2 \quad 0 \quad 0)x \quad \underbrace{+0}_{\text{obj. value}} \\ \downarrow \\ \text{s.t.} \quad & \begin{pmatrix} 1 & 1 & 1 & 1 & 0 \\ 1 & -1 & 3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 1 \end{pmatrix} \\ & x \geq 0 \end{aligned} \quad (\text{LP})$$

Canonical form for $B = \{4, 5\}$.

Corresponding BFS $\begin{matrix} x_4 = 4 \\ x_5 = 1 \end{matrix}$, $x_j = 0, \forall j \in N$

$$x = (0 \quad 0 \quad 0 \quad 4 \quad 1)^T$$

Objective value = 0

If increase x_1 or x_2 . Objective function increases.

Let's try to increase x_1 from $0 \rightarrow \theta$. (Keep $x_2 = x_3 = 0$)

$$\begin{aligned} \theta + x_4 = 4 & \iff x_4 = 4 - \theta \\ \theta + x_5 = 1 & \iff x_5 = 1 - \theta \end{aligned}$$

New objective: $0 + \theta$. However, we have

$$\begin{aligned} x_4 \geq 0 & \implies \theta \leq 4 \\ x_5 \geq 0 & \implies \theta \leq 1 \end{aligned} \implies \text{Increase } x_1 \text{ by } 1$$

x_5 will be 0 \rightarrow $\begin{matrix} x_1 \text{ enters basis} \\ x_5 \text{ leaves basis} \end{matrix}$. Then new basis $B = \{1, 4\}$.

Rewriting (LP) in canonical form for $B = \{1, 4\}$.

$$\begin{array}{l}
 \max \quad (0 \quad 4 \quad -5 \quad 0 \quad -1) x + \underbrace{1}_{\text{obj. value}} \\
 \downarrow \\
 \text{s.t.} \quad \begin{pmatrix} 1 & -1 & 3 & 0 & 1 \\ 0 & 2 & -2 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \\
 x \geq 0
 \end{array}$$

Corresponding BFS:

$$x = (1 \quad 0 \quad 0 \quad 3 \quad 0)^T$$

Obj. value = 1

Pick $j \in N$: $\bar{c}_j > 0$ ($j = 2$)

Increase x_2 to θ , keep $x_3 = x_5 = 0$

$$\begin{aligned}
 x_1 - \theta &= 1 \iff x_1 = 1 + \theta \\
 x_4 + 2\theta &= 3 \iff x_4 = 3 - 2\theta
 \end{aligned}$$

and

$$\begin{aligned}
 x_1 \geq 0 &\implies \theta \geq -1 \\
 x_4 \geq 0 &\implies \theta \leq \frac{3}{2}
 \end{aligned}$$

Set $\theta \leftarrow \frac{3}{2} \rightarrow$ x_2 enters basis
 x_4 leaves basis

New basis $B = \{1, 2\}$.

(LP) in canonical form for $B = \{1, 2\}$.

$$\begin{array}{l}
 \max \quad (0 \quad 0 \quad -1 \quad -2 \quad 1) x + 7 \\
 \downarrow \\
 \text{s.t.} \quad \begin{pmatrix} 1 & 0 & 2 & 0.5 & 0.5 \\ 0 & 1 & -1 & 0.5 & -0.5 \end{pmatrix} x = \begin{pmatrix} 2.5 \\ 1.5 \end{pmatrix} \\
 x \geq 0
 \end{array}$$

Corresponding BFS:

$$x = (2.5 \quad 1.5 \quad 0 \quad 0 \quad 0)^T$$

Obj. value = 7

Find $j \in N$, $\bar{c}_j > 0$ ($j = 5$)

$$\begin{aligned}
 x_1 = 2.5 - 0.5\theta \geq 0 &\implies \theta \leq 5 \rightarrow x_1 \text{ leaves basis} \\
 x_2 = 1.5 + 0.5\theta \geq 0 &\implies \theta \geq -3 \rightarrow x_5 \text{ enters basis}
 \end{aligned}$$

New basis $B = \{2, 5\}$

(LP) in canonical form for $B = \{2, 5\}$

$$\begin{aligned} & \max \quad (-2 \quad 0 \quad -5 \quad -3 \quad 0) x + 12 \\ & \downarrow \\ & \text{s.t.} \quad \begin{pmatrix} 1 & 1 & 1 & 1 & 0 \\ 2 & 0 & 4 & 1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 5 \end{pmatrix} \\ & \quad \quad \quad x \geq 0 \end{aligned}$$

BFS $x = (0 \quad 4 \quad 0 \quad 0 \quad 5)^T$ } Optimal Solution
 Obj. value = 12.

2.9.2 Iteration of simplex

Algorithm 2: Iteration of simplex

- 1 Start with feasible basis B
 - 2 Rewrite LP in canonical form for B
 - 3 Pick $j \in N : \bar{c}_j > 0$ (x_j enters basis)
 - 4 Let $\bar{b} = A_B^{-1}b$, $\bar{A}_N = A_B^{-1}A_N$
 Find largest θ so that $\bar{b} - \theta\bar{A}_j \geq 0$.
 Corresponding basic variable that becomes 0 (say x_k) leaves basis.
 - 5 $B \leftarrow B \setminus \{k\} \cup \{j\}$. Iterate.
-

If problem has optimal solution AND θ is always > 0 , simplex finishes.

Note

If at current BFS we have a basic variable = 0, we may have $\theta = 0$. \rightarrow May lead to cycling. (i.e. return to current basis in future iteration)

Bland's Rule

If there are multiple choices of entering or leaving variables, always pick lowest index variable.

Using Bland's Rule avoids cycling

Observations If $\bar{c}_N \leq 0$, then the (LP) obj. value in canonical form is

$$\underbrace{\bar{c}_N^T}_{\leq 0} \underbrace{x_N}_{\geq 0} + c_B^T A_B^{-1} b \leq c_B^T A_B^{-1} b$$

For any feasible solution \implies Current BFS is optimal




Figure 2.1: Simplex method

Original LP

$$\begin{aligned} \max \quad & c^T x \\ \downarrow \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

Dual

$$\begin{aligned} \min \quad & b^T y \\ \downarrow \\ \text{s.t.} \quad & A^T y \geq c \end{aligned} \quad \Longleftrightarrow \quad \begin{aligned} \min \quad & y^T b \\ \downarrow \\ \text{s.t.} \quad & y^T A \geq c^T \end{aligned}$$

$$\Longleftrightarrow \begin{aligned} \min \quad & y^T b \\ \downarrow \\ \text{s.t.} \quad & y^T A_B \geq c_B^T \\ & y^T A_N \geq c_N^T \end{aligned}$$

If satisfies C.S with BFS corresponding to B

$$\begin{aligned} y^T A_B &= c_B^T \\ \implies y^T &= c_B^T A_B^{-1} \Longleftrightarrow c_B^T A_B^{-1} A_N \geq c_N^T \Longleftrightarrow \bar{c}_N \leq 0 \\ y^T A_N &\geq c_N^T \end{aligned}$$

2.9.3 Mechanics of Simplex

Example: 1

$$\begin{aligned} \max \quad & (1 \ 3 \ -2 \ 0 \ 0) x \\ \downarrow \\ \text{s.t.} \quad & \begin{pmatrix} 1 & 1 & 1 & 1 & 0 \\ 1 & -1 & 3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 1 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

↑ enters basis ↑ j
↑ pivot row ℓ

For θ

$$\theta \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 4 \\ 1 \end{pmatrix}$$

and we have

$$\begin{pmatrix} x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 4 - \theta \\ 1 - \theta \end{pmatrix} \geq 0 \implies \boxed{\theta \leq 4}$$

We are actually picking $\min \left\{ \frac{4}{1}, \frac{1}{1} \right\}$

Pick, out of all rows $\min \left\{ \frac{\bar{b}_i}{\bar{a}_{ij}} \right\}$ where j is entering variable.

Then now in row ℓ (second row here). Make row operations so that pivot element become 1, all others in col j becomes 0.

→ Row 2×1

→ Subtract row 2 from row 1

→ subtract row 2 from objective function (with RHS multiplied by -1)

$$\begin{array}{l} \max \quad (0 \quad 4 \quad -5 \quad 0 \quad -1) x + 1 \\ \downarrow \\ \text{s.t.} \quad \begin{pmatrix} 0 & 2 & -2 & 1 & -1 \\ 1 & -1 & 3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 3 \\ 1 \end{pmatrix} \\ x \geq 0 \end{array} \quad \begin{array}{l} \text{pivot} \leftarrow \text{row } \ell \end{array}$$

$$2\theta + x_4 = 3 \iff x_4 = 3 - 2\theta \geq 0 \implies \theta \leq \frac{3}{2}$$

$$-\theta + x_1 = 1 \iff x_1 = \theta + 1 \geq 0 \implies \theta \geq -1$$

where we are finding $\min_{\bar{a}_{ij} > 0} \left\{ \frac{\bar{b}_i}{\bar{a}_{ij}} \right\}$. Now follow the similar procedure, we have

$$\begin{array}{l} \max \quad (0 \quad 0 \quad -1 \quad -2 \quad 1) x + 7 \\ \downarrow \\ \text{s.t.} \quad \begin{pmatrix} 0 & 1 & -1 & 0.5 & -0.5 \\ 1 & 0 & 2 & 0.5 & 0.5 \end{pmatrix} x = \begin{pmatrix} 1.5 \\ 2.5 \end{pmatrix} \end{array}$$

In general Pick $j \in N : \bar{c}_j > 0$.

Let $\ell = \operatorname{argmin}_{\bar{a}_{ij} > 0} \left\{ \frac{\bar{b}_i}{\bar{a}_{ij}} \right\}$ (Ratio Test)

- Multiply row ℓ by $\frac{1}{\bar{a}_{\ell j}}$
- Add $-\frac{\bar{a}_{ij}}{\bar{a}_{\ell j}}$ times row ℓ to row $i \neq \ell$.

- Add $-\frac{\bar{c}_j \cdot \bar{a}_{\ell k}}{\bar{a}_{\ell j}}$ to variable coeff in objective. $\forall k \in 1, \dots, n$
- Add $\frac{\bar{b}_\ell \cdot \bar{c}_j}{\bar{a}_{\ell j}}$ to objective value in objective function

Example: 2

$$\begin{array}{ll} \max & (2 \quad 1 \quad 1 \quad 0 \quad 0) x \\ \downarrow & \\ \text{s.t.} & \begin{array}{c} \text{pivot} \uparrow \\ \left(\begin{array}{ccccc} 1 & 2 & -1 & 1 & 0 \\ 2 & -2 & -1 & 0 & 1 \end{array} \right) x = \begin{pmatrix} 2 \\ 3 \end{pmatrix} \\ x \geq 0 \end{array} \quad \text{row } \ell \end{array}$$

Ratio Test $\min \left\{ \frac{2}{1}, \frac{3}{2} \right\} = 1.5. \ell = 2. (x_2 \text{ enters, } x_5 \text{ leaves})$

$$\begin{array}{ll} \max & (0 \quad 3 \quad 2_j \quad 0 \quad -1) x + 3 \\ \downarrow & \\ \text{s.t.} & \begin{pmatrix} 0 & 3 & -0.5 & 1 & -0.5 \\ 1 & -1 & -0.5 & 0 & 0.5 \end{pmatrix} x = \begin{pmatrix} 0.5 \\ 1.5 \end{pmatrix} \\ & x \geq 0 \end{array}$$

If we increase $x_3 \rightarrow \theta$ and keep $x_2 = x_5 = 0$

$$\begin{array}{ll} -0.5\theta + x_4 = 0.5 & \Rightarrow x_1 = 1.5 + 0.5\theta \\ -0.5\theta + x_1 = 1.5 & \Rightarrow x_4 = 0.5 + 0.5\theta \end{array} \rightarrow \text{Problem is unbounded!}$$

In general Let B be a basis

$$\begin{array}{ll} \max & \bar{c}_N^T x_N \\ \downarrow & \\ \text{s.t.} & \begin{array}{l} x_B + \bar{A}_N x_N = \bar{b} \\ x_B, x_N \geq 0 \end{array} \end{array}$$

Found $j : \bar{c}_j > 0$ AND $\bar{A}_j \leq 0$.

Construct $d \in \mathbb{R}^n$ to reflect what we are trying to do when we increase $x_j \rightarrow \theta$.

Right now, we are at BFS:

$$\begin{pmatrix} x_B \\ x_N \end{pmatrix} = \begin{pmatrix} A_B^{-1} b \\ 0 \end{pmatrix}$$

We want:

$$\begin{pmatrix} x_B \\ x_N \end{pmatrix} = \begin{pmatrix} A_B^{-1} b \\ 0 \end{pmatrix} + \theta \begin{pmatrix} d_B \\ d_N \end{pmatrix}$$

where $d_N = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \overset{j}{\uparrow} = e_j$ and $d_B = -\bar{A}_j = -A_B^{-1}A_j$.

Found d : $d \geq 0$, then

$$Ad = A_B d_B + A_N d_N = -A_B A_B^{-1} A_j + A_j = 0$$

and

$$c^T d = c_B^T d_B + c_N^T d_N = -c_B^T A_B^{-1} A_j + c_j = \bar{c}_j > 0$$

i.e.,

$$c^T d > 0$$

$$Ad = 0 \implies \text{Problem is unbounded}$$

$$d \geq 0$$

Index

B

basic feasible solution 25

C

Complementary Slackness 21
convex combination 25
convex set 24

E

extreme point 25

H

halfspace, hyperplane, polyhedron .. 7

I

infimum 5

L

line 27
line segment 24

P

pointed polyhedron 28

S

Standard Equality Form 29
supremum 5

V

vertex 25