

(a) From retrieval to fitted evidence

934

articles retrieved and screened

Elsevier and Springer, Cohort A + Cohort B

50

papers contributing fitted curves

the cohort every fitted quantity rests on

35

distinct compound labels

27 composition keys under the reduction of Sec. II.D

3,303

extracted critical-current points

digitised $J_c(T)$ and $J_c(H)$ readings

20

compounds fittable on both axes

the two-axis gate; the closed-form comparison ran on the pre-withdrawal 23

96

per-paper critical-current anchors

over 32 papers; the repair of Sec. III.F withdraws 26 and rescales 15

The retrieval corpus is the screening scope. Every fitted quantity rests on a smaller cohort, and the two highlighted rows are the cohorts the paper's claims are made on.

(b) Candidate dispatch, by family

Iron chalcogenide 11-type



0 / 29

29 refused: field below window 29 target above T_c 29 family field axis not validated 17

Iron pnictide 122-type



0 / 51

51 refused: no critical field 42 field below window 9 target above T_c 6

MgB₂-class



84 / 103

19 refused: no critical field 18 field below window 1 target above T_c 1 temperature above window 1



dispatched: the family envelope is emitted



refused: every target hit a refusal gate

a refused compound may hit several gates, so the counts above overlap

84 of 183 candidate compounds receive at least one prediction.

The refusal reason is recorded for every one of the rest.