

# Week2 Assignment Econometrics

*Pradeepta Das*

*11 November 2020*

## Questions

(a) **Prove:**

$$E(b_R) = \beta_1 + P\beta_2$$

First, it is important to express  $b_R$  in terms of  $\epsilon$ :

$$b_R = (X_1'X_1)^{-1}X_1'y = (X_1'X_1)^{-1}X_1'(X_1\beta_1 + X_2\beta_2 + \epsilon) = (X_1'X_1)^{-1}X_1'X_2\beta_2 + (X_1'X_1)^{-1}X_1'\epsilon = \beta_1 + P\beta_2 + (X_1'X_1)^{-1}X_1'\epsilon \text{ so, } E(b_R) = \beta_1 + P\beta_2$$

(b) **Prove:**

$$\text{var}(b_R) = \sigma^2(X_1'X_1)^{-1}$$

First, it is important to express  $b_R$  in terms of  $\epsilon$ :

$$b_R - E(b_R) = \beta_1 + P\beta_2 + (X_1'X_1)^{-1}X_1'\epsilon - \beta_1 - P\beta_2 = (X_1'X_1)^{-1}X_1'\epsilon \text{ var}(b_R) = E[(X_1'X_1)^{-1}X_1'\epsilon((X_1'X_1)^{-1}X_1'\epsilon)'] = E[(X_1'X_1)^{-1}X_1'\epsilon\epsilon'X_1(X_1'X_1)^{-1}] = \sigma^2(X_1'X_1)^{-1}$$

(c) **Prove:**

$$b_R = b_1 + Pb_2$$

$$b_R = (X_1'X_1)^{-1}X_1'y = (X_1'X_1)^{-1}X_1'(X_1b_1 + X_2b_2 + \text{res}) = b_1 + Pb_2 + (X_1'X_1)^{-1}X_1'\text{res} = b_1 + Pb_2 \text{ as } X_1'\text{res} = 0 \text{ because of orthogonality}$$

(d) **Argue that the columns of the  $(2 \times 3)$  matrix  $P$  are obtained by regressing each of the variables 'Age', 'Educ', and 'Parttime' on a constant term and the variable 'Female'.**

So,

$$P = (X_1'X_1)^{-1}X_1'(X_2)$$

is a  $(2 \times 3)$  matrix, with columns as such:

Column 1:

$$(X_1'X_1)^{-1}X_1'(\text{Age}) \text{ the OLS formula for regressing 'Age' on } X_1.$$

Column 2:

$$(X_1'X_1)^{-1}X_1'(\text{Educ}) \text{ the OLS formula for regressing 'Educ' on } X_1.$$

Column 3:

$$(X_1'X_1)^{-1}X_1'(\text{Parttime}) \text{ the OLS formula for regressing 'Parttime' on } X_1.$$

(e) P matrix value.

```
file25 = "C:\\Users\\Gannon_chef\\Documents\\Pradeepta\\Coursera\\Econometrics\\TrainExer21.txt"
new_logwage<-read.table(file25, header = TRUE, sep = ",", dec = ".")
str(new_logwage)
```

```
## 'data.frame':    500 obs. of  7 variables:
## $ Observ. : int  1 2 3 4 5 6 7 8 9 10 ...
## $ Wage    : int  66 34 70 47 107 188 123 57 42 200 ...
## $ LogWage : num  4.19 3.53 4.25 3.85 4.67 ...
## $ Female  : int  0 1 1 0 1 1 1 1 1 0 ...
## $ Age     : int  49 42 42 38 54 54 47 39 25 59 ...
## $ Educ    : int  1 1 1 1 1 1 1 1 1 1 ...
## $ Parttime: int  1 1 1 0 1 0 0 1 0 0 ...
```

```
X1 = new_logwage['Female']
X1['Constant']<-rep(1, length(new_logwage['Female']))
X1 <- X1[,c(2,1)]
X2 = new_logwage[c('Age', 'Educ', 'Parttime')]
str(X1)
```

```
## 'data.frame':    500 obs. of  2 variables:
## $ Constant: num  1 1 1 1 1 1 1 1 1 1 ...
## $ Female  : int  0 1 1 0 1 1 1 1 1 0 ...
```

```
str(X2)
```

```
## 'data.frame':    500 obs. of  3 variables:
## $ Age     : int  49 42 42 38 54 54 47 39 25 59 ...
## $ Educ    : int  1 1 1 1 1 1 1 1 1 1 ...
## $ Parttime: int  1 1 1 0 1 0 0 1 0 0 ...
```

Now, value of P matrix can be computed as:

```
x1_mat = as.matrix(sapply(X1, as.numeric))
x2_mat = as.matrix(sapply(X2, as.numeric))
P<-solve(t(x1_mat)%*%(x1_mat))%*%t(x1_mat)%*%x2_mat
P
```

```
##           Age      Educ  Parttime
## Constant 40.0506329  2.2594937 0.1962025
## Female   -0.1104155 -0.4931893 0.2494496
```

(f) Check the numerical validity of the result in part (c). Note: This equation will not hold exactly because the coefficients have been rounded to two or three decimals; preciser results would have been obtained for higher precision coefficients.

```
model <- lm(LogWage~Female+Age+Educ+Parttime, data=new_logwage)
summary(model)
```

```
##
## Call:
## lm(formula = LogWage ~ Female + Age + Educ + Parttime, data = new_logwage)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.73799 -0.15705 -0.00388  0.16495  0.77827
##
```

```
## Coefficients:
##           Estimate Std. Error t value Pr(>|t|)
## (Intercept)  3.052697   0.055333  55.170  <2e-16 ***
## Female      -0.041073   0.024710  -1.662   0.0971 .
## Age         0.030606   0.001273  24.041  <2e-16 ***
## Educ        0.233163   0.010660  21.873  <2e-16 ***
## Parttime    -0.365407   0.031570 -11.575  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2452 on 495 degrees of freedom
## Multiple R-squared:  0.704, Adjusted R-squared:  0.7016
## F-statistic: 294.3 on 4 and 495 DF, p-value: < 2.2e-16
```

```
b1=as.matrix(model$coeff[1:2])
b2=as.matrix(model$coeff[3:5])
b1+P%*%b2
```

```
##           [,1]
## (Intercept)  4.7336234
## Female      -0.2505962
```

Now, we should be able to replicate this value if we regress only using the Female and intercept! If we do that :

```
x1_model <- lm(LogWage~Female, data=new_logwage)
x1_model$coeff
```

```
## (Intercept)      Female
##    4.7336234   -0.2505962
```

We can see that the values match exactly!