

Assignment 2.1

Pradyot Prakash - 130050008 Utkarsh Mall - 130050037
Samarth Mishra - 130260018

February 11, 2017

A myIntegration()

Chosen step size $\Delta s = 1$. If a step size larger than this is chosen then information present in image will be missed as step size is larger than pixel size. Choosing step size smaller than this is computationally expensive and the results of integration will be very similar to that of $\Delta s = 1$.

Bilateral interpolation is being used here. A Nearest Neighbour interpolation will not be a good estimate of integral compared to integral evaluated by bilinear interpolation. In cases like $\Delta s = 0.66$ with integration along an axis, NN interpolation will select alternate pixels with twice, which will not be the case with bilinear interpolation.

B myRadon()




Figure 1: Shepp Logan Phantom Image and its Radon transform

C Parameter Choice

The Radon Transform image with $\Delta s = 1$ and $\Delta s = 0.5$ look smoother than the transform with $\Delta s = 3$. However there is no significant difference in images with $\Delta s = 1$ and $\Delta s = 0.5$. **write reason?**

The Radon Transform along a single θ with $\Delta s = 1$ and $\Delta s = 0.5$ are very similar and are smoother than the transform with $\Delta s = 3$. **write reason?**




Figure 2: Shepp-Logan Phantom Image's Radon transform with $\Delta s = 0.5, 1, 3$ respectively




Figure 3: Shepp-Logan Phantom Image's Radon transform Intesity along $\theta = 0^0$ and $\theta = 90^0$ with $\Delta s = 0.5, 1, 3$ respectively