

Metalimnetic oxygen minimum zones decouple diffusive methane and carbon dioxide fluxes from seasonal turnover in a eutrophic reservoir



R.P. McClure¹, M.E. Lofton¹, K.D. Hamre¹, B.R. Niederlehner¹, Z.M. Munger², S. Chen³, J.P. Doubek¹, M.E. Schreiber², and C.C. Carey¹



¹Department of Biological Sciences, Virginia Tech, ²Department of Geosciences, Virginia Tech, ³Department of Civil and Environmental Engineering, University of Sydney, Australia



Question: How do metalimnetic oxygen minimum zones alter carbon dynamics and surface efflux in a eutrophic reservoir?

Introduction

- As a result of anthropogenic change, oxygen dynamics in some waterbodies are changing, potentially resulting in metalimnetic oxygen minimum zones (OMZs)
- Metalimnetic OMZs may alter the carbon balance of a waterbody by generating reducing conditions that result in methane (CH_4) production in the water column, a process that normally occurs at the sediment-water interface
- In summer 2015, we conducted a whole-ecosystem manipulation of oxygen and thermal conditions to allow metalimnetic OMZs to form in a eutrophic reservoir

Methods


Falling Creek Reservoir (FCR)

$Z_{\max} = 9.3\text{m}$, Area = $1.19 \cdot 10^{-1}\text{ km}^2$, GLEON site; Vinton, VA USA

Thermal stratification and oxygen concentration manipulation



- Thermocline deepened using epilimnetic mixer
- Oxygenated hypolimnion with 95% supersaturated dissolved O₂
- Gas Efflux:**
 $\text{FLUX} = k([\text{GAS}_{\text{surface}}] - [\text{GAS}_{\text{atmosphere}}])$
- k values were calculated using LakeMetabolizer package in R
- Tested 5 separate k models:
Cole Model, Read Model, Soloviev Model, MacIntyre Model, Vachon Model

A Metalimnetic OMZ



What is the fate of elevated metalimnetic CH₄ and hypolimnetic CO₂?


Temperature (A): FCR maintained thermal stratification as the OMZ developed
Dissolved Oxygen (B): An OMZ developed between 3 and 5 meters from 1 Jul - 4 Oct



Dissolved CH₄ (A): 1 month after the OMZ developed, CH₄ concentrations rose sharply
Dissolved CO₂ (B): Hypolimnetic oxygenation increased CO₂ in the hypolimnion

CO₂ and CH₄ diffusive emissions

Regardless the k model, the largest efflux of CH₄ (A) into the atmosphere occurred 20 days before fall turnover during a strong wind event, while the largest CO₂ efflux (B) from the hypolimnion occurred during turnover



Next Steps

- Apply observed FCR data from 2015 to the general lake model (GLM-AED)
- Use GRAPLER to generate different climate scenarios for FCR guided by IPCC predictions for the region, and explore how reservoir GHG dynamics may change in response to climate
- Applying GLM-AED and GRAPLER to freshwater reservoirs is critical to assess how GHG production and emissions in waterbodies across different regions will respond to climate change**

Acknowledgements

Hilary Dugan, Paul Hanson, Paul Gantzer, Alex Gerling, Charlotte Harrell, Spencer Klepatzki, John Little, Madeline Ryan, Nicole Ward, GLEON, SAVI, University of Wisconsin, Virginia Tech Reservoir Group and Virginia Tech Stream Team. NSF PRAGMA 1234983