



Understanding Postpartum Depression

A Silent Struggle

Postpartum depression (PPD) is a complex, often misunderstood mood disorder that affects mothers following childbirth. PPD is a serious mental health condition that can impair a mother's ability to care for herself and her baby. It typically emerges within the first few weeks postpartum but can develop anytime within the first year. The condition manifests as persistent sadness, fatigue, anxiety, irritability, difficulty bonding with the newborn, and sometimes, thoughts of self-harm or harming the baby.

Biological, psychological, and social factors all contribute to PPD. Hormonal fluctuations, sleep deprivation, and physical recovery from childbirth can act as catalysts, especially in women with a history of mood disorders. Social isolation, lack of support, financial stress, and unrealistic expectations of motherhood further compound the risk. In some cultures, stigma surrounding mental illness discourages new mothers from seeking help, leading to delayed diagnosis and worsening symptoms.

Early detection is crucial. Routine postpartum screening during obstetric or paediatric visits can aid in timely intervention. Treatment options include counselling (particularly cognitive-behavioural therapy), support groups, and when necessary, pharmacotherapy with antidepressants that are safe in breastfeeding.

PPD not only affects the mother's well-being but also has long-term consequences on the infant's emotional and cognitive development. Therefore, raising awareness, integrating mental health into maternal care, and fostering supportive environments for new mothers is imperative. Combating postpartum depression is not just a medical task — it is a societal responsibility to nurture both mother and child during this vulnerable period.



Dr. Amrita Singh

MD, Obstetrician and Gynaecologist
(Assistant Professor, AIIMS Nagpur)



Content: Dr. Arun Upadhyay, Dr. Amrita Singh
Design: Isha Parlikar, Dr. Somya Mishra
Know more: arun@iitbhlai.ac.in

