
Towards comparing different radiation methods used in astrophysical codes

in the context of **protoplanetary disks**

Prakruti "Kruti" Sudarshan (PhD, Max Planck Institute for Astronomy)
sudarshan@mpia.de, prakrutisudarshan.github.io


M1, half-moment simulations in this presentation are by David Melon Fuksman
thanks to Alex Ziampras, Stanley Baronett, Thomas Pfeil, Zhaohuan Zhu, Mario Flock

Illuminating the Cosmos, MPIA, November 9 2026




Miotello+2023

Radiation and planet formation scales — drives disk evolution!



van Cappeleveen+ 25, Lawlor+26

Substructured disks in scattered light



Benisty+2023 PPVII Chapter

The radiative transfer equation

Specific intensity of radiation

$$\frac{1}{c}\frac{\partial I_\nu}{\partial t} + \mathbf{n}\cdot\nabla I_\nu = \eta_\nu - \chi_\nu I_\nu$$




time evolution

propagation

Emission and absorption

- radiation propagates in multiple spatial dimensions
- travels in different directions and interacts with matter at different frequencies
- and is coupled with the temperature and density structure of the medium

the computational challenge $I_\nu = I_\nu(x, \mathbf{n}, t)$

Category	Particle	Ray	Fluid	Ray-fluid hybrid
				
Method name	Monte Carlo	Ray-tracing Characteristics	Flux-limited diffusion M1 closure	Variable Eddington tensor Discrete ordinates
Physics				
Accurate direction of radiation?	✓	✓	× ($\tau \lesssim 1$)	✓
Accurate at different optical depths τ?	× ($\tau \gg 1$)	✓	× ($\tau \lesssim 1$)	✓
Time-and-velocity-dependent?	✓	×	✓	✓
Detailed microphysics (e.g. anisotropic scattering)?	✓	×	×	×
Known difficulties	Poisson noise; Slow transport ($\tau \gg 1$)	Ray-effect* ($\tau \ll 1$)	Filling shadows; Merging crossing beams; Colliding radiation fronts	Ray-effect* ($\tau \ll 1$)
Computational cost for a sophisticated grey 3D simulation (1= comparable to hydro, order-of-magnitude estimate)				
Relative cost	×100	×10	×1	×10

Ma et al. 2025

Moment methods: FLD and M1

Radiation intensity (freq. integrated)

$$\frac{1}{c}\frac{\partial I}{\partial t} + \mathbf{n}\cdot\nabla I = \eta - \chi I$$

Moments:

$$E_r = \frac{1}{c}\int I\, d\Omega \qquad \mathbf{F}_r = \int I\,\mathbf{n}\, d\Omega$$

Radiation energy — Radiation flux

$$\frac{\partial E_r}{\partial t} + \nabla\cdot\mathbf{F}_r = \text{source terms}$$

$$\frac{1}{c^2}\frac{\partial \mathbf{F}_r}{\partial t} + \nabla\cdot\mathbf{P}_r = \text{source terms}$$

$$\mathbf{F}_r \propto -\nabla E_r$$ FLD!

Kuiper et al. 2011, Flock et al. 2013, Robinson et al. 2024, Ziampras et al. 2024

Moment methods: FLD and M1

Radiation intensity (freq. integrated)

$$\frac{1}{c}\frac{\partial I}{\partial t} + \mathbf{n}\cdot\nabla I = \eta - \chi I$$

Moments:

$$E_r = \frac{1}{c}\int I\, d\Omega \qquad \mathbf{F}_r = \int I\,\mathbf{n}\, d\Omega$$

Radiation energy — Radiation flux

$$\frac{\partial E_r}{\partial t} + \nabla\cdot\mathbf{F}_r = \text{source terms}$$

$$\frac{1}{c^2}\frac{\partial \mathbf{F}_r}{\partial t} + \nabla\cdot\mathbf{P}_r = \text{source terms}$$

evolves both $E_r, \mathbf{F}_r$ M1!

closure relation for $\mathbf{P}_r$

Melon Fuksman et al. 2021, Muley et al. 2023


Discrete ordinates

$$\frac{1}{c}\frac{\partial I_i}{\partial t} + \mathbf{n}_i \cdot \nabla I_i = \eta - \chi I_i$$

solve for intensity in a finite set of directions $\mathbf{n}_i$

Jiang et al. 2021, Baronett et al. 2026


Radiation Shadow



two diffuse spherical sources of radiation shining on an opaque clump in an optically thin medium

Menon et al. 2022

Crossing beams



FLD: no beam crossing, they smear and diffuse out

M1: can handle a single beam, but they don't cross

Menon et al. 2022, Melon Fuksman et al. 2025

numerical methods are improving! (these are ones from recent years)

Frequency-dependent discrete ordinates (Baronett et al. 2026)

Half-moment closure (Melon Fuksman et al. 2025)

Multi-group FLD (Robinson et al. 2024)


Three-temperature (Muley et al. 2023)

More physics ↑


More cost ↑ ↑ ↑ ↑

Coupling with dust dynamics, coagulation, and additional physics still requires you to work with simpler moments.


Radiation controls a lot of processes in protoplanetary disks.



Zhang et al. 2024




Ziampras et al. 2025




Chrenko et al. 2024

Thermally driven substructures:

Su & Bai 2024, Zhang+24, Ziampras+25






Radiation hydrodynamics is key!


But… we are all using different radiation methods.

Q: what are the differences, in realistic disk setups?

Motivation from my PhD work: thermal wave propagation and shadowing


Thermal instability in the flared-disk geomtry of disks






Benisty et al. 2023 PPVII Chapter

PhD work: thermal wave propagation and shadowing






Ueda et al. 2021 (also Dullemond 2000, Watanabe & Lin 2008)

My PhD work: radiation hydrostatics and dynamics of self-shadowed disks


Optically thick disks, d2g = 10^{-2}



Sudarshan et al. 2026

My PhD work: radiation hydrostatics and dynamics of self-shadowed disks


- hydrostatics: thermal wave instability, inwardly moving thermal waves
- hydrodynamics: steady flared-staircase structure
- Ring and shadow pairs in NIR/IR emission



Hydrodynamics + radiative transfer + wave propagation is complicated: where do the differences lie?

Sudarshan et al. 2026


M1 vs FLD: thermal wave propagation, hydrostatics



- great agreement at the **surface** (Chiang & Goldreich 1997)
- temperature bumps: **lower amplitudes** with **FLD** but overall not treating (**vertical + in-plane**) radiation transport resulted in "exaggerated" wave amplitudes
- Both FLD and M1 show **higher midplane** temperatures compared to MCRT

Sudarshan et al. 2026 (with M1 data from Melon Fuksman & Klahr 2022)


M1 vs FLD: thermal wave propagation, hydrostatics



- great agreement at the **surface** (Chiang & Goldreich 1997)
- temperature bumps: **lower amplitudes** with **FLD** but overall not treating (**vertical + in-plane**) radiation transport resulted in "exaggerated" wave amplitudes
- Both FLD and M1 show **higher midplane** temperatures compared to MCRT

Sudarshan et al. 2026 (with M1 data from Melon Fuksman & Klahr 2022)


Going back to more robust, simpler setups
— a planet gap in a disk: with different methods



van Cappeleveen+ 25, Lawlor+26

Comparison: temperatures around a planet gap


Control: RADMC setup, standard disk with a duffel gap (10 $M_{\rm jup}$, at 10 au)



- The gap region in the mid plane is warmer, than the background disk.

Comparison: temperatures around a planet gap


Control: RADMC setup, standard disk with a duffel gap (10 $M_{\rm jup}$, at 10 au)



- By about 40%. This result is also consistent, with/without scattering in RADMC-3D (scattering_mode=2, anisotropic) for our opacity model.

with moment methods: FLD

Corresponding FLD setup with pyFLD (Ziampras+26), standard disk with a duffel gap (10 $M_{\rm jup}$, at 10 au)




- the atmosphere matches the standard Chiang & Goldreich optically thin solution
- differences at the disk midplane, FLD (and the other moment methods) struggle to capture the warm gap

RADMC vs FLD: temperatures around a planet gap


- The likely story for what FLD misses:
  - re-emission from the gap edge

Heated gap edge → emits IR over wavelengths where opacity is different → radiation leaks through and hence you get a warm gap



Comparison against M1 and half-moment methods


- Gray (1 frequency group) RT with moment methods seem to all show a “colder” gap
- Things start improving as you go towards multi-group methods!



M1 and HM simulations by David Melon Fuksman!

FLD, M1, HM and RADMC3D

- M1 with 3 groups seems to be the best match within the gap, but it worsens for the rest of the disk



We will also be including the Athena++ discrete ordinate method
- thanks to Stanley Baronett and Zhaohuan Zhu!

Outlook

- It can be difficult to disentangle which effects change with different radiation methods; comparisons are essential

- Standard tests are useful, but we can now start comparing realistic “disk” setups between the methods

- First results: moment methods don’t capture the warmer gap temperatures in disks, but this improves if you start including frequency-dependent (multi-group) methods, crucial for temperature-sensitive processes in the disks (for the planet-disk case: torques, RWI, spiral substructures)

- More diagnostics underway; we’re also including more codes: frequency-dependent FLD (cuDisc), discrete ordinates (Athena++)