

Counting finite semirings

J. D. Mitchell and P. Ragavan

July 4, 2025

Abstract

In this short note we count the finite semirings up to isomorphism and up to anti-isomorphism for some small values of n ; for which we utilise the existing library of small semigroups in the GAP [13] package SMALLSEMI [7].

1 Introduction

Enumeration of algebra and combinatorial structures of finite order up to isomorphism is a classical topic. Among the algebraic structures considered are groups [3, 35], rings [4, 11, 21, 37], near-rings [1, 5, 36], semigroups and monoids [8, 9, 12, 14, 15, 18, 26, 31, 32, 33, 34, 38, 41], inverse semigroups and monoids [23, 27, 28, 29, 30], and many more too numerous to mention¹. In this short note we count the number of finite semirings up to isomorphism and up to isomorphism and anti-isomorphism for $n \leq 6$. We also count several special classes of semirings for (slightly) larger values of n .

This short note was initiated by an email from M. Volkov to the first author in February of 2025 asking if it was possible to verify with GAP [13] that the number of ai-semirings up to isomorphism with 4 elements is 866 in the paper [39]; see also [43] where it is shown that there are 61 ai-semirings of order 3. After some initial missteps it was relatively straightforward to verify that this number is correct, by using the library of small semigroups in the GAP [13] package SMALLSEMI [7]. This short note arose out from these first steps. In contrast to groups or rings, where the numbers of non-isomorphic objects of order n is known for relatively large values of n , the number of semigroups of order 11 (up to isomorphism) is not known exactly. Given that 99.4% of the semigroups of order 8 are 3-nilpotent, that the number of 3-nilpotent semigroups of order 11 is approximately 10^{26} [10], this number is likely close to the exact value; see also [20]. Perhaps unsurprisingly, from the perspective of counting up to isomorphism, it seems that semirings have more in common with semigroups than with rings or groups. Roughly speaking, rings and groups are highly structured, providing strong constraints that enable their enumeration. On the other hand, semigroups, and seemingly semirings also, are less structured, more numerous, and consequently harder to enumerate.

We begin with the definition of a semiring; which is a natural generalisation of the notion of a ring.

Definition 1.1 (Semiring). A *semiring* is a set S equipped with two binary operations $+$ and \times such that:

- (a) $(S, +)$ is a commutative semigroup ($(x + y) + z = x + (y + z)$ and $x + y = y + x$ for all $x, y, z \in S$);
- (b) (S, \times) is a semigroup ($x \times (y \times z) = (x \times y) \times z$ for all $x, y, z \in S$); and
- (c) multiplication \times distributes over addition $+$ ($x \times (y + z) = (x \times y) + (x \times z)$ and $(y + z) \times x = (y \times x) + (z \times x)$ for all $x, y, z \in S$).

We note that some authors require that $(S, +)$ is a commutative monoid with (additive) identity denoted 0 where $x \times 0 = 0 \times x = 0$ for all $x \in S$; see, for example, [22, 40]. We do not add this requirement, and refer to such objects as *semirings with zero*. The numbers of semirings with zero are discussed in [42] and in [6]; see also [2] which we will discuss a little further below.

Recall that if (S, \times) and (T, \otimes) are semigroups, then $\phi : S \rightarrow T$ is a (*semigroup*) *homomorphism* if $(x \times y)\phi = (x)\phi \otimes (y)\phi$ for all $x, y \in S$. Note that we write mappings to the right of their arguments and compose them from left to right. If $\phi : S \rightarrow T$ is a semigroup homomorphism and ϕ is bijective, then ϕ is an (*semigroup*) *isomorphism*. If a semigroup isomorphism from a semigroup (S, \times) to itself is called an *automorphism* and the group of all such automorphisms is denoted $\text{Aut}(S, \times)$.

In this short note we are concerned with counting semirings up to isomorphism, and so our next definition is that of an isomorphism.

¹The disparity in the number of references for semigroups and monoids and the other algebraic structures is a consequence of the authors familiarity with the literature for semigroups and monoids, and there are likely many other references that could have been included were it not for us not knowing about them.

Definition 1.2 (Semiring isomorphism). We say that two semirings $(S, +, \times)$ and (S, \oplus, \otimes) are *isomorphic* if there exists a bijection $\phi : S \rightarrow S$ which is simultaneously a semigroup isomorphism from $(S, +)$ to (S, \oplus) and from (S, \times) to (S, \otimes) . We refer to ϕ as a (*semiring*) *isomorphism*.

Since a semiring is comprised of two semigroups, enumerating semirings is equivalent to enumerating those pairs consisting of an additive semigroup $(S, +)$ and a multiplicative semigroup (S, \times) such that \times distributes over $+$. The next theorem indicates which (S, \times) we should consider for each of the additive semigroups $(S, +)$.

We denote the symmetric group on the set S by $\text{Sym}(S)$. If $\sigma \in \text{Sym}(S)$ and $\cdot : S \times S \rightarrow S$ is a binary operation, then we define the binary operation $\cdot^\sigma : S \times S \rightarrow S$ by

$$x \cdot^\sigma y = ((x)\sigma^{-1} \cdot (y)\sigma^{-1})\sigma. \quad (1)$$

It is straightforward to verify that (1) is a (right, group) action of $\text{Sym}(S)$ on the set of all binary operations on S . Clearly if \cdot is associative, then so too is \cdot^σ for every $\sigma \in \text{Sym}(S)$. The group of automorphisms $\text{Aut}(S, \cdot)$ of a semigroup (S, \cdot) coincides with the stabiliser of the operation \cdot under the action of $\text{Sym}(S)$ defined in (1).

Recall that if H and K are subgroups of a group G , then the *double cosets* $H \backslash G / K$ are the sets of the form $\{h g k \mid h \in H, k \in K\}$ for $g \in G$. The next theorem is key to our approach for counting semirings.

Theorem 1.3. *Let $(S, +)$ be a commutative semigroup, let (S, \times) be a semigroup, and let $\sigma, \tau \in \text{Sym}(S)$ be such that $(S, +, \times^\sigma)$ and $(S, +, \times^\tau)$ are semirings. Then $(S, +, \times^\sigma)$ and $(S, +, \times^\tau)$ are isomorphic if and only if σ and τ belong to the same double coset of $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$.*

Proof. (\Rightarrow) Suppose that ϕ is a semiring isomorphism from $(S, +, \times^\sigma)$ to $(S, +, \times^\tau)$. Then $\phi \in \text{Aut}(S, +)$ and $(x \times^\sigma y)\phi = (x)\phi \times^\tau (y)\phi$ for all $x, y \in S$. It follows that

$$((x)\sigma^{-1} \times (y)\sigma^{-1})\sigma\phi\tau^{-1} = (x \times^\sigma y)\phi\tau^{-1} = ((x)\phi \times^\tau (y)\phi)\tau^{-1} = (x)\phi\tau^{-1} \times (y)\phi\tau^{-1}.$$

If we set $\gamma = \sigma\phi\tau^{-1}$, $p = (x)\sigma^{-1}$, and $q = (y)\sigma^{-1}$, then $(p \times q)\gamma = (p)\gamma \times (q)\gamma$ and so $\gamma, \gamma^{-1} \in \text{Aut}(S, \times)$. Rearranging we obtain $\tau = \gamma^{-1}\sigma\phi$. Since $\phi \in \text{Aut}(S, +)$, we conclude that τ and σ lie in the same double coset of $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$.

(\Leftarrow) Suppose that σ and τ are in the same double coset of $\text{Aut}(S, +) \backslash \text{Sym}(S) / \text{Aut}(S, \times)$. Then there exists $\alpha \in \text{Aut}(S, \times)$ and $\beta \in \text{Aut}(S, +)$ such that $\tau = \alpha\sigma\beta$.

We will show that β is a semiring isomorphism from $(S, +, \times^\sigma)$ to $(S, +, \times^\tau)$. Since $\beta \in \text{Aut}(S, +)$, it suffices to show that β is an isomorphism from (S, \times^σ) and (S, \times^τ) :

$$\begin{aligned} (x \times^\sigma y)\beta &= (x\sigma^{-1} \times y\sigma^{-1})\sigma\beta = (x\sigma^{-1} \times y\sigma^{-1})\alpha^{-1}\tau = (x\sigma^{-1}\alpha^{-1} \times y\sigma^{-1}\alpha^{-1})\tau & \alpha^{-1} \in \text{Aut}(S, \times) \\ &= (x\beta\tau^{-1} \times y\beta\tau^{-1})\tau = (x\beta \times^\tau y\beta). & \square \end{aligned}$$

If (S, \times) and (S, \otimes) are semigroups, then (S, \times) is said to be *anti-isomorphic* to (S, \otimes) if there exists a bijection $\phi : S \rightarrow S$ such that $(x \times y)\phi = y \otimes x$ for all $x, y \in S$. The bijection ϕ is referred to as an *anti-isomorphism*. If the operations \times and \otimes coincide, then ϕ is an anti-isomorphism. Clearly the composition of two anti-isomorphisms is an automorphism, and the composition of an anti-isomorphism and an isomorphism is an anti-isomorphism. As such the set of all automorphisms or anti-automorphisms forms a group under composition of functions; we denote this group by $\text{Aut}^*(S, \times)$. Note that $\text{Aut}(S, \times)$ is an index two subgroup of $\text{Aut}^*(S, \times)$.

Similarly, using the obvious analogue of Definition 1.2, we can define anti-isomorphic semirings. It is routine to adapt the proof of Definition 1.3 to prove the following.

Corollary 1.4. *Let $(S, +)$ be a commutative semigroup, let (S, \times) be a semigroup, and let $\sigma, \tau \in \text{Sym}(S)$ be such that $(S, +, \times^\sigma)$ and $(S, +, \times^\tau)$ are semirings. Then $(S, +, \times^\sigma)$ and $(S, +, \times^\tau)$ are isomorphic or anti-isomorphic if and only if σ and τ belong to the same double coset of $\text{Aut}^*(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$.*

2 Tables of results

It follows from Definition 1.3 and Definition 1.4 that to enumerate the semirings on a set S up to isomorphism (or up to isomorphism and anti-isomorphism) it suffices to consider every commutative semigroup $(S, +)$ and semigroup (S, \times) and to compute representatives of the double cosets $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ (or $\text{Aut}^*(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$). The semigroups up to isomorphism and anti-isomorphism are available in SMALLSEMI [7]. Since S is small, it is relatively straightforward to can compute $\text{Aut}(S, \times)$ and $\text{Aut}(S, +)$ (using the SEMIGROUPS [24] package for GAP [13], which reduces

the problem to that of computing the automorphisms of a graph associated to $\text{Aut}(S, \times)$ using BLISS [16, 17]). GAP [13] contains functionality for computing double coset representatives based. This is the approach implemented by the authors of the current paper in the GAP [13] package SEMIRINGS [25] to compute the numbers in this section.

Below are some tables listing the numbers of semirings (Definition 1.1), up to isomorphism, and up to isomorphism or anti-isomorphism, with various properties for some small values of $n \in \mathbb{N}$. As far as we know, many of the numbers in these tables were not previously known. In particular, we are not aware of any results in the literature about the number of semirings up to isomorphism and anti-isomorphism. Some of the numbers in the tables below can be found using [2], although this approach is considerably slower than the approach described here, largely because the precomputed data for small semigroups available in SMALLSEMI [7] does a lot of heavy lifting.

These tables are only a sample of the results that can be obtained using SEMIRINGS [25].

n	up to isomorphism	up to isomorphism or anti-isomorphism
1	1	1
2	10	9
3	132	106
4	2,341	1,713
5	57,427 ²	38,247
6	7,571,579	4,102,358

Table 1: Numbers of semirings (Definition 1.1) with n elements up to isomorphism and up to isomorphism or anti-isomorphism.

n	up to isomorphism				up to isomorphism or anti-isomorphism			
	ai	ai with 0	ai with 1	ai with 0 + 1	ai	ai with 0	ai with 1	ai with 0 + 1
1								
2								
3								
4								
5								
6								
7								
8								

Table 2: Numbers of semirings (Definition 1.1) with n elements up to isomorphism and up to isomorphism or anti-isomorphism.

n	up to isomorphism				up to isomorphism or anti-isomorphism			
	ai	ai with 0	ai with 1	ai with 0 + 1	ai	ai with 0	ai with 1	ai with 0 + 1
1								
2								
3								
4								
5								
6								
7								
8								

Table 3: Numbers of commutative semirings ($x \times y = y \times x$ for all $x, y \in S$) with n elements.

n	up to isomorphism				up to isomorphism or anti-isomorphism			
	ai	ai with 0	ai with 1	ai with 0 + 1	ai	ai with 0	ai with 1	ai with 0 + 1
1	1			1	1			1
2	6			1	5			1
3	61			3	45			3
4	866			20	581			18
5	15,751			149	9,750			125
6	354,409			1,488	205,744			1,150
7	9,908,909			18,554	5,470,437			13,171
8				295,292				116,274

Table 4: Numbers of ai-semirings ($x + x = x$ for all $x \in S$) with n elements.

n	up to isomorphism				up to isomorphism or anti-isomorphism			
	ai	ai with 0	ai with 1	ai with 0 + 1	ai	ai with 0	ai with 1	ai with 0 + 1
1								
2								
3								
4								
5								
6								
7								
8								

Table 5: Numbers of commutative ai-semirings ($x \times y = y \times x$ and $x + x = x$ for all $x, y \in S$) with n elements.

References

- [1] E. Aichinger et al. *SONATA, System of nearrings and their applications, Version 2.9.6*. Refereed GAP package. Dec. 2022. URL: <https://gap-packages.github.io/sonata/>.
- [2] A. Bauer. *alg: Algebraic library for constructive mathematics*. <https://github.com/andrejbauer/alg>. Accessed: 2025-07-03. 2019.
- [3] H. U. BESCHE, B. EICK, and E. A. O'BRIEN. "A millennium project: constructing small groups". In: *International Journal of Algebra and Computation* 12.05 (Oct. 2002), pp. 623–644. ISSN: 1793-6500. DOI: [10.1142/S0218196702001115](https://doi.org/10.1142/S0218196702001115). URL: <http://dx.doi.org/10.1142/S0218196702001115>.
- [4] S. R. Blackburn and K. R. McLean. "The enumeration of finite rings". In: *Journal of the London Mathematical Society* 106.4 (July 2022), pp. 3240–3262. ISSN: 1469-7750. DOI: [10.1112/jlms.12661](https://doi.org/10.1112/jlms.12661). URL: <http://dx.doi.org/10.1112/jlms.12661>.
- [5] C. Chow, M. Janota, and J. Araújo. "Cube-Based Isomorph-Free Finite Model Finding". In: *ECAI 2024*. IOS Press, Oct. 2024. ISBN: 9781643685489. DOI: [10.3233/faia240980](https://doi.org/10.3233/faia240980). URL: <http://dx.doi.org/10.3233/faia240980>.
- [6] A. Contributors. *MathStructures*. [Online; accessed 3-July-2025]. 2025. URL: [%5Curl%7Bhttps://math.chapman.edu/~jipsen/structures/doku.php?id=start&rev=1728801324%7D](https://math.chapman.edu/~jipsen/structures/doku.php?id=start&rev=1728801324%7D).
- [7] A. Distler and J. Mitchell. *Smallsemi, A library of small semigroups, Version 0.7.2*. GAP package. Feb. 2025. URL: <https://gap-packages.github.io/smallsemi/>.
- [8] A. Distler. "Classification and enumeration of finite semigroups". Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported. PhD thesis. University of St Andrews, May 2010. URL: <https://hdl.handle.net/10023/945>.
- [9] A. Distler and T. Kelsey. "The semigroups of order 9 and their automorphism groups". In: *Semigroup Forum* 88.1 (June 2013), pp. 93–112. ISSN: 1432-2137. DOI: [10.1007/s00233-013-9504-9](https://doi.org/10.1007/s00233-013-9504-9). URL: <http://dx.doi.org/10.1007/s00233-013-9504-9>.
- [10] A. Distler and J. D. Mitchell. "The Number of Nilpotent Semigroups of Degree 3". In: *The Electronic Journal of Combinatorics* 19.2 (June 2012). ISSN: 1077-8926. DOI: [10.37236/2441](https://doi.org/10.37236/2441). URL: <http://dx.doi.org/10.37236/2441>.

n	up to isomorphism	up to isomorphism or anti-isomorphism
1	1	1
2	2	2
3	6	6
4	40	38
5	295	262
6	3,246	2,681
7	59,314	43,331

Table 6: Numbers of semirings with one and zero with n elements up to isomorphism, and up to isomorphism or anti-isomorphism.

n	up to isomorphism	up to isomorphism or anti-isomorphism
1 2 3 4 5 6 7 8		

Table 7: Numbers of ai-semirings with one and zero with n elements up to isomorphism and up to isomorphism or anti-isomorphism.

- [11] B. Fine. “Classification of Finite Rings of Order p^2 ”. In: *Mathematics Magazine* 66.4 (Oct. 1993), p. 248. ISSN: 0025-570X. DOI: [10.2307/2690742](https://doi.org/10.2307/2690742). URL: <http://dx.doi.org/10.2307/2690742>.
- [12] G. E. Forsythe. “SWAC Computes 126 Distinct Semigroups of Order 4”. In: *Proceedings of the American Mathematical Society* 6.3 (June 1955), p. 443. ISSN: 0002-9939. DOI: [10.2307/2032786](https://doi.org/10.2307/2032786). URL: <http://dx.doi.org/10.2307/2032786>.
- [13] *GAP – Groups, Algorithms, and Programming, Version 4.14.0*. The GAP Group. 2024. URL: <https://www.gap-system.org>.
- [14] P. A. Grillet. “Computing finite commutative semigroups”. In: *Semigroup Forum* 53.1 (Dec. 1996), pp. 140–154. ISSN: 1432-2137. DOI: [10.1007/bf02574129](https://doi.org/10.1007/bf02574129). URL: <http://dx.doi.org/10.1007/BF02574129>.
- [15] P. A. Grillet. “Counting Semigroups”. In: *Communications in Algebra* 43.2 (Aug. 2014), pp. 574–596. ISSN: 1532-4125. DOI: [10.1080/00927872.2013.790036](https://doi.org/10.1080/00927872.2013.790036). URL: <http://dx.doi.org/10.1080/00927872.2013.790036>.
- [16] T. Junttila and P. Kaski. *bliss: A Tool for Computing Automorphism Groups and Canonical Labelings of Graphs*. Version as of July 2025. 2007. URL: <https://users.aalto.fi/~tjunttil/bliss/>.
- [17] T. Junttila and P. Kaski. “Engineering an efficient canonical labeling tool for large and sparse graphs”. In: *Proceedings of the Ninth Workshop on Algorithm Engineering and Experiments (ALENEX)* (2007), pp. 135–149. DOI: [10.1137/1.9781611972870.13](https://doi.org/10.1137/1.9781611972870.13). URL: <https://epubs.siam.org/doi/10.1137/1.9781611972870.13>.
- [18] H. Jürgensen, P. Wick, and V. von Št. Schwarz. “Die halbgruppen der ordnungen ≤ 7 ”. In: *Semigroup Forum* 14.1 (Dec. 1977), pp. 69–79. ISSN: 1432-2137. DOI: [10.1007/bf02194655](https://doi.org/10.1007/bf02194655). URL: <http://dx.doi.org/10.1007/BF02194655>.
- [19] V. L. Klee Jr. “The November meeting in Los Angeles”. In: *Bull. Amer. Math. Soc.* 64.2 (1958), p. 56. ISSN: 0002-9904. DOI: [10.1090/S0002-9904-1958-10159-7](https://doi.org/10.1090/S0002-9904-1958-10159-7). URL: <https://doi.org/10.1090/S0002-9904-1958-10159-7>.
- [20] D. J. Kleitman, B. R. Rothschild, and J. H. Spencer. “The Number of Semigroups of Order n ”. In: *Proceedings of the American Mathematical Society* 55.1 (Feb. 1976), p. 227. ISSN: 0002-9939. DOI: [10.2307/2041879](https://doi.org/10.2307/2041879). URL: <http://dx.doi.org/10.2307/2041879>.
- [21] R. L. Kruse and D. T. Price. “Enumerating Finite Rings”. In: *Journal of the London Mathematical Society* s2-2.1 (Jan. 1970), pp. 149–159. ISSN: 0024-6107. DOI: [10.1112/jlms/s2-2.1.149](https://doi.org/10.1112/jlms/s2-2.1.149). URL: <http://dx.doi.org/10.1112/jlms/s2-2.1.149>.
- [22] M. Lothaire. *Applied Combinatorics on Words*. Cambridge University Press, July 2005. ISBN: 9781107341005. DOI: [10.1017/cbo9781107341005](https://doi.org/10.1017/cbo9781107341005). URL: <http://dx.doi.org/10.1017/CBO9781107341005>.
- [23] M. E. Malandro. “Enumeration of finite inverse semigroups”. In: *Semigroup Forum* 99.3 (Sept. 2019), pp. 679–723. ISSN: 1432-2137. DOI: [10.1007/s00233-019-10054-9](https://doi.org/10.1007/s00233-019-10054-9). URL: <http://dx.doi.org/10.1007/s00233-019-10054-9>.

n	up to isomorphism	up to isomorphism or anti-isomorphism
1	1	1
2	4	4
3	22	21
4	169	155
5	1,819	1,561
6	41,104	30,112

Table 8: Numbers of semirings with one (unital semirings) with n elements up to isomorphism and up to isomorphism or anti-isomorphism.

- [24] J. D. Mitchell et al. *Semigroups - GAP package, Version 5.5.1*. June 2025. DOI: [10.5281/zenodo.592893](https://doi.org/10.5281/zenodo.592893). URL: <http://dx.doi.org/10.5281/zenodo.592893>.
- [25] J. D. Mitchell and P. Ragavan. *Semirings, Counting and enumerating semirings Version 0.2.0*. GAP package. July 2025. URL: [%7Dhttps://gap-packages.github.io/semirings/%7D](https://gap-packages.github.io/semirings/%7D).
- [26] T. S. Motzkin and J. L. Selfridge. “Semigroups of Order Five”. In: Presented in [19].
- [27] OEIS Foundation Inc. *Number of commutative inverse monoids of order n* . Entry A234845 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234845>.
- [28] OEIS Foundation Inc. *Number of commutative inverse semigroups of order n* . Entry A234843 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234843>.
- [29] OEIS Foundation Inc. *Number of inverse monoids of order n* . Entry A234844 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234844>.
- [30] OEIS Foundation Inc. *Number of inverse semigroups of order n , considered to be equivalent when they are isomorphic or anti-isomorphic (by reversal of the operator)*, Entry A001428 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001428>.
- [31] OEIS Foundation Inc. *Number of nonisomorphic monoids (semigroups with identity) of order n* , Entry A058129 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A058129>.
- [32] OEIS Foundation Inc. *Number of nonisomorphic semigroups of order n* , Entry A027851 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A027851>.
- [33] OEIS Foundation Inc. *Number of semigroups of order n , considered to be equivalent when they are isomorphic or anti-isomorphic (by reversal of the operator)*, Entry A001423 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001423>.
- [34] OEIS Foundation Inc. *The number of commutative semigroups of order n* , Entry A001426 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001426>.
- [35] OEIS Foundation Inc. *The number of groups of order n* , Entry A000001 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A000001>.
- [36] OEIS Foundation Inc. *The number of near-rings with n elements*, Entry A305858 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A305858>.
- [37] OEIS Foundation Inc. *The number of rings with n elements*, Entry A027623 in *The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A027623>.
- [38] R. J. Plemmons. “There are 15973 semigroups of order 6”. In: *Math. Algorithms* 2 (1967), pp. 2–17. ISSN: 0025-5548.
- [39] M. Ren et al. “The finite basis problem for additively idempotent semirings of order four, I”. In: *Semigroup Forum* 110.2 (Apr. 2025), pp. 422–457. ISSN: 1432-2137. DOI: [10.1007/s00233-025-10520-7](https://doi.org/10.1007/s00233-025-10520-7). URL: <http://dx.doi.org/10.1007/s00233-025-10520-7>.
- [40] J. Sakarovitch. *Elements of Automata Theory*. Ed. by R. Thomas. Cambridge University Press, Oct. 2009. ISBN: 9781139195218. DOI: [10.1017/cbo9781139195218](https://doi.org/10.1017/cbo9781139195218). URL: <http://dx.doi.org/10.1017/CB09781139195218>.
- [41] S. Satoh, K. Yama, and M. Tokizawa. “Semigroups of order 8”. In: *Semigroup Forum* 49.1 (Dec. 1994), pp. 7–29. ISSN: 1432-2137. DOI: [10.1007/bf02573467](https://doi.org/10.1007/bf02573467). URL: <http://dx.doi.org/10.1007/BF02573467>.

- [42] *Semirings of small orders*. <https://math.stackexchange.com/questions/3961014/semirings-of-small-orders>. Viewed Apr 20, 2025; answer by Alex J Best on Dec 25, 2020. 2020.
- [43] X. Zhao et al. “The variety generated by an ai-semiring of order three”. In: *Ural Mathematical Journal* 6.2 (Dec. 2020), p. 117. ISSN: 2414-3952. DOI: [10.15826/umj.2020.2.012](https://doi.org/10.15826/umj.2020.2.012). URL: <http://dx.doi.org/10.15826/umj.2020.2.012>.