

# Counting finite semirings

J. Edwards, J. D. Mitchell, and P. Ragavan

June 19, 2026

## Abstract

In this short note we count the finite semirings up to isomorphism, and up to isomorphism or anti-isomorphism for some small values of  $n$ ; for which we utilise the existing library of small semigroups in the GAP [19] package SMALLSEMI [11].

## 1 Introduction

Enumeration of algebraic and combinatorial structures of finite order up to isomorphism is a classical topic. Among the algebraic structures considered are groups [4, 42], rings [5, 17, 28, 44], near-rings [1, 6, 43], semigroups and monoids [12, 13, 15, 18, 20, 21, 25, 26, 33, 38, 39, 40, 41, 45, 48], inverse semigroups and monoids [30, 34, 35, 36, 37], and many more too numerous to mention<sup>1</sup>. In this short note we count the number of finite semirings up to isomorphism and up to isomorphism or anti-isomorphism for  $n \leq 6$ ; see Tables 1 to 4. We also count several special classes of semirings for (slightly) larger values of  $n$ .

This short note was initiated by an email from M. Volkov to the second author in February of 2025 asking if it was possible to verify the claim in [46] that the number of ai-semirings (additively idempotent) up to isomorphism with 4 elements is 866 (see the caption of Table 3 for the definition). After some initial missteps it was relatively straightforward to verify that this number is correct, by using the library of small semigroups in the GAP [19] package SMALLSEMI [11]. This short note arose from these first steps. In contrast to groups or rings, where the numbers of non-isomorphic objects of order  $n$  are known for relatively large values of  $n$ , the number of semigroups of order 10 (up to isomorphism) is apparently not known exactly (although the number up to isomorphism or anti-isomorphism is known [15]). The paper [27] purports to show that almost all semigroups of order  $n$  (as  $n$  tends to infinity) up to isomorphism or anti-isomorphism are 3-nilpotent (and indeed 99.4% of the semigroups of order 8 are 3-nilpotent), and it is shown in [14] that the number of 3-nilpotent semigroups of order 10 is approximately  $10^{19}$ . This number is likely close to the exact value. Perhaps unsurprisingly, from the perspective of counting up to isomorphism, it seems that semirings have more in common with semigroups than with rings or groups. Very roughly speaking, rings and groups are highly structured, providing strong constraints that facilitate their enumeration. On the other hand, semigroups, and seemingly semirings also, are less structured, more numerous, and harder to enumerate.

We begin with the definition of a semiring, which is a natural generalisation of the notion of a ring.

**Definition 1.1** (Semiring). A *semiring* is a set  $S$  equipped with two binary operations  $+$  and  $\times$  such that:

- (a)  $(S, +)$  is a commutative semigroup ( $(x + y) + z = x + (y + z)$  and  $x + y = y + x$  for all  $x, y, z \in S$ );
- (b)  $(S, \times)$  is a semigroup ( $x \times (y \times z) = (x \times y) \times z$  for all  $x, y, z \in S$ ); and
- (c) multiplication  $\times$  distributes over addition  $+$  ( $x \times (y + z) = (x \times y) + (x \times z)$  and  $(y + z) \times x = (y \times x) + (z \times x)$  for all  $x, y, z \in S$ ).

We note that some authors require that  $(S, +)$  is a commutative monoid with (additive) identity 0 such that  $0 \times x = x \times 0 = 0$  for all  $x \in S$ ; see, for example, [29, 47]. We do not add this requirement, and refer to such objects as *semirings with zero*. The numbers of semirings with zero are discussed in [49] and in [7]; see also ALG [2] which we will discuss a little further below. An *ai-semiring* is a semiring  $S$  where  $x + x = x$  for all  $x \in S$ . In [50] it was shown that there are 61 ai-semirings of order 3.

Recall that if  $(S, \times)$  and  $(T, \otimes)$  are semigroups, then  $\phi : S \rightarrow T$  is a (*semigroup*) *homomorphism* if  $(x \times y)\phi = (x)\phi \otimes (y)\phi$  for all  $x, y \in S$ . Note that we write mappings to the right of their arguments and compose them from left to right. If  $\phi : S \rightarrow T$  is a semigroup homomorphism and  $\phi$  is bijective, then  $\phi$  is an (*semigroup*) *isomorphism*. A semigroup

<sup>1</sup>The disparity in the number of references for semigroups and monoids and the other algebraic structures is a consequence of the authors' familiarity with the literature for semigroups and monoids, and there are likely many other references that could have been included were it not for us not knowing about them.

isomorphism from a semigroup  $(S, \times)$  to itself is called an *automorphism* and the group of all such automorphisms is denoted  $\text{Aut}(S, \times)$ .

In this short note we are concerned with counting semirings up to isomorphism, and so our next definition is that of an isomorphism.

**Definition 1.2** (Semiring isomorphism). We say that two semirings  $(S, +, \times)$  and  $(T, \oplus, \otimes)$  are *isomorphic* if there exists a bijection  $\phi : S \rightarrow T$  which is simultaneously a semigroup isomorphism from  $(S, +)$  to  $(T, \oplus)$  and from  $(S, \times)$  to  $(T, \otimes)$ . We refer to  $\phi$  as a (*semiring*) *isomorphism*.

Since a semiring is comprised of two semigroups, enumerating semirings is equivalent to enumerating those pairs consisting of an additive semigroup  $(S, +)$  and a multiplicative semigroup  $(S, \times)$  such that  $\times$  distributes over  $+$ . The next theorem indicates which  $(S, \times)$  we should consider for each of the additive semigroups  $(S, +)$ .

We denote the symmetric group on the set  $S$  by  $\text{Sym}(S)$ . If  $\sigma \in \text{Sym}(S)$  and  $\cdot : S \times S \rightarrow S$  is a binary operation, then we define the binary operation  $\cdot^\sigma : S \times S \rightarrow S$  by

$$x \cdot^\sigma y = ((x)\sigma^{-1} \cdot (y)\sigma^{-1})\sigma. \quad (1)$$

It is straightforward to verify that (1) is a (right, group) action of  $\text{Sym}(S)$  on the set of all binary operations on  $S$ . Clearly if  $\cdot$  is associative, then so too is  $\cdot^\sigma$  for every  $\sigma \in \text{Sym}(S)$ . The group of automorphisms  $\text{Aut}(S, \cdot)$  of a semigroup  $(S, \cdot)$  coincides with the stabiliser of the operation  $\cdot$  under the action of  $\text{Sym}(S)$  defined in (1).

Recall that if  $H$  and  $K$  are subgroups of a group  $G$ , then the *double cosets*  $H \backslash G / K$  are the sets of the form  $\{h g k \mid h \in H, k \in K\}$  for  $g \in G$ . The next theorem is key to our approach for counting semirings.

**Theorem 1.3.** Let  $(S, +)$  be a commutative semigroup, let  $(S, \times)$  be a semigroup, and let  $\sigma, \tau \in \text{Sym}(S)$ . Then there exists a semigroup isomorphism  $\phi \in \text{Aut}(S, +)$  from  $(S, \times^\sigma)$  to  $(S, \times^\tau)$  if and only if  $\sigma$  and  $\tau$  belong to the same double coset in  $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ .

*Proof.* ( $\Rightarrow$ ) Suppose that  $\phi \in \text{Aut}(S, +)$  is a semigroup isomorphism from  $(S, \times^\sigma)$  to  $(S, \times^\tau)$ . Then  $(x \times^\sigma y)\phi = (x)\phi \times^\tau (y)\phi$  for all  $x, y \in S$ . If  $x, y \in S$  are arbitrary, then

$$((x)\sigma^{-1} \times (y)\sigma^{-1})\sigma\phi\tau^{-1} = (x \times^\sigma y)\phi\tau^{-1} = ((x)\phi \times^\tau (y)\phi)\tau^{-1} = ((x)\phi\tau^{-1} \times (y)\phi\tau^{-1})\tau\tau^{-1} = (x)\phi\tau^{-1} \times (y)\phi\tau^{-1}.$$

If we set  $\gamma = \sigma\phi\tau^{-1}$ ,  $p = (x)\sigma^{-1}$ , and  $q = (y)\sigma^{-1}$ , then  $(p \times q)\gamma = (p)\gamma \times (q)\gamma$  and so  $\gamma, \gamma^{-1} \in \text{Aut}(S, \times)$ . Rearranging we obtain  $\tau = \gamma^{-1}\sigma\phi$ . Since  $\phi \in \text{Aut}(S, +)$ , we conclude that  $\tau$  and  $\sigma$  lie in the same double coset of  $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ .

( $\Leftarrow$ ) Suppose that  $\sigma$  and  $\tau$  are in the same double coset of  $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ . Then there exists  $\alpha \in \text{Aut}(S, \times)$  and  $\beta \in \text{Aut}(S, +)$  such that  $\tau = \alpha\sigma\beta$ .

We will show that  $\beta$  is a semigroup isomorphism from  $(S, \times^\sigma)$  to  $(S, \times^\tau)$ :

$$\begin{aligned} (x \times^\sigma y)\beta &= (x\sigma^{-1} \times y\sigma^{-1})\sigma\beta = (x\sigma^{-1} \times y\sigma^{-1})\alpha^{-1}\tau = (x\sigma^{-1}\alpha^{-1} \times y\sigma^{-1}\alpha^{-1})\tau & \alpha^{-1} \in \text{Aut}(S, \times) \\ &= (x\beta\tau^{-1} \times y\beta\tau^{-1})\tau = (x\beta \times^\tau y\beta). & \square \end{aligned}$$

An immediate corollary of Theorem 1.3 is the following.

**Corollary 1.4.** Let  $(S, +)$  be a commutative semigroup, let  $(S, \times)$  be a semigroup, and let  $\sigma$  and  $\tau$  belong to the same double coset in  $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ . Then the following hold:

- (a)  $(S, +, \times^\sigma)$  is a semiring if and only if  $(S, +, \times^\tau)$  is a semiring;
- (b) if  $(S, +, \times^\sigma)$  and  $(S, +, \times^\tau)$  are semirings, then they are isomorphic.

If  $(S, \times)$  and  $(S, \otimes)$  are semigroups, then  $(S, \times)$  is said to be *anti-isomorphic* to  $(S, \otimes)$  if there exists a bijection  $\phi : S \rightarrow S$  such that  $(x \times y)\phi = (y)\phi \otimes (x)\phi$  for all  $x, y \in S$ . The bijection  $\phi$  is referred to as an *anti-isomorphism*. If the operations  $\times$  and  $\otimes$  coincide, then  $\phi$  is an *anti-automorphism*. Clearly the composition of two anti-automorphisms is an automorphism, and the composition of an anti-automorphism and an automorphism is an anti-automorphism. As such the set of all automorphisms or anti-automorphisms forms a group under composition of functions; we denote this group by  $\text{Aut}^*(S, \times)$ .

Similarly, using the obvious analogue of Definition 1.2, we can define anti-isomorphic semirings. It is routine to adapt the proof of Theorem 1.3 to prove the following.

**Theorem 1.5.** Let  $(S, +)$  be a commutative semigroup, let  $(S, \times)$  be a semigroup, and let  $\sigma, \tau \in \text{Sym}(S)$ . Then there exists a semigroup isomorphism or anti-isomorphism  $\phi \in \text{Aut}(S, +)$  from  $(S, \times^\sigma)$  to  $(S, \times^\tau)$  if and only if  $\sigma$  and  $\tau$  belong to the same double coset in  $\text{Aut}^*(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ .

We also obtain a corollary of [Theorem 1.5](#) analogous to [Corollary 1.4](#).

**Corollary 1.6.** *Let  $(S, +)$  be a commutative semigroup, let  $(S, \times)$  be a semigroup, and let  $\sigma$  and  $\tau$  belong to the same double coset in  $\text{Aut}^*(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ . Then the following hold:*

- (a)  *$(S, +, \times^\sigma)$  is a semiring if and only if  $(S, +, \times^\tau)$  is a semiring;*
- (b) *if  $(S, +, \times^\sigma)$  and  $(S, +, \times^\tau)$  are semirings, then they are isomorphic or anti-isomorphic.*

## 2 Tables of results

The approach we implemented in the GAP [19] package SEMIRINGS [16] to compute the numbers in this section is as follows. It follows from [Corollaries 1.4](#) and [1.6](#) that to enumerate the semirings on a set  $S$  up to isomorphism (or up to isomorphism or anti-isomorphism) it suffices to consider every commutative semigroup  $(S, +)$  and semigroup  $(S, \times)$  and to check distributivity of  $+$  and  $\times^\sigma$  for one representative  $\sigma$  of every double coset in  $\text{Aut}(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$  (or in  $\text{Aut}^*(S, \times) \backslash \text{Sym}(S) / \text{Aut}(S, +)$ ). The semigroups with order at most 8 up to isomorphism and anti-isomorphism are available in SMALLSEMI [11]. Since  $S$  is small, it is relatively straightforward to compute  $\text{Aut}(S, \times)$  and  $\text{Aut}(S, +)$ : the problem of computing  $\text{Aut}(S, \times)$  reduces to that of computing the automorphism group of a graph associated to  $(S, \times)$ ; see [31] for details. The automorphisms of  $(S, +)$  can also be computed in this way. There are several software tools for computing automorphism groups of graphs, for example, BLISS [23, 24]. This is the approach used by the SEMIGROUPS [32] package for GAP [19], which the authors used to compute  $\text{Aut}(S, \times)$  and  $\text{Aut}(S, +)$  in this paper. Finally, GAP [19] contains functionality for computing double coset representatives; see [22, Section 4.6.8] for details.

Below are some tables listing the numbers of semirings ([Definition 1.1](#)) up to isomorphism, and up to isomorphism or anti-isomorphism, with various properties for some small values of  $n \in \mathbb{N}$ . As far as we know, many of the numbers in these tables were not previously known. In particular, we are not aware of any results in the literature about the number of semirings up to isomorphism or anti-isomorphism. Some of the numbers in the tables below can be found using ALG [2], although using ALG [2] is considerably slower than the approach described here, largely because the precomputed data for small semigroups available in SMALLSEMI [11] does a lot of the heavy lifting.

As a sanity check, where possible, we have checked the numbers produced by SEMIRINGS [16] against those in the literature, and those we could produce using ALG [2]. With one exception, these numbers agreed with each other. ALG [2] computes that there are 57,443 semirings (using [Definition 1.1](#)) of order 5 up to isomorphism, whereas SEMIRINGS [16] computes 57,427. Using GAP [19] and SEMIRINGS [16] we attempted to verify whether or not the output of ALG [2] was correct. Unfortunately, it appears that 16 of the pairs of multiplication tables output by ALG [2] do not satisfy one or the other of the distributivity conditions from [Definition 1.1\(c\)](#); see [3] for more details. This precisely accounts for the difference between the number output by SEMIRINGS [16] and the number output by ALG [2].

The numbers present in the tables below were computed using a variety of desktop and laptop computers with the following specs: a 2025 MacBook Pro M4 Pro with 14 threads and 48GB of RAM; an Intel(R) Core(TM) i7-8700 CPU @ 3.20GHz processor with 12 threads and 64GB of RAM; an Intel(R) Core(TM) i7-12700KF @ 5GHz with 20 threads and 64GB of RAM. We mostly include this information to indicate that we did not make use of particularly powerful computers. The longest of the computation in the tables below took approximately 2400 CPU hours. This was the time taken to count the semirings with 0 up to isomorphism or anti-isomorphism with 7 elements.

These tables are only a sample of the results that can be obtained using SEMIRINGS [16].

| $n$ | up to isomorphism         |           |            |            | up to isomorphism or anti-isomorphism |           |           |            |
|-----|---------------------------|-----------|------------|------------|---------------------------------------|-----------|-----------|------------|
|     | no additional constraints | with 0    | with 1     | with 0 + 1 | no additional constraints             | with 0    | with 1    | with 0 + 1 |
| 1   | 1                         | 1         | 1          | 1          | 1                                     | 1         | 1         | 1          |
| 2   | 10                        | 4         | 4          | 2          | 9                                     | 4         | 4         | 2          |
| 3   | 132                       | 22        | 22         | 6          | 106                                   | 20        | 21        | 6          |
| 4   | 2,341                     | 283       | 169        | 40         | 1,713                                 | 226       | 155       | 38         |
| 5   | 57,427                    | 4,717     | 1,819      | 295        | 38,247                                | 3,365     | 1,561     | 262        |
| 6   | 7,571,579                 | 108,992   | 41,104     | 3,246      | 4,102,358                             | 71,138    | 30,112    | 2,681      |
| 7   | -                         | 8,925,672 | 11,679,328 | 59,314     | 48,152,448,707                        | 4,910,824 | 6,268,858 | 43,331     |

Table 1: Numbers of semirings ([Definition 1.1](#)) with  $n$  elements up to isomorphism and up to isomorphism or anti-isomorphism. See [8, 9, 10] for some of these numbers up to isomorphism.

| $n$ | up to isomorphism         |         |         |            |
|-----|---------------------------|---------|---------|------------|
|     | no additional constraints | with 0  | with 1  | with 0 + 1 |
| 1   | 1                         | 1       | 1       | 1          |
| 2   | 8                         | 4       | 4       | 2          |
| 3   | 80                        | 18      | 20      | 6          |
| 4   | 1,067                     | 169     | 141     | 36         |
| 5   | 18,188                    | 1,990   | 1,276   | 228        |
| 6   | 543,458                   | 32,212  | 17,621  | 2,075      |
| 7   | 162,744,745               | 799,354 | 690,924 | 25,640     |
| 8   | -                         | -       | -       | 791,061    |

Table 2: Numbers of commutative semirings (i.e. those satisfying  $x \times y = y \times x$  for all  $x, y \in S$ ) with  $n$  elements.

| $n$ | up to isomorphism         |         |         |            | up to isomorphism or anti-isomorphism |         |         |            |
|-----|---------------------------|---------|---------|------------|---------------------------------------|---------|---------|------------|
|     | no additional constraints | with 0  | with 1  | with 0 + 1 | no additional constraints             | with 0  | with 1  | with 0 + 1 |
| 1   | 1                         | 1       | 1       | 1          | 1                                     | 1       | 1       | 1          |
| 2   | 6                         | 2       | 2       | 1          | 5                                     | 2       | 2       | 1          |
| 3   | 61                        | 12      | 11      | 3          | 45                                    | 10      | 10      | 3          |
| 4   | 866                       | 129     | 73      | 20         | 581                                   | 93      | 64      | 18         |
| 5   | 15,751                    | 1,852   | 703     | 149        | 9,750                                 | 1,207   | 574     | 125        |
| 6   | 354,409                   | 33,391  | 9,195   | 1,488      | 205,744                               | 20,142  | 6,835   | 1,150      |
| 7   | 9,908,909                 | 729,629 | 164,163 | 18,554     | 5,470,437                             | 415,527 | 109,880 | 13,171     |
| 8   | -                         | -       | -       | 295,292    | -                                     | -       | -       | 116,274    |

Table 3: Numbers of ai-semirings (i.e. those satisfying  $x + x = x$  for all  $x \in S$ , where “ai” stands for “additively idempotent”) with  $n$  elements.

## Acknowledgements

The authors would like to thank the anonymous referees for their helpful comments.

## References

- [1] E. Aichinger et al. *SONATA, System of nearrings and their applications, Version 2.9.6*. Refereed GAP package. Dec. 2022. URL: <https://gap-packages.github.io/sonata/>.
- [2] A. Bauer. *alg: Algebraic library for constructive mathematics*. <https://github.com/andrejbauer/alg>. Accessed: 2025-07-03. 2019.
- [3] A. Bauer. *Issue #16: Incorrect results*. <https://github.com/andrejbauer/alg/issues/16>. GitHub issue, opened Jul 4, 2025. 2025.
- [4] H. U. Besche, B. Eick, and E. A. O’Brien. “A millennium project: constructing small groups”. In: *International Journal of Algebra and Computation* 12.05 (Oct. 2002), pp. 623–644. ISSN: 1793-6500. DOI: [10.1142/S0218196702001115](https://doi.org/10.1142/S0218196702001115). URL: <http://dx.doi.org/10.1142/S0218196702001115>.
- [5] S. R. Blackburn and K. R. McLean. “The enumeration of finite rings”. In: *Journal of the London Mathematical Society* 106.4 (July 2022), pp. 3240–3262. ISSN: 1469-7750. DOI: [10.1112/jlms.12661](https://doi.org/10.1112/jlms.12661). URL: <http://dx.doi.org/10.1112/jlms.12661>.
- [6] C. Chow, M. Janota, and J. Araújo. “Cube-Based Isomorph-Free Finite Model Finding”. In: *ECAI 2024*. IOS Press, Oct. 2024. ISBN: 9781643685489. DOI: [10.3233/faia240980](https://doi.org/10.3233/faia240980). URL: <http://dx.doi.org/10.3233/FAIA240980>.
- [7] A. Contributors. *MathStructures*. [Online; accessed 3-July-2025]. 2025. URL: <https://math.chapman.edu/~jipsen/structures/doku.php?id=start&rev=1728801324>.
- [8] A. Contributors. *semirings — MathStructures*. [Online; accessed 4-July-2025]. 2021. URL: <https://math.chapman.edu/~jipsen/structures/doku.php?id=semirings&rev=1614028296>.

| $n$ | up to isomorphism         |           |         |            |
|-----|---------------------------|-----------|---------|------------|
|     | no additional constraints | with 0    | with 1  | with 0 + 1 |
| 1   | 1                         | 1         | 1       | 1          |
| 2   | 4                         | 2         | 2       | 1          |
| 3   | 29                        | 8         | 9       | 3          |
| 4   | 289                       | 57        | 55      | 16         |
| 5   | 3,589                     | 550       | 437     | 100        |
| 6   | 53,661                    | 6,639     | 4,296   | 794        |
| 7   | 949,843                   | 96,264    | 52,043  | 7,493      |
| 8   | 20,054,643                | 1,639,905 | 764,329 | 84,961     |

Table 4: Numbers of commutative ai-semirings (i.e. those satisfying  $x \times y = y \times x$  and  $x + x = x$  for all  $x, y \in S$ ) with  $n$  elements.

- [9] A. Contributors. *semirings\_with\_identity* — *MathStructures*. [Online; accessed 4-July-2025]. 2021. URL: [https://math.chapman.edu/~jipsen/structures/doku.php?id=semirings\\_with\\_identity&rev=1614028296](https://math.chapman.edu/~jipsen/structures/doku.php?id=semirings_with_identity&rev=1614028296).
- [10] A. Contributors. *semirings\_with\_identity\_and\_zero* — *MathStructures*. [Online; accessed 4-July-2025]. 2021. URL: [https://math.chapman.edu/~jipsen/structures/doku.php?id=semirings\\_with\\_identity\\_and\\_zero&rev=1614028296](https://math.chapman.edu/~jipsen/structures/doku.php?id=semirings_with_identity_and_zero&rev=1614028296).
- [11] A. Distler and J. Mitchell. *Smallsemi, A library of small semigroups, Version 0.7.2*. GAP package. Feb. 2025. URL: <https://gap-packages.github.io/smallsemi/>.
- [12] A. Distler. “Classification and enumeration of finite semigroups”. CC BY-NC-SA 3.0. PhD thesis. University of St Andrews, May 2010. URL: <https://hdl.handle.net/10023/945>.
- [13] A. Distler and T. Kelsey. “The semigroups of order 9 and their automorphism groups”. In: *Semigroup Forum* 88.1 (June 2013), pp. 93–112. ISSN: 1432-2137. DOI: [10.1007/s00233-013-9504-9](https://doi.org/10.1007/s00233-013-9504-9). URL: <http://dx.doi.org/10.1007/s00233-013-9504-9>.
- [14] A. Distler and J. D. Mitchell. “The Number of Nilpotent Semigroups of Degree 3”. In: *The Electronic Journal of Combinatorics* 19.2 (June 2012). ISSN: 1077-8926. DOI: [10.37236/2441](https://doi.org/10.37236/2441). URL: <http://dx.doi.org/10.37236/2441>.
- [15] A. Distler et al. “The Semigroups of Order 10”. In: *Principles and Practice of Constraint Programming*. Springer Berlin Heidelberg, 2012, pp. 883–899. ISBN: 9783642335587. DOI: [10.1007/978-3-642-33558-7\\_63](https://doi.org/10.1007/978-3-642-33558-7_63). URL: [http://dx.doi.org/10.1007/978-3-642-33558-7\\_63](http://dx.doi.org/10.1007/978-3-642-33558-7_63).
- [16] J. Edwards, J. D. Mitchell, and P. Ragavan. *Semirings, Counting and enumerating semirings Version 0.2.0*. GAP package. July 2025. URL: <https://github.com/pramothagavan/semirings>.
- [17] B. Fine. “Classification of Finite Rings of Order  $p^2$ ”. In: *Mathematics Magazine* 66.4 (Oct. 1993), p. 248. ISSN: 0025-570X. DOI: [10.2307/2690742](https://doi.org/10.2307/2690742). URL: <http://dx.doi.org/10.2307/2690742>.
- [18] G. E. Forsythe. “SWAC Computes 126 Distinct Semigroups of Order 4”. In: *Proceedings of the American Mathematical Society* 6.3 (June 1955), p. 443. ISSN: 0002-9939. DOI: [10.2307/2032786](https://doi.org/10.2307/2032786). URL: <http://dx.doi.org/10.2307/2032786>.
- [19] *GAP – Groups, Algorithms, and Programming, Version 4.14.0*. The GAP Group. 2024. URL: <https://www.gap-system.org>.
- [20] P. A. Grillet. “Computing finite commutative semigroups”. In: *Semigroup Forum* 53.1 (Dec. 1996), pp. 140–154. ISSN: 1432-2137. DOI: [10.1007/bf02574129](https://doi.org/10.1007/bf02574129). URL: <http://dx.doi.org/10.1007/BF02574129>.
- [21] P. A. Grillet. “Counting Semigroups”. In: *Communications in Algebra* 43.2 (Aug. 2014), pp. 574–596. ISSN: 1532-4125. DOI: [10.1080/00927872.2013.790036](https://doi.org/10.1080/00927872.2013.790036). URL: <http://dx.doi.org/10.1080/00927872.2013.790036>.
- [22] D. F. Holt, B. Eick, and E. A. O’Brien. *Handbook of Computational Group Theory*. Chapman and Hall/CRC, Jan. 2005. ISBN: 9780429147944. DOI: [10.1201/9781420035216](https://doi.org/10.1201/9781420035216). URL: <http://dx.doi.org/10.1201/9781420035216>.
- [23] T. Junttila and P. Kaski. *bliss: A Tool for Computing Automorphism Groups and Canonical Labelings of Graphs*. Version as of July 2025. 2007. URL: <https://users.aalto.fi/~tjunttil/bliss/>.
- [24] T. Junttila and P. Kaski. “Engineering an efficient canonical labeling tool for large and sparse graphs”. In: *Proceedings of the Ninth Workshop on Algorithm Engineering and Experiments (ALENEX)* (2007), pp. 135–149. DOI: [10.1137/1.9781611972870.13](https://doi.org/10.1137/1.9781611972870.13). URL: <https://epubs.siam.org/doi/10.1137/1.9781611972870.13>.



- [25] H. Jürgensen, P. Wick, and V. von Št. Schwarz. “Die halbgruppen der ordnungen  $\leq 7$ ”. In: *Semigroup Forum* 14.1 (Dec. 1977), pp. 69–79. ISSN: 1432-2137. DOI: [10.1007/bf02194655](https://doi.org/10.1007/bf02194655). URL: <http://dx.doi.org/10.1007/BF02194655>.
- [26] V. L. Klee Jr. “The November meeting in Los Angeles”. In: *Bull. Amer. Math. Soc.* 64.2 (1958), p. 56. ISSN: 0002-9904. DOI: [10.1090/S0002-9904-1958-10159-7](https://doi.org/10.1090/S0002-9904-1958-10159-7). URL: <https://doi.org/10.1090/S0002-9904-1958-10159-7>.
- [27] D. J. Kleitman, B. R. Rothschild, and J. H. Spencer. “The Number of Semigroups of Order  $n$ ”. In: *Proceedings of the American Mathematical Society* 55.1 (Feb. 1976), p. 227. ISSN: 0002-9939. DOI: [10.2307/2041879](https://doi.org/10.2307/2041879). URL: <http://dx.doi.org/10.2307/2041879>.
- [28] R. L. Kruse and D. T. Price. “Enumerating Finite Rings”. In: *Journal of the London Mathematical Society* s2-2.1 (Jan. 1970), pp. 149–159. ISSN: 0024-6107. DOI: [10.1112/jlms/s2-2.1.149](https://doi.org/10.1112/jlms/s2-2.1.149). URL: <http://dx.doi.org/10.1112/jlms/s2-2.1.149>.
- [29] M. Lothaire. *Applied Combinatorics on Words*. Cambridge University Press, July 2005. ISBN: 9781107341005. DOI: [10.1017/cbo9781107341005](https://doi.org/10.1017/cbo9781107341005). URL: <http://dx.doi.org/10.1017/CB09781107341005>.
- [30] M. E. Malandro. “Enumeration of finite inverse semigroups”. In: *Semigroup Forum* 99.3 (Sept. 2019), pp. 679–723. ISSN: 1432-2137. DOI: [10.1007/s00233-019-10054-9](https://doi.org/10.1007/s00233-019-10054-9). URL: <http://dx.doi.org/10.1007/s00233-019-10054-9>.
- [31] G. L. Miller. “Graph isomorphism, general remarks”. In: *Proceedings of the ninth annual ACM symposium on Theory of computing - STOC '77*. STOC '77. ACM Press, 1977, pp. 143–150. DOI: [10.1145/800105.803404](https://doi.org/10.1145/800105.803404). URL: <http://dx.doi.org/10.1145/800105.803404>.
- [32] J. D. Mitchell et al. *Semigroups - GAP package, Version 5.5.1*. June 2025. DOI: [10.5281/zenodo.592893](https://doi.org/10.5281/zenodo.592893). URL: <http://dx.doi.org/10.5281/zenodo.592893>.
- [33] T. S. Motzkin and J. L. Selfridge. “Semigroups of Order Five”. In: Presented in [26].
- [34] OEIS Foundation Inc. *Number of commutative inverse monoids of order  $n$ . Entry A234845 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234845>.
- [35] OEIS Foundation Inc. *Number of commutative inverse semigroups of order  $n$ . Entry A234843 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234843>.
- [36] OEIS Foundation Inc. *Number of inverse monoids of order  $n$ . Entry A234844 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A234844>.
- [37] OEIS Foundation Inc. *Number of inverse semigroups of order  $n$ , considered to be equivalent when they are isomorphic or anti-isomorphic (by reversal of the operator), Entry A001428 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001428>.
- [38] OEIS Foundation Inc. *Number of nonisomorphic monoids (semigroups with identity) of order  $n$ , Entry A058129 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A058129>.
- [39] OEIS Foundation Inc. *Number of nonisomorphic semigroups of order  $n$ , Entry A027851 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A027851>.
- [40] OEIS Foundation Inc. *Number of semigroups of order  $n$ , considered to be equivalent when they are isomorphic or anti-isomorphic (by reversal of the operator), Entry A001423 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001423>.
- [41] OEIS Foundation Inc. *The number of commutative semigroups of order  $n$ , Entry A001426 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A001426>.
- [42] OEIS Foundation Inc. *The number of groups of order  $n$ , Entry A000001 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A000001>.
- [43] OEIS Foundation Inc. *The number of near-rings with  $n$  elements, Entry A305858 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A305858>.
- [44] OEIS Foundation Inc. *The number of rings with  $n$  elements, Entry A027623 in The On-Line Encyclopedia of Integer Sequences*. 2025. URL: <https://oeis.org/A027623>.
- [45] R. J. Plemmons. “There are 15973 semigroups of order 6”. In: *Math. Algorithms* 2 (1967), pp. 2–17. ISSN: 0025-5548.
- [46] M. Ren et al. “The finite basis problem for additively idempotent semirings of order four, I”. In: *Semigroup Forum* 110.2 (Apr. 2025), pp. 422–457. ISSN: 1432-2137. DOI: [10.1007/s00233-025-10520-7](https://doi.org/10.1007/s00233-025-10520-7). URL: <http://dx.doi.org/10.1007/s00233-025-10520-7>.

- [47] J. Sakarovitch. *Elements of Automata Theory*. Ed. by R. Thomas. Cambridge University Press, Oct. 2009. ISBN: 9781139195218. DOI: [10.1017/cbo9781139195218](https://doi.org/10.1017/cbo9781139195218). URL: <http://dx.doi.org/10.1017/CB09781139195218>.
- [48] S. Satoh, K. Yama, and M. Tokizawa. “Semigroups of order 8”. In: *Semigroup Forum* 49.1 (Dec. 1994), pp. 7–29. ISSN: 1432-2137. DOI: [10.1007/bf02573467](https://doi.org/10.1007/bf02573467). URL: <http://dx.doi.org/10.1007/BF02573467>.
- [49] *Semirings of small orders*. <https://math.stackexchange.com/questions/3961014/semirings-of-small-orders>. Viewed Apr 20, 2025; answer by Alex J Best on Dec 25, 2020. 2020.
- [50] X. Zhao et al. “The variety generated by an ai-semiring of order three”. In: *Ural Mathematical Journal* 6.2 (Dec. 2020), p. 117. ISSN: 2414-3952. DOI: [10.15826/umj.2020.2.012](https://doi.org/10.15826/umj.2020.2.012). URL: <http://dx.doi.org/10.15826/umj.2020.2.012>.