

School of Computing
National University of Singapore
CS5340: Uncertainty Modeling in AI
Semester 1, AY 2018/19

Exercise 1

Question 1

a)




Fig. 1.1

Fig. 1.1 shows a mobile robot that traverses from pose X_t to X_{t+1} over time t to $t + 1$. The robot is equipped with an 1-dimensional range sensor that returns the distances Z_t and Z_{t+1} of a landmark structure L in the environment from the poses X_t and X_{t+1} respectively. Let U_t denotes the control command given by the user to move the robot from X_t to X_{t+1} .

- (i) Taking $\{U_t, L, X_t, X_{t+1}, Z_t, Z_{t+1}\}$ as random variables, state whether each of these random variables is an observed or latent/hidden random variable. Explain your answers.
- (ii) Given the following conditional independencies:

$$L \perp U_t \mid \emptyset, \quad X_t \perp L \mid U_t, \quad X_{t+1} \perp \{L, U_t\} \mid X_t, \\ Z_t \perp \{U_t, X_{t+1}\} \mid \{X_t, L\}, \quad Z_{t+1} \perp \{U_t, X_t, Z_t\} \mid \{L, X_{t+1}\}.$$

Write the factorized probability and draw the Bayesian network that represents the joint distribution $p(u_t, l, x_t, x_{t+1}, z_t, z_{t+1})$ assuming the following topological ordering of the random variables:

$$\{U_t, L, X_t, X_{t+1}, Z_t, Z_{t+1}\}.$$

Show all your workings clearly.

(iii) Write the following probability distribution $p(z_t, z_{t+1} | l)$ in terms of the factorized probability obtained in (ii). Simplify your answer.

b)





Fig. 1.2

For each of the Bayesian networks shown in Fig. 1.2, determine the largest set of nodes X_B such that $X_1 \perp X_B \mid X_2$. Explain your answers.

Question 2

Consider the graph shown in Fig. 2.1:




Fig 2.1

- a) What is the corresponding moral graph?
- b) What is the reconstituted graph from the UNDIRECTEDGRAPHELIMINATE algorithm on the moral graph with the ordering $\{8,7,6,5,4,3,2,1\}$?
- c) What is the reconstituted graph from the UNDIRECTEDGRAPHELIMINATE algorithm on the moral graph with the ordering $\{8,5,6,7,4,3,2,1\}$?
- d) Suppose you wish to calculate $p(x_1|x_8)$. Which ordering is preferable? Why?

Question 3

What is the treewidth of the graph below?




Fig 3.1

Question 4

Consider the following random variables. X_1 and X_2 represent the outcomes of two independent fair coin tosses. X_3 is the indicator function of the event that the outcomes are identical.

- Specify a directed graphical model that describes the joint probability distribution (i.e. specify the graph and the conditional distributions).
- Specify an undirected graphical model that describes the joint probability distribution (i.e. give the graph and specify the clique potentials).
- In both cases, list all conditional independencies that are implied by the graph.
- In both cases, list any additional conditional dependencies that are displayed by this probability distribution but are not implied by the graph.

Question 5




Fig. 5.1

The graphical model shown above describes some relationships among variables associated with chest abnormality. Answer the following questions based on the graphical model.

- True or False. Justify your choice. $Smoke \perp Dyspnea \mid Bronchitis$.
- True or False. Justify your choice. $Bronchitis \perp X-ray \mid Cancer$.
- True or False. Justify your choice. $Smoke \perp Risky\ Visit \mid Dyspnea$.
- True or False. Justify your choice. $X-ray \perp Smoke \mid \{Cancer, Bronchitis\}$.

Question 6

Evaluate (give the distribution tables) the following probabilities:

$$p(x_1 | x_5), \quad p(x_2 | x_4), \quad p(x_3 | x_2), \quad p(x_4 | x_3), \quad p(x_5)$$

for the Bayesian network shown in Fig. 6.1, where each random variable takes a binary state, i.e. $x_i \in \{T, F\}$. Show all your workings clearly.




Fig. 6.1

Question 7

Give the junction tree of the Bayesian network shown in Fig. 7.1 using the following elimination order: $\{X_7, X_6, X_5, X_4, X_3, X_2, X_1\}$. Show all your workings clearly.




Fig. 7.1

--End--