GEORGIA DEPARTMENT OF COMMUNITY HEALTH

GEORGIA MEDICAID FEE-FOR-SERVICE
HEPATITIS C AGENTS PA SUMMARY

Preferred	Non-Preferred
Direct Inhibitors	
Mavyret (glecaprevir/pibrentasvir) Sofosbuvir/velpatasvir 400/100 mg generic Vosevi (sofosbuvir/velpatasvir/voxilaprevir)	Epclusa Paks (sofosbuvir/velpatasvir 200/50, 150/37.5 mg) Harvoni 45/200 mg and Paks (ledipasvir/sofosbuvir) Ledipasvir/sofosbuvir 90/400 mg generic Sovaldi (sofosbuvir) Zepatier (elbasvir/grazoprevir)
Ribavirin	
Ribavirin generic	n/a
Pegylated Interferons	
Pegasys (peginterferon alfa-2a)	

LENGTH OF AUTHORIZATION: Varies

PA CRITERIA:

Epclusa Paks

- Approvable for members 3 years of age or older with a diagnosis of chronic hepatitis C virus infection (CHC, HCV) who are treatment naïve, have been previously treated with a regimen containing peginterferon and ribavirin or have been previously treated with sofosbuvir (Sovaldi)- or NS5A inhibitor–based regimen.
- Members with decompensated cirrhosis must take in combination with ribavirin when clinically appropriate.

Harvoni 45/200 mg and Paks and Ledipasvir/Sofosbuvir 90/400 mg Generic

- Approvable for members who are treatment naïve, have been previously treated with a peginterferon-based regimen or have been previously treated with sofosbuvir (Sovaldi)- or NS5A inhibitor–based regimen.
- Members 18 years of age or older must have a diagnosis of chronic hepatitis C virus infection (CHC, HCV) and
  - genotype 1 (1a or 1b), 4, 5 or 6 without liver transplant and without cirrhosis or with compensated cirrhosis and must be unable to take sofosbuvir/velpatasvir and Mavyret.
  - genotype 1 (1a or 1b), 4, 5 or 6 without liver transplant with decompensated cirrhosis and must be unable to take sofosbuvir/velpatasvir.
  - genotype 1 (1a or 1b), 4, 5 or 6 with liver transplant without cirrhosis or with compensated cirrhosis and must be unable to take sofosbuvir/velpatasvir and Mavyret.

- Members 3 to 17 years of age must have a diagnosis of genotype 1 (1a or 1b), 4, 5 or 6 chronic HCV infection without cirrhosis or with compensated cirrhosis and must be unable to take sofosbuvir/velpatasvir and Mavyret.
- Members must take Harvoni or ledipasvir/sofosbuvir in combination with ribavirin when clinically appropriate.

Sovaldi

- Approvable for members who are treatment naïve, have been previously treated with an interferon-based regimen or have been previously treated with glecaprevir/pibrentasvir (Mavyret), sofosbuvir/velpatasvir/voxilaprevir (Vosevi) or sofosbuvir (Sovaldi) + glecaprevir/pibrentasvir (Mavyret).
- Members 18 years of age or older must have a diagnosis of chronic hepatitis C virus infection (CHC, HCV) without cirrhosis or with compensated cirrhosis and
    - genotype 1 (1a or 1b) and must be unable to take sofosbuvir/velpatasvir and Mavyret. Members must take Sovaldi with peginterferon/ribavirin or ribavirin.
    - genotype 2 or 3 and must be unable to take sofosbuvir/velpatasvir and Mavyret. Members must take Sovaldi with ribavirin.
    - genotype 3 with compensated cirrhosis and have previously failed treatment with peginterferon/ribavirin and must be unable to take Vosevi. Members must take Sovaldi with Zepatier.
    - genotype 4 and must be unable to take sofosbuvir/velpatasvir and Mavyret. Members must take Sovaldi with peginterferon/ribavirin.
    - have hepatocellular carcinoma awaiting liver transplantation. Members must take Sovaldi with ribavirin.
- Members 3 to 17 years of age must have a diagnosis of genotype 2 or 3 chronic HCV infection without cirrhosis or with compensated cirrhosis and must be unable to take sofosbuvir/velpatasvir and Mavyret.
- Members must take Sovaldi in combination with ribavirin when clinically appropriate.

Zepatier

- Approvable for members who are treatment naïve, have been previously treated with peginterferon and ribavirin or have been previously treated with peginterferon, ribavirin and protease inhibitor.
- Members must be 12 years of age or older who weigh 30 kg or more with a diagnosis of genotype 3 chronic hepatitis C virus infection (CHC, HCV) with compensated cirrhosis and have previously failed treatment with peginterferon/ribavirin and must be unable to take Vosevi. Members must take Zepatier with Sovaldi.
- Members must be 12 years of age or older who weigh 30 kg or more with a diagnosis of genotype 1 (1a or 1b) or genotype 4 chronic hepatitis C virus infection (CHC, HCV) and members with genotype 1a must have NS5A resistance testing conducted and must be unable to take sofosbuvir/velpatasvir and Mavyret. Members must take Zepatier in combination with ribavirin if
    - genotype 1a with baseline NS5A polymorphisms.
    - genotype 4 and previous treatment failure with peginterferon/ribavirin regimen.
- Members must take Zepatier in combination with ribavirin when clinically appropriate.

EXCEPTIONS:

- Exceptions to these conditions of coverage are considered through the prior authorization process.
- The Prior Authorization process may be initiated by calling **OptumRx at 1-866-525-5827.**

PREFERRED DRUG LIST:

- For online access to the Preferred Drug List (PDL), please go to http://dch.georgia.gov/preferred-drug-lists.

PA AND APPEAL PROCESS:

- For online access to the PA process, please go to www.dch.georgia.gov/prior-authorization-process-and-criteria and click on Prior Authorization (PA) Request Process Guide.

QUANTITY LEVEL LIMITATIONS:

- For online access to the current Quantity Level Limits (QLL), please go to www.mmis.georgia.gov/portal, highlight Pharmacy and click on Other Documents, then select the most recent quarters QLL list.