AETNA BETTER HEALTH®
Coverage Policy/Guideline

| | | | |
|---|---|---|---|
| Name: | Cibinqo (abrocitinib) | Page: | 1 of 5 |
| Effective Date: | 6/1/2026 | Last Review Date: | 3/25/2026 |
| Applies to: | ☒Illinois | | |

## Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Cibinqo under the patient's prescription drug benefit.

## Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met, and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication
Cibinqo is indicated for the treatment of adults and pediatric patients 12 years of age and older with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is inadvisable.

All other indications are considered experimental/investigational and not medically necessary.

## Applicable Drug List:

Cibinqo

## Policy/Guideline:

**Documentation for all indications**:
The patient is unable to take Dupixent, where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.

**Documentation**

Submission of the following information is necessary to initiate the prior authorization review:

Initial requests
- Chart notes or medical record documentation showing affected area(s) and body surface area.
- Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy (where applicable).

Continuation requests
Chart notes or medical record documentation supporting positive clinical response.

**Prescriber Specialties**
This medication must be prescribed by or in consultation with a dermatologist or allergist/immunologist.

**Coverage Criteria**

Atopic Dermatitis
Authorization of 4 months may be granted for treatment of moderate-to-severe atopic dermatitis in members 12 years of age or older when ALL the following criteria are met:

- Affected body surface is greater than or equal to 10% body surface area OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
- Member meets EITHER of the following:
  - Member has had an inadequate treatment response with ONE of the following:
    - A medium potency to super-high potency topical corticosteroid (see Appendix)
    - A topical calcineurin inhibitor (e.g., pimecrolimus, tacrolimus)
    - A topical Janus kinase (JAK) inhibitor (e.g., Opzelura)
    - A topical phosphodiesterase-4 (PDE-4) inhibitor (e.g., Eucrisa, Zoryve)
    - A topical aryl hydrocarbon receptor agonist (e.g., Vtama)
  - The use of medium potency to super-high potency topical corticosteroid, topical calcineurin inhibitor, topical JAK inhibitor, topical PDE-4 inhibitor, and topical aryl hydrocarbon receptor agonist are not advisable for the member (e.g., due to contraindications, prior intolerances).
- Member has had an inadequate response, intolerance or contraindication to treatment with other systemic drug products, including biologics (e.g., Adbry, Dupixent, Ebglyss, Nemluvio, Rinvoq) indicated for the treatment of atopic dermatitis in the past 12 months.

**Continuation of Therapy**
Authorization of 12 months may be granted for members 12 years of age or older (including new members) who are using the requested medication for moderate-to-severe atopic dermatitis when the member has achieved or maintained a positive clinical response as evidenced by low disease activity (i.e., clear or almost clear skin), or improvement in signs and symptoms of atopic dermatitis (e.g., redness, itching, oozing/crusting).

**Other**
Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members

with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

Member cannot use the requested medication concomitantly with any other biologic drug, targeted synthetic drug, or potent immunosuppressant such as azathioprine or cyclosporine.

## Approval Duration and Quantity Restrictions:

**Approval:**

- **Initial Duration:** 4 months
- **Renewal Duration:** 12 months

**Quantity Level Limit:** 30 tablets per 30 days

**Appendix**:

**Table. Relative potency of select topical corticosteroid products**

| Potency | Drug | Dosage form | Strength |
|---|---|---|---|
| I. Super-high potency (group 1) | Augmented betamethasone dipropionate | Ointment, Lotion, Gel | 0.05% |
| | Clobetasol propionate | Cream, Gel, Ointment, Solution, Cream (emollient), Lotion, Shampoo, Foam, Spray | 0.05% |
| | Fluocinonide | Cream | 0.1% |
| | Flurandrenolide | Tape | 4 mcg/cm$^2$ |
| | Halobetasol propionate | Cream, Lotion, Ointment, Foam | 0.05% |
| II. High potency (group 2) | Amcinonide | Ointment | 0.1% |
| | Augmented betamethasone dipropionate | Cream | 0.05% |
| | Betamethasone dipropionate | Ointment | 0.05% |
| | Clobetasol propionate | Cream | 0.025% |
| | Desoximetasone | Cream, Ointment, Spray | 0.25% |
| | | Gel | 0.05% |
| | Diflorasone diacetate | Ointment, Cream (emollient) | 0.05% |
| | Fluocinonide | Cream, Ointment, Gel, Solution | 0.05% |
| | Halcinonide | Cream, Ointment | 0.1% |
| | Halobetasol propionate | Lotion | 0.01% |
| **Potency** | **Drug** | **Dosage form** | **Strength** |
| III. High potency (group 3) | Amcinonide | Cream, Lotion | 0.1% |
| | Betamethasone dipropionate | Cream, hydrophilic emollient | 0.05% |
| | Betamethasone valerate | Ointment | 0.1% |
| | | Foam | 0.12% |
| | Desoximetasone | Cream, Ointment | 0.05% |
| | Diflorasone diacetate | Cream | 0.05% |

| Potency | Drug | Dosage form | Strength |
|---|---|---|---|
| | Fluocinonide | Cream, aqueous emollient | 0.05% |
| | Fluticasone propionate | Ointment | 0.005% |
| | Mometasone furoate | Ointment | 0.1% |
| | Triamcinolone acetonide | Cream, Ointment | 0.5% |
| IV. Medium potency (group 4) | Betamethasone dipropionate | Spray | 0.05% |
| | Clocortolone pivalate | Cream | 0.1% |
| | Fluocinolone acetonide | Ointment | 0.025% |
| | Flurandrenolide | Ointment | 0.05% |
| | Hydrocortisone valerate | Ointment | 0.2% |
| | Mometasone furoate | Cream, Lotion, Solution | 0.1% |
| | Triamcinolone acetonide | Cream | 0.1% |
| | | Ointment | 0.05% and 0.1% |
| | | Aerosol Spray | 0.2 mg per 2-second spray |
| V. Lower-mid potency (group 5) | Betamethasone dipropionate | Lotion | 0.05% |
| | Betamethasone valerate | Cream | 0.1% |
| | Desonide | Ointment, Gel | 0.05% |
| | Fluocinolone acetonide | Cream | 0.025% |
| | Flurandrenolide | Cream, Lotion | 0.05% |
| | Fluticasone propionate | Cream, Lotion | 0.05% |
| | Hydrocortisone butyrate | Cream, Lotion, Ointment, Solution | 0.1% |
| | Hydrocortisone probutate | Cream | 0.1% |
| | Hydrocortisone valerate | Cream | 0.2% |
| | Prednicarbate | Cream (emollient), Ointment | 0.1% |
| | Triamcinolone acetonide | Lotion | 0.1% |
| | | Ointment | 0.025% |
| VI. Low potency (group 6) | Alclometasone dipropionate | Cream, Ointment | 0.05% |
| | Betamethasone valerate | Lotion | 0.1% |
| | Desonide | Cream, Lotion, Foam | 0.05% |
| | Fluocinolone acetonide | Cream, Solution, Shampoo, Oil | 0.01% |
| | Triamcinolone acetonide | Cream, lotion | 0.025% |
| VII. Least potent (group 7) | Hydrocortisone (base, greater than or equal to 2%) | Cream, Ointment, Solution | 2.5% |
| | | Lotion | 2% |
| | Hydrocortisone (base, less than 2%) | Cream, Ointment, Gel, Lotion, Spray, Solution | 1% |
| | | Cream, Ointment | 0.5% |
| | Hydrocortisone acetate | Cream | 2.5% |

| Potency | Drug | Dosage form | Strength |
|---|---|---|---|
| | | Lotion | 2% |
| | | Cream | 1% |

## References:

1. Cibinqo [package insert]. New York, NY: Pfizer Inc.; December 2023.
2. Eichenfield LF, Tom WL, Chamlin SL, et. al. Guidelines of care for the management of atopic dermatitis: Section 1. Diagnosis and assessment of atopic dermatitis. J Am Acad Dermatol. 2014;70:338-51.
3. Sidbury R, Alikhan A, Bercovitch L, et. al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. J Am Acad Dermatol. 2023;89(1):e1-e20.
4. Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. J Am Acad Dermatol. 2024;90(2):e43-e56.
5. Davis DMR, Frazer-Green L, Alikhan A, et al. Focused update: Guidelines of care for the management of atopic dermatitis in adults. J Am Acad Dermatol. 2025;93(3):745.e1-745.e7.
6. Sidbury R, Tom WL, Bergman JN, et al. Guidelines of care for the management of atopic dermatitis: Section 4. Prevention of disease flares and use of adjunctive therapies and approaches. J Am Acad Dermatol. 2014;71:1218-1233.
7. Comparison of Representative Topical Corticosteroid Preparations (Classified According to the United States System). Drug Facts and Comparisons. Facts and Comparisons. UpToDate Inc. https://fco.factsandcomparisons.com. Accessed November 7, 2025.
8. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on November 7, 2025 from: https://www.cdc.gov/tb/testing/index.html.