Clinical Criteria, Step Therapy, and Quantity Limits for TennCare Preferred Drug List (PDL)

November 2, 2026

To view the full **Preferred Drug List (PDL)** and other Program Information, please visit the Optum Rx/TennCare **website**.

With the exception of the "**Branded Drugs to be Classified as Generics**" list, TennCare is a mandatory generic program in accordance with state law (TCA 53-10-205). Approval of a branded product when a generic is available requires documentation of a serious adverse reaction from the generic via an FDA MedWatch form **OR** contraindication to an inactive ingredient in the AB-rated generic equivalent. Therapeutic Failure of an AB-rated generic equivalent may be considered for approval of branded products in the following high-risk medication classes: Anticonvulsants, Atypical Antipsychotics, HIV antivirals, Immunosuppressants, and Oncology Agents.

Faxed PA submissions require the use of the **General PA Form**, unless a drug/class specific form is noted for the drug/drug class.

PA approval duration length is one (1) year unless otherwise indicated.

Please note the following:

- All agents must be prescribed by a provider with a Tennessee Medicaid Provider ID.
- Approval of Non-Preferred (NP) agents requires trial and failure, contraindication, or intolerance of two (2) preferred agents, unless otherwise indicated. This is the Standard NP Criteria.
- Agents not listed on the PDL may be considered non-preferred unless otherwise indicated.
- Agents that must be administered by a healthcare provider or in a healthcare facility may by covered under medical benefits. Providers should refer to the patient's Managed Care Organization (MCO) for coverage information. Please refer to the Covered Injectable Drugs list for exceptions.

Abbreviations

Abbreviation	Definition
B	Budgetary Reduction edit for utilization control
I/DD	Intellectual and Developmental Disabilities
N/A	Not applicable
NP	Non-Preferred Agent
P	Preferred Agent
PA	Prior Authorization
PAC	Pharmacy Advisory Committee
PDL	Preferred Drug List
QL	Quantity Limit

Abbreviation	Definition
Caps	Capsules
Conc	Concentrate
Cr	Cream
Inh	Inhalation
Inj	Injection
Oint	Ointment
Soln	Solution
Supp	Suppository
Susp	Suspension
Tabs	Tablets

I. ANALGESICS

Agents for Dependency

Lucemyra® tablets

PDL Status	P
Quantity Limit	16/day; 1 treatment course (up to 14 days) per 6-months
PA Criteria	See below
PA Form	General PA Form

Lucemyra® tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescriber attestation that patient will be monitored if at risk for QT interval prolongation, CV effects, or CNS effects; **AND**
- Patient has tried and failed, had a contraindication to, or experienced an adverse reaction/intolerance to clonidine; **AND**
- Prescriber attestation of a comprehensive treatment plan between provider and patient; **AND**
- In the case of opioid use disorder (OUD), provide verbal attestation that patient:
- Has a referral to **OR** active involvement in substance abuse counseling; **OR**
- Is unable to have counseling **AND** provides verbal attestation that patient has been offered medication-assisted treatment (MAT) as part of a comprehensive treatment plan; **AND**

Lucemyra® tablets Renewal PA Criteria:

- If request documents that member has been started and stabilized on Lucemyra®, continuation of therapy will be allowed to complete a total duration not to exceed 14 days

Vivitrol® Injection

PDL Status	P
Quantity Limit	1 vial per 28 days
PA Criteria	None
PA Form	General PA Form

Lofexidine tablets

PDL Status	NP
Quantity Limit	16/day; 1 treatment course (up to 14 days) per 6-months
PA Criteria	See below
PA Form	General PA Form

Lofexidine tablets PA Criteria:

- Clinically valid reason why Lucemyra® cannot be used

Buprenorphine and Buprenorphine/Naloxone

Please submit faxed PA requests using the Buprenorphine Products PA Form.

Buprenorphine Enhanced Supportive Medication-Assisted Recovery and Treatment (BESMART) Network Providers Only

BESMART Quantity Limit Criteria for Requests > 32 mg/day

- Provider is a MD or DO; **AND**
- The patient is currently taking 32 mg/day or has been titrated to the requested dose without evidence of over-medication side effects (e.g., sedation, cognitive fog) (confirmed via paid claims or submitted chart notes); **AND**
- Submission of medical records (e.g. chart notes) documenting ONE of the following:
  - Patient demonstrates current active substance use, as evidenced by history, recent hospitalization or relapse, and/or recent drug screening results (PA approval duration: 6 months); **OR**
  - The patient is pregnant (must provide estimated due date) (PA approval duration: 3 months past due date); **OR**
  - The patient is less than 12 months postpartum from delivery date (must provide delivery date) (PA approval duration: 12 months from delivery date); **OR**
  - The patient is tolerating 32 mg/day but continues to experience significant cravings, withdrawal symptoms, or both, which impairs daily functioning (PA approval duration: 6 months); **AND**
- Provider has submitted clinical literature to support efficacy of the requested dose above 32 mg

Note: Nurse Practitioner (NP), Physician Assistant (PA), and other mid-level prescribers in the BESMART network are limited to max daily dose (MDD) 16mg/day per TN state law.

Please submit faxed PA requests using the Buprenorphine Products PA Form.

Buprenorphine/naloxone tablets

PDL Status	P
Quantity Limit	8/2 mg: 4/day for MD/DO prescribers, 2/day for NP/PA prescribers*; 2/0.5 mg: 3 /day ^
PA Criteria	None
PA Form	Buprenorphine Products PA Form
Notes	*NP/PA prescribers are limited to 16 mg of buprenorphine per day. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine/naloxone film

PDL Status	P
Quantity Limit	12/3 mg: 2/day for MD/DO prescribers, 1/day for NP/PA prescribers*; 8/2 mg: 4 /day for MD/DO prescribers, 2/day for NP/PA prescribers*; 4/1 mg: 2/day; 2/0.5 mg: 3/day ^
PA Criteria	None
PA Form	Buprenorphine Products PA Form
Notes	*NP/PA prescribers are limited to 16 mg of buprenorphine per day. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine tablets

PDL Status	NP
Quantity Limit	8 mg: 4 /day for MD/DO prescribers, 2/day for NP/PA prescribers*; 2 mg: 3 /day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Pregnancy: 3 months past due date Breastfeeding: 6 months (maximum 4 approvals) Contraindication to naloxone: initial authorization: 6 months; reauthorization: 12 months
Notes	The PA for buprenorphine monotherapy sublingual tablets in pregnant or breastfeeding patients diagnosed with opioid use disorder can be bypassed using Professional Pharmacy Service (PPS) Codes. Please refer to Provider Notice Buprenorphine Updates 06.01.25 posted on the Optum Rx/TennCare website for details. Mild rash, itching, and GI intolerance are not accepted as intolerance to naloxone. *NP/PA prescribers are limited to 16 mg of buprenorphine per day. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine tablets PA Criteria:

- Diagnosis of opiate addiction; **AND**
- Prescriber is enrolled and in good standing in the BESMART program; **AND**
- Buprenorphine will not be approved for treatment of depression or pain; **AND**
- ONE of the following:
  - Patient is actively pregnant (must provide estimated due date)
  - Patient is actively breastfeeding (must provide delivery date)
  - Request is for a two-day induction for patients transitioning off methadone
  - Patient is unable to take naloxone containing products due to a contraindication, drug to drug interaction, or history of toxic side effects that caused immediate or long-term damage (DOCUMENTATION REQUIRED). Note: Mild rash, itching, and GI intolerance are not accepted as intolerance to naloxone.

Suboxone® films

PDL Status	NP
Quantity Limit	12/3 mg: 2/day for MD/DO prescribers, 1/day for NP/PA prescribers*; 8/2 mg: 4/day for MD/DO prescribers, 2/day for NP/PA prescribers*; 4/1 mg: 2/day; 2/0.5 mg: 3/day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months
Notes	*NP/PA prescribers are limited to 16 mg of buprenorphine per day. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Suboxone® films PA Criteria:

- Diagnosis of opiate addiction; **AND**
- Prescriber is enrolled and in good standing in the BESMART program; **AND**
- Buprenorphine will not be approved for treatment of depression or pain; **AND**
- Patient has a documented allergy to inactive ingredient in preferred product that is not in requested product (DOCUMENTATION REQUIRED)

Zubsolv® tablets

PDL Status	NP
Quantity Limit	11.4/2.9 mg: 2/day for MD/DO prescribers, 1/day for NP/PA prescribers*; 8.6/2.1 mg: 2/day for MD/DO prescribers, 1/day for NP/PA prescribers*; 5.7/1.4 mg: 4/day for MD/DO prescribers, 2/day for NP/PA prescribers*; 2.9/0.71 mg: 2/day; 1.4/0.36 mg: 3/day^; 0.7/0.18 mg: 3/day^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months
Notes	*NP/PA prescribers are limited to 16 mg of buprenorphine equivalent per day. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Zubsolv® tablets PA Criteria:
See Suboxone® film prior authorization criteria.

All Other TennCare Providers (Non-BESMART)

Please submit faxed PA requests using the **Buprenorphine Products PA Form**.

Nurse Practitioners (NP), Physician Assistants (PA)s, or other mid-level providers must be in the BESMART Network to prescribe buprenorphine products.

Non-BESMART Quantity Limit Criteria for Requests for > 16 mg to ≤ 32 mg

- Provider is a MD or DO; **AND**
- If the request is for ≥ 24 mg/day, the patient is currently taking 24 mg/day without evidence of over-medication side effects (e.g., sedation, cognitive fog) (confirmed via paid claims or submitted chart notes); **AND**
- Patient meets ONE of the following:
  - The patient is pregnant (must provide estimated due date) (PA approval duration: 3 months past due date)
  - The patient is less than 12 months postpartum from delivery date (must provide delivery date) (PA approval duration: 12 months from delivery date)
  - Recent IV drug use, or fentanyl use, or recent OUD-related hospitalization or relapse confirmed by prescriber attestation (PA approval duration: 12 months)
  - Current users receiving greater than 50 mg of methadone for OUD treatment transitioning to buprenorphine agonist therapy demonstrated by paid TennCare (or previous insurer if new to TennCare) claims data from the enrollee's health insurer, provider attestation, or medical records. (PA approval duration: 12 months)
  - Newly eligible TennCare enrollees who are current users of > 16 mg per day of buprenorphine demonstrated by paid claims data from the enrollee's previous health insurer (PA duration: 12 months);

Note: Criteria applies to adult (≥ 21 years old) enrollees only. Children have access to 32 mg of buprenorphine daily across both networks if prescribed by MD or DO.

Buprenorphine/naloxone tablets

PDL Status	P
Quantity Limit	8/2 mg: 2/day for 6 months, then 1/day; 2/0.5 mg: 3 /day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months
Notes	^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine/naloxone tablet PA Criteria:

- Diagnosis of opiate addiction; **AND**
- Prescriber is NOT a nurse practitioner or physician assistant; **AND**
- Physician attests they have reviewed the Tennessee Controlled Substance Monitoring Database for this patient on the date of the prior authorization request to ensure that concomitant narcotic or benzodiazepine use is not occurring.

Additional Information:

- Buprenorphine will not be approved for treatment of depression or pain.
- Buprenorphine will not be approved for recipients whose medication history indicates use of concomitant narcotics or benzodiazepines without a clinically valid reason and drug tapering plan.
- Quantity limit is as a single daily dose. Twice daily dosing may be approved as clinically necessary.
- Physicians will be asked to provide an anticipated treatment plan for the patient (including anticipated dosing for induction & maintenance phases, anticipated frequency of office visits, & anticipated plan for psychosocial counseling).
- The "Here to Help" program as an exclusive provider of counseling will not be accepted.
- Prior Authorizations will be assigned to the prescribing physician.
- Requests for buprenorphine from a different physician will require a new prior authorization request and documentation that the previous prescribing physician has communicated transfer of care.

Buprenorphine tablets

PDL Status	NP
Quantity Limit	8 mg: 2/day for 6 months, then 1/day; 2 mg: 3 /day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Pregnancy: 3 months past due date; Breastfeeding: 6 months (maximum 4 approvals); Contraindication to naloxone: initial authorization: 6 months; reauthorization: 12 months
Notes	Mild rash, itching, and GI intolerance are not accepted as intolerance to naloxone. ^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine tablets PA Criteria:

See buprenorphine/naloxone tab prior authorization criteria; **AND**

- ONE of the following:
  - Patient is actively pregnant (must provide estimated due date)
  - Patient is actively breastfeeding (must provide delivery date)
  - Request is for a two-day induction for patients transitioning off methadone
  - Patient is unable to take naloxone containing products due to a contraindication, drug to drug interaction, or history of toxic side effects that caused immediate or long-term damage (DOCUMENTATION REQUIRED).

Buprenorphine/naloxone film

PDL Status	NP
Quantity Limit	12/3 mg and 8/2 mg: 2/day for 6 months, then 1/day; 4/1 mg: 2/day; 2/0.5 mg: 3/day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months;
Notes	^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Buprenorphine/naloxone film PA Criteria:

See buprenorphine/naloxone tab prior authorization criteria; **AND**

- Patient has a documented allergy to inactive ingredient in preferred product that is not in requested product (DOCUMENTATION REQUIRED)

Suboxone® film

PDL Status	NP
Quantity Limit	12/3 mg and 8/2 mg: 2/day for 6 months, then 1/day; 4/1 mg: 2/day; 2/0.5 mg: 3/day ^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months;
Notes	^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Suboxone® film PA Criteria:

See buprenorphine/naloxone tab prior authorization criteria; **AND**

- Patient has a documented allergy to inactive ingredient in preferred product that is not in requested product (DOCUMENTATION REQUIRED)

Zubsolv®

PDL Status	NP
Quantity Limit	11.4/2.9 mg and 8.6/2.1 mg: 1/day; 5.7/1.4 mg: 2/day for 6 months, then 1/day; 2.9/0.71 mg: 2/day; 1.4/0.36 mg: 3/day^; 0.7/0.18 mg: 3/day^
PA Criteria	See below
PA Form	Buprenorphine Products PA Form
PA Approval Duration	Initial authorization: 6 months; reauthorization: 12 months;
Notes	^Requests for 4/day will only be approved if dose is being titrated or patient's condition is too unstable

Zubsolv® PA Criteria:

See buprenorphine/naloxone tab prior authorization criteria; **AND**

- Patient has a documented allergy to inactive ingredient in preferred product that is not in requested product (DOCUMENTATION REQUIRED)

Narcotic Agonists/Antagonists

Nalbuphine

PDL Status	P
Quantity Limit	10 mg/mL: 4 mL/day; 20 mg/mL: 8 mL/day
PA Criteria	See below
PA Form	General PA Form

Nalbuphine PA Criteria:

- Trial and failure of at least 2 short acting narcotics; **OR**
- Documented contraindication, or intolerance to short acting narcotics; **AND**
- Unable to swallow, **OR** Unable to absorb medications through the GI tract.

Butorphanol nasal spray

PDL Status	NP
Quantity Limit	2.5 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Butorphanol nasal spray PA Criteria:

- Documented inability to swallow or absorb PO narcotics, **OR**
- For the treatment of migraines; **AND**
- Recipient MUST be receiving prophylactic therapy for migraines, **AND**
  - Trial and failure, intolerance, or contraindication to at least ONE agent in EACH of the following categories:
    - 5-HT1 receptor antagonist (triptans)
    - Anti-migraine combinations
    - NSAIDs

Pentazocine/naloxone

PDL Status	NP
Quantity Limit	12/day
PA Criteria	See below
PA Form	General PA Form

Pentazocine/naloxone PA Criteria:

- Contraindication, or intolerance to ALL short acting narcotics
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database for this patient within the last 30 days

Narcotics, Long-Acting

Approval of non-preferred agents in the Long-Acting Narcotics class requires contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (NOTE: this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated.

Edits on agents in the Short-Acting and Long-Acting Narcotics classes of the PDL impact all first-time (acute) and non-chronic opioid users. For details, visit: **Opioid Criteria - Acute, Chronic, and Exceptions.**

For agents in this class, please submit one of the following PA forms with all PA submissions: **Acute Opioid PA Form**, **Chronic Opioid PA Form**, or **Exceptions Opioid PA Form.**

***^Morphine Milligram Equivalent (MME) Criteria:**

- Indication or diagnosis is Cancer pain or Hospice
  - Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days (document date); **AND**
  - Document prescriber's specialty; **AND**
  - Patient has a written treatment plan with established objectives; **AND**
  - Patient has a signed Pain Management Agreement; **AND**
  - If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
    - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
    - Has an intrauterine device (IUD) or implant; **OR**
    - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
    - If the patient is pregnant, provider attests to ALL the following:
      - Alternative treatment options are ineffective, or not appropriate; **AND**
      - Lowest effective dose and shortest duration will be prescribed; **AND**
      - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS)

Fentanyl patch (12, 25, 50, 75, & 100 mcg/HR)

PDL Status	P
Quantity Limit	10 patches/30 days; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Fentanyl patch (12, 25, 50, 75, & 100 mcg/hr) PA Criteria:

- Management of severe pain with need for around-the-clock analgesia for an extended period; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- Requests for strengths ≥ 90 MME per day: (Please refer to the TennCare**MME Conversion Chart**)
  - Recipient must be opioid tolerant (as demonstrated by at least a week or longer history of morphine ≥ 60 mg/day, oral oxycodone ≥ 30 mg/day, oral hydromorphone ≥ 8 mg/day, or an equianalgesic dose of another opioid); **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The provider attests to investigating ALL of the following before submitting a PA:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids

Morphine ER tablets

PDL Status	P
Quantity Limit	1/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Morphine ER tablets PA Criteria:

See fentanyl patch (12, 25, 50, 75, & 100 mcg) PA criteria

Tramadol ER tablets

PDL Status	P
Quantity Limit	1/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Tramadol ER tablets PA Criteria:
See fentanyl patch (12, 25, 50, 75, & 100 mcg) PA criteria

Belbuca®

PDL Status	NP
Quantity Limit	2/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Belbuca® PA Criteria:

- Management of severe pain with need for around-the-clock analgesia for an extended period; **AND**
- Prescriber has checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Patients who have not been titrated down to no more than 30 mg morphine (or morphine equivalents) per day will NOT be approved; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
  - Alternative treatment options are ineffective, or not appropriate; **AND**
  - Lowest effective dose and shortest duration will be prescribed; **AND**
  - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The prescriber attests to investigating all of following before submitting a PA:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids; **AND**
- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated

Buprenorphine patch

PDL Status	NP
Quantity Limit	4 patches/28 days; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary. Additionally, buprenorphine patch 7.5, 10, 15, and 20 mcg/hr will be approved for opioid-experienced patients only.

Buprenorphine patch PA Criteria:
See Belbuca® prior authorization criteria

Butrans® patch

PDL Status	NP
Quantity Limit	4 patches/28 days; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary. Additionally, Butrans® patch 7.5, 10, 15, and 20 mcg/hr will be approved for opioid-experienced patients only.

Butrans® patch PA Criteria:
See Belbuca® prior authorization criteria

ConZip®

PDL Status	NP
Quantity Limit	1/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Notes	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

ConZip® PA Criteria:

- Management of severe pain with need for around-the-clock analgesia for an extended period; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The prescriber attests to investigating ALL of the following before submitting a PA:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids; **AND**
- If patient is 12 to 18 years of age: (For patients less than 12 years of age, approval will not be granted)
  - Patient does not have any of the following:
  - Obesity (BMI ≥ 30)
  - Obstructive Sleep Apnea
  - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, pulmonary hypertension, etc.)
  - Recent adenectomy/tonsillectomy; **AND**
  - Trial and failure or contraindication to acetaminophen; **AND**
  - Trial and failure or contraindication to ALL NSAIDs; **AND**
  - Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated

Fentanyl patch (37.5, 62.5, & 87.5 mcg/HR)

PDL Status	NP
Quantity Limit	10 patches/30 days; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Fentanyl patch (37.5, 62.5, & 87.5 mcg/HR) PA Criteria:

- Management of severe pain with need for around-the-clock analgesia for an extended period; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- Requests for strengths ≥ 90 MME per day: (Please refer to the TennCare**MME Conversion Chart**)
  - Recipient must be opioid tolerant (as demonstrated by at least a week or longer history of morphine ≥ 60 mg/day, oral oxycodone ≥ 30 mg/day, oral hydromorphone ≥ 8 mg/day, or an equianalgesic dose of another opioid); **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The provider attests to investigating ALL of the following before submitting a PA:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids; **AND**
- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated

Hydrocodone ER capsules & tablets

PDL Status	NP
Quantity Limit	Tabs: 1 tab/day; Caps: 2 caps/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hydrocodone ER PA Criteria:

- The prescriber has checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- Approval of non-preferred agents requires: Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (NOTE: this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated.; **AND**
- The following should be investigated before a PA is granted:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids; **AND**
- Requests for strengths ≥ 90 MME per day: (Please refer to the TennCare **MME Conversion Chart**)
  - Recipient must be opioid tolerant (as demonstrated by at least a week or longer history of morphine ≥ 60 mg/day, oral oxycodone ≥ 30 mg/day, oral hydromorphone ≥ 8 mg/day, or an equianalgesic dose of another opioid)

Hydromorphone ER tablets

PDL Status	NP
Quantity Limit	Tabs: 1 tab/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hydromorphone ER tablets PA Criteria:

- Management of severe pain with need for around-the-clock analgesia for an extended period; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- Requests for strengths ≥ 90 mg: (Please refer to the TennCare **MME Conversion Chart**)
  - Recipient must be opioid tolerant (as demonstrated by at least a week or longer history of morphine ≥ 60 mg/day, oral oxycodone ≥ 30 mg/day, oral hydromorphone ≥ 8 mg/day, or an equianalgesic dose of another opioid); **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The provider attests to investigating ALL of the following before submitting a PA:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids; **AND**
- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated

Hysingla® ER tablets

PDL Status	NP
Quantity Limit	Tabs: 1 tab/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hysingla® ER tablets PA Criteria:

See hydromorphone ER prior authorization criteria.

Methadone concentrate, tablets

PDL Status	NP
Quantity Limit	5 mg tabs: 8/day; 10 mg tabs: 4/day 5 mg/5 mL: 40 mL/day 10 mg/5 mL: 20 mL/day 10 mg/mL: 4 mL/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary. TennCare does not cover any form of methadone for the treatment of opioid addiction.

Methadone concentrate, tablets PA Criteria:

- One of the following:
  - Diagnosis of Metastatic Neoplasia
  - Infants up to 1 year of age who are discharged from hospital on a methadone taper will be approved for up to 30 days
  - Management of severe pain with need for around-the-clock analgesia for an extended period **AND** patient has contraindication to all other long-acting opioids; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines & opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- Requests for strengths ≥ 90 MME per day: (Please refer to the TennCare **MME Conversion Chart**)
  - Recipient must be opioid tolerant (as demonstrated by at least a week or longer history of morphine ≥ 60 mg/day, oral oxycodone ≥ 30 mg/day, oral hydromorphone ≥ 8 mg/day, or an equianalgesic dose of another opioid); **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS); **AND**
- The following should be investigated before a PA is granted:
  - History of substance abuse
  - Frequent requests for early refills
  - Reported frequent instances of lost tablets
  - Requests for odd quantities which requires fractional dosing
  - Requests for short-term or prn usage
  - Medication history indicates concurrent use of other extended-release opioids

Methadose® concentrate, tablet for oral suspension

PDL Status	NP
Quantity Limit	40 mg tab for oral susp: 1 tab/day 10 mg/mL: 4 mL/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary. TennCare does not cover any form of methadone for the treatment of opioid addiction.

Methadose® concentrate, tablet for oral suspension PA Criteria:
See methadone prior authorization criteria.

Morphine ER capsules

PDL Status	NP
Quantity Limit	Beads caps: 1/day; Caps: 2/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Morphine ER capsules PA Criteria:
See hydromorphone ER prior authorization criteria.

MS Contin® tablets

PDL Status	NP
Quantity Limit	15 mg, 30 mg, 60 mg tabs: 3/day; 100 mg tabs: 2/day; 200 mg tabs: 1/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

MS Contin® tablets PA Criteria:
See hydromorphone ER prior authorization criteria.

Oxycodone ER tablets

PDL Status	NP
Quantity Limit	Tabs: 2/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Oxycodone ER tablets PA Criteria:
See hydromorphone ER prior authorization criteria.

Oxycontin® tablets

PDL Status	NP
Quantity Limit	Tabs: 2/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Oxycontin® tablets PA Criteria:
See hydromorphone ER prior authorization criteria.

Oxymorphone ER tablets

PDL Status	NP
Quantity Limit	Tabs: 2/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary. Due to cross-reactivity with morphine, oxymorphone ER will not be approved for patients with immune-mediated morphine allergy.

1. Oxymorphone ER tablets PA Criteria:

See hydromorphone ER prior authorization criteria.

Tapendatol ER tablets

PDL Status	NP
Quantity Limit	2 tabs/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Tapendatol ER tablets PA Criteria:
See hydromorphone ER prior authorization criteria.

Tramadol ER capsules

PDL Status	NP
Quantity Limit	1 cap/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Tramadol ER capsules PA Criteria:
See hydromorphone ER prior authorization criteria.

Narcotics, Short-Acting

Approval of non-preferred agents in the Short-Acting Narcotics class requires contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (NOTE: this does not include GI intolerance) with ALL preferred agents, unless otherwise indicated.

Edits on agents in the Short-Acting and Long-Acting Narcotics classes of the PDL impact all first-time (acute) and non-chronic opioid users. For details, visit: **Opioid Criteria - Acute, Chronic, and Exceptions.**

For agents in this class, please submit one of the following PA forms with all PA submissions: **Acute Opioid PA Form**, **Chronic Opioid PA Form**, or **Exceptions Opioid PA Form.**

****Quantity Limit Override Criteria for Butalbital-Containing products:**
Requests for butalbital-containing products for quantities greater than 20 per 30 days will be approved for patients meeting the following criteria:

- Trial and failure of at least 2 prophylactic headache treatments: a tricyclic antidepressant (unless contraindicated) PLUS at least one of the following:
  - Divalproex sodium
  - Sodium valproate
  - Topiramate
  - Frovatriptan
  - Beta-blocker

***^Morphine Milligram Equivalent (MME) Criteria:**

- Indication or diagnosis is Cancer pain or Hospice
  - Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days (document date); **AND**
  - Document prescriber's specialty; **AND**
  - Patient has a written treatment plan with established objectives; **AND**
  - Patient has a signed Pain Management Agreement; **AND**
  - If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
    - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
    - Has an intrauterine device (IUD) or implant; **OR**
    - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
    - If the patient is pregnant, provider attests to ALL the following:
      - Alternative treatment options are ineffective, or not appropriate; **AND**
      - Lowest effective dose and shortest duration will be prescribed; **AND**

Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS)

APAP/Codeine soln, tabs

PDL Status	P
Quantity Limit	120-12 mg/5 mL soln: 15 mL/day 300-15 mg, 300-30 mg tabs: 12/day 300-60 mg tabs: 6/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

APAP/Codeine PA Criteria:

- Patient is ≥ 12 years of age and < 18 years of age; **AND**
- Trial and failure of acetaminophen; **AND**
- Contraindication to ALL NSAIDs; **AND**
- Patient does not have any of the following:
  - Obesity
  - Obstructive Sleep Apnea
  - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, etc.)
  - Recent adenectomy/tonsillectomy

Endocet®

PDL Status	P
Quantity Limit	2.5/325 mg tab: 12/day; All other tabs: 8/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Hydrocodone/APAP 325 mg soln, tabs

PDL Status	P
Quantity Limit	5/325 mg tabs: 12/day; 7.5/325, 10/325 mg tabs: 8/day; 7.5-325 mg/15 mL soln: 120 mL/day; 10-325 mg/15 mL soln: 89 mL/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Hydrocodone/ibuprofen

PDL Status	P
Quantity Limit	5/200 mg tabs: 12/day; 7.5/200 mg tabs: 8/day; 10/200 mg tabs: 6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Hydromorphone tabs

PDL Status	P
Quantity Limit	2 mg tabs: 7/day; 4 mg tabs: 3/day; 8 mg tabs: 1/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Morphine IR tabs

PDL Status	P
Quantity Limit	Tabs: 6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Morphine solution

PDL Status	P
Quantity Limit	*^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Morphine solution PA Criteria:

- Prescriber has checked the Tennessee Controlled Substance Database for this patient within the last 7 days (document date); **OR** request is for a hospice patient, HIV/AIDS patient, active cancer patient, **OR** long-term care facility resident (document name of facility); **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider; **AND**
- Pain agreement required. Please refer to the Opioid and Controlled Substance Agreement; **AND**
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS)

Oxycodone/APAP 325 mg soln, tablets

PDL Status	P
Quantity Limit	2.5/325 mg tab: 12/day; All other tabs: 8/day; Soln: 40 mL/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Oxycodone concentrate (100 mg/5 mL or 20 mg/mL)

PDL Status	P
Quantity Limit	*^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Oxycodone concentrate PA Criteria:

See morphine solution prior authorization criteria; **AND**

- Patient must be opioid tolerant by having at least 32 MME per day for at least 1 week

Oxycodone tablets

PDL Status	P
Quantity Limit	5 & 10 mg tab: 8/day; 15, 20, & 30 mg: 4/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Oxycodone soln (5 mg/5 mL)

PDL Status	P
Quantity Limit	*^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Tramadol soln, tabs (50 and 100 mg)

PDL Status	P
Quantity Limit	50 mg and 100 mg Tabs: 8/day; Soln: 80 mL/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill. Patients 18 years and older will only be subject to the quantity limit and opioid criteria

Tramadol PA Criteria:

- Patient is > 12 years of age and < 18 years of age; **AND**
- Patient does not have any of the following:
  - Obesity (BMI ≥ 30)
  - Obstructive Sleep Apnea
  - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, etc.)
  - Recent adenectomy/tonsillectomy; **AND**
- Trial and failure or contraindication to acetaminophen; **AND**
- Trial and failure or contraindication to ALL NSAIDs

Tramadol/APAP

PDL Status	P
Quantity Limit	Tabs: 12/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Prior Authorization is required for all subsequent acute prescriptions after the first fill.

Tramadol/APAP PA Criteria:
See tramadol prior authorization criteria.

Butalbital/APAP/caffeine/codeine

PDL Status	NP
Quantity Limit	**Butalbital containing products: 20/30 days Max: 4 g APAP/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.

Butalbital/APAP/caffeine/codeine PA Criteria:

- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long- term damage (NOTE: this does not include GI intolerance) with ALL preferred short-acting narcotic agents; **AND**
- One of the following:
  - Patients ≥ 18 years of age
  - Patient is > 12 years of age and < 18 years of age; **AND**
    - Trial and failure of acetaminophen; **AND**
    - Contraindication to ALL NSAIDs; **AND**
    - Patient does not have any of the following:
      - Obesity
      - Obstructive Sleep Apnea
      - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, etc.)
      - Recent adenectomy/tonsillectomy

Butalbital/ASA/caffeine/codeine

PDL Status	NP
Quantity Limit	**Butalbital containing products: 20/30 days
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.

Butalbital/ASA/caffeine/codeine PA Criteria:
See butalbital/APAP/caffeine/codeine prior authorization criteria.

Codeine

PDL Status	NP
Quantity Limit	15 & 30 mg tabs: 12/day; 60 mg tabs: 6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.

Codeine PA Criteria:

- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long- term damage (NOTE: this does not include GI intolerance) with ALL preferred short-acting narcotic agents; **AND**
- One of the following:
  - Patients ≥ 18 years of age
  - Patient is > 12 years of age and < 18 years of age; **AND**
    - Trial and failure of acetaminophen; **AND**
    - Contraindication to ALL NSAIDs; **AND**
    - Patient does not have any of the following:
      - Obesity
      - Obstructive Sleep Apnea
      - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, etc.)
      - Recent adenectomy/tonsillectomy

Dihydrocodeine/APAP/caffeine

PDL Status	NP
Quantity Limit	Tabs: 8/day Max: 4 g APAP/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	

Dihydrocodeine/APAP/caffeine PA Criteria:

See butalbital/APAP/caffeine/codeine prior authorization criteria.

Dilaudid® tablets

PDL Status	NP
Quantity Limit	2 mg tabs: 7/day; 4 mg tabs: 3/day; 8 mg tabs: 1/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Dilaudid® PA Criteria:

- Contraindication, drug to drug interaction, or history of toxic side effects that cause immediate or long-term damage (NOTE: this does not include GI intolerance) with ALL preferred agents; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 7 days; **AND**
- Pain agreement required. Please refer to the **Opioid and Controlled Substance Agreement**; **AND**
- Concomitant use of benzodiazepines and opioids will only be approved under the care of, or referral to, a mental health provider.
- If request is for a female of child-bearing age (14-44 years), patient meets ONE of the following:
  - Using contraception (e.g., barrier, oral contraceptive, rhythm method); **OR**
  - Has an intrauterine device (IUD) or implant; **OR**
  - Has history of hysterectomy, tubal ligation, or endometrial ablation; **OR**
  - If the patient is pregnant, provider attests to ALL the following:
    - Alternative treatment options are ineffective, or not appropriate; **AND**
    - Lowest effective dose and shortest duration will be prescribed; **AND**
    - Patient has been counseled on the risk associated with Neonatal Abstinence Syndrome (NAS)

Fioricet® with Codeine

PDL Status	NP
Quantity Limit	**Butalbital containing products: 20/30 days Max: 4 g APAP/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.

Fioricet® with Codeine PA Criteria:

See butalbital/APAP/caffeine/codeine prior authorization criteria.

Hydrocodone/APAP 300 mg

PDL Status	NP
Quantity Limit	5/300 mg tab: 12/day; 10/300 mg tab: 6/day; Soln: 89 mL/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hydrocodone/APAP 300 mg PA Criteria:
See Dilaudid® prior authorization criteria.

Hydromorphone liquid

PDL Status	NP
Quantity Limit	Liquid: 15 mL/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hydromorphone liquid PA Criteria:
See Dilaudid® prior authorization criteria.

Hydromorphone suppositories

PDL Status	NP
Quantity Limit	Suppositories: 5/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Hydromorphone suppositories PA Criteria:
See Dilaudid® prior authorization criteria.

Levorphanol tablets

PDL Status	NP
Quantity Limit	Tablets: 6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Levorphanol tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Meperidine solution, tablets

PDL Status	NP
Quantity Limit	Solution: 60 mL/day Tablets: 12/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Meperidine solution, tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Morphine suppositories

PDL Status	NP
Quantity Limit	5 mg suppositories: 12/day; All other strength suppositories: 6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Morphine suppositories PA Criteria:
See Dilaudid® prior authorization criteria.

Nalocet® tablets

PDL Status	NP
Quantity Limit	Tablets: 12/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Nalocet® tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Oxycodone/APAP 300 mg solution, tablets

PDL Status	NP
Quantity Limit	2.5/325 mg tabs: 12/day; All other strength tabs: 8/day; Soln: 40 mL/day *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Oxycodone/APAP 300 mg solution, tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Oxycodone capsules

PDL Status	NP
Quantity Limit	Capsules: 8/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Oxycodone capsules PA Criteria:
See Dilaudid® prior authorization criteria.

Oxymorphone tablets

PDL Status	NP
Quantity Limit	Tablets: 4/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Oxymorphone tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Percocet® tablets

PDL Status	NP
Quantity Limit	2.5/325 mg tablets: 12/day; All other strength tablets: 8/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Percocet® tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Prolate® solution, tablets

PDL Status	NP
Quantity Limit	Solution: 40 mL/day; Tablets: 8/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Prolate® solution, tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Qdolo® solution

PDL Status	NP
Quantity Limit	Solution: 40 mL/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Qdolo® solution PA Criteria:
See Dilaudid® prior authorization criteria.

Roxicodone® tablets

PDL Status	NP
Quantity Limit	Tablets: 4/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Roxicodone® tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Roxybond® tablets

PDL Status	NP
Quantity Limit	5 mg tablets: 8/day 10, 15, & 30 mg tablets: 4/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Roxybond® tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Tapendatol IR tablets

PDL Status	NP
Quantity Limit	6/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	See below
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Tapendatol IR tablets PA Criteria:
See Dilaudid® prior authorization criteria.

Tramadol tabs (25 mg and 75 mg)

PDL Status	NP
Quantity Limit	25 mg and 75 Tabs: 8/day; *^Max Total: Non-Chronic: 60 MME/day; Chronic: 200 MME/day
PA Criteria	None
PA Form	Acute Opioid PA Form, Chronic Opioid PA Form, or Exceptions Opioid PA Form.
Note	Use of opioid analgesics during pregnancy has been associated with Neonatal Opioid Withdrawal Syndrome. Providers MUST counsel women of childbearing age on the risks of becoming pregnant while receiving opioids, including the risk of Neonatal Opioid Withdrawal Syndrome. Providers should offer access to contraceptive services when necessary.

Tramadol PA Criteria:
See Dilaudid® prior authorization criteria; **AND**

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- One of the following:
  - Patient is ≥ 18 years of age; **OR**
  - Patient is > 12 years of age and < 18 years of age; **AND**
    - Patient does not have any of the following:
      - Obesity (BMI ≥ 30)
      - Obstructive Sleep Apnea
      - Severe Lung Disease (acute or severe asthma, COPD, Cystic Fibrosis, hypoxemia, hypercapnia, pneumonia, etc.)
      - Recent adenectomy/tonsillectomy; **AND**
    - Trial and failure or contraindication to acetaminophen; **AND**
    - Trial and failure or contraindication to ALL NSAIDs

NSAIDs

Please refer to the OTC Drug Formulary for coverage of OTC products.

Celecoxib capsules

PDL Status	P
Quantity Limit	2 capsules/day
PA Criteria	None

Diclofenac 1% gel

PDL Status	P
Quantity Limit	10 g/day
PA Criteria	None

Diclofenac 2% solution

PDL Status	P
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Diclofenac 2% solution PA Criteria:

- Diagnosis of osteoarthritis pain of the knee

Ketorolac tablets

PDL Status	P
Quantity Limit	20 tablets/60 days
PA Criteria	None

Voltaren® 1% gel

PDL Status	P
Quantity Limit	Gel: 10 g/day
PA Criteria	None

Anaprox DS®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Anaprox DS® PA Criteria:

Standard NP Criteria.

Celebrex® capsules

PDL Status	NP
Quantity Limit	2 capsules/day
PA Criteria	See below
PA Form	General PA Form

Celebrex® PA Criteria:

Standard NP Criteria.

Coxanto® capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See below
PA Form	General PA Form

Coxanto® capsules PA Criteria:

- Diagnosis of ONE of the following:
  - Juvenile rheumatoid arthritis
  - Osteoarthritis
  - Rheumatoid arthritis; **AND**
- Clinically valid reason why oxaprozin tablets cannot be used; **AND**
- Clinically valid reason why generic oxaprozin capsules cannot be used; **AND**
- Inadequate response or intolerance to TWO preferred NSAIDs

Daypro® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Daypro® PA Criteria:

Standard NP Criteria.

Diclofenac potassium capsules, diclofenac potassium packets

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Diclofenac potassium capsules, diclofenac potassium packets PA Criteria:

- Clinically valid reason why the preferred NSAIDs cannot be used

Diclofenac patch

PDL Status	NP
Quantity Limit	Patches: 2/day
PA Criteria	See below
PA Form	General PA Form

Diclofenac patch PA Criteria:

- Clinically valid reason why the preferred NSAIDs cannot be used

Diclofenac potassium 25 mg tablets

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Diclofenac patch PA Criteria:

- Clinically valid reason why the preferred diclofenac products cannot be used

Diclofenac 1.5% solution

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Diclofenac 1.5% solution PA Criteria:

- Clinically valid reason why the preferred NSAIDs cannot be used

EC-Naprosyn®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

EC-Naprosyn® PA Criteria:

Standard NP Criteria.

Elyxyb® solution

PDL Status	NP
Quantity Limit	6 bottles (28.8 mL)/30 days
PA Criteria	See below
PA Form	General PA Form

Elyxyb® solution PA Criteria:

- Diagnosis of migraine; **AND**
- Patient is unable to swallow solid dosage forms; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Etodolac capsules, tablets, ER tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Etodolac capsules, tablets, ER tablets PA Criteria:

Standard NP Criteria.

Fenoprofen capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See below
PA Form	General PA Form

Fenoprofen capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Fenoprofen tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Fenoprofen tablets PA Criteria:
Standard NP Criteria.

Fenopron® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Fenopron® capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Flurbiprofen tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Flurbiprofen tablets PA Criteria:
Standard NP Criteria.

Ibuprofen 300 mg tablets

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Ibuprofen 300 mg tablets PA Criteria:

- Clinically valid reason why the preferred NSAIDs cannot be used

Lofena® tablets

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Lofena® tablets PA Criteria:

- Clinically valid reason why the preferred diclofenac products cannot be used

Ketoprofen capsules, ER capsules

PDL Status	NP
Quantity Limit	25 mg caps: 3/day; 50 mg caps: 6/day
PA Criteria	See below
PA Form	General PA Form

Ketoprofen capsules, ER capsules PA Criteria:
Standard NP Criteria.

Ketorolac nasal spray

PDL Status	NP
Quantity Limit	5 bottles/60 days
PA Criteria	See below
PA Form	General PA Form

Ketorolac nasal spray PA Criteria:

- Trial and failure, contraindication, or intolerance of oral ketorolac; **OR**
- Patient is unable to swallow solid dosage forms

Meloxicam capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Meloxicam capsules PA Criteria:

- Clinically valid reason why the preferred meloxicam tablets cannot be used

Naprosyn® suspension, tablets, naproxen suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Naprosyn® suspension, tablets, naproxen suspension PA Criteria:
Standard NP Criteria.

Naproxen CR/ER tabs (375, 500, 750 mg)

PDL Status	NP
Quantity Limit	Not applicable
PA Criteria	See below
PA Form	General PA Form

Naproxen CR/ER tabs PA Criteria:

- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications

Oxaprozin capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See below
PA Form	General PA Form

Oxaprozin capsules PA Criteria:

- Diagnosis of ONE of the following:
  - Juvenile rheumatoid arthritis
  - Osteoarthritis
  - Rheumatoid arthritis; **AND**
- Clinically valid reason why oxaprozin tablets cannot be used; **AND**
- Inadequate response or intolerance to TWO preferred NSAIDs

Oxaprozin tablets, piroxicam capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Oxaprozin tablets, piroxicam capsules PA Criteria:

Standard NP Criteria.

Relafen DS® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Relafen DS® tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Tolectin DS® capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See below
PA Form	General PA Form

Tolectin DS® capsules PA Criteria:

- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred NSAIDs

Tolectin® tablets

PDL Status	NP
Quantity Limit	3 tablets/day
PA Criteria	See below
PA Form	General PA Form

Tolectin DS® tablets PA Criteria:

- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred NSAIDs

Tolmetin capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See below
PA Form	General PA Form

Tolmetin capsules PA Criteria:

- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred NSAIDs

Tolmetin tablets

PDL Status	NP
Quantity Limit	3 tablets/day
PA Criteria	See below
PA Form	General PA Form

Tolmetin tablets PA Criteria:

- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred NSAIDs

Vyscoxa® suspension

PDL Status	NP
Quantity Limit	40 mL/day
PA Criteria	See below
PA Form	General PA Form

Vyscoxa® suspension PA Criteria:

- Diagnosis of one of the following:
  - Osteoarthritis (OA)
  - Rheumatoid arthritis (RA)
  - Ankylosing Spondylitis (AS); **OR**
  - Juvenile Rheumatoid Arthritis; **AND**
- Clinically valid reason why preferred celecoxib capsules or brand Celebrex capsules cannot be used; **AND**
- Medical documentation that patient is not at an increased risk for GI bleeds (e.g., prior GI ulcer, bleeding, perforation, etc.) that would prevent use of other NSAID agents; **AND**
- Patient is unable to swallow solid dosage forms

NSAIDs/Anti-Ulcer Agents

Diclofenac/misoprostol tablets

PDL Status	P
Quantity Limit	50 mg/200 mcg tablets: 4/day; 75 mg/200 mcg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Diclofenac/misoprostol tablets PA Criteria:

- Patient is > 60 years old; **OR**
- Patients < 60 years old and is at high risk for GI side effects as indicated by ANY of the following:
  - History of peptic ulcer disease/GI bleed/NSAID gastropathy
  - GERD (gastroesophageal reflux disease) due to conventional NSAIDS
  - Patient on anticoagulants
  - Patient on chronic corticosteroids
  - History of platelet dysfunction or coagulopathy, including use of clopidogrel or aspirin
  - Patient on methotrexate

Ibuprofen/famotidine tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Ibuprofen/famotidine tablets PA Criteria:

- Patient is at high risk for GI side effects as indicated by ANY of the following:
  - History of peptic ulcer disease/GI bleed/NSAID gastropathy
  - GERD (gastroesophageal reflux disease) due to conventional NSAIDS
  - Patient on anticoagulants
  - Patient on chronic corticosteroids
  - History of platelet dysfunction or coagulopathy, including use of clopidogrel or aspirin
  - Patient on methotrexate

Arthrotec® tablets

PDL Status	NP
Quantity Limit	50 mg/200 mcg tablets: 4/day; 75 mg/200 mcg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Arthrotec® PA Criteria:

Standard NP Criteria.

Naproxen/esomeprazole tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Naproxen/esomeprazole tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Non-Opioid Analgesics

Journavx® tablets

PDL Status	P
Quantity Limit	30 tablets/180 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	14 days
Note	Journavx is available without prior authorization for up to 30 tablets per 180 days. Requests exceeding this quantity limit will require prior authorization.

Journavx® tablets PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of moderate to severe acute pain; **AND**
- Medication will not be used for longer than 14 days for any one acute pain occurrence; **AND**
- Prescriber attests that pain cannot be controlled with conventional pharmacological therapies such as acetaminophen, NSAIDS, and local anesthetics; **AND**
- Patient does not have severe hepatic impairment (Child Pugh Class-C); **AND**
- Medication will not be used in combination with opioid products (e.g., oxycodone, hydrocodone); **AND**
- Medication will not be used in combination with strong CYP3A inhibitors; **AND**
- For female patients with reproductive potential, prescriber attest to BOTH of the following:
  - Patient is not pregnant prior to initiation of therapy
  - Patients taking concomitant hormonal contraceptives containing progestins other than levonorgestrel and norethindrone have been advised to use alternative contraceptives during Journavx treatment and for 28 days after discontinuation of therapy

Opioid Reversal Agents

Please refer to the OTC Drug Formulary for coverage of OTC products.

Kloxxado® spray, naloxone nasal spray, Narcan® nasal spray, Opvee® spray, Rextovy® spray

PDL Status	P
Quantity Limit	2 sprayers/30 days
PA Criteria	N/A
Note	Please refer to the OTC Drug Formulary for coverage of OTC products.

Naloxone injection, Zurnai® injection

PDL Status	P
Quantity Limit	2 injections/30 days
PA Criteria	N/A

Salicylates

Salsalate tablets

PDL Status	P
Quantity Limit	500 mg tablets: 6/day; 750 mg tablets: 4/day
PA Criteria	None

Diflunisal tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	Standard NP Criteria
PA Form	General PA Form

Diflunisal tablets PA Criteria:
Standard NP Criteria.

Dolobid® 250 and 375 mg tablets

PDL Status	NP
Quantity Limit	250 mg tablets: 6/day; 375 mg tablets: 4/day
PA Criteria	See below
PA Form	General PA Form

Dolobid® 250 and 375 mg tablets PA Criteria:

- Clinically valid reason why generic diflunisal cannot be used; **AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice

Transmucosal Fentanyl Products

Fentanyl lozenge

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form
Note	Prescription should be written by, or in consultation with, an oncologist or pain management specialist unless patient is enrolled in or eligible for hospice care.

Fentanyl lozenge PA Criteria:

- Medication is ordered for the treatment of breakthrough cancer pain
- Recipient must be receiving around-the-clock scheduled long-acting opioids
- Recipient must be tolerant to opioids, defined as one of the following:
  - ≥ 60 mg oral morphine per day for at least one week without adequate pain relief
  - ≥ 25 mcg/hr transdermal fentanyl for at least one week without adequate pain relief
  - ≥ 30 mg oral oxycodone/day for at least one week without adequate pain relief
  - ≥ 8 mg oral hydromorphone/day for at least one week without adequate pain relief
  - ≥ 25 mg oral oxymorphone/day for at least one week without adequate pain relief
  - Equianalgesic dose of another opioid for at least one week without adequate pain relief
- Trial and failure, contraindication, intolerance, or drug-to-drug interaction with at least 2 immediate release opioid products

II. ANTI-INFECTIVES

Antibiotics: Agents for Diarrhea

Vancomycin solution

PDL Status	P
Quantity Limit	2,000 mg/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 days

Vancomycin solution PA Criteria:

- Patient is unable to swallow sold dosage forms; **OR**
- Patient is < 12 years of age

Aemcolo®

PDL Status	NP
Quantity Limit	Tabs: 12/Rx, max 24 tabs/year.
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 days

Aemcolo® PA Criteria:

- Patient is being treated for traveler's diarrhea; **AND**
- Trial and failure, contraindication, intolerance, drug-drug interaction or resistance to a fluoroquinolone or azithromycin

Firvanq®

PDL Status	NP
Quantity Limit	2,000 mg/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 days

Firvanq® PA Criteria:

- Trial and failure, contraindication, or intolerance to generic vancomycin solution

Vancocin® capsules

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 days

Vancocin® capsules PA Criteria:

- Trial and failure, contraindication, or intolerance to generic vancomycin capsules

Antibiotics: Aminoglycosides, Oral

Arikayce®

PDL Status	NP
Quantity Limit	8.4 mL/day
PA Criteria	See Below
PA Form	General PA Form

Arikayce® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of Mycobacterium avium complex (MAC) lung disease as determined by the following:
  - Chest radiography or high-resolution computed tomography (HRCT) scan; **AND**
  - At least two positive sputum cultures; **AND**
  - Other conditions such as tuberculosis and lung malignancy have been ruled out; **AND**
- Patient has failed a multi-drug regimen with a macrolide (clarithromycin or azithromycin), rifampin, and ethambutol. (Failure is defined as continual positive sputum cultures for MAC while adhering to a multi-drug treatment regimen for a minimum duration of 6-months); **AND**
- Prescribed in conjunction with a multi-drug antimycobacterial regimen

Arikayce® Renewal PA Criteria:

- Patient has demonstrated response to therapy defined as having three consecutive monthly negative sputum cultures by month six of treatment; **AND**
- Patient has not experienced toxicity to amikacin treatment (e.g., ototoxicity, renal toxicity, neuromuscular blockade)

Antibiotics: Anti-Tuberculosis, Oral

Cycloserine capsules, Priftin® tablets

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Cycloserine capsules, Priftin® tablets PA Criteria:

Standard NP Criteria

Rifapentine tablets

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Rifapentine tablets PA Criteria:

- Clinically valid reason why brand Priftin® tablets cannot be used

Sirturo®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	9 months

Sirturo® PA Criteria:

- Patient is ≥ 2 years of age and weighs ≥ 8 kg; **AND**
- Patient has a diagnosis of pulmonary multi-drug-resistant tuberculosis (MDR-TB); **AND**
- Sirturo is prescribed as part of a combination regimen with at least 3 other drugs to which the patient's MDRTB isolate has been shown to be susceptible; **AND**
- Sirturo is prescribed by, or in consultation with, an infectious disease specialist

Antibiotics: Cephalosporins, First Generation

Cefadroxil tablets, cephalexin tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cefadroxil tablets, cephalexin tablets PA Criteria:
Standard NP Criteria

Antibiotics: Cephalosporins, Second Generation

Cefaclor ER tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cefaclor ER tablets PA Criteria:
Standard NP Criteria

Antibiotics: Lincosamides, Oral

Clindamycin pediatric solution

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	No PA required for patients less than 12 years of age

Clindamycin pediatric solution PA Criteria:

- Patient less than 12 years of age; **OR**
- Patient is unable to swallow solid dosage forms

Cleocin® Pediatric Granules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cleocin® Pediatric Granules PA Criteria:

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why generic clindamycin solution cannot be used

Antibiotics: Macrolides

Azithromycin packets, tablets

PDL Status	P
Quantity Limit	1 g packet: 2/Rx 250 and 500 mg tablets: 12/Rx; 600 mg tablets: 8/month
PA Criteria	None

Dificid® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Dificid® tablets PA Criteria:

- Diagnosis of Clostridium difficile (C. diff) associated diarrhea

Clarithromycin ER/XL tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	None

Dificid® suspension

PDL Status	NP
Quantity Limit	1 bottle/Rx
PA Criteria	See Below
PA Form	General PA Form
Note	Individuals started on Dificid® therapy in the hospital will be approved for this agent following hospital discharge to allow for completion of the course of therapy.

Dificid® tablets PA Criteria:

- Diagnosis of *Clostridium difficile (C. diff)* associated diarrhea

E.E.S. Granules®, Ery-Tab®, Eryped®, erythromycin base tablets, erythromycin stearate tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

E.E.S. Granules®, Ery-Tab®, Eryped®, erythromycin base tablets, erythromycin stearate tablets PA Criteria:
Standard NP Criteria

Fidaxomicin tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Fidaxomicin tablets PA Criteria:

- Diagnosis of Clostridium difficile (C. diff) associated diarrhea; **AND**
- Clinically valid reason why Dificid® tablets cannot be used

Zithromax® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Zithromax® suspension PA Criteria:
Standard NP Criteria

Zithromax® tablets

PDL Status	NP
Quantity Limit	250 and 500 mg tablets: 12/Rx; 600 mg tablets: 8/month
PA Criteria	See below
PA Form	General PA Form

Zithromax® tablets PA Criteria:
Standard NP Criteria

Antibiotics: Methenamines

Hiprex®

PDL Status	NP
Quantity Limit	None
PA Criteria	None

Hiprex® PA Criteria:
Standard NP Criteria

Antibiotics: Nitrofurans, Oral

Nitrofurantoin suspension (25 mg/5 mL and 50 mg/10 mL)

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	No PA required for patients less than 12 years of age

Nitrofurantoin suspension (25 mg/5 mL and 50 mg/10 mL) PA Criteria:

- Patient is unable to swallow solid dosage forms

Macrobid® capsules, Macrodantin® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Macrobid® capsules, Macrodantin® capsules PA Criteria:
Standard NP Criteria

Nitrofurantoin suspension (50 mg/5 mL)

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	No PA required for patients less than 12 years of age

Nitrofurantoin suspension (50 mg/5 mL) PA Criteria:

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why preferred nitrofurantoin suspension cannot be used

Antibiotics: Oxazolidinones

Linezolid tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Linezolid tablets PA Criteria:

- Treatment is for ONE of the following:
  - Vancomycin Resistant Enterococcus faecalis infections
  - Healthcare-associated Methicillin-Resistant Staph Aureus (MRSA) infections or community-acquired MRSA with polyresistance
  - Community-acquired pneumonia (CAP) caused by S. pneumoniae or S. aureus (MSSA)
  - Nosocomial pneumonia caused by S. pneumoniae or S. aureus (including MSSA and MRSA)
  - Complicated skin and skin structure infections (SSSI) caused by S. aureus (MSSA and MRSA), S. pyogenes, or S. agalactiae.
  - Uncomplicated SSTI caused by S. aureus (MSSA only) or S. pyogenes
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Linezolid suspension

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Linezolid suspension PA Criteria:

- One of the following:
  - Patient is less than 12 years of age
  - Patient is unable to swallow oral dosage forms
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Sivextro®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Sivextro® PA Criteria:

- Diagnosis of acute bacterial skin and skin structure infection; **AND**
  - Patient must be resistant to or have a contraindication, or intolerance, to all other treatment options; **OR**
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Zyvox® suspension, tablets

PDL Status	NP
Quantity Limit	Susp: 60 mL/day; Tabs: 2/day
PA Criteria	See Below
PA Form	General PA Form

Zyvox® suspension, tablets PA Criteria:

Standard NP Criteria

Antibiotics: Penicillins

All brand penicillins, amoxicillin/clavulanic acid XR tablets, Augmentin® suspension, Augmentin ES® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

All brand penicillins, amoxicillin/clavulanic acid XR tablets, Augmentin® suspension, Augmentin ES® suspension PA Criteria:
Standard NP Criteria

Antibiotics: Quinolones

Baxdela® tablets

PDL Status	NP
Quantity Limit	2/day; Max 14-day supply
PA Criteria	See Below
PA Form	General PA Form

Baxdela® tablets PA Criteria:

- Patient age ≥ 18 years of age; **AND**
- ONE of the following:
  - Diagnosis of acute bacterial skin and skin structure infection (ABSSSI); **AND**
    - Trial and failure to, contraindication, or resistance to ONE preferred standard of care agents for ABSSSI (e.g., linezolid, clindamycin, doxycycline, SMX-TMP, vancomycin, cephalosporin, a preferred fluoroquinolone)
  - Diagnosis of community-acquired bacterial pneumonia (CABP); **AND**
    - Trial and failure to, contraindication, or resistance to TWO preferred standard of care agents for CABP (e.g., macrolide, doxycycline, a preferred fluoroquinolone, beta-lactam, linezolid)
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Cipro® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cipro® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Cipro® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cipro® tablets PA Criteria:
Standard NP Criteria

Ciprofloxacin suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Ciprofloxacin suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Levofloxacin suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Ciprofloxacin suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Moxifloxacin tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	See Appropriate Diagnosis PA bypass list for patients needing moxifloxacin tablets for *Mycoplasma genitalium* infections

Moxifloxacin tablets PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred agents; **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy); **OR**
- Diagnosis of *Mycoplasma genitalium*

Ofloxacin tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Ofloxacin tablets PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred agents; **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Antibiotics: Sulfonamides, Folate Agonist

Bactrim® tablets, Bactrim DS® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Bactrim® tablets, Bactrim DS® tablets PA Criteria:

Standard NP Criteria

Antibiotics: Tetracyclines

Doxycycline hyclate capsules (50 and 100 mg)

PDL Status	P
Quantity Limit	50 mg caps: 3/day; 100 mg caps: 2/day
PA Criteria	None

Doxycycline hyclate tablets (50 and 100 mg)

PDL Status	P
Quantity Limit	50 mg tabs: 3/day; 100 mg tabs: 2/day
PA Criteria	None

Doxycycline monohydrate capsules (50 and 100 mg)

PDL Status	P
Quantity Limit	50 mg caps: 3/day; 100 mg caps: 2/day
PA Criteria	None

Demeclocycline tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Demeclocycline tablets PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred agents; **OR**
- Treatment is for syndrome of inappropriate antidiuretic hormone secretion (SIADH)

Doryx® DR tablets

PDL Status	NP
Quantity Limit	50 mg: 3/day; All other strengths: 2/day
PA Criteria	See below
PA Form	General PA Form

Doryx® DR tablets PA Criteria:

Standard NP Criteria

Doxycycline DR capsules, tablets

PDL Status	NP
Quantity Limit	50 mg: 3/day; All other strengths: 2/day
PA Criteria	See below
PA Form	General PA Form

Doxycycline DR capsules, tablets PA Criteria:

Standard NP Criteria

Doxycycline hyclate capsules (75 mg)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Doxycycline hyclate capsules (75 mg) PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Doxycycline hyclate tablets (20, 75, and 150 mg)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Doxycycline hyclate tablets (20, 75, and 150 mg) PA Criteria:

- Agent is used as an adjunct to scaling and root planting to promote attachment level gain and to reduce pocket depth for adult periodontitis

Doxycycline monohydrate capsules (75 and 150 mg)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Doxycycline monohydrate capsules (75 and 150 mg) PA Criteria:

Standard NP Criteria

Doxycycline monohydrate suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Doxycycline monohydrate suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Doxycycline monohydrate tablets (50, 75, 100, and 150 mg)

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Doxycycline monohydrate tablets PA Criteria:

Standard NP Criteria

Minocycline ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Minocycline ER tablets PA Criteria:

- Patient is < 21 years old; **AND**
- Diagnosis of non-nodular moderate to severe acne vulgaris with inflammatory lesions; **AND**
- Patient requires long-term therapy with an oral tetracycline; **AND**
- Trial and failure, contraindication, or intolerance of TWO of the following topical agents:
  - Metronidazole (Metrogel®)
  - Azelaic acid (Azelex®, Finacea®)
  - Erythromycin (A/T/S® solution, gel)
  - Clindamycin (Cleocin T®)
  - Topical keratolytic agents (such as benzoyl peroxide, salicylic acid preparations); **AND**
- Clinically valid reason why the preferred minocycline capsules cannot be used

Minocycline tablets (50, 75, 100 mg)

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Minocycline tablets PA Criteria:

Standard NP Criteria

Nuzyra® tablets

PDL Status	NP
Quantity Limit	3/day; Max 14-day supply
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	14 days

Nuzyra® tablets PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- One of the following:
  - Community-acquired bacterial pneumonia (CABP); **AND**
    - Trial and failure to, contraindication, or resistance to TWO preferred standard of care agents for CABP (e.g., macrolide, doxycycline, a preferred fluoroquinolone, beta-lactam, linezolid)
  - Diagnosis of acute bacterial skin and skin structure infections (ABSSSI); **AND**
    - Trial and failure to, contraindication, or resistance to ONE preferred standard of care agents for ABSSSI (e.g., linezolid, clindamycin, doxycycline, SMX-TMP, vancomycin, cephalosporin, a preferred fluoroquinolone)
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Oracea® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Oracea® capsules PA Criteria:

- Diagnosis of inflammatory lesions (papules and pustules) of rosacea; **AND**
- Patient is < 21 years of age; **AND**
- Patient requires long-term therapy (greater than 3 months) with an oral antibiotic; **AND**
- Trial and failure, contraindication, or intolerance to ONE of the following topical agents:
  - Metronidazole (e.g., MetroGel®, MetroCream®)
  - Azelaic Acid (e.g., Azelex®, Finacea®)
  - Erythromycin gel or solution

Targadox® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Targadox® tablets PA Criteria:

Standard NP Criteria

Tetracycline tablets (250 and 500 mg)

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tetracycline tablets (250 and 500 mg) PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Antibiotics: UTI Agents, Miscellaneous

Fosfomycin powder packets

PDL Status	P
Quantity Limit	1 packet (3 g) per course of therapy
PA Criteria	See below
PA Form	General PA Form

Fosfomycin PA Criteria:

- Trial and failure, contraindication, intolerance, or resistance to at least 2 of the following agents:
  - Sulfamethoxazole/trimethoprim
  - Quinolones
  - Nitrofurantoin

Blujepa® tablets

PDL Status	NP
Quantity Limit	UTI: 20 tablets/course of therapy; Gonorrhea: 4 tablets/course of therapy
PA Criteria	See below
PA Form	General PA Form
Notes	Agent will not be approved for needle phobia

Uncomplicated Urinary Tract Infection

Blujepa® tablets PA Criteria:

- Diagnosis of uncomplicated urinary tract infection (uUTI); **AND**
- Patient is 12 years of age or older and weighs at least 40 kg; **AND**
- Infection has been confirmed to have been caused by one of the following:
  - Escherichia coli
  - Klebsiella pneumoniae
  - Citrobacter freundii complex
  - Staphylococcus saprophyticus
  - Enterococcus faecalis; **AND**
- Trial and failure, contraindication, intolerance, or resistance to at least 2 of the following agents:
  - Sulfamethoxazole/trimethoprim
  - Quinolones
  - Nitrofurantoin; **AND**
- Trial and failure, contraindication, intolerance, or resistance to the preferred fosfomycin agent; **AND**
- Prescribed by, or in consultation with, an infectious disease specialist

Uncomplicated Urogenital Gonorrhea

Blujepa® tablets PA Criteria:

- Diagnosis of uncomplicated urogenital gonorrhea; **AND**
- Patient is 12 years of age or older and weighs at least 45kg; **AND**
- Clinically valid reason why patient cannot use ONE of the following regimens:
  - IM ceftriaxone
  - IM cefoxitin
  - Oral cefixime
  - An alternative combination regimen with IM spectinomycin or gentamicin plus oral azithromycin; **AND**
- Prescribed by, or in consultation with, an infectious disease specialist; **AND**
- Prescriber attests that patient has no alternative options for the treatment of uncomplicated urogenital gonorrhea caused by susceptible strains of Neisseria gonorrhoeae

Orlynvah® tablets

PDL Status	NP
Quantity Limit	10 tablets/course of therapy
PA Criteria	See below
PA Form	General PA Form
Notes	Reserve use for infections caused by drug-resistant pathogens with limited treatment options.

Orlynvah® tablets PA Criteria:

- Diagnosis of uncomplicated urinary tract infection (uUTI); **AND**
- Patient is 18 years of age or older; **AND**
- Infection has been confirmed to have been caused by one of the following:
  - Escherichia coli
  - Klebsiella pneumoniae
  - Proteus mirabilis; **AND**
- Trial and failure, contraindication, intolerance, or resistance to at least 2 of the following agents:
  - Sulfamethoxazole/trimethoprim
  - Quinolones
  - Nitrofurantoin; **AND**
- Trial and failure, contraindication, intolerance, or resistance to the preferred fosfomycin agent; **AND**
- Prescribed by, or in consultation with, an infectious disease specialist

Pivya® tablets

PDL Status	NP
Quantity Limit	21 tabs/course of therapy
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	1 week

Pivya® tablets Initial PA Criteria:

- Patient is female and at least 18 years of age or older; **AND**
- Diagnosis of uncomplicated urinary tract infection (uUTI); **AND**
- Infection is caused by one of the following:
  - E. coli
  - Proteus mirabilis
  - Staphylococcus saprophyticus; **AND**
- Trial and failure, contraindication, intolerance, or submission of lab documented resistance to at least 1 of the following agents:
  - Sulfamethoxazole/trimethoprim
  - Quinolones
  - Nitrofurantoin; **AND**
- Trial and failure, contraindication, intolerance, or submission of lab documented resistance to the preferred fosfomycin agent; **AND**
- Lab documentation demonstrating both sensitivity to Pivya

Antibiotics, Vaginal

Clindamycin phosphate 2% vaginal cream

PDL Status	P
Quantity Limit	40 g/Rx
PA Criteria	None

Metronidazole 0.75% vaginal gel

PDL Status	P
Quantity Limit	70 g/Rx
PA Criteria	None

Nuvessa®

PDL Status	P
Quantity Limit	5 g/Rx
PA Criteria	None

Vandazole®

PDL Status	P
Quantity Limit	70 g/Rx
PA Criteria	None

Cleocin® vaginal cream

PDL Status	NP
Quantity Limit	40 g/Rx
PA Criteria	See below
PA Form	General PA Form

Cleocin® vaginal cream PA Criteria:

- Clinically valid reason why generic clindamycin vaginal cream cannot be used

Clindesse® vaginal cream

PDL Status	NP
Quantity Limit	5 g/Rx
PA Criteria	See below
PA Form	General PA Form

Clindesse® vaginal cream PA Criteria:

Standard NP Criteria

Xaciato®

PDL Status	NP
Quantity Limit	8 g/Rx
PA Criteria	See below
PA Form	General PA Form

Xaciato® PA Criteria:

Standard NP Criteria

Antifungals, Oral

Fluconazole suspension

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	PA not required for members < 20 years old

Fluconazole suspension PA Criteria:

- Patient is unable to swallow solid dosage forms; **OR**
- Patient ≤ 20 years of age

Fluconazole tablets

PDL Status	P
Quantity Limit	150 mg tablets: 4/28 days No quantity limit on other strengths
PA Criteria	Not applicable

Itraconazole capsules

PDL Status	P
Quantity Limit	4/day
PA Criteria	Not applicable

Itraconazole solution

PDL Status	P
Quantity Limit	40 mL/day
PA Criteria	See below
PA Form	General PA Form

Itraconazole solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Terbinafine tablets

PDL Status	P
Quantity Limit	84/year
PA Criteria	Not applicable

Ancobon® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Ancobon® capsules PA Criteria:

- Diagnosis of systemic candidiasis or cryptococcosis; **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Brexafemme® tablets

PDL Status	NP
Quantity Limit	4 tabs/Rx
PA Criteria	See below
PA Form	General PA Form

Brexafemme® tablets PA Criteria:

- Diagnosis of vulvovaginal candidiasis; **AND**
- One of the following:
  - Patient is ≥ 18 years of age
  - Patient is a post-menarchal female; **AND**
- Patient is not pregnant; **AND**
- Trial and failure, contraindication, or intolerance to 1 preferred oral agent (fluconazole tablets) **OR** 1 preferred topical agent (miconazole-3 or terconazole)

Cresemba® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Cresemba® capsules PA Criteria:

- Patient is > 6 years of age; **AND**
- Diagnosis of one of the following:
  - Invasive aspergillosis; **AND**
    - Trial and failure, contraindication, or intolerance to voriconazole **OR** posaconazole
  - Invasive mucormycosis; **AND**
- A fungal culture and relevant laboratory study (including histopathology) has been obtained to isolate and identify the causative organism(s); **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Diflucan® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Diflucan® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Diflucan® tablets

PDL Status	NP
Quantity Limit	150 mg tablets: 4/28 days No quantity limit on other strengths
PA Criteria	See below
PA Form	General PA Form

Diflucan® tablets PA Criteria:

Standard NP Criteria.

Flucytosine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Flucytosine capsules PA Criteria:

- Diagnosis of systemic candidiasis or cryptococcosis; **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Fulvicin P/G® 165 mg tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Fulvicin P/G® 165 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Griseofulvin ultramicrosize 165 mg tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Griseofulvin ultramicrosize 165 mg tablets PA Criteria:

- Diagnosis of ONE of the following:
  - Tinea corporis
  - Tinea pedis
  - Tinea cruris
  - Tinea barbae
  - Tinea capitis
  - Tinea unguium; **AND**
- Clinically valid reason why griseofulvin ultramicrosize 125 or 250 mg tablets cannot be used; **AND**
- Trial and failure of two preferred oral antifungal products

Ketoconazole tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Ketoconazole tablets PA Criteria:

- Trial and failure, contraindication, or intolerance to TWO preferred agents; **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Noxafil® suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Noxafil® suspension, tablets PA Criteria:

- ONE of the following:
  - As indicated for the prophylaxis of invasive aspergillus and/or candida in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with Graft versus Host Disease (GVHD), recipients with hematologic malignancies (leukemia, lymphoma, myelodysplastic syndromes) with prolonged neutropenia from chemotherapy, or recipients with AIDS.
  - Treatment of Fusariosis disease
  - Treatment of Zygomycetes disease
  - Treatment of other fungal infections or molds that are refractory or resistant to, or in patient who have a contraindication, or intolerance to itraconazole or voriconazole
  - Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Noxafil® Pak

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	If started as an inpatient hospital regimen and this is a continuation of therapy, then the drug is approvable.

Noxafil® Pak PA Criteria:

- Indication is prophylaxis of invasive aspergillus and/or candida in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with Graft versus Host Disease (GVHD), recipients with hematologic malignancies (leukemia, lymphoma, myelodysplastic syndromes) with prolonged neutropenia from chemotherapy, or recipients with AIDS; **AND**
- Patient is $\geq$ 2 years old and weighs $\leq$ 40 kg

Oravig®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Oravig® PA Criteria:

- Patient is 18 years of age or older; **AND**
- Patient has a diagnosis of oropharyngeal candidiasis; **AND**
- Patient has a contraindication, allergic reaction, or drug-drug interaction to clotrimazole troche and nystatin

Posaconazole suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Posaconazole suspension, tablets PA Criteria:

- See Noxafil® suspension/tablet prior authorization criteria.

Tolsura®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Tolsura® PA Criteria:

- Diagnosis ONE of the following:
  - Aspergillosis (pulmonary and extrapulmonary)
  - Blastomycosis (pulmonary and extrapulmonary)
  - Histoplasmosis (including chronic cavitary pulmonary disease, disseminated, or nonmeningeal); **AND**
- Clinically valid reason why the patient cannot use the other itraconazole capsules or solution

Vfend®

PDL Status	NP
Quantity Limit	18/84 days
PA Criteria	See below
PA Form	General PA Form

Vfend® PA Criteria:

- Treatment is for ONE of the following:
  - Candidemia (in non-neutropenic patients)
  - Esophageal candidiasis
  - Invasive aspergillosis
  - Serious fungal infections caused by *S. apiospermum* and *Fusarium* species including *F. solani*
  - Part of standard anti-fungal regimen in febrile neutropenic patients
  - Other fungal infections that are refractory or resistant to other oral triazole agents (i.e., fluconazole, ketoconazole); **OR**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate completion of therapy)

Vivjoa®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vivjoa® PA Criteria:

- Diagnosis of recurrent vulvovaginal candidiasis (RVCC); **AND**
- Provider attests patient is NOT of reproductive potential; **AND**
- The member has experienced ≥ 3 episodes of VVC in less than one year; **AND**
- Failure of a maintenance course of oral fluconazole defined as 100-mg, 150-mg, or 200-mg taken weekly for 6-months

Voriconazole

PDL Status	NP
Quantity Limit	18/84 days
PA Criteria	See below
PA Form	General PA Form

Voriconazole PA Criteria:

- See Vfend® prior authorization criteria

Antifungals, Vaginal

Please refer to the OTC Drug Formulary for coverage of OTC products.

Gynazole-1®

PDL Status	P
Quantity Limit	5 g/day
PA Criteria	Not applicable

Miconazole-3 vaginal suppository

PDL Status	P
Quantity Limit	1 box/Rx
PA Criteria	Not applicable

Terconazole vaginal cream, suppository

PDL Status	P
Quantity Limit	1 box/Rx
PA Criteria	Not applicable

Anti-Infectives: Amebicides

Paromomycin

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Paromomycin PA Criteria:
Standard NP Criteria

Anti-Infectives: Anthelmintics

Please refer to the OTC Drug Formulary for coverage of OTC products.

Albendazole tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Albendazole tablets PA Criteria:

- Treatment of neurocysticercosis caused by Taenia solium; **AND**
- Prescribed by, or in consultation with, an Infectious Disease specialist; **OR**
- Treatment of cystic hydatid disease caused by Echinococcus granulosus; **OR**
- Treatment of hookworm

Ivermectin tablets

PDL Status	P
Quantity Limit	20/90 days
PA Criteria	Not applicable

Emverm® chewable tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	PA approval duration will be based on FDA indication.

Emverm® chewable tablets PA Criteria:

- Treatment of Enterobius vermicularis (pinworm) in single or mixed infections; **AND**
  - Patient has tried and failed, has an intolerance, **OR** contraindication to pyrantel pamoate; **OR**
- Treatment of Ancylostoma duodenale (common hookworm) or Necator americanus (American hookworm); **AND**
  - Patient has tried and failed, has an intolerance, **OR** contraindication to albendazole; **OR**
- Treatment of Trichuris trichiura (whipworm) or Ascaris lumbricoides (common roundworm); **AND**
  - Patient has tried and failed, has an intolerance, **OR** contraindication to ivermectin

Praziquantel tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Praziquantel tablets PA Criteria:

- Diagnosis of infection with Schistosoma species; **OR**
- Trial and failure, contraindication, or intolerance of two of the following:
  - Albendazole
  - Ivermectin tablets
  - Pyrantel pamoate (OTC)

Stromectol® tablets

PDL Status	NP
Quantity Limit	20/90 days
PA Criteria	See below
PA Form	General PA Form

Stromectol® tablets PA Criteria:

Standard NP Criteria.

Anti-Infectives: Antimalarials

Coartem®, Krintafel®, Malarone®, Qualaquin®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Coartem®, Krintafel®, Malarone®, Qualaquin® PA Criteria:

Standard NP Criteria

Anti-Infectives: Antiprotozoals, Oral

Atovaquone suspension

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Atovaquone suspension PA Criteria:

- Treatment is for Pneumocystis pneumonia (PCP) prevention or treatment; **AND**
  - Trial and failure, contraindication, intolerance to sulfamethoxazole/trimethoprim; **OR**
- Diagnosis of Toxoplasmosis gondii encephalitis; **AND**
  - Trial and failure, contraindication, intolerance to sulfamethoxazole/trimethoprim; **OR**
- Diagnosis of Babesiosis

Benznidazole tablets

PDL Status	NP
Quantity Limit	12.5 mg tabs: 6/day; 100 mg tabs: 4/day
PA Criteria	See below
PA Form	General PA Form

Benznidazole tablets PA Criteria:

- Diagnosis of American trypanosomiasis (Chagas disease) caused by Trypanosoma cruzi

Daraprim® tablets, Flagyl® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Daraprim® tablets, Flagyl® capsules PA Criteria:

Standard NP Criteria

Lampit® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lampit® tablets PA Criteria:

- Diagnosis of American trypanosomiasis (Chagas disease) caused by Trypanosoma cruzi

Likmez® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Likmez® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms; **OR**
- Patients less than 12 years of age

Mepron® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Mepron® suspension PA Criteria:

- See atovaquone prior authorization criteria; **AND**
- Trial and failure, contraindication, intolerance, or drug-drug interaction to sulfamethoxazole/trimethoprim

Metronidazole capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Metronidazole capsules PA Criteria:

Standard NP Criteria

Metronidazole tablets (125 mg)

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Metronidazole tablets (125 mg) PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Nitazoxanide tablets

PDL Status	NP
Quantity Limit	6/Rx
PA Criteria	See below
PA Form	General PA Form

Nitazoxanide tablets PA Criteria:

- Patient is > 12 years of age or older; **AND**
- One of the following:
- Treatment of diarrhea caused by Cryptosporidium parvum (Note: Will not be approved for the treatment of diarrhea caused by C. parvum in HIV-infected or immunodeficient patients); **OR**
- Treatment of diarrhea caused by Giardia lamblia; **AND**
- Patient has failed and failed, or has a contraindication, intolerance, or adverse drug reaction to tinidazole and metronidazole

Pyrimethamine tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pyrimethamine tablets PA Criteria:

- Treatment of toxoplasmosis when used in combination with a sulfonamide

Solosec® packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Solosec® packets PA Criteria:

- Patient is 12 years of age or older; **AND**
- One of the following:
  - Diagnosis of bacterial vaginosis; **AND**
  - Trial and failure, contraindication, or intolerance to one of the following:
    - Cleocin® vaginal cream
    - Cleocin® Ovules
    - clindamycin capsules
    - metronidazole tablets
    - metronidazole vaginal gel
  - Diagnosis of trichomoniasis caused by *Trichomonas vaginalis* (*T. vaginalis*);
    - Trial and failure, contraindication, or intolerance to preferred metronidazole tablets

Sulfadiazine tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sulfadiazine tablets PA Criteria:

- Treatment of Toxoplasma gondii encephalitis in combination with pyrimethamine; **OR**
- Rheumatic fever prophylaxis in patients who have a contraindication or intolerance to penicillin

Tinidazole tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tinidazole tablets PA Criteria:

Standard NP Criteria

Antivirals: COVID Treatment

Lagevrio®

PDL Status	P
Quantity Limit	40/5 days
PA Criteria	See below
PA Form	General PA Form

Lagevrio® PA Criteria:

- Patient is ≥ 18 years of age and older

Paxlovid®

PDL Status	P
Quantity Limit	30/5 days
PA Criteria	See below
PA Form	General PA Form

Paxlovid® PA Criteria:

- Patient is ≥ 12 years of age and older

Antivirals: Cytomegalovirus Agents

Livtencity® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Livtencity® tablets PA Criteria:

- Patient is ≥ 12 years of age and weighs ≥ 35kg; **AND**
- Diagnosis of post-transplant cytomegalovirus (CMV) infection; **AND**
- Infection is refractory to prior treatment with at least one of the following:
  - Ganciclovir, valganciclovir, cidofovir or foscarnet

Prevymis® PAK, tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
Notes	When co-administered with cyclosporine the recommended dose of Prevymis is 240 mg daily.

Prevymis® PAK, tablets PA Criteria:

- Patient is scheduled or has received an allogeneic hematopoietic stem cell transplant (HSCT) and ONE of the following:
- Patient is 6 months of age or older and weights at least 6 kg; **AND**
- One of the following:
  - Patient is seropositive for CMV; **OR**
  - Treatment is for prophylaxis against CMV disease; **OR**
- Patient is a kidney transplant recipient and **BOTH** of the following:

- Must be prescribed by, or in consultation with, an oncology, hematology, infectious disease, or transplant specialist; **AND**
- Patient is NOT receiving concurrent therapy with pimozide, ergot alkaloids, or cyclosporine in conjunction with either pitavastatin or simvastatin

Valcyte® solution, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Valcyte® solution, tablets PA Criteria:

Standard NP Criteria

Antivirals: Hepatitis B

Entecavir tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Lamivudine-HBV tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Tenofovir disoproxil fumarate tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Adefovir tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Adefovir tablets PA Criteria:
Standard NP Criteria.

Baraclude® solution

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See below
PA Form	General PA Form
Notes	Prior authorization is not required for patients 2 through 11 years of age.

Baraclude® solution PA Criteria:

- Diagnosis of chronic hepatitis B virus infection with evidence of active viral replication and either evidence of persistent elevations in serum aminotransferases (ALT or AST) or histologically active disease; **AND**
- Patient is unable to swallow solid dosage forms

Baraclude® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Baraclude® tablets PA Criteria:
Standard NP Criteria.

Vemlidy® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Vemlidy® tablets PA Criteria:

- Patient is 6 years of age and older; **AND**
- Diagnosis of Chronic Hepatitis B virus (HBV) infection in adults with compensated liver disease; **AND**
- Inadequate treatment response (detectable HBV DNA level after 24 weeks of therapy), virologic breakthrough, resistance, intolerance, or contraindication to entecavir; **AND**
- Patient has ONE of the following:
  - History of osteoporosis or osteopenia; **OR**
  - Creatinine clearance 15-50 mL/min or on hemodialysis; **OR**
  - Clinically valid reason as to why the preferred tenofovir disoproxil fumarate (TDF) cannot be used; **AND**
- Patient is not using Vemlidy® as monotherapy if (HIV)-1 positive (must have additional antiviral therapy if HIV-1 positive for coverage of both disease states)

Viread® powder

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	See Appropriate Diagnosis PA bypass list for patients ages 2-6 years (wt ≥ 10 kg) old who need Viread powder for PEP.

Viread® powder PA Criteria:

- Patient has had a trial and failure, contraindication, or intolerance to 2 preferred agents; **OR**
- Patient cannot swallow solid dosage forms; **OR**
- Patient is 2 to 6 years of age; **AND**
- Patient is being treated for post-exposure prophylaxis (PEP)

Viread® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Viread® tablets PA Criteria:

Standard NP Criteria.

Antivirals: Hepatitis C Antivirals

Epclusa® 200-50 mg tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Epclusa & Sofosbuvir/Velpatasvir PA Form
PA Approval Duration	12 weeks unless indicated otherwise below
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Chronic Hepatitis C Virus, Treatment Naïve, No Cirrhosis

Epclusa® 200-50 mg tablets PA Criteria:

- Diagnosis of Chronic Hepatitis C; **AND**
- Patient does not have cirrhosis; **AND**
- Patient is treatment naïve; **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Chronic Hepatitis C Virus, Treatment Naïve, Compensated Cirrhosis (Child Pugh A)

Epclusa® 200-50 mg tablets PA Criteria:

- Diagnosis of Chronic Hepatitis C; **AND**
- Patient is treatment naïve; **AND**
- Patient has compensated cirrhosis (Child Pugh A); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Chronic Hepatitis C Virus, Treatment Naïve, Decompensated Cirrhosis (Child Pugh B or C)

Epclusa® 200-50 mg tablets PA Criteria:

- Diagnosis of Chronic Hepatitis C; **AND**
- Patient has decompensated Cirrhosis (Child Pugh B or C); **AND**
- Prescribed by or in consultation with a physician specialist with the experience in the treatment of Hepatitis C (e.g. hepatologist, Infectious Disease, or gastroenterology); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary; **AND**
- One of the following:
  - Will be given in combination with ribavirin (Total duration – 12 weeks); **OR**
  - Patient is ribavirin ineligible (Total duration – 24 weeks);

Chronic Hepatitis C Virus~~,~~ Treatment Experienced

Epclusa® 200-50 mg tablets PA Criteria:

- If patient has a history of HBV co-infection, prior history with direct acting hepatitis C antivirals, or decompensated cirrhosis, authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease, or Gastroenterology); **AND**
- Patients requiring retreatment of HCV or 2nd course of therapy with Hepatitis C Direct Acting Antiviral, requires escalation and documentation of ALL the following:
  - Requested HCV treatment regimen is recommended by the AASLD/IDSA guidelines for treatment-experienced patients (HCV Guidance - Treatment Experienced);
  - Current quantitative HCV RNA levels;
  - Quantitative HCV RNA level measured 12 weeks after completion of previous treatment;
  - Previous treatment history;
  - Genotype testing from current and previous (if available, from previous) infections;
    - If patient has Genotype 3 HCV, then documentation is submitted showing ONE of the following:
      - Without baseline resistance to NS5A RAS Y93H (Total duration – 12 weeks); **OR**
      - With baseline resistance to NS5A RAS Y93H AND given in combination with ribavirin (Total duration – 12 weeks); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Ledipasvir/sofosbuvir 90/400 mg tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Harvoni PA Form
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Ledipasvir/sofosbuvir 90/400 mg tablets PA Criteria:

- One of the following:
  - Diagnosis of Chronic Hepatitis C, genotype 1
    - Patients without cirrhosis:
      - Treatment naïve patients with documentation of pre-treatment HCV RNA < 6 million IU/mL (Total duration – 8 weeks)
      - Treatment naïve patients with documentation of pre-treatment HCV RNA > 6 million IU/mL (Total duration – 12 weeks)
      - Liver or kidney transplant patient (Total duration – 12 weeks); **OR**
    - Patients with compensated cirrhosis (Child-Pugh A):
      - Treatment naïve patients (Total duration – 12 weeks)
      - Liver or kidney transplant patient (Total duration – 12 weeks); **OR**
    - Patients with decompensated cirrhosis (Child-Pugh B or C):
      - Given in combination with ribavirin (Total duration – 12 weeks)
      - If ribavirin ineligible, may take as monotherapy (Total duration – 24 weeks); **OR**
  - Diagnosis of Chronic Hepatitis C, genotype 4, 5, 6
    - Treatment naïve patients with **OR** without compensated cirrhosis (Child-Pugh A) (Total Duration- 12 weeks)
    - Liver or kidney transplant patient with or without compensated cirrhosis (Child-Pugh A) (Total duration – 12 weeks)
    - Patients with decompensated cirrhosis (Child-Pugh B or C)
      - Given in combination with ribavirin (Total duration – 12 weeks)
      - If ribavirin ineligible, may take as monotherapy (Total duration – 24 weeks); **AND**
- If patient has a history of HBV co-infection, prior history with direct acting hepatitis C antivirals, or decompensated cirrhosis, authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease or Gastroenterology); **AND**
- Patients requiring retreatment of HCV or 2nd course of therapy with Hepatitis C Direct Acting Antiviral, requires escalation and documentation of ALL the following:
  - Requested HCV treatment regimen is recommended by the AASLD/IDSA guidelines for treatment-experienced patients (**HCV Guidance - Treatment Experienced**)
  - Current quantitative HCV RNA levels
  - Quantitative HCV RNA level measured 12 weeks after completion of previous treatment
  - Previous treatment history
  - Genotype testing from current and previous infections; **AND**
- Patient has been screened for Hepatitis B prior to treatment with a direct-acting antiviral agent for Chronic Hepatitis C

Mavyret® tablets (100/40 mg)

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	Mavyret PA Form
PA Approval Duration	8 weeks unless indicated otherwise below
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Acute or Chronic Hepatitis C Virus, Treatment Naïve, No Cirrhosis

Mavyret® tablets PA Criteria:

- Diagnosis of Chronic or Acute Hepatitis C; **AND**
- Patient is treatment naïve; **AND**
- Patient does not have cirrhosis; **AND**
- ONE of the following:
  - Patient has not received a liver or kidney transplant (Total duration- 8 weeks); **OR**
  - Liver or kidney transplant recipients (Total duration – 12 weeks); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Acute or Chronic Hepatitis C Virus, Treatment Naïve, Compensated Cirrhosis (Child Pugh A)

Mavyret® tablets PA Criteria:

- Diagnosis of Chronic or Acute Hepatitis C; **AND**
- Patient is treatment naïve; **AND**
- Patient has compensated cirrhosis (Child Pugh A); **AND**
- ONE of the following:
  - Patient has not received a liver or kidney transplant (Total duration- 8 weeks); **OR**
  - Liver or kidney transplant recipients (Total duration – 12 weeks); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Acute or Chronic Hepatitis C Virus, Treatment Naïve, Decompensated Cirrhosis (Child Pugh B or C)

Mavyret® tablets PA Criteria:

- Diagnosis of Chronic or Acute Hepatitis C; **AND**
- Patient is treatment naïve; **AND**
- Patient has decompensated cirrhosis (Child Pugh B or C); **AND**
- ONE of the following:
  - Patient has not received a liver or kidney transplant (Total duration- 8 weeks); **OR**
  - Liver or kidney transplant recipients (Total duration – 12 weeks); **AND**
- Authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease, or Gastroenterology); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Acute or Chronic Hepatitis C Virus, Treatment Experienced

Mavyret® tablets PA Criteria:

- Diagnosis of Chronic or Acute Hepatitis C; **AND**
- Patient is treatment experienced; **AND**
- Patients requiring retreatment of HCV or 2nd course of therapy with Hepatitis C Direct Acting Antiviral, requires escalation and documentation of ALL the following:
  - Requested HCV treatment regimen is recommended by the AASLD/IDSA guidelines for treatment-experienced patients (**HCV Guidance - Treatment Experienced**)
  - Current quantitative HCV RNA levels
  - Quantitative HCV RNA level measured 12 weeks after completion of previous treatment
  - Previous treatment history
  - Genotype testing from current and previous (if available from previous) infections;
  - Authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease or Gastroenterology); **AND**
- Prescriber attests that patient has been or will be screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C and treated appropriately if necessary

Mavyret® pellet (50/20 mg)

PDL Status	P
Quantity Limit	5/day
PA Criteria	See below
PA Form	**Mavyret PA Form**
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Mavyret® pellet PA Criteria:

See Mavyret® tablet prior authorization criteria; **AND**

- Patient is unable to swallow tablets

Sofosbuvir/velpatasvir tablets (400/100 mg)

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	**Epclusa & Sofosbuvir/Velpatasvir PA Form**
PA Approval Duration	12 weeks unless indicated otherwise below
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Sofosbuvir/velpatasvir tablets PA Criteria:

See Epclusa® 200-50 mg tablets prior authorization criteria.

Epclusa® pellets (150/37.5 mg and 200/50 mg)

PDL Status	NP
Quantity Limit	150 mg pellet: 1/day; 200 mg pellet: 2/day
PA Criteria	See below
PA Form	Epclusa & Sofosbuvir/Velpatasvir PA Form
PA Approval Duration	12 weeks unless indicated otherwise below
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Epclusa® pellets PA Criteria:

See Epclusa® 200-50 mg tablets prior authorization criteria; **AND**

- Patient is unable to swallow tablets

Epclusa® 400-100 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Epclusa & Sofosbuvir/Velpatasvir PA Form
PA Approval Duration	12 weeks unless indicated otherwise below
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Epclusa® 400-100 mg tablets PA Criteria:

See Epclusa® 200-50 mg tablets prior authorization criteria; **AND**

- Clinically valid reason why sofosbuvir/velpatasvir 400/100 mg tablets cannot be used

Harvoni® pellets (33.75/150 mg and 45/200 mg)

PDL Status	NP
Quantity Limit	1 pack/28 days
PA Criteria	See below
PA Form	Harvoni PA Form
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Harvoni® pellets PA Criteria:

See ledipasvir/sofosbuvir tablet prior authorization criteria; **AND**

- Patient is unable to swallow tablets

Harvoni® tablets (90/400 mg and 45/200 mg)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Harvoni PA Form
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Harvoni® tablets PA Criteria:

See ledipasvir/sofosbuvir tablet prior authorization criteria; **AND**

- Clinically valid reason why ledipasvir/sofosbuvir tablet cannot be used

Sovaldi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Sovaldi PA Form
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Sovaldi® tablets PA Criteria:

- One of the following:
  - Diagnosis of Chronic Hepatitis C, genotype 1 or 4 (Total duration – 12 weeks)
    - Used in combination with ribavirin and peginterferon alfa; **OR**
    - Patient must have a contraindication or drug-drug interaction with two preferred agents;**OR**
    - Patients must be treatment naïve to all HCV therapy (including therapies with pegylated interferon or ribavirin); **OR**
    - If patient has a documented contraindication to interferon; may use in combination with ribavirin alone (Total duration – 24 weeks); **AND**
  - Diagnosis of Chronic Hepatitis C, genotype 2 (Total duration – 12 weeks):
    - Treatment-naïve and treatment-experienced with or without cirrhosis (Child-Pugh A); **AND**
    - Requires contraindication or drug-drug interaction with two preferred agents; **AND**
    - Used in combination with ribavirin
  - Diagnosis of Chronic Hepatitis C, genotype 3 (Total duration – 24 weeks):
    - Treatment-naïve and treatment-experienced without cirrhosis or with compensated cirrhosis (Child-Pugh A); **AND**
    - Requires contraindication or drug-drug interaction with Mavyret and Epclusa; **AND**
    - Used in combination with ribavirin
- If patient has a history of HBV co-infection, prior history with direct acting hepatitis C antivirals, or decompensated cirrhosis, authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease, or Gastroenterology); **AND**
- Patients requiring retreatment of HCV or 2nd course of therapy with Hepatitis C Direct Acting Antiviral, requires escalation and documentation of ALL the following:
  - Requested HCV treatment regimen is recommended by the AASLD/IDSA guidelines for treatment-experienced patients (**HCV Guidance - Treatment Experienced**)
  - Current quantitative HCV RNA levels
  - Quantitative HCV RNA level measured 12 weeks after completion of previous treatment
  - Previous treatment history
  - Genotype testing from current and previous infections; **AND**
- Patient has been screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C; **AND**
- If request is for diagnosis of Hepatocellular Carcinoma awaiting liver transplant (Length of authorization: 48 weeks), must be used in combination with ribavirin; **AND**
  - Must meet ALL Milan criteria, defined as:
    - The presence of a tumor 5 cm or less in diameter in subjects with single hepatocellular carcinoma
    - No more than three tumor nodules, each 3 cm or less in diameter, in subjects with multiple tumors
    - No extrahepatic manifestations of the cancer and no evidence of vascular invasion of the tumor

Sovaldi® pellets

PDL Status	NP
Quantity Limit	1 pack/28 days
PA Criteria	See below
PA Form	Sovaldi PA Form
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Sovaldi® pellets PA Criteria:

See Sovaldi® tablets prior authorization criteria; **AND**

- Patient is unable to swallow tablets

Vosevi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Vosevi PA Form.
Notes	Patients previously treated with one of the following are considered treatment-naïve: sofosbuvir+ daclatasvir, peginterferon alfa + ribavirin, paritaprevir/ritonavir/ombitasvir/dasabuvir, and telaprevir or boceprevir + pegylated interferon, ribavirin

Vosevi® tablets PA Criteria:

- Diagnosis of chronic Hepatitis C, genotype 1–6
  - Sofosbuvir- based treatment failures, with or without compensated cirrhosis (Total duration – 12 weeks); **OR**
  - Glecaprevir/Pibrentasvir treatment failure with or without compensated cirrhosis (Total duration – 12 weeks); **OR**
  - Multiple Direct-Acting Antiviral (DAA) treatment failures in combination with weight-based ribavirin (Total duration- 24 weeks); **AND**
- If patient has a history of HBV co-infection, prior history with direct acting hepatitis C antivirals, or decompensated cirrhosis, authorization is being requested by or in consultation with a physician specialist with experience in the treatment of hepatitis C infection (e.g., Hepatology, Infectious Disease or Gastroenterology); **AND**
- Patients requiring retreatment of HCV or 2nd course of therapy with Hepatitis C Direct Acting Antiviral, requires escalation and documentation of all the following:
  - Requested HCV treatment regimen is recommended by the AASLD/IDSA guidelines for treatment-experienced patients (**HCV Guidance - Treatment Experienced**)
  - Current quantitative HCV RNA levels
  - Quantitative HCV RNA level measured 12 weeks after completion of previous treatment
  - Previous treatment history
  - Genotype testing from current and previous infections; **AND**
- Patient does not have, nor has ever had, decompensated cirrhosis [Child-Pugh score greater than 6 (class B or C)]; **AND**
- Patient has been screened for Hepatitis B prior to treatment with any direct-acting antiviral agent for Chronic Hepatitis C

Antivirals: Hepatitis C Pegylated Interferons

Pegasys® syringes

PDL Status	P
Quantity Limit	4/24 days
PA Criteria	See below
PA Form	General PA Form
Notes	Prior authorization will be required after 24 weeks of therapy.

Pegasys® syringes PA Criteria:

- Diagnosis of ONE of the following:
- Chronic Hepatitis C and one of the following:
  - Adult Patients: In combination therapy with other hepatitis C virus drugs for adults with compensated liver disease. Pegasys monotherapy is indicated only if patient has contraindication or significant intolerance to other Hepatitis C drugs
  - Pediatric Patients: In combination with ribavirin for pediatric patients 5 years of age and older with compensated liver disease
- Chronic Hepatitis B and one of the following:
- Adult Patients: Treatment of adults with HBeAg-positive and HBeAg-negative chronic hepatitis B (CHB) infection who have compensated liver disease and evidence of viral replication and liver inflammation; **OR**
- Pediatric Patients: Treatment of non-cirrhotic pediatric patients 3 years of age and older with HBeAg-positive CHB and evidence of viral replication and elevations in serum alanine aminotransferase (ALT)

Pegasys® vials

PDL Status	P
Quantity Limit	4/24 days
PA Criteria	See below
PA Form	General PA Form
Notes	Prior authorization will be required after 24 weeks of therapy.

Pegasys® vials PA Criteria:

See Pegasys® syringes prior authorization criteria.

Antivirals: Heptatitis Delta

Hepcludex® injection

PDL Status	NP
Quantity Limit	1 vial/day
PA Criteria	See below
PA Form	General PA Form

Hepcludex® injection Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of Chronic Hepatitis Delta virus (HDV) infection in adults without cirrhosis OR with compensated liver disease; **AND**
- Patient exhibits detectable serum HDV RNA levels as confirmed by quantitative assay in past 6 months; **AND**
- Prescriber attests that patient will be given the first dose under healthcare provider supervision; **AND**
- Prescriber attests that patient or caregiver will be counseled on proper dilution and administration technique; **AND**
- Prescribed by or in consultation with a hepatologist, gastroenterologist, or infectious disease specialist

Hepcludex® vials Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g. undetectable HDV RNA or ≥ 2 log10 IU/mL decline from baseline, ALT normalization); **AND**
- Patient is not experiencing any unacceptable toxicities from the drug (e.g., severe hypersensitivity reactions, headache, pruritis, abdominal pain, injection site reactions, etc.)

Antivirals: Herpes Agents, Oral

Famciclovir tablets

PDL Status	P
Quantity Limit	125 mg tablets: 20/30 days; 250 mg tablets: 60/30 days; 500 mg tablets: 3/day and 21/Rx
PA Criteria	None

Valacyclovir tablets

PDL Status	P
Quantity Limit	500 mg tablets: 60/30 days; 1000 mg tablets: 30/Rx
PA Criteria	None

Sitavig® buccal tablets

PDL Status	NP
Quantity Limit	2/Rx
PA Criteria	See below
PA Form	General PA Form

Sitavig® buccal tablets Criteria:

Standard NP Criteria.

Valtrex® tablets

PDL Status	NP
Quantity Limit	500 mg tablets: 60/30 days; 1000 mg tablets: 30/Rx
PA Criteria	See below
PA Form	General PA Form

Valtrex® tablets Criteria:
Standard NP Criteria.

Antivirals: HIV Attachment Inhibitors

Rukobia® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Rukobia® tablets Initial Criteria:

- Diagnosis of treatment-experienced multidrug-resistant HIV-1 infection; **AND**
- HIV-1 RNA levels > 200 copies/mL; **AND**
- Prescriber attests that the patient lacks sufficient treatment options due to resistance, intolerability, contraindication, or other safety concerns to construct a fully suppressive antiretroviral regimen; **AND**
- Will not be used with strong cytochrome P450 (CYP)3A inducers; **AND**
- Prescribed by, or in consultation with or by an infectious disease specialist

Rukobia® tablets Renewal Criteria:

- Patient demonstrates documented efficacy (e.g., reduced viral load/improved CD4, remain virologically suppressed)

Antivirals: HIV CCR5 Antagonists

Maraviroc tablets

PDL Status	P
Quantity Limit	150 mg tablets: 2/day; 300 mg tablets: 4/day
PA Criteria	See below
PA Form	General PA Form

Maraviroc tablets PA Criteria:

- Diagnosis of CCR5-tropic HIV-1 via a co-receptor tropism; **AND**
- Verification that agent will be administered in combination with other antiretroviral agents

Selzentry® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Selzentry® solution PA Criteria:

- Diagnosis of CCR5-tropic HIV-1 via a co-receptor tropism; **AND**
- Verification that agent will be administered in combination with other antiretroviral agents; **AND**
- Patient is 11 years of age or younger; **OR**
- Patient is unable to swallow tablets

Selzentry® tablets

PDL Status	NP
Quantity Limit	150 mg tablets: 2/day; 300 mg tablets: 4/day
PA Criteria	See below
PA Form	General PA Form

Selzentry® tablets PA Criteria:

- Diagnosis of CCR5-tropic HIV-1 via a co-receptor tropism; **AND**
- Verification that agent will be administered in combination with other antiretroviral agents; **AND**
- Trial and failure, contraindication, or intolerance to maraviroc tablets

Antivirals: HIV Fusion Inhibitors

Fuzeon®

PDL Status	P
Quantity Limit	1 kit/30 days (2 vials/day)
PA Criteria	See below
PA Form	General PA Form

Fuzeon® Initial Criteria:

- Diagnosis of treatment-experienced multidrug-resistant HIV-1 infection; **AND**
- HIV-1 RNA levels > 200 copies/mL; **AND**
- Prescriber attests that the patient lacks sufficient treatment options due to resistance, intolerability, contraindication, or other safety concerns to construct a fully suppressive antiretroviral regimen; **AND**
- Agent will be used in combination with an optimized antiretroviral regimen therapy (ART); **AND**
- Prescribed by, or in consultation with or by an infectious disease specialist

Fuzeon® Renewal Criteria:

- Patient demonstrates documented efficacy (e.g., reduced viral load/improved CD4, remain virologically suppressed)

Antivirals: HIV Integrase Inhibitors and Combinations

Isentress® chews, granules, tablets

PDL Status	P
Quantity Limit	Chews: 6/day; Granules: 2 packs/day; Tablets: 2/day
PA Criteria	None

Tivicay®

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Tivicay PD®

PDL Status	P
Quantity Limit	3 bottles/30 days
PA Criteria	See below
PA Form	General PA Form

Tivicay PD® PA Criteria:

- Patient is < 6 years of age; **OR**
- Patient is unable to swallow solid dosage forms; **OR**
- Clinically valid reason why the patient cannot use Tivicay tablets

Isentress HD®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Isentress HD® PA Criteria:

- Verification that agent will be administered in combination with other antiretroviral agents; **AND**
- Clinically valid reason why the patient cannot use the preferred agents

Juluca®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Juluca® PA Criteria:

- Patient has a diagnosis of HIV; **AND**
- Patient does not have any prior history of treatment failure to other HIV agents **OR** known resistance to the individual components (dolutegravir/rilpivirine); **AND**
- Patient is virologically suppressed (HIV-1 RNA < 50 copies/mL) on a current ART regimen for ≥ 6-months

Antivirals: HIV NNRTIs

Efavirenz capsules, tablets

PDL Status	P
Quantity Limit	50 mg: 7/day; 200 mg: 2/day; 600 mg: 1/day
PA Criteria	None

Intelence® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Intelence® tablets PA Criteria:

- Patient is treatment-experienced; **AND**
- Patient will concomitantly take at least two additional antiretroviral agents; **AND**
- Patient has documented non-nucleoside reverse transcriptase inhibitor (NNRTI) resistance

Nevirapine suspension, tablets

PDL Status	P
Quantity Limit	Suspension: 40 mL/day 200 mg tablets: 2/day;
PA Criteria	None

Pifeltro® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Etravirine tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Etravirine tablets PA Criteria:

See Intelence® prior authorization criteria; **AND**

- Clinically valid reason why preferred Intelence® cannot be used

Nevirapine ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Nevirapine ER tablets PA Criteria:

Standard NP Criteria

Rilpivirine tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Rilpivirine tablets PA Criteria:

- Clinically valid reason why preferred Edurant® tablets cannot be used

Antivirals: HIV NRTIs

Abacavir solution, tablets

PDL Status	P
Quantity Limit	Solution: 30 mL/day Tablets: 2/day;
PA Criteria	None

Emtricitabine

PDL Status	P
Quantity Limit	1/day;
PA Criteria	None

Lamivudine solution, tablets

PDL Status	P
Quantity Limit	Solution: 30 mL/day 100 & 300 mg tablets: 1/day 150 mg tablets: 2/day
PA Criteria	None

Zidovudine syrup, tablets

PDL Status	P
Quantity Limit	Syrup: 60 mL/day 100 mg tablets: 6/day 300 mg tablets: 2/day
PA Criteria	None

Emtriva® capsules, solution

PDL Status	NP
Quantity Limit	Caps: 1/day; Solution: 24 mL/day
PA Criteria	See below
PA Form	General PA Form

Emtriva® capsules, solution PA Criteria:
Standard NP Criteria

Emtriva® capsules, solution

PDL Status	NP
Quantity Limit	Caps: 1/day; Solution: 24 mL/day
PA Criteria	See below
PA Form	General PA Form

Emtriva® capsules, solution PA Criteria:
Standard NP Criteria

Epivir® capsules, solution

PDL Status	NP
Quantity Limit	150 mg caps: 2/day; 300 mg caps: 1/day Solution: 30 mL/day
PA Criteria	See below
PA Form	General PA Form

Epivir® capsules, solution PA Criteria:
Standard NP Criteria

Retrovir® syrup, tablet

PDL Status	NP
Quantity Limit	Syrup: 60 mL/day; 100 mg tabs: 6/day
PA Criteria	See below
PA Form	General PA Form

Retrovir® syrup, tablet PA Criteria:
Standard NP Criteria

Ziagen® solution, tablet

PDL Status	NP
Quantity Limit	Solution: 30 mL/day; Tabs: 2/day
PA Criteria	See below
PA Form	General PA Form

Ziagen® solution, tablet PA Criteria:
Standard NP Criteria

Antivirals: HIV NRTI Combos

Abacavir/lamivudine tablets, Biktarvy® tablets, Complera® tablets, Delstrigo® tablets, Descovy® tablets, Dovato® tablets, emtricitabine/tenofovir tablets, efavirenz/emtricitabine/tenofovir tablets, Genvoya® tablets

PDL Status	P
Quantity Limit	1/day;
PA Criteria	None

Lamivudine/zidovudine tablets

PDL Status	P
Quantity Limit	2/day;
PA Criteria	None

Odefsy® tablets, Stribild® tablets

PDL Status	P
Quantity Limit	1/day;
PA Criteria	None

Symtuza® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Symtuza® Initial PA Criteria:

- Patient has a diagnosis of HIV-1; **AND**
- Patient has no known substitutions associated with resistance to darunavir or tenofovir; **AND**
- One of the following:
  - Patient is ARV treatment-naïve; **OR**
  - Patient is ARV treatment-experienced and meets the following requirements:
    - Virologically suppressed (HIV-1 RNA < 50 copies/mL) on a stable ARV regimen for ≥ 6-months; **OR**
    - Patient is switching medication due to adverse effects or documented compliance issues due to pill burden or dose frequency

Symtuza® Renewal PA Criteria:

- Patient demonstrates documented efficacy (e.g., reduced viral load/improved CD4, remaining virologically suppressed)

Triumeq® tablets

PDL Status	P
Quantity Limit	1/day;
PA Criteria	None

Cimduo® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Cimduo® tablets PA Criteria:
Standard NP Criteria

Efavirenz/lamivudine/tenofovir tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Efavirenz/lamivudine/tenofovir tablets PA Criteria:
Standard NP Criteria

Rilpivirine/emtricitabine/tenofovir tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Rilpivirine/emtricitabine/tenofovir tablets PA Criteria:

- Clinically valid reason why Complera® cannot be used

Symfi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Symfi® tablets PA Criteria:
Standard NP Criteria

Triumeq PD® tablets for oral suspension

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Triumeq PD® tablets for oral suspension PA Criteria:
Standard NP Criteria

Truvada® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Truvada® tablets PA Criteria:
Standard NP Criteria

Antivirals: HIV Pharmacokinetic Enhancers

Ritonavir tablet

PDL Status	P
Quantity Limit	12/day;
PA Criteria	None

Norvir® packet

PDL Status	NP
Quantity Limit	12/day
PA Criteria	See below
PA Form	General PA Form
Notes	Norvir oral powder is dosed in increments of 100 mg; prescription should not be written for <100 mg increments

Norvir® packet PA Criteria:

- Patient has a diagnosis of HIV-1; **AND**
- The requested will be used in combination with other antiretroviral agents; **AND**
- One of the following:
  - Patient is ≤ 18 years of age; **OR**
  - Clinically valid reason why the preferred ritonavir tablets cannot be used

Norvir® tablet

PDL Status	NP
Quantity Limit	12/day;
PA Criteria	None

Norvir® tablet PA Criteria:

Standard NP Criteria.

Tybost®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
Notes	

Tybost® PA Criteria:

- Verification that agent will be administered in combination with Prezista® (darunavir) **OR** atazanavir; **AND**
- Patient has a contraindication or intolerance to ritonavir; **AND**
- Patient is not pregnant; **AND**
- Patient does not have renal impairment

Antivirals: HIV Protease Inhibitors

Atazanavir capsules

PDL Status	P
Quantity Limit	150 and 200 mg caps: 2/day; 300 mg: 1/day
PA Criteria	None

Darunavir tablets

PDL Status	P
Quantity Limit	800 mg tablets: 1/day; All other strengths: 2/day
PA Criteria	None

Evotaz® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Fosamprenavir tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Lopinavir/ritonavir tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Prezcobix® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Prezista® suspension

PDL Status	P
Quantity Limit	12 mL/day
PA Criteria	None

Reyataz® powder

PDL Status	P
Quantity Limit	5/day
PA Criteria	None

Viracept® tablets

PDL Status	P
Quantity Limit	250 mg tablets: 9/day; 625 mg tablets: 4/day
PA Criteria	None

Aptivus® capsules

PDL Status	P
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Aptivus® capsules PA Criteria:

- Confirmation patient has had previous exposure to at least one PI indicated for first-line therapy

Kaletra® solution, tablets

PDL Status	P
Quantity Limit	Solution: 15 mL/day Tablets: 6/day
PA Criteria	See below
PA Form	General PA Form

Kaletra® solution, tablets PA Criteria:
Standard NP Criteria

Prezista® tablets

PDL Status	P
Quantity Limit	800 mg tabs: 1/day All other strengths: 2/day
PA Criteria	See below
PA Form	General PA Form

Prezista® tablets PA Criteria:
Standard NP Criteria

Reyataz® capsules

PDL Status	P
Quantity Limit	200 mg caps: 2/day; 300 mg caps: 1/day
PA Criteria	See below
PA Form	General PA Form

Reyataz® capsules PA Criteria:
Standard NP Criteria

Antivirals: Influenza

Oseltamivir capsules, suspension

PDL Status	P
Quantity Limit	Caps: 20/180 days; Susp: 300 mL/180 days
PA Criteria	None

Relenza®

PDL Status	P
Quantity Limit	40/180 days
PA Criteria	None

Tamiflu® capsules, suspension

PDL Status	NP
Quantity Limit	Caps: 20/180 days; Susp: 300 mL/180 days
PA Criteria	See below
PA Form	Influenza Antiviral PA Form.

Tamiflu® capsules, suspension PA Criteria:
Standard NP Criteria

Xofluza® tablets

PDL Status	NP
Quantity Limit	2/Rx
PA Criteria	See below
PA Form	Influenza Antiviral PA Form.

Xofluza® tablets PA Criteria:

- Agent is being used for treatment of influenza **OR** post-exposure prophylaxis of influenza; **AND**
- Treatment is being used for ONE of the following:
  - Acute uncomplicated influenza in patients ≥ 5 years of age who have been symptomatic for no more than 48 hours and who are otherwise healthy
  - Acute uncomplicated influenza in patients ≥ 5 years of age who are at high risk of developing influenza-related complications
  - Post-exposure prophylaxis of influenza in patients > 5 years of age; **AND**
- One of the following:
  - Contraindication to both Relenza® and Tamiflu® that is not associated with requested agent
  - Area surveillance data that indicates an oseltamivir resistant strain
  - Recurrent documented influenza in the same flu season that was previously treated with a preferred agent

III. CARDIOVASCULAR

ACE Inhibitors (ACEIs)

Ramipril capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Altace® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Altace® capsules PA Criteria:

Standard NP Criteria

Captopril tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	PA is not required for members 18 years of age and younger

Captopril tablets PA Criteria:

Standard NP Criteria

Enalapril solution, Epaned® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	PA is not required for members 8 years of age and younger

Enalapril solution, Epaned® solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Fosinopril tablets, Lotensin® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Fosinopril tablets, Lotensin® tablets PA Criteria:

Standard NP Criteria

Moexipril tablets

PDL Status	NP
Quantity Limit	7.5 mg: 1/day; 15 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Moexipril tablets PA Criteria:

Standard NP Criteria

Perindopril tablets

PDL Status	NP
Quantity Limit	2 mg, 4 mg: 1/day; 8 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Perindopril tablets PA Criteria:

Standard NP Criteria

Qbrelis® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	PA is not required for members 7 years of age and younger

Qbrelis® solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Quinapril tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Quinapril tablets PA Criteria:
Standard NP Criteria

Trandolapril tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Trandolapril tablets PA Criteria:
Standard NP Criteria

Zestril® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Zestril® tablets PA Criteria:
Standard NP Criteria

ACEIs + Calcium Channel Blockers

Benazepril/amlodipine capsules

PDL Status	P
Quantity Limit	5/40 mg: 2/day; All others: 1/day
PA Criteria	Not applicable

Lotrel® capsules

PDL Status	NP
Quantity Limit	5/40 mg: 2/day; All others: 1/day
PA Criteria	See Below
PA Form	General PA Form

Lotrel® capsules PA Criteria:

- Patient is unable to take the two components separately

Prestalia® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Prestalia® tablets PA Criteria:

- Patient is unable to take the two components separately

Trandolapril/verapamil tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Trandolapril/verapamil tablets PA Criteria:

- Patient is unable to take the two components separately

ACEIs + Diuretics

Accuretic® tablets, captopril/HCTZ tablets, fosinopril/HCTZ tablets, Lotensin HCT® tablets, quinapril/HCTZ tablets, Zestoretic® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Accuretic®, captopril/HCTZ, fosinopril/HCTZ, Lotensin HCT®, quinapril/HCTZ, Zestoretic® PA Criteria:
Standard NP Criteria

Alpha Blockers

Cardura® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cardura® tablets PA Criteria:
Standard NP Criteria

Alpha-Beta Blockers

Carvedilol

PDL Status	P
Quantity Limit	2/day
PA Criteria	Not applicable

Carvedilol ER

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Carvedilol ER PA Criteria:
Standard NP Criteria

Angiotensin II Receptor Blockers (ARBs)

Irbesartan tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	Not applicable

Losartan tablets

PDL Status	P
Quantity Limit	25 mg, 100 mg: 1/day; 50 mg: 2/day
PA Criteria	Not applicable

Olmesartan tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	Not applicable

Valsartan tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	Not applicable

Atacand® tablets, Avapro® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Atacand® tablets, Avapro® tablets PA Criteria:
Standard NP Criteria

Azilsartan tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Azilsartan tablets PA Criteria:
Standard NP Criteria; **AND**

- Clinically valid reason why brand Edarbi® tablets cannot be used

Benicar® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Benicar® tablets PA Criteria:
Standard NP Criteria

Candesartan tablets

PDL Status	NP
Quantity Limit	4 & 32 mg: 1/day; 8 & 16 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Candesartan tablets PA Criteria:
Standard NP Criteria

Cozaar® tablets

PDL Status	NP
Quantity Limit	25 mg, 100 mg: 1/day; 50 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Cozaar® tablets PA Criteria:
Standard NP Criteria

Diovan® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Diovan® tablets PA Criteria:
Standard NP Criteria

Edarbi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Edarbi® tablets PA Criteria:
Standard NP Criteria

Micardis® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Micardis® tablets PA Criteria:
Standard NP Criteria

Telmisartan tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Telmisartan tablets PA Criteria:
Standard NP Criteria

Valsartan solution

PDL Status	NP
Quantity Limit	80 mL/day
PA Criteria	See Below
PA Form	General PA Form

Valsartan solution PA Criteria:

- Patient is unable to swallow solid dosage forms

ARB + Calcium Channel Blocker

Valsartan/amlodipine, valsartan/amlodipine/HCTZ

PDL Status	P
Quantity Limit	1/day
PA Criteria	Not applicable

Azor®, Exforge®, Exforge HCT®, olmesartan/amlodipine

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Azor®, Exforge®, Exforge HCT®, olmesartan/amlodipine PA Criteria:
Standard NP Criteria

Olmesartan/amlodipine/HCTZ

PDL Status	NP
Quantity Limit	20/5/12.5 mg: 2/day; All others: 1/day
PA Criteria	See Below
PA Form	General PA Form

Olmesartan/amlodipine/HCTZ PA Criteria:
Standard NP Criteria

Telmisartan/amlodipine

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Telmisartan/amlodipine PA Criteria:
Standard NP Criteria

Tribenzor®

PDL Status	NP
Quantity Limit	20/5/12.5 mg: 2/day; All others: 1/day
PA Criteria	See Below
PA Form	General PA Form

Tribenzor® PA Criteria:
Standard NP Criteria

ARB + Diuretic

Irbesartan/HCTZ, Losartan/HCTZ, Olmesartan/HCTZ, & Valsartan/HCTZ

PDL Status	P
Quantity Limit	1/day
PA Criteria	Not applicable

Atacand HCT®, Avalide®, Benicar HCT®, candesartan/HCTZ, Diovan HCT®, Edarbyclor®, Hyzaar®, Micardis HCT®, telmisartan/HCTZ

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Atacand HCT®, Avalide®, Benicar HCT®, candesartan/HCTZ, Diovan HCT®, Edarbyclor®, Hyzaar®, Micardis HCT®, &telmisartan/HCTZ PA Criteria:
Standard NP Criteria

ARB + Neprilysin Inhibitor

Sacubitril/valsartan tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Sacubitril/valsartan® PA Criteria:

- Diagnosis of chronic heart failure (NYHA Class II-IV)

Entresto® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Entresto® tablets PA Criteria:

- Diagnosis of chronic heart failure (NYHA Class II-IV); **AND**
- Clinically valid reason why Entresto tablets cannot be used

Entresto® sprinkles

PDL Status	NP
Quantity Limit	8/day
PA Criteria	See Below
PA Form	General PA Form

Entresto® sprinkles PA Criteria:

- Diagnosis of chronic heart failure (NYHA Class II-IV); **AND**
- Clinically valid reason why sacubitril/valsartan tablets cannot be used

Antianginals: Nitrates

Rectiv®

PDL Status	P
Quantity Limit	1 tube/Rx
PA Criteria	See Below
PA Form	General PA Form

Rectiv® PA Criteria:

- Diagnosis of history of anal fissure; **AND**
- Patient is a candidate for surgery

Isordil®, isosorbide dinitrate 10 mg tablets Nitro-Bid®, Nitro-Dur®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Isordil®, isosorbide dinitrate 10 mg tablets Nitro-Bid®, Nitro-Dur® PA Criteria:
Standard NP Criteria

Nitroglycerin 0.4% ointment

PDL Status	NP
Quantity Limit	1 tube/Rx
PA Criteria	See Below
PA Form	General PA Form

Nitroglycerin 0.4% oint PA Criteria:

- Diagnosis of history of anal fissure; **AND**
- Patient is a candidate for surgery; **AND**
- Clinically valid reason why brand Rectiv® cannot be used

Nitroglycerin spray

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Nitroglycerin spray PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred agents; **OR**
- Clinically valid reason why the preferred agent cannot be used

Antiarrhythmics, Oral

Dofetilide

PDL Status	P
Quantity Limit	2/day
PA Criteria	None
PA Form	General PA Form

Betapace®, Betapace-AF®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Betapace®, Betapace-AF® PA Criteria:
Standard NP Criteria

Multaq®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Multaq® PA Criteria:

- Not on concurrent Class I or III anti-arrhythmic agent; **AND**
- Not hospitalized for exacerbation of heart failure in past 30 days; **AND**
- Patient does not have NYHA class IIIb or IV heart failure; **AND**
- Trial and failure, contraindication, or intolerance of TWO of the following preferred antiarrhythmic agents: (Note: Requirement is waived if patient has structural heart disease)
  - amiodarone
  - flecainide
  - propafenone
  - sotalol

Norpace®, Pacerone®, propafenone ER, quinidine gluconate & sulfate

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Norpace®, Pacerone®, propafenone ER, quinidine gluconate & sulfate PA Criteria:
Standard NP Criteria

Sotylize®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	PA is not required for patients 8 years of age and younger

Sotylize® PA Criteria:

- Patient is unable to swallow tablets and capsules

Tikosyn®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Tikosyn® PA Criteria:
Standard NP Criteria

Anticoagulants, Injectable

Enoxaparin

PDL Status	P
Quantity Limit	2 injections/day
PA Criteria	None

Fondaparinux

PDL Status	P
Quantity Limit	1 injection/day
PA Criteria	None

Arixtra®

PDL Status	NP
Quantity Limit	1 injection/day
PA Criteria	See Below
PA Form	General PA Form

Arixtra® PA Criteria:
Standard NP Criteria

Fragmin®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Fragmin® PA Criteria:
Standard NP Criteria

Lovenox®

PDL Status	NP
Quantity Limit	2 syringes/day
PA Criteria	See Below
PA Form	General PA Form

Lovenox® PA Criteria:
Standard NP Criteria

Anticoagulants, Oral

Dabigatran capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Eliquis® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Rivaroxaban tablets (2.5 mg)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below

Xarelto® tablets (10 & 15 mg and starter pack)

PDL Status	P
Quantity Limit	2.5 & 15 mg: 2/day 10 & 20 mg: 1/day
PA Criteria	None

Eliquis Sprinkle®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Eliquis Sprinkle® PA Criteria:

- One of the following:
  - Patient has diagnosis of Venous Thromboembolism (VTE);
  - Recurrent VTE risk reduction; **AND**
- Prescriber attests patient has received > 5 days of initial anticoagulation therapy; **AND**
  - Patient weighs 2.6 kg to less than 4 kg

Eliquis® tablets for oral susp

PDL Status	NP
Quantity Limit	0.5 mg: 4 tablets/day; 1.5 mg: 12 tablets/day; 2 mg: 16 tablets/day
PA Criteria	See Below
PA Form	General PA Form

Eliquis® tablets for oral suspension PA Criteria:

- One of the following:
  - Patient has diagnosis of Venous Thromboembolism (VTE);
  - Recurrent VTE risk reduction; **AND**
- Prescriber attests patient has received > 5 days of initial anticoagulation therapy; **AND**
  - Patient weighs 4 kg to less than 35 kg

Pradaxa® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Pradaxa® capsules PA Criteria:

Standard NP Criteria

Pradaxa® packs

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Pradaxa® packs PA Criteria:

- Patient is unable to swallow sold dosage forms; **OR**
- Clinically valid reason why the patient cannot use Pradaxa® capsules

Savaysa®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Savaysa® PA Criteria:

- One of the following:
  - Diagnosis of non-valvular atrial fibrillation; **AND**
    - Documentation that CrCl NOT ≥ 95 mL/min as calculated by Cockcroft-Gault equation
  - Diagnosis of deep vein thrombosis or pulmonary embolism; **AND**
- Trial and failure, intolerance, or contraindication to Xarelto® and Pradaxa®

Xarelto® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Xarelto® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Xarelto® tablets (2.5 mg)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Xarelto® tablets PA Criteria:

- Clinically valid reason why generic rivaroxaban tablets cannot be used

Antihypertensives, Miscellaneous

Clonidine 0.05 mg tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Clonidine weekly patch

PDL Status	P
Quantity Limit	0.1, 0.2 mg: 4/28 days; 0.3 mg: Age ≤21 years: 4/28 days; Age >21 years: 8/28 days
PA Criteria	None

Clonidine 24hr ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Clonidine 24hr ER PA Criteria:
Standard NP Criteria

Javadin® solution

PDL Status	NP
Quantity Limit	4 bottles/month
PA Criteria	See Below
PA Form	General PA Form

Javadin® solution PA Criteria:

- One of the following:
  - Documentation of need for dosing that will not allow the use of a preferred clonidine product; **OR**
  - Patient is unable to swallow solid oral dosage forms

Minoxidil tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Note	Minoxidil will not be approved for alopecia.

Minoxidil tablets PA Criteria:

- Diagnosis of severe hypertension (symptomatic or associated with target organ damage only); **AND**
- Trial and failure, contraindication, or intolerance to TWO of the following:
  - ACEI or ARBs
  - Calcium channel blockers
  - Methyldopa
  - Clonidine; **AND**
- Patient is concomitantly taking a diuretic (e.g., hydrochlorothiazide, chlorthalidone, furosemide, etc.) unless contraindicated; **AND**
- Patient does not have diagnosis of pheochromocytoma (minoxidil may stimulate secretions of catecholamines from the tumor)

Beta Blockers

Metoprolol succinate ER

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Acebutolol capsules, Betapace®, betaxolol tablets, Bystolic®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Acebutolol capsules, Betapace®, betaxolol tablets, Bystolic® PA Criteria:
Standard NP Criteria

Hemangeol®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Hemangeol® ER PA Criteria:

- Diagnosis of Infantile Hemangioma; **AND**
- Clinically valid reason why the preferred propranolol solution cannot be used

Inderal LA®, Inderal XL®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Inderal LA®, Inderal XL® PA Criteria:
Standard NP Criteria

InnoPran XL®

PDL Status	NP
Quantity Limit	80 mg: 2/day; 120 mg: 1/day
PA Criteria	See Below
PA Form	General PA Form

InnoPran XL® solution PA Criteria:
Standard NP Criteria

Kapspargo Sprinkle®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Kapspargo Sprinkle® PA Criteria:

- Diagnosis of ONE of the following:
  - Heart Failure or LVEF ≤ 40%
  - Hypertension
  - Angina Pectoris; **AND**
- Patient is unable to swallow tablets and capsules

Lopressor® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Lopressor® tablets PA Criteria:
Standard NP Criteria

Lopressor® Oral Solution

PDL Status	NP
Quantity Limit	45 mL/day
PA Criteria	See Below
PA Form	General PA Form

Lopressor® Oral solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Pindolol, Tenormin®, timolol maleate tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Pindolol, Tenormin®, timolol maleate tablets PA Criteria:
Standard NP Criteria

Toprol XL®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Toprol XL® PA Criteria:
Standard NP Criteria

Beta Blockers + Diuretics

Tenoretic®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tenoretic® PA Criteria:
Standard NP Criteria

Calcium Channel Blockers (Dihydropyridine (DHP))

Amlodipine tablets

PDL Status	P
Quantity Limit	2.5 & 5 mg (1.5/day); 10 mg (1/day)
PA Criteria	None

Nifedipine ER 24HR tabs, nifedipine ER 24HR osmotic tabs

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Norliqva® solution

PDL Status	P
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Norliqva® solution PA Criteria:

- Diagnosis of one of the following:
  - Hypertension
  - Chronic stable angina or treatment
  - Vasospastic Angina (Prinzmetal's or Variant Angina)
  - Confirmed or suspected vasospastic angina
  - Angiographically documented Coronary Artery Disease in patients without heart failure or an ejection fraction < 40%; **AND**
- One of the following:
  - Patient is unable to swallow solid dosage forms; **OR**
  - Clinically valid reason why amlodipine tablets cannot be used

Cardamyst® nasal spray

PDL Status	Awaiting PAC Review
Quantity Limit	2 devices/Rx and 4 devices /month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	6 months

Cardamyst® nasal spray Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of paroxysmal supraventricular tachycardia (PSVT); **AND**
- Confirmation of failure, intolerance, or contraindication to oral preventative therapy (e.g., diltiazem, verapamil, beta-blockers), OR must be used concurrently; **AND**
- Patient has had prior sustained symptomatic episodes lasting greater than or equal to 20 minutes; **AND**
- Drug will be used for conversion of acute symptomatic episodes of PSVT to sinus rhythm; **AND**
- Patient does not have any contraindication to etripamil (e.g., heart failure, Wolff-Parkinson-White (WPW), Lown-Ganong-Levine (LGL) syndromes); **AND**
- Prescribed by, or in consultation with, a cardiologist or electrophysiologist

Isradipine capsules

PDL Status	NP
Quantity Limit	2.5 mg: 2/day; 5 mg: 4/day
PA Criteria	See Below
PA Form	General PA Form

Isradipine capsules PA Criteria:

Standard NP Criteria

Katerzia® suspension

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Katerzia® suspension PA Criteria:
See Norliqva prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to Norliqva®

Nimodipine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Nimodipine capsules PA Criteria:

- Diagnosis of subarachnoid hemorrhage (SAH)

Nisoldipine ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Nisoldipine ER tablets PA Criteria:
Standard NP Criteria

Norvasc® tablets

PDL Status	NP
Quantity Limit	2.5 & 5 mg: 1.5/day; 10 mg: 1/day
PA Criteria	See Below
PA Form	General PA Form

Norvasc® tablets PA Criteria:
Standard NP Criteria

Nymalize® solution

PDL Status	NP
Quantity Limit	120 mL/day
PA Criteria	See Below
PA Form	General PA Form

Nymalize® solution PA Criteria:
Standard NP Criteria

Procardia XL® 24HR osmotic tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Procardia XL® 24HR osmotic tablets PA Criteria:
Standard NP Criteria

Sdamlo® solution

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Sdamlo® solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Sular® ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Sular® ER tablets PA Criteria:
Standard NP Criteria

Calcium Channel Blockers (Non-DHP)

Verapamil ER/SR 24HR capsules (except 100, 200, & 300 mg)

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Diltiazem ER tablets & Matzim LA®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Diltiazem ER tablets & Matzim LA® PA Criteria:
Standard NP Criteria

Verapamil ER PM 100, 200, & 300 mg capsules, Verelan® capsules, Verelan PM® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Verapamil ER PM 100, 200, & 300 mg capsules, Verelan® capsules, Verelan PM® capsules PA Criteria:
Standard NP Criteria

Cardiac Agents, Miscellaneous

Ranolazine ER tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Ivabradine tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Ivabradine tablets PA Criteria:

- Diagnosis of Congestive Heart Failure (NYHA class II to IV) and documentation of the following:
  - Left ventricular ejection fraction ≤ 35%; **AND**
  - In sinus rhythm with resting heart rate ≥ 70 beats per minute; **AND**
  - One of the following:
    - Currently taking a maximum tolerated dose of a beta-blocker and still experiencing heart failure symptoms; **OR**
    - Patient has a contraindication, adverse reaction, or drug-drug interaction to a beta-blocker; **OR**
- Diagnosis of Congestive Heart Failure (NYHA class II to IV) due to dilated cardiomyopathy (DCM); **AND**
  - Left ventricular ejection fraction ≤ 45%; **AND**
  - Patient is in sinus rhythm with elevated heart rate; **AND**
- Will NOT be approved for patients with any of the following:
  - Concomitant use of potent CYP3A inhibitors or inducers
  - Acute decompensated heart failure
  - Clinically significant hypotension or bradycardia
  - Sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present
  - Severe hepatic impairment
  - Pacemaker dependence (heart rate maintained exclusively by the pacemaker)

Aspruzyo Sprinkle® packets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Aspruzyo Sprinkle® packets PA Criteria:

- Diagnosis of chronic angina; **AND**
- Patient is unable to swallow solid dosage form

Camzyos® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Camzyos® capsules Initial PA Criteria:

- Diagnosis of obstructive hypertrophic cardiomyopathy (HCM); **AND**
- Left ventricular hypertrophy (LVH) confirmed by cardiac imaging (i.e., echocardiography, cardiac MRI); **AND**
- Heart failure is classified New York Heart Association (NYHA) class II or III Patient has New York Heart Association (NYHA) Class II or III symptoms (e.g., shortness of breath, chest pain); **AND**
- Patient has left ventricular outflow tract (LVOT) peak gradient > 50 mmHg at rest or with provocation; **AND**
- Patient has a left ventricular ejection fraction > 55% (for initiation of therapy); **AND**
- Prescribed by or in consultation with a cardiologist; **AND**
- Trial and failure, contraindication, or intolerance to ONE of the following at a maximally tolerated dose:
  - Non-vasodilating beta blocker (e.g., bisoprolol, propranolol)
  - Calcium channel blocker (e.g., verapamil, diltiazem)

Camzyos® capsules Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., NHYA class remains stable or improves improved symptom relief, improvement of LVOT gradient); **AND**
- Patient has a left ventricular ejection fraction > 50%; **AND**
- Prescribed by, or in consultation with, a cardiologist

Corlanor® solution, tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Corlanor® solution, tablets PA Criteria:

- See ivabradine prior authorization criteria; **AND**
- Clinically valid reason why ivabradine cannot be used

Lodoco® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Lodoco® tablets PA Criteria:

- Patient has established atherosclerotic disease or with multiple risk factors for CV disease; **AND**
- Treatment is being requested to reduce the risk of myocardial infarction (MI), stroke, coronary revascularization, and cardiovascular (CV) death; **AND**
- Patient is on optimized guideline-directed therapy such as beta-blockers, RAS inhibitors, and lipid lowering agents to ensure reduced cardiovascular risk unless contraindicated (supported by medical documentation or claims history); **AND**
- Prescriber attest patient remains at very high risk for recurrent ASCVD events despite maximally tolerated GDMT

Tryvio® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Tryvio® tablets Initial PA Criteria:

- Diagnosis of resistant hypertension; **AND**
- Patient is 18 years of age or older; **AND**
- Submission of medical records (e.g. chart notes) documenting patient has not achieved target blood pressure (e.g. systolic <130/80) following concurrent maximumly tolerated treatment with at least THREE of the following antihypertensive classes unless contraindicated or intolerance:
  - ACE inhibitors or Angiotensin II receptor blockers (ARBs)
  - Beta blockers
  - Calcium channel blockers
  - Mineralocorticoid Receptor Antagonists
  - Thiazide diuretics; **AND**
- Tryvio will be used in combination with at least one other hypertensive drug

Tryvio® tablets Renewal PA Criteria:

- Submission of medical records (e.g. chart notes) documenting a positive clinical response to therapy (e.g. decrease of systolic blood pressure from baseline); **AND**
  - Patient is receiving concomitant therapy with other hypertensive drugs (documented by claims or medical records)

Verquvo® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Verquvo® tablets PA Criteria:

- Diagnosis of symptomatic chronic heart failure (NYHA class II-IV) with reduced ejection fraction (≤45%); **AND**
- Prescribed by, or in consultation with, a cardiologist (initial approval only); **AND**
- Patient has had a heart failure hospitalization in the last 6 months **OR** has received outpatient IV diuretics for heart failure in the last 3 months; **AND**
- Patient is 18 years of age or older; **AND**
- Patient is currently being treated with guideline directed therapy (e.g., SGLT2 inhibitors, ACEI, ARB, etc.); **AND**
- Patient is not pregnant or breastfeeding; **AND**
- Female patients of reproductive potential will be counseled to use effective contraception during treatment with therapy and for at least one month after the last dose; **AND**
- Patient does not meet any of the following:
  - Concomitant use with another soluble guanylate cyclase (sGC) stimulator (e.g., Adempas)
  - Concomitant use with a PDE-5 inhibitor (e.g., tadalafil, sildenafil)

Cardiac Glycosides

Lanoxin® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Lanoxin® tablets PA Criteria:
Standard NP Criteria

Direct Renin Inhibitors

Aliskiren

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Aliskiren PA Criteria:

- Patient has a diagnosis of hypertension; **AND**
- Trial and failure, contraindication, or intolerance to an agent from at least TWO of the following drug classes:
  - ACEI/ARB
  - Calcium channel blocker
  - Thiazide diuretic

Tekturna®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Tekturna® PA Criteria:
See aliskiren prior authorization criteria

Tekturna HCT®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Tekturna HCT® PA Criteria:

- Trial and failure, contraindication, or intolerance to an agent from at least TWO of the following drug classes:
  - ACEI/ARB
  - Calcium channel blocker
  - Thiazide diuretic; **AND**
- Patient is unable to take the individual components

Diuretics: Carbonic Anhydrase

Dichlorphenamide

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Dichlorphenamide PA Criteria:

- See Keveyis criteria**; AND**
- Clinically valid reason why Keveyis® cannot be used

Keveyis®

PDL Status	NP
Quantity Limit	2/DAY
PA Criteria	See Below
PA Form	General PA Form

Keveyis® Initial PA Criteria:

- Diagnosis of Primary Hypokalemic/Hyperkalemic Periodic Paralysis, and related variants; **AND**
- Patient does not have any of the following:
  - Hepatic insufficiency
  - Severe pulmonary disease
  - Hypersensitivity to dichlorphenamide or other sulfonamides
  - Concomitant use with high dose aspirin

Keveyis® Renewal PA Criteria:

- Clinical documentation that patient has exhibited a reduction in symptoms or attacks; **AND**
- Patient's serum potassium and bicarbonate levels are being monitored

Ormalvi®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Ormalvi® PA Criteria:

See Keveyis criteria**; AND**

- Clinically valid reason why Keveyis® cannot be used

Diuretics: Loop

Enbumyst® nasal spray

PDL Status	NP
Quantity Limit	120 units/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	2 months

Enbumyst® nasal spray Initial PA Criteria:

- Diagnosis of one of the following:
  - Chronic heart failure
  - Hepatic disease
  - Kidney disease, including nephrotic syndrome; **AND**
- Prescribed by, or in consultation with, a cardiologist, nephrologist, hepatologist, or gastroenterologist; **AND**
- Prescriber attests being used for short-term treatment, when the use of IV or oral bumetanide is not feasible (e.g., access issues, resistance, poor gastrointestinal absorption, delayed onset)

Furoscix® On-Body Infusor

PDL Status	NP
Quantity Limit	4 devices/30 days; 8 devices/180 days
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 months

Furoscix® On-Body Infusor PA Criteria:

- Diagnosis of chronic heart failure or chronic kidney disease, including nephrotic syndrome; **AND**
- Patient has signs and symptoms of fluid overload; **AND**
- The patient is currently receiving maximal oral diuretic therapy; **AND**
- Prescriber attests to BOTH of the following:
  - Additional oral diuretic therapy would be ineffective;
  - The requested agent will be for short-term outpatient use only and patient will be transitioned back to oral diuretics as soon as clinically feasible; **AND**
- Prescribed by, or in verbal consultation with, a cardiologist or nephrologist; **AND**
- Prescriber has demonstrated appropriate administration use of the On-Body Infusor®

Lasix® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Lasix® tablets PA Criteria:

Standard NP Criteria

Diuretics: Potassium Sparing

Aldactone® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Aldactone® tablets PA Criteria:
Standard NP Criteria

CaroSpir®

PDL Status	NP
Quantity Limit	15 mL/day
PA Criteria	See Below
PA Form	General PA Form

CaroSpir® PA Criteria:

- One of the following:
  - Diagnosis of hypertension
  - Diagnosis of heart failure
  - Diagnosis of edema associated with hepatic cirrhosis; **AND**
- Patient is unable to swallow solid dosage forms
- Clinically valid reason why generic spironolactone suspension cannot be used

Dyrenium® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Dyrenium® capsules PA Criteria:
Standard NP Criteria

Eplerenone tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Eplerenone tablets PA Criteria:

- One of the following:
  - Patient has a diagnosis of hypertension
  - Patient has a diagnosis of congestive heart failure
  - Patient has a diagnosis of Duchenne muscular dystrophy (DMD); **AND**
- Trial and failure, contraindication, or intolerance of spironolactone

Kerendia®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Kerendia® PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescriber attests that the patient be monitored for Hyperkalemia while on therapy; **AND**
- One of the following:
- Diagnosis of chronic kidney disease (CKD) associated with type 2 diabetes (T2D); **AND**
  - Currently taking the maximum tolerated dose of an ACE inhibitor or ARB, unless contraindicated or intolerant; **AND**
  - Currently taking an antidiabetic agent (e.g., insulin, metformin, GLP-1 receptor agonist, SGLT2 inhibitor); **OR**
- Diagnosis of chronic heart failure (NYHA Class II-IV); **AND**
  - Left ventricular ejection fraction (LVEF) ≥ 40%

Spironolactone Susp

PDL Status	NP
Quantity Limit	15 mL/day
PA Criteria	See Below
PA Form	General PA Form
Note	PA not required for patients < 6 years of age

Spironolactone Suspension PA Criteria:

- One of the following:
  - Diagnosis of hypertension
  - Diagnosis of heart failure
  - Diagnosis of edema associated with hepatic cirrhosis; **AND**
- Patient is unable to swallow solid dosage forms

Triamterene capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Triamterene capsules PA Criteria:
Standard NP Criteria

Diuretics: Thiazide and Related

Hemiclor®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Hemiclor® PA Criteria:

- Clinically valid reason why the preferred generic chlorthalidone cannot be used (NOTE: Patient convenience is NOT an approvable reason)

Inzirqo®

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Inzirqo® PA Criteria:

- Documentation of need for dosing that will not allow the use of a preferred HCTZ tablet; **OR**
- Patient is unable to swallow solid oral dosage forms

Thalitone®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Thalitone® PA Criteria:

Standard NP Criteria

Hemostatics, Oral

Tranexamic Acid

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Tranexamic Acid PA Criteria:

- Diagnosis of acute uterine or cyclic heavy menstrual bleeding; **AND**
- Trial and failure, contraindication, or intolerance to ALL the following:
  - Two other forms of hormone therapy (oral, vaginal, topical, or injectable estrogen and/or progesterone)
  - Levonorgestrel-releasing IUD; **OR**
- All other diagnoses require trial and failure, intolerance, or contraindication to aminocaproic acid.

Lipotropics: Antihyperlipidemic Agents

Repatha®

PDL Status	P
Quantity Limit	Repatha: 2/28 days; Repatha Pushtronex: 1/28 days
PA Criteria	See Below
PA Form	PCSK9 Inhibitors PA Form
PA Approval Duration	6 months

Cardiovascular disease (CVD) Prevention

Repatha® Initial PA Criteria:

- Treatment is for the prevention of cardiovascular disease; **AND**
- Patient age ≥ 18 years; **AND**
- Documented current LDL-C value (within 3 months); **AND**
- Patient specific target LDL-C value is provided; **AND**
- One of the following:
  - Failure to reach patient specific LDL target despite a ≥ 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
    - High-intensity statin (atorvastatin/rosuvastatin)
    - Ezetimibe; **OR**
  - Patient requires > 25% additional LDL-C lowering to meet LDL target after a > 3-month trial (supported by claims history or clinical documentation) of therapy with a high-intensity statin, unless contraindicated or intolerance; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance

Repatha® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Heterozygous familial hypercholesterolemia (HeFH) or homozygous familial hypercholesterolemia (HoFH)

Repatha® Initial PA Criteria:

- Diagnosis of HeFH or HoFH confirmed by one of the following:
  - Presence of a mutation in LDL receptor, ApoB, PCSK9 gene
  - Clinical criteria using either the Simon Broome or WHO/Dutch Lipid Network criteria; **AND**
- Patient age is >10 years; **AND**
- One of the following:
  - Failure to reach patient specific LDL target despite a ≥ 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
  - High-intensity statin (atorvastatin/rosuvastatin)
  - Ezetimibe; **OR**
  - Patient requires > 25% additional LDL-C lowering to meet LDL target after a > 3-month trial (supported by claims history or clinical documentation) of therapy with a high-intensity statin, unless contraindicated or intolerance; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance

Repatha® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Juxtapid®

PDL Status	NP
Quantity Limit	5 mg, 10 mg: 1/day 20 mg: 3/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Juxtapid® Initial PA Criteria:

- Diagnosis of homozygous familial hypercholesterolemia (HoFH) confirmed by one of the following:
  - Presence of a mutation in LDL receptor, ApoB, PCSK9 gene
  - Clinical criteria using either the Simon Broome or WHO/Dutch Lipid Network criteria; **AND**
- Patient age is appropriate according to package labeling (i.e., Praluent is indicated for age >18 years, Repatha is indicated for age >10 years; **AND**
- Documented current LDL-C value (within 3 months); **AND**
- Patient specific target LDL-C value is provided; **AND**
- Failure to reach patient specific LDL target despite a ≥ 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
  - High-intensity statin (atorvastatin/rosuvastatin)
  - Ezetimibe; **AND**
- Trial and failure, contraindication, or intolerance to Repatha; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance; **AND**
- If female, documentation patient is not currently pregnant; **AND**
- Patient is not concomitantly taking strong or moderate inhibitors of cytochrome P450 (CYP) 3A4

Juxtapid® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Nexletol®, Nexlizet®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Nexletol®, Nexlizet® Initial PA Criteria:

- Age ≥ 18 years; **AND**
- Diagnosis of ONE of the following:
  - Primary prevention of cardiovascular disease
  - Primary hyperlipidemia
  - Atherosclerotic cardiovascular disease (ASCVD)
  - Heterozygous familial hypercholesterolemia (HeFH)) confirmed by one of the following:
    - Presence of a mutation in LDLR, ApoB, PCSK9 or ARH adaptor protein (LDLRAP1) gene
    - Clinical criteria is met using either the Simon Broome or WHO/Dutch Lipid Network criteria; **AND**
- Documented current LDL-C value (within 3 months); **AND**
- Patient specific target LDL-C value is provided; **AND**
- Failure to reach patient specific LDL target despite a ≥ 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
  - High-intensity statin (atorvastatin/rosuvastatin)
  - Ezetimibe; **AND**
- Trial and failure, contraindication, or intolerance to Repatha; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance

Nexletol®, Nexlizet® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Praluent®

PDL Status	NP
Quantity Limit	2 pens/28 days
PA Criteria	See Below
PA Form	PCSK9 Inhibitors PA Form
PA Approval Duration	6 months

Cardiovascular Disease (CVD) Prevention

Praluent® Initial PA Criteria:

- Treatment is for the prevention of cardiovascular disease; **AND**
- Patient age ≥ 18 years; **AND**
- Patient has history of ONE of the following:
  - MI, unstable angina, or symptomatic peripheral artery disease
  - Stroke
  - Primary Hyperlipidemia (including heterozygous familial hypercholesterolemia (HeFH))
  - Baseline LDL-C ≥ 190; **AND**
- Documented current LDL-C value (within 3 months); **AND**
- Patient specific target LDL-C value is provided; **AND**
- One of the following:
  - Failure to reach patient specific LDL target despite a ≥ 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
    - High-intensity statin (atorvastatin/rosuvastatin)
    - Ezetimibe; **OR**
  - Patient requires > 25% additional LDL-C lowering to meet LDL target after a > 3-month trial (supported by claims history or clinical documentation) of therapy with a high-intensity statin, unless contraindicated or intolerance; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance
- Trial and failure, contraindication, or intolerance to a preferred agent

Praluent® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Heterozygous familial hypercholesterolemia (HeFH) or homozygous familial hypercholesterolemia (HoFH)

Praluent® Initial PA Criteria:

- Diagnosis of HeFH or HoFH confirmed by one of the following:
  - Presence of a mutation in LDL receptor, ApoB, PCSK9 gene
  - Clinical criteria using either the Simon Broome or WHO/Dutch Lipid Network criteria; **AND**
- Patient age is appropriate according to package labeling (i.e., Praluent is indicated for age ≥8 years, Repatha is indicated for age ≥10 years); **AND**
- One of the following:
  - Failure to reach patient specific LDL target despite a > 3-month trial (supported by claims history or clinical documentation) of concurrent therapy with BOTH the following, unless contraindicated or intolerance:
    - High-intensity statin (atorvastatin/rosuvastatin)
    - Ezetimibe; **OR**
  - Patient requires > 25% additional LDL-C lowering to meet LDL target after a > 3-month trial (supported by claims history or clinical documentation) of therapy with a high-intensity statin, unless contraindicated or intolerance; **AND**
- Agent will be used in combination with other lipid lowering therapies, unless documented intolerance
- Trial and failure, contraindication, or intolerance to a preferred agent

Praluent® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., LDL reduction from baseline, patient has meet LDL-C target)

Lipotropics: Bile Acid Sequestrants

Colestid® granules, tablets

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Colestid® granules, tablets PA Criteria:
Standard NP Criteria

Colesevelam packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Colesevelam packets PA Criteria:

- Patient is unable to swallow solid dosage forms

Colestipol granules, tablets, Questran® powder, Questran Light® powder

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Colestipol granules, tablets, Questran® powder, Questran Light® powder PA Criteria:
Standard NP Criteria

Welchol® packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Welchol® packets PA Criteria:

- Patient is unable to swallow solid dosage forms

Welchol® tablets

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Welchol® tablets PA Criteria:
Standard NP Criteria

Lipotropics: Cholesterol Absorption Inhibitors

Zetia® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Zetia® tablets PA Criteria:

- One of the following:
  - Patient is currently taking a high-intensity statin and has experienced less than anticipated therapeutic response
  - Patient is unable to tolerate lower doses of high-intensity therapy
  - Use in combination with a bile acid sequestrant, fibrate, or niacin will be approved.
  - For requests as monotherapy, recipients must have been intolerant to, or have a contraindication to, a statin

Lipotropics: Combination Agents

Ezetimibe/simvastatin tablets, Vytorin® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Ezetimibe/simvastatin tablets, Vytorin® tablets PA Criteria:

- For patients that require ≤45% LDL reduction: 4-week trial and failure of both atorvastatin and simvastatin; **OR**
- For patients that require >45% LDL reduction: 4-week trial and failure of atorvastatin

Lipotropics: Fibric Acid Derivatives

Fenofibrate capsules, Fenofibrate tablets 40, 54, & 120 mg, Fenofibric acid, Fibricor® tablets, Lipofen® capsules, Tricor® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Fenofibrate capsules, Fenofibrate tablets 40, 54, & 120 mg, Fenofibric acid, Fibricor® tablets, Lipofen® capsules, Tricor® tablets PA Criteria:

- Patient will take fenofibrate concomitantly with a sulfonylurea, thiazolidinedione, repaglinide, or a statin; **OR**
- Clinically valid reason why a preferred agent cannot be used (e.g., gemfibrozil, fenofibrate tablets 48, 145, & 160 mg)

Lipotropics: Niacin Derivatives

Niacin ER

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Niacin ER PA Criteria:

- One of the following:
  - Triglycerides > 500 mg/dL; **AND**
  - Diagnosis of hyperlipidemia; **AND**
    - Trial and failure, contraindication, or intolerance to a statin

Lipotropics: Omega-3 Fatty Acids

Omega-3 acid ethyl esters

PDL Status	P
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Omega-3 acid ethyl esters Initial PA Criteria:

- Diagnosis of Severe Hypertriglyceridemia (TG level is above 500 mg/dl)

Omega-3 acid ethyl esters Renewal PA Criteria:

- Documentation of positive clinical response (e.g., reduction in TG from baseline)

Icosapent ethyl

PDL Status	NP
Quantity Limit	0.5 g: 2/day; 1 g: 4/day
PA Criteria	See Below
PA Form	General PA Form

Initial Icosapent ethyl Initial PA Criteria:

- Diagnosis of Severe Hypertriglyceridemia (TG level is above 500 mg/dl); **OR**
- Patient is on maximally tolerated statin **AND** has triglyceride levels $\geq$ 135

Renewal Icosapent ethyl PA Criteria:

- Documentation of positive clinical response (e.g., reduction in TG from baseline)

Lipotropics: Statins

Atorvastatin tablets, Lovastatin tablets, Pravastatin tablets, Rosuvastatin tablets, Simvastatin 5 to 40 mg tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Simvastatin 80 mg

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
Notes	FDA labeling states that due to the increased risk of myopathy, including rhabdomyolysis, associated with the 80 mg dose of simvastatin, patients unable to achieve their LDL-C goal utilizing the 40 mg dose of simvastatin should not be titrated to the 80 mg dose, but should be placed on alternative LDL-C lowering treatment(s) that provides greater LDL-C lowering. Atorvastatin and rosuvastatin are available without PA.

Simvastatin 80 mg PA Criteria:

- Patient has previously received simvastatin 80 mg for 12 months or longer with no evidence of myopathy; **AND**
- Clinically valid reason why the patient cannot use high-intensity doses of atorvastatin or rosuvastatin (for patients requiring > 50% LDL-C reduction or have LDL-C goal of < 70)

Altoprev ER® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Altoprev ER® tablets PA Criteria:

Standard NP Criteria

Atorvaliq® suspension

PDL Status	NP
Quantity Limit	80 mg/day
PA Criteria	See Below
PA Form	General PA Form

Atorvaliq® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Crestor® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Crestor® tablets PA Criteria:

Standard NP Criteria

Ezallor Sprinkles® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Ezallor Sprinkles® capsules PA Criteria:

- Patient is unable to swallow solid dosage forms

Flolipid® suspension

PDL Status	NP
Quantity Limit	40 mg/day
PA Criteria	See Below
PA Form	General PA Form

Flolipid® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms

Fluvastatin tablets, Fluvastatin ER tablets, Lipitor® tablets, Livalo® tablets, Pitavastatin tablets, Zocor® 5 to 40 mg tablets, & Zypitamag® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Fluvastatin tablets, Fluvastatin ER tablets, Lipitor® tablets, Livalo® tablets, Pitavastatin tablets, Zocor® 5 to 40 mg tablets, & Zypitamag® tablets PA Criteria:

Standard NP Criteria

Lipotropics: Statin + Calcium Channel Blocker

Amlodipine/atorvastatin tablets, Caduet® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Amlodipine/atorvastatin tablets, Caduet® tablets PA Criteria:

- Patient is unable to take the 2 components separately

Pheochromocytoma Agents

Metyrosine

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Metyrosine PA Criteria:

- Documentation of pheochromocytoma diagnosis; **AND**
- Trial and failure of an alpha and beta blocker

Phenoxybenzamine

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Phenoxybenzamine PA Criteria:

- Diagnosis of pheochromocytoma

Platelet Inhibitors

Prasugrel tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Prasugrel tablets PA Criteria:

- Patients has unstable angina, NSTEMI, or STEMI; **AND**
- PCI will be performed, or PCI is planned; **AND**
- Age < 75 years; **AND**
- Weight ≥ 60 kg; **AND**
- No history of stroke or TIA

Ticagrelor tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Note	Will NOT be approved if patient is receiving aspirin doses > 100 mg/day (includes Rx & OTC aspirin containing products)

Ticagrelor tablets PA Criteria:

- History of Myocardial Infarction (MI); **OR**
- ACS initial event (USA, NSTEMI or STEMI) or recurrence within previous 12 months; **OR**
- Patient has diagnosis of coronary artery disease (CAD) and is at high risk for myocardial infarction (MI) or stroke, **OR**
- Acute ischemic stroke or transient ischemic attack (TIA) risk reduction

Agrylin® capsules

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Agrylin® capsules PA Criteria:

Standard NP Criteria

Brilinta® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Note	Will NOT be approved if patient is receiving aspirin doses > 100 mg/day (includes Rx & OTC aspirin containing products)

Brilinta® PA Criteria:

- History of Myocardial Infarction (MI); **OR**
- ACS initial event (USA, NSTEMI or STEMI) or recurrence within previous 12 months; **OR**
- Patient has diagnosis of coronary artery disease (CAD) and is at high risk for myocardial infarction (MI) or stroke, **OR**
- Acute ischemic stroke or transient ischemic attack (TIA) risk reduction

Cablivi® kit for injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	2 months
Note	If started as an inpatient hospital regimen and this is continuation of therapy, Cablivi® will be approved

Cablivi® kit for injection PA Criteria:

- Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP); **AND**
- Used in combination with both of the following:
  - Plasma exchange until at least 2 days after normalization of the platelet count
  - Immunosuppressive therapy (e.g., corticosteroids); **AND**
- Date Cablivi IV was initiated/administered by a healthcare provider; **AND**
- Total treatment duration will be limited to 58 days beyond the last therapeutic plasma exchange; **AND**
- The patient has not experienced more than two recurrences of aTTP while on Cablivi

Clopidogrel 300 mg tablets

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Clopidogrel 300 mg tablets PA Criteria:

Standard NP Criteria

Effient® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Effient® tablets PA Criteria:

- Patients has unstable angina, NSTEMI, or STEMI; **AND**
- PCI has been performed, or PCI is planned; **AND**
- Age < 75 years; **AND**
- Weight ≥ 60 kg; **AND**
- No history of stroke or TIA; **AND**
- Trial and failure of prasugrel

Plavix® tablets

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Plavix® tablets PA Criteria:

Standard NP Criteria

Yosprala® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Note	Will NOT be approved if patient is receiving aspirin doses > 100 mg/day (includes Rx & OTC aspirin containing products)

Yosprala® tablets PA Criteria:

- Diagnosis of one of the following:
  - Ischemic stroke,
  - Transient ischemia of the brain,
  - Previous myocardial infarction,
  - Unstable angina pectoris,
  - Chronic stable angina pectoris; **OR**
- Patient has had **ONE** of the following:
  - Coronary Artery Bypass Graft (CABG)
  - Percutaneous Transluminal Coronary Angioplasty (PTCA); **AND**
- Patient meets **ALL** the following:
  - Patient is considered a high-risk candidate for aspirin-associated gastric ulcers due to **ONE** of the following:
    - Age ≥ 55, **OR**
    - Documented history of gastric ulcers; **AND**
  - Patient had an inadequate treatment response, or intolerance to use of aspirin and omeprazole separately

Pulmonary Arterial Hypertension (PAH) Agents

Alyq® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Alyq® tablets PA Criteria:

- Diagnosis of Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension (PPH); **OR**
- Diagnosis of Congenital heart disease with elevated pulmonary vascular resistance

Ambrisentan tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Ambrisentan tablets PA Criteria:

See Alyq® prior authorization criteria

Bosentan tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Bosentan PA Criteria:

See Alyq® prior authorization criteria

Sildenafil tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Sildenafil PA Criteria:
See Alyq® prior authorization criteria

Tadalafil tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Tadalafil PA Criteria:
See Alyq® prior authorization criteria

Ventavis® solution

PDL Status	P
Quantity Limit	3 mL/day
PA Criteria	See Below
PA Form	General PA Form

Ventavis® solution PA Criteria:
See Alyq® prior authorization criteria

Adcirca® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Adcirca® tablets PA Criteria:

- Diagnosis of one of the following:
- Pulmonary arterial hypertension (PAH)/elevated pulmonary vascular resistance or primary pulmonary hypertension
- Congenital heart disease with elevated pulmonary vascular resistance; **AND**
- Clinically valid reason why the preferred generic cannot be used

Adempas® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form
Note	Use of Adempas® is contraindicated in patients also taking PDE-5 inhibitors

Adempas® tablets PA Criteria:

- One of the following:
  - Diagnosis of pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension (PPH); **AND**
    - Trial of ONE preferred agent with persistent signs or symptoms
  - Diagnosis of congenital heart disease with elevated pulmonary vascular resistance; **AND**
    - Trial of ONE preferred agent with persistent signs or symptoms
  - Diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH) with one of the following:
    - Patient has disease that is inoperable; **OR**
    - Patient has residual post-pulmonary endarterectomy hypertension

Bosentan tablets for oral susp

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Bosentan tablets for oral suspension PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension (PPH)
  - Diagnosis of congenital heart disease with elevated pulmonary vascular resistance; **AND**
- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why brand Tracleer® tablets for oral suspension cannot be used

Letairis® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Letairis® tablets PA Criteria:

See Adcirca® prior authorization criteria

Macitentan tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Macitentan tablets PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension
  - Congenital heart disease with elevated pulmonary vascular resistance; **AND**
- Trial of one preferred agent with persistent signs or symptoms

Opsumit® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Opsumit® tablets PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension
  - Congenital heart disease with elevated pulmonary vascular resistance; **AND**
- Trial of one preferred agent with persistent signs or symptoms; **AND**
- Clinically valid reason why generic macitentan tablets cannot be used

Opsynvi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Opsynvi® tablets PA Criteria:

- Diagnosis of Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension; **AND**
- Clinically valid reason as to why the patient is unable to take components of Opsynvi® individually

Orenitram® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Orenitram® tablets PA Criteria:

See Opsumit® prior authorization criteria

Revatio® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Revatio® PA Criteria:

See Adcirca® prior authorization criteria

Riociguat tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Riociguat tablets PA Criteria:

- See Adempas® prior authorization criteria; **AND**
- Clinically valid reason why brand Adempas® cannot be used

Sildenafil suspension

PDL Status	NP
Quantity Limit	6 mL/day; Max day supply=60
PA Criteria	See Below
PA Form	General PA Form

Sildenafil suspension PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension
  - Congenital heart disease with elevated pulmonary vascular resistance; **AND**
- One of the following:
  - Patient is unable to swallow tablets
  - Patient is < 6 years of age
- Clinically valid reason why a preferred tablet formulation cannot be used

Tadliq® suspension

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Tadliq® suspension PA Criteria:

See sildenafil suspension prior authorization criteria

Tracleer® soluble tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Tracleer® soluble tablets PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension (PPH)
  - Diagnosis of congenital heart disease with elevated pulmonary vascular resistance; **AND**
- Patient is unable to swallow solid dosage forms

Tracleer® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Tracleer® tablets PA Criteria:

See Adcirca® prior authorization criteria

Tyvaso® solution

PDL Status	NP
Quantity Limit	2.9 mL/day
PA Criteria	See Below
PA Form	General PA Form

Tyvaso® solution PA Criteria:

- Diagnosis of pulmonary hypertension associated with interstitial lung disease (PH-ILD; WHO Group 3) to improve exercise ability; **OR**
- Diagnosis of pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension (PPH); **AND**
- Clinically valid reason why a preferred agent cannot be used; **AND**
- Diagnosis of congenital heart disease with elevated pulmonary vascular resistance; **OR**
- Clinically valid reason why a preferred agent cannot be used

Tyvaso DPI®

PDL Status	NP
Quantity Limit	Single cartridges: 4/day; Combo cartridges: 8/day; Kits: 2/year
PA Criteria	See Below
PA Form	General PA Form

Tyvaso DPI® PA Criteria:

- Diagnosis of one of the following:
  - Pulmonary arterial hypertension (PAH)/elevated pulmonary vascular resistance or primary pulmonary hypertension
  - Pulmonary hypertension associated with interstitial lung disease; **AND**
- Clinically valid reason why Tyvaso inhalation solution cannot be used

Uptravi® tablets

PDL Status	NP
Quantity Limit	Tablets: 2 /day; Pack: 1 /Rx
PA Criteria	See Below
PA Form	General PA Form

Uptravi® tablets PA Criteria:

See Opsumit® prior authorization criteria

Winrevair® injection

PDL Status	NP
Quantity Limit	1 kit/21 days
PA Criteria	See Below
PA Form	General PA Form

Winrevair® injection PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of Pulmonary arterial hypertension (PAH)/ elevated pulmonary vascular resistance or primary pulmonary hypertension; **AND**
- Patient is Functional Class II or III; **AND**
- Trial and failure of 1 agent with persistent signs and symptoms from TWO different classes for PAH such as:
- Endothelin receptor antagonist (e.g. ambrisentan, bosentan)
- Phosphodiesterade-5 inhibitors (e.g. sildenafil, tadalafil)
- Prostacyclin analogue or receptor agonist (e.g., treprostinil, epoprostenol, Uptravi, Ventavis); **AND**
- Winrevair® will be used in combination with other PAH therapies

Yutrepia® caps for inhalation

PDL Status	NP
Quantity Limit	5/day
PA Criteria	See Below
PA Form	General PA Form

Yutrepia® caps for inhalation PA Criteria:

- Diagnosis of one of the following:
- Pulmonary arterial hypertension (PAH)/elevated pulmonary vascular resistance or primary pulmonary hypertension
- Pulmonary hypertension associated with interstitial lung disease; **AND**
- Clinically valid reason why Tyvaso inhalation solution cannot be used

Pulmonary Fibrosis Agents

Nintendanib capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Nintedanib capsules PA Criteria:

- Diagnosis of one of the following:
  - Idiopathic pulmonary fibrosis
  - Interstitial Lung Disease Associated with Systemic Sclerosis- associated interstitial lung disease (SSc-ILD)
  - Chronic Fibrosing Interstitial Lunch Diseases (ILDs) with a progressive phenotype (at least 10% of the lungs show presence of fibrotic ILD); **AND**
- Prescribed by, or in consultation with, a pulmonologist (initial approval only)

Nintendanib capsules Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improved lung function, improved measured FVC, decrease in hospitalization for respiratory cause); **AND**
- Prescribed by, or in consultation with, a pulmonologist;

Pirfenidone tablets

PDL Status	P
Quantity Limit	534 mg, 801 mg: 3/day; 267 mg: 9/day
PA Criteria	See Below
PA Form	General PA Form

Pirfenidone tablets PA Criteria:

- Patient has a diagnosis of idiopathic pulmonary fibrosis; **AND**
- Prescribed by, or in consultation with, a pulmonologist (initial approval only)

Esbriet® capsules, tablets

PDL Status	NP
Quantity Limit	801 mg: 3/day; 267 mg: 9/day
PA Criteria	See Below
PA Form	General PA Form

Esbriet® capsules, tablets PA Criteria:

- Patient has a diagnosis of idiopathic pulmonary fibrosis; **AND**
- Prescribed by, or in consultation with, a pulmonologist (initial approval only); **AND**
- Clinically valid reason as to why the preferred pirfenidone cannot be used

Jascayd® tablets

PDL Status	NP
Quantity Limit	2 tabs/day
PA Criteria	See Below
PA Form	General PA Form

Jascayd® tablets Initial PA Criteria:

- Diagnosis of one of the following:
  - Idiopathic pulmonary fibrosis
  - Progressive pulmonary fibrosis; **AND**
- Prescribed by, or in consultation with, a pulmonologist; **AND**
- If patient is using concomitantly with pirfenidone patient should be receiving 18 mg twice daily; **AND**
- Clinically valid reason as to why the preferred nintendanib cannot be used

Jascayd® tablets Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g. improved lung function, improved measured FVC, decrease in acute ILD exacerbation, decrease in hospitalization for respiratory causes); **AND**
- If patient is using concomitantly with pirfenidone, patient should be receiving 18mg twice daily; **AND**
- Prescribed by, or in consultation with, a pulmonologist; **AND**
- Patient has absence of unacceptable side effects or toxicities (e.g. diarrhea, upper respiratory infections, depression)

Ofev® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Ofev® capsules Initial PA Criteria:

- Diagnosis of one of the following:
  - Idiopathic pulmonary fibrosis
  - Interstitial Lung Disease Associated with Systemic Sclerosis- associated interstitial lung disease (SSc-ILD)
  - Chronic Fibrosing Interstitial Lunch Diseases (ILDs) with a progressive phenotype (at least 10% of the lungs show presence of fibrotic ILD); **AND**
- Prescribed by, or in consultation with, a pulmonologist (initial approval only)

Ofev® capsules Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improved lung function, improved measured FVC, decrease in hospitalization for respiratory cause); **AND**
- Prescribed by, or in consultation with, a pulmonologist; **AND**
- If the request is for brand Ofev®, clinically valid reason why the generic nintedanib cannot be used.

Pirfenidone capsules

PDL Status	NP
Quantity Limit	267 mg: 9/day
PA Criteria	See Below
PA Form	General PA Form

Pirfenidone capsules PA Criteria:

See Esbriet® prior authorization criteria

Thrombopoietin Agonists, Orals

Alvaiz® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Alvaiz® tablets PA Criteria:

- Diagnosis of persistent or chronic thrombocytopenia purpura (ITP) in patients ≥6 years of age;**AND**
  - Documentation of failure or insufficient response to adequate treatment with corticosteroids **AND** immunoglobulins, **OR** ITP related splenectomy; **AND**
  - Documentation that patient's thrombocytopenia and clinical condition puts the patient at increased risk of bleeding; **OR**
- Diagnosis of thrombocytopenia in patient with chronic hepatitis C in patients ≥ 18 years of age;**AND** Patient receiving (or planning to initiate) interferon-based anti-viral therapy; **OR**
- Diagnosis of severe aplastic anemia in patients ≥ 18 years of age; **AND**
  - Patient has tried and failed or has intolerance to immunosuppressive therapy; **AND**
- Prescribed by, or in consultation with, a hematologist or hepatologist

Promacta® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Promacta® tablets PA Criteria:

- Diagnosis of persistent or chronic thrombocytopenia purpura (ITP) in patients ≥1 year of age; **AND**
  - Documentation of failure or insufficient response to adequate treatment with corticosteroids **AND** immunoglobulins, **OR** ITP related splenectomy; **AND**
  - Documentation that patient's thrombocytopenia and clinical condition puts the patient at increased risk of bleeding; **OR**
- Diagnosis of thrombocytopenia in patient with chronic hepatitis C; **AND**
  - Patient receiving (or planning to initiate) interferon-based anti-viral therapy; **OR**
- Diagnosis of severe aplastic anemia in patients 2 years of age or older; **AND**
  - Patient will use in combination with standard immunosuppressive therapy for first-line treatment; **OR**
- Diagnosis of severe aplastic anemia; **AND**
- Patient has tried and failed or has intolerance to immunosuppressive therapy; **AND**
- Prescribed by, or in consultation with, a hematologist or hepatologist

Doptelet Sprinkle® capsules

PDL Status	NP
Quantity Limit	2 capsules/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	See Criteria

Doptelet Sprinkle® capsules Initial PA Criteria:

- Patient is 1 year to less than 6 years old; **AND**
- Patient must have a diagnosis of Chronic Immune Thrombocytopenia (ITP); **AND**
- Patient had an insufficient response to a previous treatment; **AND**
- Patient has a platelet count of $< 50 \times 10^9/L$; **AND**
- Prescribed by, or in consultation with, a hematologist

Doptelet Sprinkle® Capsules Renewal PA Criteria:

- Patient has laboratory values documenting platelet response to therapy (platelet count $\geq 50 \times 10^9/L$); **AND**
- Patient has not experienced severe adverse effects as a result of therapy.

Doptelet® tablets

PDL Status	NP
Quantity Limit	Thrombocytopenia with Chronic Liver Disease: 15 tabs/treatment; Chronic Immune Thrombocytopenia (ITP): 2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	See Criteria

Thrombocytopenia with Chronic Liver Disease (PA Duration: single course of treatment per scheduled procedure)

Doptelet® tablets Initial PA Criteria:

- Patient is ≥ 6 years old or labeled age minimum; **AND**
- Patient must have diagnosis of thrombocytopenia and chronic liver disease and scheduled to undergo a medical procedure; **AND**
- One of the following:
  - Patient is scheduled to take the requested agent 10 to 13 days prior to the procedure, with the procedure occurring 5 to 8 days following the last dose of Doptelet; **OR**
  - Prescribed dose is according to baseline platelet count (10 tabs per 5 day ≥ 40 x 10^9/L or 15 tabs per 5 days platelets < 40 x 10^9/L); **AND**
- Prescribed by, or in consultation with, a hematologist or hepatologist

Doptelet® tablets Renewal PA Criteria

- Patient has laboratory values documenting platelet response to therapy (platelet count ≥ 50 x 10^9/L); **AND**
- Patient has not experienced severe adverse effect as a result of therapy

Chronic Immune Thrombocytopenia (ITP) (PA duration: 1 year)

Doptelet® tablets Initial PA Criteria

- Patient is ≥ 6 years old *or labeled age minimum*; **AND**
- Patient must have a diagnosis of thrombocytopenia; **AND**
- Patient has had an insufficient response to a previous treatment; **AND**
  - Patient has a platelet count of < 50 x 10^9/L**;AND**
- Prescribed by, or in consultation with, a hematologist

Doptelet® tablets Renewal PA Criteria

- Patient has laboratory values documenting platelet response to therapy (platelet count ≥ 50 x 10^9/L); **AND**
- Patient has not experienced severe adverse effect as a result of therapy

Eltrombopag powder for suspension

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Eltrombopag powder for suspension PA Criteria:

See Promacta prior authorization criteria; **AND**

- Clinically valid reason why Promacta® tablets cannot be used; **AND**
- Patient is unable to swallow solid dosage forms

Eltrombopag tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Eltrombopag powder for suspension and tablets PA Criteria:

See Promacta prior authorization criteria; **AND**

- Clinically valid reason why Promacta® cannot be used

Mulpleta® tablets

PDL Status	NP
Quantity Limit	7 tablets/Rx
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Single course of treatment per scheduled procedure

Mulpleta® tablets PA Criteria:

- Patient is ≥ 18 years old; **AND**
- Patient has a diagnosis of Chronic Liver Disease (CLD); **AND**
- Patient does NOT have Child-Pugh class C liver disease, absence of hepatopetal blood flow, a prothrombotic condition other than CLD nor a history of splenectomy, partial splenic embolization, or thrombosis; **AND**
- Patient has a platelet count of < 50 x 10^9/L; **AND**
- Patient has an upcoming invasive procedure scheduled; **AND**
- Patient is scheduled to take the requested agent 8 to 14 days prior to the procedure, with the procedure occurring 2 to 8 days following the last dose of Mulpleta®; **AND**
- Patient is NOT scheduled for a thoracotomy, laparotomy, open-heart surgery, craniotomy, or organ resection; **AND**
- Prescribed by, or in consultation with, a hematologist or hepatologist

Promacta® powder for susp

PDL Status	NP
Quantity Limit	2 packets/day
PA Criteria	See Below
PA Form	General PA Form

Promacta® powder for suspension PA Criteria:

See Promacta® prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms

Tavalisse® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Tavalisse® tablets Initial PA Criteria:

- Patient has a diagnosis of chronic immune thrombocytopenia; **AND**
- Trial and failure (platelet count ≥ 50 x 10^9/L not achieved) of ONE of the following:
  - Corticosteroids
  - Thrombopoietin receptor antagonists (e.g., Promacta)
  - Splenectomy
  - Azathioprine (Azasan, Imuran), cyclosporine (Neoral, Sandimmune), cyclophosphamide (Cytoxan), mycophenolate mofetil (CellCept), danazol, or rituximab (Rituxan); **AND**
- Patient is not on concomitant therapy with a strong CYP3A4 inducer; **AND**
- Patient has received a baseline and will receive ongoing routine monitoring that includes:
  - Neutropenia (measure ANC monthly)
  - Hepatotoxicity (measure LFTs monthly)
  - Hypertension (measure blood pressure every 2 weeks until stable dose established, then monthly); **AND**
- Prescribed by, or in consultation with, a hematologist

Tavalisse® tablets Renewal PA Criteria:

- Patient has laboratory values documenting platelet response to therapy (platelet count ≥ 50 x 10^9/L); **AND**
- Patient has not experienced severe adverse effect as a result of fostamatinib therapy

Wayrilz® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Wayrilz® tablets Initial PA Criteria:

- Patient is ≥ 18 years old; **AND**
- Patient has a diagnosis of chronic OR persistent immune thrombocytopenia; **AND**
- Trial and failure (platelet count ≥ 50 x 109/L not achieved) of ONE of the following:
  - Corticosteroids
  - Thrombopoietin receptor antagonists (e.g., Promacta)
  - Splenectomy
  - Azathioprine (Azasan, Imuran), cyclosporine (Neoral, Sandimmune), cyclophosphamide (Cytoxan), mycophenolate mofetil (CellCept), danazol, or rituximab (Rituxan); **AND**
- Patient is not on concomitant therapy with a strong CYP3A4 inducer; **AND**
- Females must use effective contraception due to embryo-fetal toxicity; **AND**
- Prescribed by, or in consultation with, a hematologist; **AND**
- Patient has received a baseline and will receive ongoing routine monitoring that includes:
  - Neutropenia (measure ANC monthly)
  - Hepatotoxicity (measure LFTs monthly)

Wayrilz® tablets Renewal PA Criteria:

- Patient has laboratory values documenting platelet response to therapy (platelet count ≥ 50 x 109/L);**AND**
- Patient has not experienced severe adverse effect as a result of Wayrilz® therapy

Vasodilator + Nitrates

BiDil® tablets

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

BiDil® tablets PA Criteria:
Standard NP Criteria

Vasopressors

Droxidopa, Northera® capsules

PDL Status	NP
Quantity Limit	100 mg & 200 mg: 3/day; 300 mg: 6/day
PA Criteria	See Below
PA Form	General PA Form

Droxidopa, Northera® capsules PA Criteria:

- Diagnosis of symptomatic neurogenic orthostatic hypotension secondary to primary autonomic failure, dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy; **AND**
- Trial and failure, contraindication, or intolerance to midodrine **OR** fludrocortisone

IV. CENTRAL NERVOUS SYSTEM

Agents for Narcolepsy

For agents in this class requiring PA, please submit the Narcolepsy Agents PA Form on faxed PA submissions.

Modafinil tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Modafinil tablets PA Criteria:

- Diagnosis of ADD/ADHD; **AND**
  - Contraindication, adverse reaction, or drug-drug interaction to **ALL** preferred antihyperkinesis agents; **OR**
- Daytime sleepiness/hypersomnolence occurring for at least 3 months; **AND**
  - Diagnosis is associated with ONE of the following:
    - Idiopathic hypersomnia
    - Narcolepsy
    - Obstructive sleep apnea/hypopnea syndrome supported by a documented sleep study, **AND**
      - Trial and failure (minimum duration 3 months with documented compliance) of Continuous Positive Airway Pressure (CPAP) or BiPAP device, unless contraindications
    - Shift Work Sleep Disorder; **AND**
      - Statement of patient's work schedule showing a minimum of 6 hours work between 10 pm and 8 am; **AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out

Provigil® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Provigil® tablets PA Criteria:
See modafinil prior authorization criteria.

Xyrem® solution

PDL Status	P
Quantity Limit	9 g/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Xyrem® PA Criteria:

- Enrolled in the Xyrem Program (1-866-997-3688); **AND**
- One of the following:
  - Diagnosis of cataplexy associated with narcolepsy
  - Diagnosis of excessive daytime sleepiness/hypersomnolence associated with narcolepsy occurring > 3 months; **AND**
    - Trial and failure, intolerance, or contraindication to modafinil; **AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out

Armodafinil tablets

PDL Status	NP
Quantity Limit	50 mg tablets: 2/day; 150, 200, & 250 mg tablets: 1/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Armodafinil tablets PA Criteria:

- Daytime sleepiness/hypersomnolence occurring for at least 3 months; **AND**
- Diagnosis is associated with ONE of the following:
  - Narcolepsy
  - Obstructive sleep apnea/hypopnea syndrome supported by a documented sleep study, **AND**
    - Trial and failure (minimum duration 3 months with documented compliance) of Continuous Positive Airway Pressure (CPAP) or BiPAP device, unless contraindications
  - Shift Work Sleep Disorder; **AND**
    - Statement of patient's work schedule showing a minimum of 6 hours work between 10 pm and 8 am; **AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out; **AND**
- Trial and failure, contraindication, or intolerance to modafinil

Nuvigil® tablets

PDL Status	NP
Quantity Limit	50 mg tablets: 2/day; 150, 200, & 250 mg tablets: 1/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Nuvigil® tablets PA Criteria:
See armodafinil tablets prior authorization criteria.

Sodium oxybate solution

PDL Status	NP
Quantity Limit	9 g/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Sodium oxybate solution PA Criteria:
See Xyrem® prior authorization criteria; **AND**

- Trial and failure of Xyrem®

Sunosi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Sunosi® tablets PA Criteria:

- Daytime sleepiness/hypersomnolence occurring for at least 3 months; **AND**
- Diagnosis is associated with ONE of the following:
  - Narcolepsy
  - Obstructive sleep apnea/hypopnea syndrome supported by a documented sleep study, **AND**
    - Trial and failure (minimum duration 3 months with documented compliance) of Continuous Positive Airway Pressure (CPAP) or BiPAP device, unless contraindications; **AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out; **AND**
- Trial and failure, contraindication, or intolerance to modafinil

Wakix®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Wakix® PA Criteria:

- Daytime sleepiness/hypersomnolence occurring for at least 3 months; **AND**
- ONE of the following:
  - Diagnosis of cataplexy associated with narcolepsy; **AND**
    - Trial and failure, contraindication, or intolerance to Xyrem
  - Diagnosis of excessive daytime sleepiness (EDS) associated with Narcolepsy; **AND**
    - Trial and failure, contraindication, or intolerance to modafinil; **AND**
  - Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out

Xywav®

PDL Status	NP
Quantity Limit	18 mL/day
PA Criteria	See below
PA Form	Narcolepsy Agents PA Form

Xywav® PA Criteria:

- Enrolled in the Xywav Program (1-866-997-3688); **AND**
- One of the following:
  - Diagnosis of cataplexy associated with narcolepsy; **AND**
    - Clinically valid reason is given why the patient requires Xywav over Xyrem
  - Diagnosis of excessive daytime sleepiness/hypersomnolence associated with narcolepsy occurring > 3 months; **AND**
    - Trial and failure, intolerance, or contraindication to modafinil; **AND**
    - Clinically valid reason is given why the patient requires Xywav over Xyrem
  - Diagnosis of idiopathic hypersomnia (IH) in patients ≥ 18 years of age; **AND**
    - Trial and failure, intolerance, or contraindication to modafinil; **AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out

Agents for Neuropathic Pain and Fibromyalgia

Duloxetine 20, 30, and 60 mg capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Gabapentin 100, 300, and 400 mg capsules

PDL Status	P
Quantity Limit	100 mg caps: 6/day; 300 mg caps: 12/day; 400 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	None

Gabapentin solution

PDL Status	P
Quantity Limit	72 mL/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Gabapentin solution PA Criteria:

- One of the following:
  - Patient is less than 12 years of age; **OR**
  - Patient is unable to swallow solid dosage forms

Gralise® once daily tablets

PDL Status	P
Quantity Limit	300, 450, and 600 mg tablets: 3/day; 750 and 900 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Gralise® once daily tablets PA Criteria:

- Diagnosis of post-herpetic neuralgia; **AND**
- Clinically valid reason why the preferred gabapentin agents cannot be used

Horizant® ER tablets

PDL Status	P
Quantity Limit	1/day; Gabapentin max daily dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Horizant® ER tabletsPA Criteria:

- Diagnosis of post-herpetic neuralgia; **OR**
- Diagnosis of Restless Leg Syndrome

Lidocaine 5% patches

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Lidocaine 5% patches PA Criteria:

- Diagnosis of post-herpetic neuralgia;

Pregabalin capsules

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pregabalin capsules PA Criteria:

- Diagnosis of post-herpetic neuralgia; **OR**
- Diagnosis of fibromyalgia; **OR**
- Diagnosis of seizure disorder

Pregabalin solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pregabalin solution PA Criteria:

See pregabalin capsules prior authorization criteria; **AND**

- Patient is less than 12 years of age; **OR**
- Patient is unable to swallow solid dosage forms

Cymbalta® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Cymbalta® PA Criteria:
Standard NP Criteria

Duloxetine 40 mg capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Duloxetine 40 mg capsules PA Criteria:

- Clinically valid reason as to why the preferred duloxetine strengths (20mg, 30mg, 60mg) cannot be used

Duloxetine 80, 90, and 120 mg capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Duloxetine 80, 90, and 120 mg capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Drizalma Sprinkle™ capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Drizalma Sprinkle™ capsules PA Criteria:

- Patient is unable to swallow solid dosage forms

Gabapentin 200 mg capsules

PDL Status	NP
Quantity Limit	9/day
PA Criteria	See below
PA Form	General PA Form

Gabapentin 200 mg capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Gabapentin tablets

PDL Status	NP
Quantity Limit	600 mg tabs: 6/day; 800 mg tabs: 4.5/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Gabapentin tablets PA Criteria:

- Documented allergy or contraindication to an inactive ingredient in the capsules that is NOT present in the tablets

Gabapentin once daily tablets

PDL Status	NP
Quantity Limit	300, 450, and 600 mg tablets: 3/day; 750 and 900 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Gabapentin once daily tablets PA Criteria:

- Diagnosis of post-herpetic neuralgia; **AND**
- Clinically valid reason why the preferred gabapentin agents cannot be used

Gabarone® tablets

PDL Status	NP
Quantity Limit	100 mg tablets: 6/day; 400 mg tablets: 9/day
PA Criteria	See below
PA Form	General PA Form

Gabarone® tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Lyrica® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lyrica® capsules PA Criteria:

See pregabalin capsules prior authorization criteria; **AND**

- Trial and failure of preferred pregabalin capsules

Lyrica® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lyrica® solution PA Criteria:

See pregabalin solution prior authorization criteria; **AND**

- Trial and failure of preferred pregabalin solution

Lyrica CR® tablets

PDL Status	NP
Quantity Limit	82.5 & 165 mg tabs: 1/day; 330 mg tabs: 2/day
PA Criteria	See below
PA Form	General PA Form

Lyrica CR® tablets PA Criteria:

- Diagnosis of postherpetic neuralgia **OR** neuropathic pain associated with-diabetic peripheral neuropathy; **AND**
- Trial and failure, contraindication, or intolerance to a tricyclic antidepressant **OR** gabapentin; **AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus immediate-release pregabalin

Milnacipran tablets and titration packs

PDL Status	NP
Quantity Limit	Tablets: 2 tabs/day; Titration packs: 1/year
PA Criteria	See below
PA Form	General PA Form

Milnacipran tablets and titration packs PA Criteria:

- Patient has a diagnosis of fibromyalgia accompanied by fatigue; **AND**
- Patient is 18 years of age or older; **AND**
- Patient MUST have tried and failed, or have contraindication, or intolerance to duloxetine; **AND**
- Clinically valid reason why brand Savella® cannot be used

Neurontin® capsules

PDL Status	NP
Quantity Limit	100 mg caps: 6/day; 300 mg caps: 12/day; 400 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® capsules PA Criteria:

Standard NP Criteria

Neurontin® solution

PDL Status	NP
Quantity Limit	72 mL/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® solution PA Criteria:

See gabapentin solution prior authorization criteria

Neurontin® tablets

PDL Status	NP
Quantity Limit	600 mg tabs: 6/day; 800 mg tabs: 4.5/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® tablets PA Criteria:
Standard NP Criteria

Pregabalin CR tablets

PDL Status	NP
Quantity Limit	82.5 & 165 mg tabs: 1/day; 330 mg tabs: 2/day
PA Criteria	See below
PA Form	General PA Form

Pregabalin CR tablets PA Criteria:
See Lyrica CR® prior authorization criteria

Relgaabi® 200 mg capsules

PDL Status	NP
Quantity Limit	200 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Relgaabi® 200 mg capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Savella® tablets and titration packs

PDL Status	NP
Quantity Limit	Tablets: 2 tabs/day; Titration packs: 1/year
PA Criteria	See below
PA Form	General PA Form

Savella® tablets and titration packs PA Criteria:

- Patient has a diagnosis of fibromyalgia accompanied by fatigue; **AND**
- Patient is 18 years of age or older; **AND**
- Patient MUST have tried and failed, or have contraindication, or intolerance to duloxetine.

Tonmya® SL tablets

PDL Status	NP
Quantity Limit	2 tabs/day
PA Criteria	See below
PA Form	General PA Form

Tonmya® SL tablets packs PA Criteria:

- Patient has a diagnosis of fibromyalgia; **AND**
- Patient is 18 years of age or older; **AND**

- Patient must have tried and failed, or have contraindication or intolerance to duloxetine AND Savella® (milnacipran); **OR**
- Patient is unable to swallow solid dosage forms

Agents for RLS (Restless Leg Syndrome)

Horizant® ER tablets

PDL Status	P
Quantity Limit	1 tab/day; Gabapentin max daily dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Horizant® PA Criteria:

- Diagnosis of post-herpetic neuralgia; **OR**
- Diagnosis of Restless Leg Syndrome

Pramipexole tablets

PDL Status	P
Quantity Limit	3 tabs/day
PA Criteria	None

Neupro® patches

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Neupro® patches PA Criteria:

- Diagnosis of Parkinson's Disease or Restless Leg Syndrome, **AND**
- Trial and failure, contraindication, or intolerance to Horizant, pramipexole, and ropinirole, **OR**
- Inability to swallow

Alzheimer's: Cholinesterase Inhibitors

Donepezil tablets (excluding 23 mg)

PDL Status	P
Quantity Limit	1 tab/day
PA Criteria	None

Donepezil ODT

PDL Status	P
Quantity Limit	1 ODT/day
PA Criteria	See below
PA Form	General PA Form

Donepezil ODT PA Criteria:

- Inability to swallow; **OR**
- Unable to absorb medications through the GI tract

Exelon® patches

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Aricept® tablets (except 23 mg)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Aricept® PA Criteria:
Standard NP Criteria.

Aricept® 23 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Aricept® 23 mg tablets PA Criteria:

- Patient has been established (at least 3 months) on therapy with Aricept 10 mg daily

Donepezil 23 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Donepezil 23 mg tablets PA Criteria:

- Patient has been established (at least 3 months) on therapy with Aricept 10 mg daily

Galantamine ER capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Galantamine ER capsules PA Criteria:
Standard NP Criteria.

Rivastigmine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Rivastigmine capsules PA Criteria:
Standard NP Criteria.

Rivastigmine patch

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Rivastigmine patch PA Criteria:
Standard NP Criteria.

Zunveyl® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Zunvey® tablets PA Criteria:
Standard NP Criteria.

Alzheimer's: NMDA Receptor Antagonists

Memantine tablets

PDL Status	P
Quantity Limit	5 and 10 mg tabs: 2/day; Titration pack: 1/Rx
PA Criteria	None

Memantine/donepezil caps

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Memantine/donepezil PA Criteria:

- Diagnosis of moderate to severe dementia associated with Alzheimer's disease; **AND**
- Concomitantly taking donepezil and memantine (immediate release or extended release) [≥10 mg/day on both agents]; **AND**
- Clinical reason why recipient is unable to take the components individually; **AND**
- Clinically valid reason why brand Namzaric® cannot be used

Memantine ER capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Memantine ER capsulesPA Criteria:

- Diagnosis of moderate to severe Alzheimer's disease

Memantine solution

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See below
PA Form	General PA Form

Memantine solution PA Criteria:

- Diagnosis of moderate to severe Alzheimer's disease

Namenda® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Namenda® tablets PA Criteria:
Standard NP Criteria

Namzaric® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Namzaric® capsules PA Criteria:

- Diagnosis of moderate to severe dementia associated with Alzheimer's disease; **AND**
- Concomitantly taking donepezil and memantine (immediate release or extended release) [≥10 mg/day on both agents]; **AND**
- Clinical reason why recipient is unable to take the components individually

Analeptics

Caffeine citrate solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	2 months

Caffeine citrate solution PA Criteria:

- Diagnosis of apnea in premature infants (born between 28 and <33 weeks gestational age); **AND**
- Patient is continuing therapy from an inpatient hospital stay (to facilitate transition to outpatient for completion of therapy); **AND**
- Infant does not have renal impairment, hepatic impairment, or cardiovascular disease; **AND**
- Prescriber must attest that they are aware of the risks of fatal necrotizing enterocolitis in premature infants and will monitor patient for efficacy and to avoid serious toxicity; **AND**
- Prescribed by, or in consultation with a board-certified neonatologist

Antiparkinson Agents: Adenosine Antagonists

Nourianz® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Nourianz® tablets Initial PA Criteria:

- Diagnosis of Parkinson's disease; **AND**
  - Patient is experiencing "off" episode; **AND**
- Patient is 18 years of age or older; **AND**
- Patient is currently being treated with a stable dosage of levodopa/carbidopa; **AND**
- Prescriber advises women of childbearing potential to use contraception during treatment; **AND**
- Prescriber agrees to monitor the following:
  - Patients with moderate hepatic impairment (Child-Pugh B) for adverse reactions
  - Exacerbation of pre-existing dyskinesia
  - Presence of hallucinations/psychotic behavior
  - Presence of impulse control/compulsive behaviors; **AND**
- Trial and failure, intolerance, or contraindication to ONE agent in TWO different antiparkinson classes (e.g., Dopamine Agents, Decarboxylase Inhibitors, COMT Inhibitors, MAO-B inhibitors, NMDA Antagonists)

Nourianz® tablets Renewal PA Criteria:

- Patient is currently being treated with levodopa/carbidopa; **AND**
- Patient has a positive clinical response to therapy (e.g., reduction in number or total daily hours of "off" episodes, increase "on" time without troublesome dyskinesia)

Antiparkinson Agents: COMT Inhibitors

Ongentys®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Ongentys® Initial PA Criteria:

- Diagnosis of Parkinson's disease; **AND**
- Patient is experiencing "off" episodes; **AND**
- Patient is currently being treated with a stable dose of carbidopa/levodopa; **AND**
- Trial and failure, intolerance, or contraindication to ONE agent in TWO different antiparkinson classes (e.g., dopamine agents, decarboxylase inhibitors, COMT inhibitors, MAO-B inhibitors, NMDA antagonists); **AND**
- Will not be taken concomitantly with a non-selective monoamine oxidase inhibitor (MAOI); **AND**
- Patient does not have a history of pheochromocytoma, paraganglioma, or other catecholamine secreting neoplasms

Ongentys® Renewal PA Criteria:

- Patient is currently being treated with levodopa/carbidopa; **AND**
- Patient has a positive clinical response to therapy (e.g., reduction in number or total daily hours of "off" episodes, increase "on" time without troublesome dyskinesia)

Tolcapone tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tolcapone tablets PA Criteria:
Standard NP Criteria

Antiparkinson Agents: Decarboxylase Inhibitors

Lodosyn® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lodosyn® tablets PA Criteria:
Standard NP Criteria

Antiparkinson Agents: Dopamine Agents

Pramipexole tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Apokyn® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Apokyn® injection PA Criteria:

- Patient has a diagnosis of Parkinson's disease; **AND**
- Patient is experiencing acute, intermittent treatment of "off" episodes; **AND**
- Must be 18 years of age or older; **AND**
- Patient is currently being treated with a carbidopa/levodopa agent; **AND**
- Patient has had a trial and failure, contraindication, or intolerance of TWO of the following preferred adjunct drugs prescribed in combination with levodopa/carbidopa, each from different classes:
  - MAO-B inhibitor: selegiline
  - COMT inhibitor: entacapone, carbidopa/levodopa/entacapone, Stalevo
  - Dopamine agonist: pramipexole, ropinirole; **AND**
- Patient must not meet any of the following:
  - Patient is on concomitant 5HT3 antagonist
  - Patient is pregnant
  - Patient has a sensitivity to sulfites

Apomorphine injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Apomorphine injection PA Criteria:
See Apokyn® prior authorization criteria; **AND**

- Clinically valid reason why brand Apokyn® cannot be used

Bromocriptine capsules, tablets, Cycloset® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Bromocriptine capsules, tablets, Cycloset® tablets PA Criteria:
Standard NP Criteria

Neupro® patch

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Neupro® patch PA Criteria:

- Diagnosis of Parkinson's Disease **OR** Restless Leg Syndrome, **AND**
  - Trial and failure, contraindication, or intolerance to BOTH pramipexole **AND** ropinirole, **OR**
  - Inability to swallow

Parlodel® capsules, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Parlodel® capsules, tablets PA Criteria:
Standard NP Criteria

Pramipexole ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Pramipexole ER tablets PA Criteria:
Standard NP Criteria

Ropinirole ER tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Ropinirole ER tablets PA Criteria:
Standard NP Criteria

Antiparkinson Agents: Levodopa and Combinations

Crexont® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Crexont® capsules PA Criteria:
Standard NP Criteria

Carbidopa/levodopa ER capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carbidopa/levodopa ER capsules PA Criteria:

- Clinically valid reason why brand Rytary® cannot be used

Dhivy® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dhivy® tablets PA Criteria:

- Clinically valid reason why all preferred carbidopa/levodopa agents cannot be used

Inbrija® capsules

PDL Status	NP
Quantity Limit	60 blisters/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Inbrija® capsules Initial PA Criteria:

- Diagnosis of Parkinson's disease; **AND**
- Experiencing "off" episodes; **AND**
- Patient is currently being treated with a stable dose of carbidopa/levodopa; **AND**
- Trial and failure, intolerance, or contraindication to ONE agent in TWO different antiparkinson classes (e.g., dopamine agents, decarboxylase inhibitors, COMT inhibitors, MAO-B inhibitors, NMDA antagonists); **AND**
- Will not be taken concomitantly with a non-selective monoamine oxidase inhibitor (MAOI); **AND**
- Patient does not have asthma, COPD, or other chronic lung disease

Inbrija® capsules Renewal PA Criteria:

- Patient is currently being treated with levodopa/carbidopa; **AND**
- Patient has a positive clinical response to therapy (e.g., reduction in number or total daily hours of "off" episodes, increase "on" time without troublesome dyskinesia)

Rytary® capsules, Sinemet® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Rytary® capsules, Sinemet® tablets PA Criteria:
Standard NP Criteria

Antiparkinson Agents: MAOI-Bs

Azilect®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Azilect® PA Criteria:
Standard NP Criteria

Xadago®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Xadago® PA Criteria:
Standard NP Criteria

Antiparkinson Agents: NMDA Antagonists

Gocovri®

PDL Status	NP
Quantity Limit	68.5 mg: 1/day; 127 mg: 2/day
PA Criteria	See below
PA Form	General PA Form

Gocovri® Initial PA Criteria:

- One of the following:
  - Patient has a diagnosis of dyskinesia associated with Parkinson's disease
  - Patient is experiencing "off" episodes; **AND**
- Patient must be on concomitant levodopa-based therapy; **AND**
- Patient has tried/failed an adequate trial of or is intolerant to amantadine immediate release; **AND**
- Patient does not have end-stage renal disease (creatinine clearance < 15 mL/min/1.73 m_2)

Gocovri® Renewal PA Criteria:

- Patient is currently being treated with levodopa/carbidopa; **AND**
- Patient has a positive clinical response to therapy (e.g., reduction in number or total daily hours of "off" episodes, increase "on" time without troublesome dyskinesia)

Anti-Anxiety and Anti-Panic Agents

For agents in this class requiring PA, please submit the Anti-Anxiety PA Form on faxed PA submissions.

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Anti-anxiety agents prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** I/DD Prior Authorization Form

Alprazolam tablet

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Alprazolam tablet PA Criteria:

- Diagnosis of one of the following:
  - Anxiety disorder
  - Panic disorder with or without agoraphobia; **AND**
- Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., cognitive behavioral therapy, worry exposure, applied relaxation, muscle relaxation, short-term psychodynamic psychotherapy, mindfulness-based therapy); **AND**
- Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
  - SSRI (minimum trial duration of 4 weeks)
  - SNRI (minimum trial duration of 4 weeks)
  - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse; **AND**
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Alprazolam ER tablet

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Alprazolam ER tablet PA Criteria:

See alprazolam tablets prior authorization criteria

Buspirone tablets (5 mg, 10 mg, 15 mg, 30 mg)

PDL Status	P
Quantity Limit	30 mg tabs: 2/day; All other strengths: 3/day
PA Criteria	None

Chlordiazepoxide capsules

PDL Status	P
Quantity Limit	4/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Chlordiazepoxide capsules PA Criteria:

- Diagnosis of acute alcohol withdrawal syndrome; **OR**
- Diagnosis of anxiety disorder; **AND**
  - Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., cognitive behavioral therapy, worry exposure, applied relaxation, muscle relaxation, mindfulness-based therapy); **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Clonazepam 0.5, 1, and 2 mg tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Clonazepam 0.5, 1, and 2 mg tablets PA Criteria:

- Diagnosis of seizure disorder; **OR**
- Diagnosis of panic disorder; **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Clorazepate tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Clorazepate tablets PA Criteria:

- Diagnosis of acute alcohol withdrawal syndrome; **OR**
- Diagnosis of seizure disorder; **AND**
  - Must be used in conjunction with another anticonvulsant; **OR**
- Diagnosis of anxiety disorder; **AND**
  - Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., cognitive behavioral therapy, worry exposure, applied relaxation, muscle relaxation, mindfulness-based therapy); **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol/meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse (does not apply to diagnosis of acute alcohol withdrawal syndrome); **AND**
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Diazepam concentrate, solution, tablets

PDL Status	P
Quantity Limit	Concentrate: 2 mL/day; Solution: 10 mL/day; Tablets: 4/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Diazepam concentrate, solution, tablets PA Criteria:

- Diagnosis of acute alcohol withdrawal syndrome; **OR**
- Diagnosis of seizure disorder; **AND**
  - Must be used in conjunction with another anticonvulsant; **OR**
- Diagnosis of muscle spasms; **AND**
  - Patient has tried and failed at least TWO preferred skeletal muscle relaxants; **OR**
- Diagnosis of anxiety disorder; **AND**
  - Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., cognitive behavioral therapy, worry exposure, applied relaxation, muscle relaxation, mindfulness-based therapy); **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse (does not apply to diagnosis of acute alcohol withdrawal syndrome); **AND**
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Lorazepam concentrate, tablets

PDL Status	P
Quantity Limit	Concentrate: 3 mL/day; Tablets: 3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Lorazepam concentrate, tablets PA Criteria:

- Patient is < 1 year of age and completing taper following inpatient hospital use for Neonatal Withdrawal symptoms; **OR**
- Diagnosis of anxiety disorder; **AND**
  - Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., cognitive behavioral therapy, worry exposure, applied relaxation, muscle relaxation, mindfulness-based therapy); **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse; **AND**
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Xanax® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Xanax® tablets PA Criteria:

See alprazolam tablets prior authorization criteria

Alprazolam concentrate, ODT

PDL Status	NP
Quantity Limit	Concentrate: 6 mL/day; ODT: 3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Alprazolam concentrate, ODT PA Criteria:

See alprazolam tablets prior authorization criteria

- Patient is unable to swallow solid dosage forms or unable to absorb medications through the GI tract; **AND**
- Trial and failure, contraindication, or intolerance to BOTH preferred concentrate solutions

Bucapsol® capsules

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Bucapsol® capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Buspirone 2.5 mg and 12.5 mg tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Buspirone 2.5 mg and 12.5 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Clonazepam 0.25 mg tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Clonazepam 0.25 mg tablets PA Criteria:

See clonazepam tablets prior authorization criteria; **AND**

- Requested dose is unattainable with preferred clonazepam 0.5mg tablets

Loreev XR®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Loreev XR® PA Criteria:

See lorazepam prior authorization criteria; **AND**

- Clinically valid reason why the preferred lorazepam tablets or concentrate cannot be used

Meprobamate

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Meprobamate PA Criteria:

See alprazolam tablets prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance of TWO preferred agents

Oxazepam

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Oxazepam PA Criteria:

See chlordiazepoxide capsule prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance of TWO preferred agents

Anticonvulsants

Clobazam tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Clobazam tablets PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant

Clonazepam 0.5, 1, and 2 mg tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-anxiety PA Form

Clonazepam 0.5, 1, and 2 mg tablets PA Criteria:

- Diagnosis of seizure disorder; **OR**
- Diagnosis of panic disorder; **AND**
    - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
        - SSRI (minimum trial duration of 4 weeks)
        - SNRI (minimum trial duration of 4 weeks)
        - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use

Diazepam rectal gel (10 and 20 mg)

PDL Status	P
Quantity Limit	2 packs/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	PA not required for patients less than 21 years of age.

Diazepam rectal gel PA Criteria:

- Will be approved for patients 21 years of age and older with a Diagnosis of Seizure Disorder or Epilepsy.

Eslicarbazepine tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Eslicarbazepine tablets PA Criteria:

- Use as monotherapy for partial onset seizures and trial and failure with ONE preferred anticonvulsant with the same indication; **OR**
- Use as adjunctive therapy for partial onset seizures when used in combination with at least ONE other anticonvulsant.

Epidiolex® solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Epidiolex solution Initial PA Criteria:

- Diagnosis of one of the following:
  - Dravet Syndrome (DS)
  - Lennox-Gastaut Syndrome (LGS)
  - Tuberous sclerosis complex (TSC)
  - Treatment-Refractory Epilepsy; **AND**
- Trial of 2 anticonvulsants within the past 12 months (documented by claims); **AND**
- Epidiolex will be used as adjunct therapy with ≥ 1 anticonvulsant (documented by claims)

Epidiolex® solution Renewal PA Criteria:

- Epidiolex will be used as adjunct therapy with ≥ 1 anticonvulsant (documented by claims)

Gabapentin 100, 200, and 300 mg capsules

PDL Status	P
Quantity Limit	100 mg caps: 6/day; 300 mg caps: 12/day; 400 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	None

Gabapentin solution

PDL Status	P
Quantity Limit	72 mL/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Gabapentin solution PA Criteria:

- Inability to swallow solid oral dosage forms; **AND**
  - Patient and caregiver are unable to open capsule and empty contents in food or drink; **OR**
- Patient is less than 12 years of age; **OR**

Lacosamide tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lacosamide tablets PA Criteria:

- Use as monotherapy for partial onset seizures requires trial and failure with at least ONE other preferred anticonvulsant for the same indication; **OR**
- Use as adjunctive therapy for partial onset seizures when used in combination with at least ONE other anticonvulsant; **OR**
- Used as adjunctive therapy in the treatment of primary generalized tonic-clonic (PGTC) seizures in patients 4 years of age and older

Libervant® buccal film

PDL Status	P
Quantity Limit	10 doses/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Libervant® buccal film PA Initial Criteria:

- Diagnosis of intermittent, stereotypic episodes of frequent seizure activity (e.g., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern; **AND**
- Patient is 2 to 5 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient is on a stable antiepileptic regimen; **AND**
- Prescriber has counseled patient on the following:
  - Risks if combined with opioids or benzodiazepines
  - Identification of a seizure cluster
  - Proper administration
  - When to seek emergency medical treatment; **AND**
- Patient is not using moderate or strong CYP2C19 and CYP 3A4 inhibitors or, if unavoidable, prescriber will monitor toxicity risk during concomitant use; **AND**
- Patient does not have acute narrow-angle glaucoma

Libervant® buccal film Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not have treatment-limiting adverse effects (e.g., treatment-limiting central nervous system depression or cognitive impairment, worsened glaucoma, respiratory depression, suicidal ideation, clinically significant changes in blood pressure or heart rate); **AND**
- Prescriber to provide verbal attestation of diazepam effectiveness (e.g., decreased typical length of repetitive seizures)

Nayzilam® nasal spray

PDL Status	P
Quantity Limit	10 doses/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Nayzilam® nasal spray Initial PA Criteria:

- Patient has diagnosis of intermittent, stereotypic episodes of frequent seizure activity (e.g., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern; **AND**
- Patient is 12 years of age or older; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient is on a stable antiepileptic regimen; **AND**
- Prescriber has counseled patient on the following:
  - Risks if combined with opioids
  - Identification of a seizure cluster
  - Proper administration
  - When to seek emergency medical treatment; **AND**
- Patient is not using moderate or strong CYP 3A4 inhibitors or, if unavoidable, prescriber will monitor toxicity risk during concomitant use; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, CNS depressants, carisoprodol, meprobamate, or barbiturates; **AND**
- Patient does not have acute narrow-angle glaucoma

Nayzilam® nasal spray Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not have treatment-limiting adverse effects (e.g., treatment-limiting central nervous system depression or cognitive impairment, worsened glaucoma, respiratory depression, suicidal ideation, clinically significant changes in blood pressure or heart rate); **AND**
- Prescriber to provide verbal attestation of midazolam effectiveness (e.g., decreased typical length of repetitive seizures)

Pregabalin capsules

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pregabalin capsules PA Criteria:

- Diagnosis of post-herpetic neuralgia; **OR**
- Diagnosis of fibromyalgia; **OR**
- Diagnosis of seizure disorder

Pregabalin solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pregabalin solution PA Criteria:

See pregabalin capsules prior authorization criteria; **AND**

- Patient is less than 12 years of age; **OR**
- Patient is unable to swallow solid dosage forms

Phenobarbital elixir

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	PA not required for patients less than 2 years of age

Phenobarbital elixir PA Criteria:

- Will be approved for use ONLY in patients with diagnosis of seizure disorder

Phenobarbital tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Phenobarbital tablets PA Criteria:

- Will be approved for use ONLY in patients with diagnosis of seizure disorder

Rufinamide suspension

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Rufinamide suspension PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant; **AND**
- Patient is unable to swallow solid dosage forms

Rufinamide tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Rufinamide tablets PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant

Trokendi XR® tablets

PDL Status	P
Quantity Limit	25, 50, & 100 mg: 1/day; 200 mg: 2/day
PA Criteria	See below
PA Form	General PA Form

Trokendi XR® tablets PA Criteria:

- Adjunctive therapy for patients with partial-onset seizures or primary generalized tonic-clonic seizures; **OR** seizures associated with Lennox-Gastaut syndrome; **AND**
  - Will be used approved in combination with at least one other anticonvulsant; **AND**
  - Trial and failure of preferred immediate release product and one additional preferred agent; **OR**
- Initial monotherapy in patients with partial-onset or primary generalized tonic-clonic seizures; **AND**
  - Trial and failure of preferred immediate release product and one additional preferred agent; **OR**
- Migraine Prophylaxis in patients ≥ 12 years of age

Valtoco® nasal spray

PDL Status	P
Quantity Limit	10 doses/30 days
PA Criteria	See below
PA Form	General PA Form

Valtoco® nasal spray Initial PA Criteria:

- Patient has diagnosis of intermittent, stereotypic episodes of frequent seizure activity (e.g., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern; **AND**
- Patient is 2 years of age or older; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient is on a stable antiepileptic regimen; **AND**
- Prescriber has counseled patient on the following:
  - Risks if combined with opioids
  - Identification of a seizure cluster
  - Proper administration
  - When to seek emergency medical treatment; **AND**
- Patient is not using CYP 2C19 and CYP 3A4 inhibitors or, if unavoidable, prescriber will monitor toxicity risk during concomitant use; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, CNS depressants, carisoprodol, meprobamate, or barbiturates; **AND**
- Patient does not have acute narrow-angle glaucoma

Valtoco® nasal spray Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not have treatment-limiting adverse effects (e.g., treatment-limiting CNS depression or cognitive impairment, worsened glaucoma, respiratory depression, suicidal ideation, clinically significant changes in blood pressure); **AND**
- Prescriber to provide verbal attestation of diazepam effectiveness (e.g., decreased typical length of repetitive seizures)

Zonisamide capsules

PDL Status	P
Quantity Limit	25 mg: 4/day; 50 mg: 2/day; 100 mg: 6/day
PA Criteria	None

Ztalmy® suspension

PDL Status	P
Quantity Limit	36 mL/day
PA Criteria	See below
PA Form	General PA Form

Ztalmy® suspension Initial PA Criteria:

- Patient is 2 years of age and older; **AND**
- Diagnosis of seizure disorder associated with cyclin-dependent kinase-like 5 deficiency disorder; **AND**
- Prescriber has confirmed that patient is not pregnant (if applicable) and counseled patient on risks of pregnancy while taking Ztalmy; **AND**
- Prescriber has confirmed member does not have hepatic disease and will monitor hepatic function (dose reductions may be required in impaired hepatic function)

Ztalmy® suspension Renewal PA Criteria:

- Prescriber has confirmed that patient is not pregnant (if applicable); **AND**
- Prescriber has confirmed member does not have hepatic disease and will monitor hepatic function (dose reductions may be required in impaired hepatic function)

Aptiom® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Aptiom® tablets PA Criteria:

- Use as monotherapy for partial onset seizures and trial and failure with ONE preferred anticonvulsant with the same indication; **OR**
- Use as adjunctive therapy for partial onset seizures when used in combination with at least ONE other anticonvulsant; **AND**
- Clinically valid reason why the generic eslicarbazepine cannot be used

Banzel® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Banzel® suspension PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant; **AND**
- Patient is unable to swallow solid dosage forms

Banzel® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Banzel® tablets PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant

Brivaracetam solution

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See below
PA Form	General PA Form

Brivaracetam solution PA Criteria:

- Patient is ≥ 1 month of age; **AND**
- Have diagnosis of partial-onset seizures; **AND**
- Have tried and failed at least 1 other medication indicated for partial-onset seizures; **AND**
- Patient is unable to swallow solid dosage forms

Brivaracetam tablets

PDL Status	NP
Quantity Limit	2 /day
PA Criteria	See below
PA Form	General PA Form

Brivaracetam tablets PA Criteria:

- Patient is ≥ 1 month of age; **AND**
- Have diagnosis of partial-onset seizures; **AND**
- Have tried and failed at least 1 other medication indicated for partial-onset seizures

Briviact® solution

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See below
PA Form	General PA Form

Briviact® solution PA Criteria:

- Patient is ≥ 1 month of age; **AND**
- Have diagnosis of partial-onset seizures; **AND**
- Have tried and failed at least 1 other medication indicated for partial-onset seizures; **AND**
- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why generic brivaracetam solution cannot be used

Briviact® tablets

PDL Status	NP
Quantity Limit	2 /day
PA Criteria	See below
PA Form	General PA Form

Briviact® tablets PA Criteria:

- Patient is ≥ 1 month of age; **AND**
- Have diagnosis of partial-onset seizures; **AND**
- Have tried and failed at least 1 other medication indicated for partial-onset seizures

Carbamazepine suspension, tablets, Carbatrol® ER capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carbamazepine suspension, tablets, Carbatrol® ER capsules PA Criteria:

Standard NP Criteria

Clobazam suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Clobazam suspension PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunct therapy with at least one other anticonvulsant; **AND**
- Patient unable to swallow solid dosage forms

Clonazepam 0.25 mg tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Clonazepam 0.25 mg tablets PA Criteria:

See clonazepam tablets prior authorization criteria; **AND**

- Requested dose is unattainable with preferred clonazepam 0.5mg tablets

Clonazepam ODT

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-Anxiety PA Form

Clonazepam ODT PA Criteria:

- Diagnosis of seizure disorder; **OR**
- Diagnosis of panic disorder; **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use; **AND**
- Patient must be unable to swallow **OR** unable to absorb medications through the GI tract

Depakote® tablets, Depakote ER® capsules, tablets, Depakote Sprinkles® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Depakote® tablets, Depakote ER® capsules, tablets, Depakote Sprinkles® capsules PA Criteria:

Standard NP Criteria

Diacomit® capsules, packs

PDL Status	NP
Quantity Limit	250 mg: 1/day; 500 mg: 6/day
PA Criteria	See below
PA Form	General PA Form

Diacomit® capsules, packs Initial PA Criteria:

- Patient is ≥ 6 months of age and weighing at least 7 kg; **AND**
- Patient must also be taking clobazam concomitantly; **AND**
- Diagnosis of Dravet syndrome (DS); **AND**
- Prescribed by neurologist or epileptologist; **AND**
- Prescriber attests that baseline serum hematologic testing has been completed; **AND**
- Prescriber attests the patient has refractory epilepsy (failed to become seizure free after trials of 2 antiepileptic drugs); **AND**
- If the oral powder for suspension is prescribed, the patient does not have phenylketonuria (PKU).

Diacomit® capsules, packs Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has no treatment-limiting adverse effects (e.g., thrombocytopenia, neutropenia, new onset or worsened depression; suicidal thoughts, worsened seizure control); **AND**
- Prescriber to provide verbal attestation of Diacomit effectiveness (e.g., reduced seizure frequency and/or duration)

Dilantin 125® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dilantin 125® suspension PA Criteria:

Standard NP Criteria

Elepsia XR® tablets

PDL Status	NP
Quantity Limit	1000 mg: 3/day; 1500 mg: 2/day
PA Criteria	See below
PA Form	General PA Form

Elepsia XR® tablets PA Criteria:

- Patient has a diagnosis or history of partial-onset seizures, juvenile myoclonic epilepsy, or primary generalized tonic-clonic seizure; **AND**
- Will be used as adjunctive therapy with another anticonvulsant; **AND**
- Patient must be 12 years of age or older; **AND**
- Clinically valid reason why the preferred levetiracetam ER cannot be used

Epitol® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Epitol® tablets PA Criteria:

Standard NP Criteria

Eprontia® solution

PDL Status	NP
Quantity Limit	16 mL/day
PA Criteria	See below
PA Form	General PA Form

Eprontia® solution PA Criteria:

- Patient is unable to swallow tablets; **AND**
- If request is for Eprontia®, clinically valid reason why the generic topiramate solution cannot be used

Felbatol® suspension, tablets, felbamate suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	PA will not be approved if there is a history of blood dyscrasia or liver disease unless the prescriber can make a compelling clinical case demonstrating that the benefits of the drug outweigh the risks.

Felbatol® suspension, tablets, felbamate suspension, tablets PA Criteria:

- Used as adjunctive therapy for the treatment of partial and generalized seizures associated with Lennox-Gastaut Syndrome in children 2-14 years of age with a contraindication to, or trial and failure of, **TWO** of the following:
  - Valproic acid/divalproex sodium
  - Lamotrigine
  - Topiramate
- Used as monotherapy or adjunctive therapy for the treatment of partial seizures with or without generalization in adults > 14 years of age with a contraindication to, or trial and failure of, **THREE** of the following:
  - Carbamazepine
  - Oxcarbazepine
  - Phenytoin
  - Gabapentin
  - Lamotrigine
  - Topiramate
  - Valproic acid/divalproex sodium
  - Levetiracetam
  - Zonisamide

Fintepla® solution

PDL Status	NP
Quantity Limit	1 bottle/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Fintepla® solution Initial PA Criteria:

- Patient must be ≥ 2 years of age; **AND**
- Diagnosis of Dravet syndrome (DS) or Lennox-Gastaut syndrome (LGS); **AND**
- Prescribed by, or in consultation with, a neurologist or epileptologist; **AND**
- Patient has not received MAOI therapy within 14 days and will not receive during Fintepla therapy; **AND**
- Prescriber attestation that baseline echocardiogram has been completed and will be monitored throughout treatment and 3 – 6 months after the final dose; **AND**
- Inadequate response to trials of 2 preferred anticonvulsant agents

Fintepla® solution Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has no treatment-limiting adverse effects (e.g., serotonin syndrome, abnormal AST/ALT, CrCl, abnormal echocardiogram); **AND**
- Patient is responding to therapy (e.g., reduced seizure frequency and/or duration)

Fycompa® suspension

PDL Status	NP
Quantity Limit	Suspension: none
PA Criteria	See below
PA Form	General PA Form

Fycompa® suspension PA Criteria:

- Diagnosis of partial onset seizures with or without secondarily generalized seizures; **AND**
  - Patient is ≥ 4 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred agents, one of which must be lacosamide **OR**
- Will be used as adjunctive therapy for the treatment of primary generalized tonic-clonic (PGTC) seizures; **AND**
  - Patient is ≥ 12 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents

Fycompa® tablets

PDL Status	NP
Quantity Limit	Tablets: 2, 4, 8, 10 & 12 mg: 1/day 6 mg: 2/day;
PA Criteria	See below
PA Form	General PA Form

Fycompa® suspension, tablets PA Criteria:

- Diagnosis of partial onset seizures with or without secondarily generalized seizures; **AND**
  - Patient is ≥ 4 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred agents, one of which must be lacosamide **OR**
- Will be used as adjunctive therapy for the treatment of primary generalized tonic-clonic (PGTC) seizures; **AND**
  - Patient is ≥ 12 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents; **AND**
- If request is for Fycompa® tablets, clinically valid reason why generic perampanel tablets cannot be used

Gabapentin 200 mg capsules

PDL Status	NP
Quantity Limit	9/day
PA Criteria	See below
PA Form	General PA Form

Gabapentin 200 mg capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Gabapentin tablets

PDL Status	NP
Quantity Limit	600 mg tabs: 6/day; 800 mg tabs: 4.5/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Gabapentin tablets PA Criteria:

- Documented allergy or contraindication to an inactive ingredient in the capsules that is NOT present in the tablets

Gabarone® tablets

PDL Status	NP
Quantity Limit	100 mg tablets: 6/day; 400 mg tablets: 9/day
PA Criteria	See below
PA Form	General PA Form

Gabarone® tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Keppra® solution, tablets, Keppra XR® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Keppra® solution, tablets, Keppra XR® tablets PA Criteria:

Standard NP Criteria

Klonopin® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	Anti-anxiety PA Form

Klonopin® tablets PA Criteria:

- Diagnosis of seizure disorder; **OR**
- Diagnosis of panic disorder; **AND**
  - Trial and failure, contraindication, or intolerance to therapy with TWO of the following:
    - SSRI (minimum trial duration of 4 weeks)
    - SNRI (minimum trial duration of 4 weeks)
    - Buspirone; **AND**
- Due to increased risk of toxicity, patient should not be pregnant **OR** concurrently taking CNS stimulants, opiates, carisoprodol, meprobamate, or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse
- Prescriber has checked the Tennessee Controlled Substance Monitoring Database (CSMD) on the date of the request for concomitant controlled substance use; **AND**
- Trial and failure of clonazepam

Lacosamide solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lacosamide solution PA Criteria:
Standard NP Criteria

Lamictal® chewable tablets, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamictal® chewable tablets, tablets PA Criteria:
Standard NP Criteria

Lamictal ODT®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamictal ODT® PA Criteria:

- Patient is unable to swallow solid dosage forms

Lamictal XR® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamictal XR® tablets PA Criteria:
Standard NP Criteria.

Lamotrigine ODT

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamotrigine ODT PA Criteria:

- Patient is unable to swallow solid dosage forms

Levetiracetam soluble tablets, Spritam® soluble tablets

PDL Status	NP
Quantity Limit	250, 500, & 1000 mg: 2/day; 750 mg: 4/day
PA Criteria	See below
PA Form	General PA Form

Levetiracetam soluble tablets, Spritam® soluble tablets PA Criteria:

- Patient is unable to swallow solid oral dosage form; **AND**
- Provider must have a clinically valid reason as to why the generic levetiracetam solution cannot be used

Lyrica® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lyrica® capsules PA Criteria:

See pregabalin capsules prior authorization criteria; **AND**

- Trial and failure of preferred pregabalin capsules

Lyrica® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lyrica® capsules PA Criteria:

See pregabalin solution prior authorization criteria; **AND**

- Trial and failure of preferred pregabalin solution

Methsuximide capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Methsuximide capsules PA Criteria:

Standard NP Criteria

Motpoly XR® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Motpoly XR® capsules PA Criteria:

- One of the following:
  - Initial monotherapy for partial onset seizures
  - Adjunctive therapy for partial onset seizures and will be used in combination with at least one other anticonvulsant; **AND**
- Trial and failure of preferred immediate release product and one additional preferred agent

Neurontin® capsules

PDL Status	NP
Quantity Limit	100 mg caps: 6/day; 300 mg caps: 12/day; 400 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® capsules PA Criteria:

Standard NP Criteria

Neurontin® solution

PDL Status	NP
Quantity Limit	72 mL/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® solution PA Criteria:

See gabapentin solution prior authorization criteria

Neurontin® tablets

PDL Status	NP
Quantity Limit	600 mg caps: 6/day; 800 mg caps: 4.5/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Neurontin® tablets PA Criteria:

Standard NP Criteria

Onfi® suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Onfi® suspension, tablets PA Criteria:

- Diagnosis of Lennox-Gastaut Syndrome; **AND**
- Used as adjunctive therapy with at least one other anticonvulsant; **AND**
- Trial and failure of the generic equivalent (clobazam); **AND**
- If request is for suspension, patient is unable to swallow solid dosage forms

Oxtellar XR® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Oxtellar XR® tablets PA Criteria:

Standard NP Criteria

Perampanel suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Perampanel suspension PA Criteria:

- Diagnosis of partial onset seizures with or without secondarily generalized seizures; **AND**
  - Patient is ≥ 4 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred agents, one of which must be lacosamide **OR**
- Will be used as adjunctive therapy for the treatment of primary generalized tonic-clonic (PGTC) seizures; **AND**
  - Patient is ≥ 12 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents; **AND**
- Clinically valid reason why brand Fycompa® cannot be used

Perampanel tablets

PDL Status	NP
Quantity Limit	2, 4, 8, 10 & 12 mg: 1 tabs/day; 6 mg: 2 tabs/day
PA Criteria	See below
PA Form	General PA Form

Perampanel tablets PA Criteria:

- Diagnosis of partial onset seizures with or without secondarily generalized seizures; **AND**
  - Patient is ≥ 4 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred agents, one of which must be lacosamide **OR**
- Will be used as adjunctive therapy for the treatment of primary generalized tonic-clonic (PGTC) seizures; **AND**
  - Patient is ≥ 12 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents

Phenytek® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Phenytek® capsules PA Criteria:
Standard NP Criteria

Relgaabi® 200 mg capsules

PDL Status	NP
Quantity Limit	200 mg caps: 9/day; Max daily gabapentin dose: 3,600 mg
PA Criteria	See below
PA Form	General PA Form

Relgaabi® 200 mg capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Sabril® powder, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	This drug is subject to REMS requirements to ensure the benefits of treatment outweigh the risks of vision loss.

Sabril® powder, tablets PA Criteria:

- Adjunctive therapy for patients with refractory complex partial seizures who have responded inadequately to several alternative treatments; **AND**
  - Patient has tried and failed 2 preferred anticonvulsants; **OR**
  - Monotherapy for patients with infantile spasms

Sympazan® oral film

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Sympazan® oral film PA Criteria:

- Patient has a diagnosis of Lennox-Gastaut syndrome (LGS); **AND**
- Requested drug will be used as adjunctive therapy in combination with at least one other anticonvulsant; **AND**
- Provider must have a clinically valid reason as to why both clobazam tablets and suspension cannot be used.

Subvenite® suspension

PDL Status	NP
Quantity Limit	40 mL/day
PA Criteria	See below
PA Form	General PA Form

Subvenite® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why lamotrigine ODT cannot be used

Tegretol XR® tablets, tiagabine tablets, Topamax® sprinkle caps, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tegretol XR® tablets, tiagabine tablets, Topamax® sprinkle caps, tablets PA Criteria:

Standard NP Criteria

Topiramate ER capsules

PDL Status	NP
Quantity Limit	200 mg: 2/day; All other strengths: 1/day
PA Criteria	See below
PA Form	General PA Form

Topiramate ER capsules PA Criteria:

- Will be used as monotherapy or adjunctive therapy in patients with focal (partial) onset or primary generalized tonic-clonic seizures; **OR**
- Will be used as adjunctive therapy in patients with seizures associated with Lennox-Gastaut syndrome; **OR**
- Migraine Prophylaxis in patients ≥ 12 years of age; **AND**
- Trial and failure of an Trokendi XR and 1 other preferred agent

Topiramate solution

PDL Status	NP
Quantity Limit	16 mL/day
PA Criteria	See below
PA Form	General PA Form

Topiramate solution PA Criteria:

See Eprontia® prior authorization criteria; **AND**

Trileptal® suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Trileptal® suspension, tablets PA Criteria:

Standard NP Criteria

Vigabatrin packs, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vigabatrin packs, tablets PA Criteria:
See Sabril® prior authorization criteria

Vigafyde® oral solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	This drug is subject to REMS requirements to ensure the benefits of treatment outweigh the risks of vision loss.

Vigafyde® oral solution PA Criteria:

- Treatment is for monotherapy for patients with infantile spasms; **AND**
- Clinically valid reason why vigabatrin 50 mg/mL powder oral solution cannot be used

Vigadrone® packs, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vigadrone® packs, tablets PA Criteria:
See Sabril® prior authorization criteria

Vimpat® solution, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vimpat® solution, tablets PA Criteria:
See lacosamide prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to lacosamide

Xcopri® packs, tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Xcopri® packs, tablets Initial PA Criteria:

- Diagnosis of partial-onset seizures; **AND**
- Prescribed by, or in consultation with, a neurologist or epileptologist; **AND**
- Must be 18 years of age and older; **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred anticonvulsants indicated for partial-onset seizures; **AND**
- Patient does not have Familial Short QT syndrome

Xcopri® packs, tablets Renewal PA Criteria:

- Patient must demonstrate disease improvement and stabilization as a result of the medication; **AND**
- Patient is absent of unacceptable toxicity from the drug; **AND**
- Patient's QT interval is being monitored

Zarontin® capsules, solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Zarontin® capsules, solution PA Criteria:

Standard NP Criteria

Zonisade® suspension

PDL Status	NP
Quantity Limit	30 mL/day
PA Criteria	See below
PA Form	General PA Form

Zonisade® suspension PA Criteria:

- Diagnosis of partial-onset seizures; **AND**
- Zonisade will be used as adjunctive therapy; **AND**
- Patient must be unable to swallow solid dosage forms

Antidepressants: MAOIs

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antidepressants prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** **I/DD Prior Authorization Form**

Phenelzine

PDL Status	P
Quantity Limit	6 tabs/day
PA Criteria	See below
PA Form	General PA Form

Phenelzine PA Criteria:

- Diagnosis of major depression; **AND**
- Trial and failure, contraindication, or intolerance to an agent from at least TWO of the following drug classes:
  - SSRIs
  - SNRIs
  - New generation antidepressants

Emsam®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Emsam® PA Criteria:

- Diagnosis of major depression; **AND**
- Trial and failure, contraindication, or intolerance to an agent from at least TWO of the following drug classes:
  - SSRIs
  - SNRIs
  - New generation antidepressants; **AND**
- Trial and failure, contraindication, or intolerance to preferred phenelzine; **AND**
- Patient must be ≥ 13 years of age

Marplan®

PDL Status	NP
Quantity Limit	6 tabs/day
PA Criteria	See below
PA Form	General PA Form

Marplan® PA Criteria:

- Diagnosis of major depression; **AND**
- Trial and failure, contraindication, or intolerance to an agent from at least TWO of the following drug classes:
  - SSRIs
  - SNRIs
  - New generation antidepressants; **AND**
  - Trial and failure, contraindication, or intolerance to preferred phenelzine

Nardil®

PDL Status	NP
Quantity Limit	6 tabs/day
PA Criteria	See below
PA Form	General PA Form

Nardil® PA Criteria:

See Marplan® prior authorization criteria.

Parnate®

PDL Status	NP
Quantity Limit	6 tabs/day
PA Criteria	See below
PA Form	General PA Form

Parnate® PA Criteria:

See Marplan® prior authorization criteria.

Tranylcypromine

PDL Status	NP
Quantity Limit	6 tabs/day
PA Criteria	See below
PA Form	General PA Form

Tranylcypromine PA Criteria:

See Marplan® prior authorization criteria.

Antidepressants: New Generation

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antidepressants prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** **I/DD Prior Authorization Form**

Bupropion IR/SR (Generic for Wellbutrin® or Wellbutrin SR®) tablets

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Bupropion XL (Generic for Wellbutrin XL®) tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Exxua® tablets

PDL Status	P
Quantity Limit	Tablets: 1 /day; Titration pack: 1/year
PA Criteria	See below
PA Form	General PA Form

Exxua® Initial PA Criteria:

- Diagnosis of Major Depression Disorder; **AND**
- Patient is the labeled age minimum; **AND**
- Prescriber attests patient does not have abnormal electrolyte lab levels or abnormal ECG prior to initiating therapy and will continue to ~~watch~~ *monitor* during dose titration; **AND**
- Patient has been screened for personal or family history of bipolar disorder, mania, or hypomania; **AND**
- Patient does not have ANY of the following:
- Prolonged QTc interval greater than or equal to 450 msec at baseline
- Congenital long QT syndrome
- Severe hepatic impairment; **AND**
- Patient will not concomitantly use *strong* CYP3A4 inhibitors (e.g., clarithromycin, cobicistat, ketoconazole, itraconazole, lopinavir, posaconazole, ritonavir) while on Exxua therapy; **AND**
- Prescriber attests patient has not received MAOI therapy within 14 days and will not receive during therapy; **AND**
- Adequate trial and failure of TWO agents at an appropriate dose (3 weeks at the maximum tolerated dose within the recommended therapeutic range) within the following drug classes:
    - SSRI,
    - SNRI, or
    - Another preferred New Generation Antidepressant;

Exxua® Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g., improvement in Hamilton Depression Rating Scale (HAM-D) scores, improved overall mood, etc.); **AND**
- Patient has absence of unacceptable toxicity (e.g., increased QTc interval, abnormal electrolytes, abnormal liver enzyme labs, hypersensitivity reactions)

Mirtazapine ODT

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Mirtazapine ODT PA Criteria:

- Patient is unable to swallow solid dosage forms

Auvelity® tablets

PDL Status	NP
Quantity Limit	Tablets: 2/day; Titration pack: 1/year
PA Criteria	See below
PA Form	General PA Form

Auvelity® tablets PA Criteria:

- Patient is 18 years of age or older; **AND**
- One of the following:
  - Diagnosis of Major Depressive Disorder (MDD); **AND**
    - Trial and failure, or contraindication, intolerance to 2 preferred antidepressants; **OR**
  - Diagnosis of dementia due to Alzheimer's disease; **AND**
    - Patient is experiencing agitation; **AND**
- Patient does not have ANY of the following:
  - Seizure disorder
  - Current or prior diagnosis of bulimia or anorexia nervosa
  - Undergoing abrupt discontinuation of alcohol, benzodiazepines, barbiturates, or antiepileptic drugs; **AND**
- Prescriber attests patient has not received MAOI therapy within 14 days and will not receive during therapy

Forfivo XL® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Forfivo XL® tablets PA Criteria:

- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Patient must currently be on a bupropion product titrated to a dose of 300 mg per day

Nefazodone tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Nefazodone tablets PA Criteria:

- Diagnosis of major depression; **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Patient does not have hepatic impairment

Raldesy® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Raldesy® solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Remeron® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Remeron® tablets PA Criteria:
Standard NP Criteria

Remeron SolTab®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Remeron SolTab® PA Criteria:

- Patient is unable to swallow solid dosage forms

Trazodone 25 and 75 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Trazodone 25 and 75 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Trazodone 300 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Trazodone 300 mg tablets PA Criteria:

- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Clinically valid reason why the preferred lower strength tablets cannot be used (i.e., trazodone 50 mg, 100 mg, 150 mg)

Wellbutrin® IR and SR tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Wellbutrin® IR and SR tablets PA Criteria:
Standard NP Criteria

Zurzuvae® capsules

PDL Status	NP
Quantity Limit	1 treatment course/year
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	3 months
Notes	I/DD members not required to meet additional I/DD criteria for this agent.

Zurzuvae® capsules PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of postpartum depression (PPD); **AND**
- Patient's symptoms began in the third trimester or within 4 weeks of delivery; **AND**
- Prescriber attests that the PPD requires rapid improvement and resolution of symptoms; **AND**
- Prescribed by, or in consultation with, a psychiatrist, psychologist, or an obstetrician-gynecologist; **AND**
- Prescriber attests to ALL of the following:
  - Patient has been advised not to drive or operate machinery until at least 12 hours after administration due central nervous system (CNS) depressant effects such as somnolence and confusion
  - Females of reproductive potential have been advised to use effective contraception during treatment and for 1 week after the final dose due to potential risk to fetus and to notify healthcare provider if they become pregnant during treatment
  - Lactating women have been counseled on risk versus benefits of breastfeeding while on treatment

Antidepressants: SNRIs

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antidepressants prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at: **I/DD Prior Authorization Form**

Desvenlafaxine ER tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Duloxetine 20, 30, and 60 mg capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Effexor XR® capsules

PDL Status	P
Quantity Limit	37.5 mg caps: 1/day 75 mg caps: 3/day 150 mg caps: 2/day For 225 and 375 mg doses: use 150 and 75 mg caps
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Venlafaxine IR tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Venlafaxine ER capsules

PDL Status	P
Quantity Limit	37.5 mg: 1/day; 75 mg: 3/day; 150 mg: 2/day; For 225 & 375 mg doses: use 150 mg & 75 mg caps
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Cymbalta® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Cymbalta® capsules PA Criteria:
Standard NP Criteria.

Duloxetine 40 mg capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Duloxetine 40 mg capsules PA Criteria:

- Clinically valid reason as to why the preferred duloxetine strengths (20mg, 30mg, 60mg) cannot be used

Drizalma Sprinkle™ capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Drizalma Sprinkle™ capsules PA Criteria:

- Patient is unable to swallow solid dosage forms

Fetzima® capsules, titration pack

PDL Status	NP
Quantity Limit	Caps: 1/day Titration pack: 1/day (56 caps/lifetime)
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Fetzima® capsules PA Criteria:

Standard NP Criteria.

Pristiq® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Pristiq® tablets PA Criteria:

Standard NP Criteria.

Venlafaxine besylate ER tablets, venlafaxine HCl ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antidepressant SNRI PA Form

Venlafaxine besylate ER tablets, venlafaxine HCl ER tablets PA Criteria:

Standard NP Criteria.

Antidepressants: SSRIs

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antidepressants prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** **I/DD Prior Authorization Form**

Citalopram tablets

PDL Status	P
Quantity Limit	10 & 20 mg tabs: 1.5/day; 40 mg tabs: 1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Citalopram solution, escitalopram solution

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Escitalopram tablets

PDL Status	P
Quantity Limit	1.5/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Fluoxetine capsules

PDL Status	P
Quantity Limit	3/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Fluoxetine solution

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Fluvoxamine tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Paroxetine IR tablets

PDL Status	P
Quantity Limit	10 & 20 mg tabs: 1/day; 30 & 40 mg tabs: 2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Sertraline concentrate

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Sertraline tablets

PDL Status	P
Quantity Limit	25 & 50 mg tabs: 1.5/day; 100 mg tabs: 2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Vilazodone tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Celexa® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Celexa® tablets PA Criteria:
Standard NP Criteria.

Citalopram capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Citalopram capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Escitalopram capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Escitalopram capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Fluoxetine DR capsules

PDL Status	NP
Quantity Limit	4/28 days
PA Criteria	See below
PA Form	General PA Form

Fluoxetine DR capsules PA Criteria:

- Stabilized at a dose of 20 mg/day of fluoxetine for > 1 month; **AND**
- Documented reason why the patient is unable to continue fluoxetine 20 mg daily

Fluoxetine tablets

PDL Status	NP
Quantity Limit	20 mg tablets: 3/day; 60 mg tablets: 1/day
PA Criteria	See below
PA Form	General PA Form

Fluoxetine tablets PA Criteria:

Standard NP Criteria.

Fluvoxamine ER capsules

PDL Status	NP
Quantity Limit	100 mg tablets: 3/day; 150 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Fluvoxamine ER capsules PA Criteria:

Standard NP Criteria.

Lexapro® tablets

PDL Status	NP
Quantity Limit	1.5/day
PA Criteria	See below
PA Form	General PA Form

Lexapro® PA Criteria:
Standard NP Criteria.

Paroxetine 7.5 mg capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Paroxetine 7.5 mg capsules PA Criteria:
Standard NP Criteria.

Paroxetine CR tablets

PDL Status	NP
Quantity Limit	12.5 & 25 mg tablets: 1/day; 37.5 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Paroxetine CR tablets PA Criteria:
Standard NP Criteria.

Paxil® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Paxil® suspension PA Criteria:
Standard NP Criteria.

Paxil® tablets

PDL Status	NP
Quantity Limit	10 & 20 mg tablets: 1/day; 30 & 40 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Paxil® tablets PA Criteria:
Standard NP Criteria.

Paxil CR® tablets

PDL Status	NP
Quantity Limit	12.5 & 25 mg tablets: 1/day; 37.5 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Paxil CR® tablets PA Criteria:
Standard NP Criteria.

Prozac® capsules

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Prozac® capsules PA Criteria:
Standard NP Criteria.

Sertraline capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Sertraline capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Trintellix® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Trintellix® tablets PA Criteria:

- Diagnosis of Major Depression Disorder; **AND**
- Adequate trial and failure of TWO agents at an appropriate dose (3 weeks at the maximum tolerated dose within the recommended therapeutic range) within the following drug classes:
  - SSRI
  - SNRI
  - New Generation Antidepressants

Viibryd® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Viibryd® tablets PA Criteria:
Standard NP Criteria.

Zoloft® concentrate

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Zoloft® concentrate PA Criteria:
Standard NP Criteria.

Zoloft® tablets

PDL Status	NP
Quantity Limit	25 & 50 mg tabs: 1.5/day; 100 mg tabs: 2/day
PA Criteria	See below
PA Form	General PA Form

Zoloft® tablets PA Criteria:
Standard NP Criteria.

Antidepressants: Tricyclics

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antidepressants prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** **I/DD Prior Authorization Form**

Amitriptyline tablets

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Doxepin capsules

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Imipramine tablets

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Nortriptyline capsules

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Amoxapine

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Amoxapine PA Criteria:
Standard NP Criteria.

Anafranil®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Anafranil® PA Criteria:

- Diagnosis of obsessive-compulsive disorder; **AND**
- Trial and failure of at least 2 unique SSRIs

Clomipramine

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Clomipramine PA Criteria:

- Diagnosis of obsessive-compulsive disorder; **AND**
- Trial and failure of at least 2 unique SSRIs

Desipramine

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Desipramine PA Criteria:
Standard NP Criteria.

Imipramine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Imipramine capsules PA Criteria:
Standard NP Criteria.

Norpramin®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Norpramin® PA Criteria:
Standard NP Criteria.

Nortriptyline solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Nortriptyline solution PA Criteria:

- Patient is unable to swallow nortriptyline capsules

Pamelor®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pamelor® PA Criteria:
Standard NP Criteria.

Protriptyline

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Protriptyline PA Criteria:
Standard NP Criteria.

Antihyperkinesis: Stimulants

For agents in this class requiring PA, please submit the Antihyperkinesis: Stimulants PA Form on faxed PA submissions.

Adderall XR® capsules

PDL Status	P
Quantity Limit	5, 10, &15 mg caps: 1/day; 25 & 30 mg caps: 2/day; 20 mg caps: 3/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old. Patients ≤ 20 years of age will be subject to the initial criteria if they exceed 80 mg/day of total amphetamine.

Adderall XR® capsules PA Criteria:

- Agent must not be prescribed by a pain clinic; **AND**
- Patient will not concurrently take a benzodiazepine, barbiturate, sedative hypnotic, opioid (excluding buprenorphine for OUD), MAOI (monoamine oxidase inhibitor) agent, or meprobamate/carisoprodol; **AND**
- One of the following:
  - Diagnosis of Attention Deficit Disorder and/or Hyperactivity Disorder (ADD/ADHD); **AND**
    - Documentation that the symptoms affect the patient's ability to function in daily life tasks or creates significant difficulties in at least 2 major settings (school, work, social settings, and/or home)
  - Diagnosis of Narcolepsy supported with documentation of polysomnography or multiple sleep latency test (MSLT)
  - Diagnosis of neurocognitive disorder (also known as organic brain disorder)
  - Diagnosis of treatment resistant Major Depressive Disorder; **AND**
    - Adequate trial and failure of 3 agents at an appropriate dose (defined as: 3 weeks at the maximum tolerated dose within the recommended therapeutic range) from at least 3 distinct drug classes:
      - SSRI
      - SNRI
      - New Generation Antidepressants
      - TCAs

Amphetamine salt ER combination capsules

PDL Status	P
Quantity Limit	5, 10, &15 mg caps: 1/day; 25 & 30 mg caps: 2/day; 20 mg caps: 3/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old. Patients ≤ 20 years of age will be subject to the initial criteria if they exceed 80 mg/day of total amphetamine.

Amphetamine salt ER combination capsules PA Criteria:

See Adderall XR® prior authorization criteria.

Amphetamine salt IR combination tablets

PDL Status	P
Quantity Limit	5, 7.5, 10, & 12.5 mg tabs: 4/day; 15 & 30 mg tabs: 2/day; 20 mg tabs: 3/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old. Patients ≤ 20 years of age will be subject to the initial criteria if they exceed 80 mg/day of total amphetamine.

Amphetamine salt IR combination tablets PA Criteria:
See Adderall XR® prior authorization criteria

Amphetamine sulfate tablets

PDL Status	P
Quantity Limit	5 mg tab: 3/day; 10 mg tab: 6/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old. Patients ≤ 20 years of age will be subject to the initial criteria if they exceed 80 mg/day of total amphetamine.

Amphetamine sulfate tablets PA Criteria:
See Adderall XR® prior authorization criteria

Aptensio XR® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Aptensio XR® capsules PA Criteria:
See Adderall XR® prior authorization criteria

Dexmethylphenidate tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Dexmethylphenidate tablets PA Criteria:
See Adderall XR® prior authorization criteria

Dexmethylphenidate XR capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Dexmethylphenidate XR capsules PA Criteria:
See Adderall XR® prior authorization criteria

Dextroamphetamine tablets

PDL Status	P
Quantity Limit	20 mg: 3/day; 30 mg: 2/day; All others: 4/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old. Patients ≤ 20 years of age will be subject to the initial criteria if they exceed 80 mg/day of total amphetamine.

Dextroamphetamine tablets PA Criteria:
See Adderall XR® prior authorization criteria

Focalin XR® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Focalin XR® capsules PA Criteria:
See Adderall XR® prior authorization criteria

Lisdexamfetamine chewables

PDL Status	P
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Lisdexamfetamine chewables PA Criteria:
See Adderall XR® prior authorization criteria

Methylphenidate tablets (generic for Ritalin®)

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Methylphenidate tablets (generic for Ritalin®) PA Criteria:
See Adderall XR® prior authorization criteria

Methylphenidate solution (generic for Methylin®)

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Methylphenidate solution PA Criteria:
See Adderall XR® prior authorization criteria

Methylphenidate ER tablets (generic for Metadate ER®)

PDL Status	P
Quantity Limit	10 mg tabs: 1/day; 20 mg tabs: 3/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Methylphenidate ER tablets (generic for Metadate ER®) PA Criteria:
See Adderall XR® prior authorization criteria

Methylphenidate ER OSM tablets (generic for Concerta®)

PDL Status	P
Quantity Limit	18, 27, & 54 mg: 1/day; 36 mg: 2/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate ER OSM tablets (generic for Concerta®) PA Criteria:
See Adderall XR® tablets prior authorization criteria

ProCentra® solution

PDL Status	P
Quantity Limit	20 mL/day Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

ProCentra® solution PA Criteria:
See Adderall XR® prior authorization criteria

Quillichew ER® chewables

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Quillichew ER® chewables PA Criteria:
See Adderall XR® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance of 2 preferred agents, unless otherwise indicated

Quillivant XR® suspension

PDL Status	P
Quantity Limit	12 mL/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Quillivant XR® suspension PA Criteria:

See Adderall XR® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance of 2 preferred agents, unless otherwise indicated

Vyvanse® capsules

PDL Status	P
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form
Notes	PA required for patients ≥ 21 years old.

Vyvanse® capsules PA Criteria:

See Adderall XR® prior authorization criteria

Adderall® tablets

PDL Status	NP
Quantity Limit	5, 7.5, 10, & 12.5 mg tabs: 4/day; 15 & 30 mg tabs: 2/day; 20 mg tabs: 3/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Adderall® tablets PA Criteria:

- Agent must not be prescribed by a pain clinic; **AND**
- Patient will not concurrently take a benzodiazepine, barbiturate, sedative hypnotic, opioid (excluding buprenorphine for OUD), MAOI (monoamine oxidase inhibitor) agent, or meprobamate/carisoprodol; **AND**
- One of the following:
  - Diagnosis of Attention Deficit Disorder and/or Hyperactivity Disorder (ADD/ADHD); **AND**
    - Documentation that the symptoms affect the patient's ability to function in daily life tasks or creates significant difficulties in at least 2 major settings (school, work, social settings, and/or home)
  - Diagnosis of Narcolepsy supported with documentation of polysomnography or multiple sleep latency test (MSLT)
  - Diagnosis of neurocognitive disorder (also known as organic brain disorder)
  - Diagnosis of treatment resistant Major Depressive Disorder; **AND**
    - Adequate trial and failure of 3 agents at an appropriate dose (defined as: 3 weeks at the maximum tolerated dose within the recommended therapeutic range) from at least 3 distinct drug classes:
      - SSRI
      - SNRI
      - New Generation Antidepressants
      - TCAs; **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents, unless otherwise indicated

Adzenys XR® ODT

PDL Status	NP
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Adzenys XR® ODT PA Criteria:
See Adderall® tablets prior authorization criteria.

Amphetamine/dextroamphetamine ER (3-bead) 24 HR caps (generic for Mydayis®

PDL Status	NP
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Amphetamine/dextroamphetamine ER (3-bead) 24 hr caps (generic for Mydayis® PA Criteria:
See Adderall® tablets prior authorization criteria.

Amphetamine ER ODT (generic for Adzenys XR® ODT)

PDL Status	NP
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Amphetamine ER ODT (generic for Adzenys XR® ODT) PA Criteria:
See Adderall® tablets prior authorization criteria.

Arynta® solution

PDL Status	NP
Quantity Limit	7 ml/Day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Arynta® solution PA Criteria:
See Adderall® tablets prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms

Azstarys® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Azstarys® capsules PA Criteria:
See Adderall® tablets prior authorization criteria.

Concerta® OSM tablets

PDL Status	NP
Quantity Limit	18, 27, & 54 mg: 1/day; 36 mg: 2/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Concerta® OSM tablets PA Criteria:
See Adderall ® prior authorization criteria

Cotempla XR® ODT

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Cotempla XR® ODT PA Criteria:

- Agent must not be prescribed by a pain clinic; **AND**
- Patient will not concurrently take a benzodiazepine, barbiturate, sedative hypnotic, opioid (excluding buprenorphine for OUD), MAOI (monoamine oxidase inhibitor) agent, or meprobamate/carisoprodol; **AND**
- Patient is 6 to 17 years of age; **AND**
- Diagnosis of Attention Deficit Disorder and/or Hyperactivity Disorder (ADD/ADHD); **AND**
    - Documentation that the symptoms affect the patient's ability to function in daily life tasks or creates significant difficulties in at least 2 major settings (school, work, social settings, and/or home); **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents, unless otherwise indicated

Dexedrine® capsules

PDL Status	NP
Quantity Limit	4/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Dexedrine® PA Criteria:
See Adderall® tablets prior authorization criteria.

Dextroamphetamine solution

PDL Status	NP
Quantity Limit	20 mL/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Dextroamphetamine solution PA Criteria:
See Adderall® tablets prior authorization criteria.

Dyanavel XR® suspension, tablets

PDL Status	NP
Quantity Limit	Suspension: 8 mL/day; Tablets: 1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Dyanavel XR® suspension, tablets PA Criteria:
See Adderall® tablets prior authorization criteria.

Evekeo® tablets

PDL Status	NP
Quantity Limit	5 mg tab: 3/day; 10 mg tab: 6/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Evekeo® tablets PA Criteria:
See Adderall XR® prior authorization criteria

Focalin® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Focalin® tablets PA Criteria:
See Adderall® tablets prior authorization criteria

Jornay PM® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Jornay PM® capsules PA Criteria:
See Adderall® tablets prior authorization criteria

Lisdexamfetamine capsules

PDL Status	NP
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Lisdexamfetamine capsules PA Criteria:
See Adderall® tablets prior authorization criteria; **AND**

- Clinically valid reason why brand Vyvanse capsules cannot be used

Methamphetamine tablets

PDL Status	NP
Quantity Limit	4/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methamphetamine tablets PA Criteria:
See Adderall® tablets prior authorization criteria.

Methylin® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylin® solution PA Criteria:
See Adderall® tablets prior authorization criteria

Methylphenidate chewables

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate chewables PA Criteria:
See Adderall® tablets prior authorization criteria

Methylphenidate patch

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate patch PA Criteria:
See Adderall® tablets prior authorization criteria

Methylphenidate ER 24hr capsules (generic for Aptensio XR®, Ritalin LA®)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate ER 24hr capsules (generic for Aptensio XR®, Ritalin LA®) PA Criteria:
See Adderall® tablets prior authorization criteria

Methylphenidate ER tablets (18, 27, 36, and 54 mg), ER diffusion tablets (27, 36, and 54 mg)

PDL Status	NP
Quantity Limit	18, 27, & 54 mg: 1/day; 36 mg: 2/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate ER tablets, ER diffusion tablets PA Criteria:
See Adderall® tablets prior authorization criteria; **AND**

- Clinically valid reason why methylphenidate OSM ER tablets (generic for Concerta®) cannot be used

Methylphenidate ER OSM tablets (45, 63, and 72 mg) (generic for Relexxii®)

PDL Status	NP
Quantity Limit	45, 63 and 72 mg tabs: 1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate ER OSM tablets (generic for Relexxii®) PA Criteria:

See Adderall® tablets prior authorization criteria

Methylphenidate XR ODT (generic for Cotempla XR® ODT)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Methylphenidate XR ODT (generic for Cotempla XR® ODT) PA Criteria:

- Agent must not be prescribed by a pain clinic; **AND**
- Patient will not concurrently take a benzodiazepine, barbiturate, sedative hypnotic, opioid (excluding buprenorphine for OUD), MAOI (monoamine oxidase inhibitor) agent, or meprobamate/carisoprodol; **AND**
- Patient is 6 to 17 years of age; **AND**
- Diagnosis of Attention Deficit Disorder and/or Hyperactivity Disorder (ADD/ADHD); **AND**
    - Documentation that the symptoms affect the patient's ability to function in daily life tasks or creates significant difficulties in at least 2 major settings (school, work, social settings, and/or home); **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents, unless otherwise indicated; **AND**
- If request is for methylphenidate XR ODT, clinically valid reason why brand Cotempla XR ODT cannot be used

Mydayis ER® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Mydayis ER® capsules PA Criteria:

See Adderall® tablets prior authorization criteria; **AND**

- Clinically valid reason why generic equivalent cannot be used

Relexxii ER® OSM tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Relexxii ER® OSM tablets PA Criteria:

See Adderall® tablets prior authorization criteria

Ritalin® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Ritalin® tablets PA Criteria:

See Adderall® tablets prior authorization criteria

Ritalin LA® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Ritalin LA® capsules PA Criteria:
See Adderall® tablets prior authorization criteria

Vyvanse® chewables

PDL Status	NP
Quantity Limit	1/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Vyvanse® chewables PA Criteria:
See Adderall® tablets prior authorization criteria

Xelstrym® patch

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Xelstrym® patch PA Criteria:
See Adderall® tablets prior authorization criteria

Zenzedi® tablets

PDL Status	NP
Quantity Limit	20 mg: 3/day; 30 mg: 2/day; All others: 4/day; Max total amphetamine dose (Age ≥ 21): 60 mg/day
PA Criteria	See below
PA Form	Antihyperkinesis: Stimulants PA Form

Zenzedi® tablets PA Criteria:
See Adderall® tablets prior authorization criteria

Antihyperkinesis: Non-Stimulants

Atomoxetine capsules

PDL Status	P
Quantity Limit	60, 80, & 100 mg caps: 1/day; All other strengths: 2/day
PA Criteria	None

Clonidine ER 12-hour tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Guanfacine ER tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Qelbree® capsules

PDL Status	P
Quantity Limit	100 & 150 mg caps: 2/day; 200 mg caps: 3/day
PA Criteria	See below
PA Form	General PA Form

Qelbree® capsules PA Criteria:

- Diagnosis of Attention Deficit Hyperactivity Disorder (ADHD); **AND**
- Patient is 6 years of age or older; **AND**
- Patient's blood pressure and heart rate will be assessed prior to therapy and monitored throughout therapy; **AND**
- Patient will not concomitantly use monoamine oxidase inhibitors (MAOIs); **AND**
- Patient will not concomitantly use CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic range; **AND**
- Patient is not pregnant; **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred antihyperkinesis stimulant and/or non-stimulant agents

Intuniv® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Intuniv® tablets PA Criteria:

- Clinically valid reason why preferred guanfacine ER cannot be used

Onyda XR® suspension

PDL Status	NP
Quantity Limit	4 mL/day
PA Criteria	See below
PA Form	General PA Form

Onyda XR® suspension PA Criteria:

- Diagnosis of attention deficit hyperactivity disorder (ADHD); **AND**
- Patient is 6 years of age or older; **AND**
- One of the following:
  - Trial & failure, contraindication, or intolerance of 2 preferred non-stimulant antihyperkinesis agents; **OR**
  - Patient is unable to swallow solid dosage forms

Strattera® capsules

PDL Status	NP
Quantity Limit	60, 80, & 100 mg caps: 1/day; All other strengths: 2/day
PA Criteria	See below
PA Form	General PA Form

Strattera® capsules PA Criteria:

Standard NP Criteria

Anti-Migraine: 5-HT1 Receptor Agonists and Combinations

Eletriptan

PDL Status	P
Quantity Limit	6/30 days
PA Criteria	None

Frovatriptan

PDL Status	P
Quantity Limit	6/30 days
PA Criteria	None

Rizatriptan tablets, ODT

PDL Status	P
Quantity Limit	12/30 days
PA Criteria	None

Sumatriptan tablets

PDL Status	P
Quantity Limit	9/30 days
PA Criteria	None

Sumatriptan vials

PDL Status	P
Quantity Limit	8 vials/30 days
PA Criteria	None

Zolmitriptan nasal spray

PDL Status	P
Quantity Limit	6/30 days
PA Criteria	None

Almotriptan

PDL Status	NP
Quantity Limit	12/30 days
PA Criteria	See below
PA Form	General PA Form

Almotriptan PA Criteria:
Standard NP Criteria

Frova®

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Frova® PA Criteria:
Standard NP Criteria

Imitrex® injectable

PDL Status	NP
Quantity Limit	8/30 days
PA Criteria	See below
PA Form	General PA Form

Imitrex® injectable PA Criteria:
Standard NP Criteria

Imitrex® Cartridge

PDL Status	NP
Quantity Limit	8/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Imitrex® Cartridge PA Criteria:

- Clinically valid reason why the injectable vials cannot be used.

Imitrex® Nasal

PDL Status	NP
Quantity Limit	6/30 days
PA Criteria	See below
PA Form	General PA Form

Imitrex® Nasal PA Criteria:
Standard NP Criteria

Imitrex® tablets

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Imitrex® tablets PA Criteria:
Standard NP Criteria

Maxalt®-MLT, Maxalt® tablets

PDL Status	NP
Quantity Limit	12/30 days
PA Criteria	See below
PA Form	General PA Form

Maxalt®- MLT, Maxalt® tablets PA Criteria:
Standard NP Criteria

Naratriptan tablets

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Naratriptan tablets PA Criteria:
Standard NP Criteria

Onzetra Xsail®

PDL Status	NP
Quantity Limit	16/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Onzetra Xsail® PA Criteria:

- Patient has an allergy to an inactive ingredient found in the preferred sumatriptan containing agents; **AND**
- Patient has a contraindication, allergic reaction, or drug-drug interaction to preferred rizatriptan containing agents; **AND**
- Clinically valid reason why the patient requires a nasal powder

Relpax® tablets

PDL Status	NP
Quantity Limit	6/30 days
PA Criteria	See below
PA Form	General PA Form

Relpax® tablets PA Criteria:

Standard NP Criteria

Reyvow®

PDL Status	NP
Quantity Limit	4/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 3 months; Renewal: 1 year

Reyvow® Initial PA Criteria:

- Agent is being used for acute treatment of migraine with or without aura; **AND**
- Patient is 18 years of age or older; **AND**
- Trial and failure, contraindication, or intolerance to TWO triptans (e.g., eletriptan, rizatriptan, sumatriptan)

Reyvow® Renewal PA Criteria:

- Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)

Sumatriptan autoinjector, cartridge

PDL Status	NP
Quantity Limit	8/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Sumatriptan autoinjector, cartridge PA Criteria:

- Clinically valid reason why the injectable vials cannot be used

Sumatriptan nasal spray

PDL Status	NP
Quantity Limit	6/30 days
PA Criteria	See below
PA Form	General PA Form

Sumatriptan nasal spray PA Criteria:

Standard NP Criteria

Sumatriptan/naproxen tablets

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Sumatriptan/naproxen tablets PA Criteria:
Standard NP Criteria

Symbravo®

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Symbravo® PA Criteria:

- Clinically valid reason as to why the patient cannot use a preferred NSAID and 5-HT receptor agonist as separate agents

Tosymra®

PDL Status	NP
Quantity Limit	12/30 days
PA Criteria	See below
PA Form	General PA Form

Tosymra® PA Criteria:
Standard NP Criteria

Treximet® tablets

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Treximet® tablets PA Criteria:
Standard NP Criteria

Zolmitriptan tablets

PDL Status	NP
Quantity Limit	9/30 days
PA Criteria	See below
PA Form	General PA Form

Zolmitriptan tablets PA Criteria:
Standard NP Criteria

Zembrace Symtouch®

PDL Status	NP
Quantity Limit	2 mL/30 days
PA Criteria	See below
PA Form	General PA Form
Notes	Patient convenience is NOT an approvable reason.

Zembrace Symtouch® PA Criteria:

- Patient has an allergy to an inactive ingredient found in the preferred sumatriptan containing agents; **AND**
- Patient has a contraindication, allergic reaction, or drug-drug interaction to preferred rizatriptan containing agents; **AND**
- Clinically valid reason why the patient requires an autoinjector device

Zomig® nasal spray, tablets

PDL Status	NP
Quantity Limit	6/30 days
PA Criteria	See below
PA Form	General PA Form

Zomig® nasal spray, tablets PA Criteria:

Standard NP Criteria

Anti-Migraine: Barbiturate Agents, Non-Narcotic

****Quantity Limit Override Criteria for Butalbital-Containing products:**

Requests for butalbital-containing products for quantities greater than 20 per 30 days will be approved for patients meeting the following criteria:

- Trial and failure of at least 2 prophylactic headache treatments: a tricyclic antidepressant (unless contraindicated) PLUS at least one of the following:
  - Divalproex sodium
  - Sodium valproate
  - Topiramate
  - Frovatriptan
  - Beta-blocker

Butalbital/APAP

PDL Status	P
Quantity Limit	20/30 days; Max APAP: 4 g/day
PA Criteria	None

Butalbital/APAP/caffeine

PDL Status	P
Quantity Limit	20/30 days; Max APAP: 4 g/day
PA Criteria	None

Allzital®

PDL Status	NP
Quantity Limit	20/30 days; Max APAP: 4 g/day
PA Criteria	See below
PA Form	General PA Form

Allzital® PA Criteria:
Standard NP Criteria

Butalbital/ASA/caffeine

PDL Status	NP
Quantity Limit	20/30 days;
PA Criteria	See below
PA Form	General PA Form

Butalbital/ASA/caffeine PA Criteria:

- Allergy or intolerance to APAP (acetaminophen)

Fioricet®

PDL Status	NP
Quantity Limit	20/30 days; Max APAP: 4 g/day
PA Criteria	See below
PA Form	General PA Form

Fioricet® PA Criteria:
Standard NP Criteria

Anti-Migraine: CGRP Antagonists

Aimovig®

PDL Status	P
Quantity Limit	1 syringe/30 days
PA Criteria	See below
PA Form	General PA Form

Aimovig® Initial PA Criteria:

- Patient has a diagnosis of migraine with or without aura; **AND**
- Patient has ≥ 4 migraine days per month; **AND**
- Patient is utilizing prophylactic interventions (e.g., behavioral therapy, physical therapy, lifestyle modifications); **AND**
- Trial (duration > 8 weeks) and failure of ONE of the following oral medication classes, unless contraindicated:
  - Antidepressants (i.e., amitriptyline, venlafaxine)
  - Beta blockers (i.e., propranolol, metoprolol, timolol, atenolol)
  - Antiepileptics (i.e., valproate, topiramate)

Aimovig® Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g., decrease in the number, frequency, and/or intensity of headaches, improved function, decreased reliance on acute treatments for migraine headaches); **AND**
- Patient has absence of unacceptable toxicity (e.g., intolerable injection site pain or constipation)

Emgality® pen, syringe

PDL Status	P
Quantity Limit	1 syringe/30 days; 120 mg for migraine, 300 mg for cluster headaches
PA Criteria	See below
PA Form	General PA Form

Emgality® pen, syringe Initial PA Criteria:

- Diagnosis of episodic cluster headache; **OR**
- Diagnosis of migraine with or without aura; **AND**
  - Patient has ≥ 4 migraine days per month; **AND**
  - Patient is utilizing prophylactic interventions (e.g., behavioral therapy, physical therapy, life-style modifications); **AND**
  - Trial (duration ≥ 8 weeks) and failure of ONE of the following oral medication classes, unless contraindicated:
    - Antidepressants (i.e., amitriptyline, venlafaxine)
    - Beta blockers (i.e., propranolol, metoprolol, timolol, atenolol)
    - Antiepileptics (i.e., valproate, topiramate); **OR**

Emgality® pen, syringe Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g., decrease in the number, frequency, and/or intensity of headaches, improved function, decreased reliance on acute treatments for migraine headaches); **AND**
- Patient has absence of unacceptable toxicity (e.g., intolerable injection site pain or constipation)

Nurtec ODT®

PDL Status	P
Quantity Limit	Acute treatment: 1 dose pack (8 tabs)/30 days; Prophylaxis: 2 dose packs (16 tabs)/30 days
PA Criteria	See below
PA Form	General PA Form

Nurtec ODT® Initial PA Criteria:

- Diagnosis of migraine with or without aura; **AND**
- One of the following:
  - Acute treatment of migraine, **AND**
    - Medication will not be used in combination with another acute CGRP inhibitor; **AND**
    - Trial and failure or intolerance to TWO triptans (e.g., eletriptan, rizatriptan, sumatriptan) **OR** contraindication to all triptans
  - Preventative treatment of migraine; **AND**
    - Patient has ≥ 4 migraine days per month; **AND**
    - Patient is utilizing prophylactic interventions (e.g., behavioral therapy, physical therapy, life-style modifications); **AND**
    - Trial (duration ≥ 8 weeks) and failure of ONE of the following oral medication classes, unless contraindicated:
      - Antidepressants (i.e., amitriptyline, venlafaxine)
      - Beta blockers (i.e., propranolol, metoprolol, timolol, atenolol)
      - Antiepileptics (i.e., valproate, topiramate); **AND**

Nurtec ODT® Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g., decrease in the number, frequency, and/or intensity of headaches, improved function, decreased reliance on acute treatments for migraine headaches)

Quilipta®

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Quilipta® Initial PA Criteria:

- Patient has a diagnosis of migraine with or without aura; **AND**
- Patient has ≥ 4 migraine days per month; **AND**
- Patient is utilizing prophylactic interventions (e.g., behavioral therapy, physical therapy, lifestyle modifications); **AND**
- Trial (duration ≥ 8 weeks) and failure of ONE of the following oral medication classes, unless contraindicated:
  - Antidepressants (i.e., amitriptyline, venlafaxine)
  - Beta blockers (i.e., propranolol, metoprolol, timolol, atenolol)
  - Antiepileptics (i.e., valproate, topiramate); **AND**

Quilipta® Renewal PA Criteria:

- Patient has experienced positive response to therapy (e.g., decrease in the number, frequency, and/or intensity of headaches, improved function, decreased reliance on acute treatments for migraine headaches)

Ubrelvy®

PDL Status	P
Quantity Limit	1 box (10 tablets)/30 days
PA Criteria	See below
PA Form	General PA Form

Ubrelvy® Initial PA Criteria:

- Diagnosis of migraine with or without aura and will be used for the acute treatment of migraine, **AND**
- Trial and failure or intolerance to TWO triptans (e.g., eletriptan, rizatriptan, sumatriptan) **OR** contraindication to all triptan; **AND**
- Medication will not be used in combination with another acute CGRP inhibitor

Ubrelvy® Renewal PA Criteria:

- Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)

Ajovy® autoinjector, prefilled syringe

PDL Status	NP
Quantity Limit	3 injections/90 days
PA Criteria	See below
PA Form	General PA Form

Ajovy® autoinjector, prefilled syringe PA Criteria:

- One of the following:
  - Patient is 18 years of age or older; **AND**
    - Patient meets Aimovig® prior authorization criteria; **AND**
    - Trial and failure of Aimovig® and Emgality®; **OR**
  - Patient is 6 to 17 years old and ≥ 45 kg; **AND**
    - ALL of the following:
      - Patient has diagnosis of episodic migraines defined as migraines for at least 6 months and < 15 headache days per month; **AND**
      - Patient has ≥ 4 migraine days per month; **AND**
      - Patient is utilizing prophylactic interventions (e.g., behavioral therapy, physical therapy, lifestyle modifications, supplements); **AND**
      - Will not be used in combination with another CGRP for preventative treatment; **AND**
      - Trial (duration > 8 weeks) and failure of ONE of the following oral medication classes, unless contraindicated:
        - Beta blockers (i.e., propranolol)
        - Anticonvulsants (i.e., topiramate); **AND**
    - Prescribed by, or in consultation with, a neurologist

Zavzpret®

PDL Status	NP
Quantity Limit	6 devices (60 mg)/30 days
PA Criteria	See below
PA Form	General PA Form

Zavzpret® Initial PA Criteria:

- Diagnosis of migraine with or without aura and will be used for the acute treatment of migraine, **AND**
- Trial and failure or intolerance to Nurtec ODT® and Ubrelvy®; **AND**
- Medication will not be used in combination with another acute CGRP inhibitor

Zavzpret® Renewal PA Criteria:

- Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)

Antimigraine: Ergotamine Derivatives

Dihydroergotamine nasal spray

PDL Status	P
Quantity Limit	8 mL/30 days
PA Criteria	None

Dihydroergotamine injection

PDL Status	P
Quantity Limit	8 mL/30 days
PA Criteria	None

Brekiya®

PDL Status	NP
Quantity Limit	8 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Brekiya® PA Criteria:

- Trial and failure of dihydroergotamine nasal spray and injection; **OR**
- Clinically valid reason why other dihydroergotamine formulations are not appropriate; **AND**
- For initial requests only: Prescriber attests that patient has received first injection in the setting of an equipped healthcare facility

Migergot®

PDL Status	NP
Quantity Limit	15/30 days
PA Criteria	See below
PA Form	General PA Form

Migergot® PA Criteria:

Standard NP Criteria

Antipsychotics: Atypical

Note: A list of ICD-10 codes to allow PA bypass for preferred oral atypical antipsychotics that require PA can be found at **Appropriate Diagnosis for PA Bypass List**.

For agents in this class requiring PA, please submit the **Atypical Antipsychotics PA Form** on faxed PA submissions.

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antipsychotics prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at: **I/DD Prior Authorization Form**

Abilify Asimtufii® injection

PDL Status	P
Quantity Limit	1 injection/60 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Abilify Asimtufii® injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Patient has documented tolerance to the oral active ingredient

Abilify Maintena® injection

PDL Status	P
Quantity Limit	1 injection/30 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Abilify Maintena® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Aripiprazole ODT, solution, tablets

PDL Status	P
Quantity Limit	ODT: 1/day; Solution: 10 mL/day; 2, 5, 10, & 15 mg tabs: 2/day; 20 & 30 mg tabs: 1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Aristada® injection

PDL Status	P
Quantity Limit	1064 mg injection: 1/60 days; All other strengths: 1/30 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Aristada® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Aristada Initio® injection

PDL Status	P
Quantity Limit	2.4 mL/60 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Aristada Initio® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Asenapine SL tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Asenapine SL tablets PA Criteria:

- Diagnosis of ONE of the following:
  - Agitation in dementia
  - Bipolar and manic disorders
  - Bipolar depression, bipolar maintenance, bipolar mania-acute, bipolar mixed states
  - Brief psychotic disorder
  - Delusional disorder
  - Depression with psychotic symptoms
  - Drug-induced psychotic disorder with hallucinations
  - Impulse control disorders, including Oppositional Defiant Disorder and Intermittent Explosive Disorder
  - Organic psychotic condition
  - Psychosis secondary to a medical condition, psychotic depression, psychotic disorders
  - Schizoaffective disorder, schizoid/schizotypal personality disorder, schizophrenia, schizophrenic disorders
  - Substance-induced psychotic disorder, substance-induced withdrawal psychotic disorder
  - Severe refractory OCD or PTSD
  - Tourette's/Severe tic disorder; **OR**
- Diagnosis of major depressive disorder (MDD)**; AND**
  - Atypical agents will be approved only as adjunctive treatment for MDD; **AND**
  - Adequate trial (4 - 6 weeks) of ONE agent from any of the following classes (unless contraindication or intolerance):
    - SSRIs
    - SNRIs
    - TCAs
    - New generation antidepressants (including bupropion, mirtazapine, etc.); **OR**
- For patients without one of the above diagnoses:
  - May be approved if the physician can provide documented clinical evidence supporting the use of the requested medication for the requested indication

Clozapine tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Erzofi® injection

PDL Status	P
Quantity Limit	1 injection/28 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Erzofi® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Invega Hafyera® injection

PDL Status	P
Quantity Limit	1 injection/168 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Invega Hafyera® injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- TennCare prescription claims history must indicate patient has been on Invega Sustenna® for 4 months **OR** Invega Trinza for at least one three-month cycle

Invega Sustenna® injection

PDL Status	P
Quantity Limit	1 injection/28 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Invega Sustenna® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Invega Trinza® injection

PDL Status	P
Quantity Limit	1 injection/76 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Invega Trinza® injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- TennCare prescription claims history must indicate patient has been on Invega Sustenna® for 4 months

Lurasidone tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Lurasidone tablets PA Criteria:

See asenapine prior authorization criteria

Olanzapine tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Olanzapine IM injection

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Olanzapine IM injection PA Criteria:

See asenapine prior authorization criteria

Olanzapine ODT

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Olanzapine ODT PA Criteria:

See asenapine prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms or absorb medications through the GI tract; **OR**
- Non-response due to non-compliance

Paliperidone ER tablets

PDL Status	P
Quantity Limit	6 mg tablets: 2/day; All other strengths: 1/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Perseris® injection

PDL Status	P
Quantity Limit	1 injection/month
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Perseris® injection PA Criteria:

See Abilify Asimtufii® prior authorization criteria

Quetiapine tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Quetiapine ER tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Quetiapine ER tablets PA Criteria:

See asenapine prior authorization criteria

Rexulti® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Rexulti® tablets PA Criteria:

See asenapine prior authorization criteria

Risperidone ODT

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Risperidone ODT PA Criteria:
See olanzapine ODT prior authorization criteria.

Risperidone solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Risperidone solution PA Criteria:
See asenapine prior authorization criteria

Risperidone tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Uzedy® injection

PDL Status	P
Quantity Limit	50, 75, 100, & 125 mg injection: 1/30 days; 150, 200, & 250 mg injection: 1/60 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Uzedy® injection PA Criteria:
See Abilify Asimtufii® prior authorization criteria

Vraylar® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Vraylar® capsules PA Criteria:
See asenapine prior authorization criteria

Ziprasidone capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Ziprasidone injection

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Ziprasidone injection PA Criteria:
See asenapine prior authorization criteria

Abilify® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Abilify® tablets PA Criteria:

- Approval of non-preferred atypical antipsychotics requires trial and failure of ONE preferred agent; **AND**
- Diagnosis of ONE of the following:
  - Agitation in dementia
  - Bipolar and manic disorders
  - Bipolar depression, bipolar maintenance, bipolar mania-acute, bipolar mixed states
  - Brief psychotic disorder
  - Delusional disorder
  - Depression with psychotic symptoms
  - Drug-induced psychotic disorder with hallucinations
  - Impulse control disorders, including Oppositional Defiant Disorder and Intermittent Explosive Disorder
  - Organic psychotic condition
  - Psychosis secondary to a medical condition, psychotic depression, psychotic disorders
  - Schizoaffective disorder, schizoid/schizotypal personality disorder, schizophrenia, schizophrenic disorders
  - Substance-induced psychotic disorder, substance-induced withdrawal psychotic disorder
  - Severe refractory OCD or PTSD
  - Tourette's/Severe tic disorder; **OR**
- Diagnosis of major depressive disorder (MDD)**; AND**
  - Atypical agents will be approved only as adjunctive treatment for MDD; **AND**
  - Adequate trial (4 - 6 weeks) of ONE agent from any of the following classes (unless contraindication or intolerance):
    - SSRIs
    - SNRIs
    - TCAs
    - New generation antidepressants (including bupropion, mirtazapine, etc.); **OR**
- For patients without one of the above diagnoses:
  - May be approved if the physician can provide documented clinical evidence supporting the use of the requested medication for the requested indication

Abilify MyCite® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Abilify MyCite® tablets PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Clinically valid reason why none of the other dosage forms of aripiprazole cannot be used.

Caplyta® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Caplyta® capsules PA Criteria:

See Abilify® tablets prior authorization criteria

Olanzapine ODT

PDL Status	NP
Quantity Limit	12.5 & 25 mg ODT: 2/day; 100 mg ODT: 9/day; 150 mg ODT: 6/day; 200 mg ODT: 4/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Olanzapine ODT PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms or absorb medications through the GI tract; **OR**
- Non-response due to noncompliance

Clozaril® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Clozaril® tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Cobenfy® capsules, starter pack

PDL Status	NP
Quantity Limit	Cap: 2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Cobenfy® capsules, starter pack PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of schizophrenia; **AND**
- Approval of non-preferred atypical antipsychotics requires trial and failure of ONE preferred agent; **AND**
- Prescriber attests patient does NOT have any of the following:
  - Urinary retention
  - Moderate (Child-Pugh Class B) or severe (Child-Pugh Class C) hepatic impairment
  - Gastric retention
  - Untreated narrow-angle glaucoma

Fanapt® tablets, tablet packs

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Fanapt® tablets, tablet packs PA Criteria:

See Abilify® tablets prior authorization criteria

Geodon® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Geodon® capsules PA Criteria:
See Abilify® tablets prior authorization criteria

Geodon® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Geodon® injection PA Criteria:
See Abilify® tablets prior authorization criteria

Invega® tablets

PDL Status	NP
Quantity Limit	6 mg tablets: 2/day; All other strengths: 1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Invega® tablets PA Criteria:
See Abilify® tablets prior authorization criteria

Latuda® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Latuda® tablets PA Criteria:
See Abilify® tablets prior authorization criteria

Lybalvi® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Lybalvi® tablets PA Criteria:

- Patient is ≥18 years of age; **AND**
- One of the following:
  - Diagnosis of schizophrenia
  - Diagnosis of Bipolar I disorder and will be used for the acute treatment of manic or mixed episodes
  - Diagnosis of Bipolar I disorder and will be used as maintenance monotherapy treatment
- Prescriber attests to ALL of the following:
  - Patient is NOT using opioids; nor has used a short-acting opioid in the last 7 days; nor used a long-acting opioid in the last 14 days
  - Patient is NOT undergoing acute opioid withdrawal; **AND**
- Approval of non-preferred atypical antipsychotics requires trial and failure of ONE preferred agent

Nuplazid® capsules, tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form
Notes	Coverage will not be approved for psychosis not related to Parkinson's disease

Nuplazid® capsules, tablets PA Criteria:

- Hallucinations and/or delusions associated with Parkinson's disease psychosis; **AND**
- Must be ≥18 years of age; **AND**
- Trial of dose adjustment or withdrawal of anti-Parkinson medications (anticholinergics, amantadine, dopamine agonists, COMT inhibitors, selegiline) prior to treatment with Nuplazid®
- Trial and failure of ONE preferred agent

Opipza® films

PDL Status	NP
Quantity Limit	2 & 5 mg: 1/day; 10 mg: 3/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Opipza® films PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Clinically valid reason why none of the other dosage forms of aripiprazole (e.g., aripiprazole ODT) cannot be used.

Risperdal Consta® injection

PDL Status	NP
Quantity Limit	2 vials/28 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Risperdal Consta® injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Documented tolerance to the oral active ingredient; **AND**
- One of the following:
  - Diagnosis of Bipolar Disorder
  - Clinically valid reason why the patient cannot use the preferred long-acting injectables

Risperdal® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Risperdal® tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Risperidone ER injection

PDL Status	NP
Quantity Limit	2 vials/28 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Risperdal ER injection PA Criteria:

See Risperdal Consta® prior authorization criteria; **AND**

- Clinically valid reason why brand Risperdal Consta® cannot be used.

Rykindo® injection

PDL Status	NP
Quantity Limit	2 injections/28 days
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Rykindo® injection PA Criteria:

See Risperdal Consta® prior authorization criteria

Saphris® sublingual tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Saphris® sublingual tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Secuado® patches

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Secuado® patches PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms or absorb medications through the GI tract; **OR**
- Non-response due to noncompliance

Seroquel® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Seroquel® tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Seroquel XR® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Seroquel XR® tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Versacloz® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Versacloz® suspension PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Allergy or intolerance to inactive ingredient in clozapine ODT tab (i.e., dye, filler, excipient, etc.); **OR**
- Dose not achievable with ODT tab

Zyprexa® IM injection

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Zyprexa® IM injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Patient has documented tolerance to the oral active ingredient; **AND**
- Trial and failure of ONE preferred atypical antipsychotic

Zyprexa® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Zyprexa® tablets PA Criteria:

See Abilify® tablets prior authorization criteria

Zyprexa Relprevv® injection

PDL Status	NP
Quantity Limit	210 & 300 mg injection: 1 injection/2 weeks; 450 mg injection: 1 injection/month
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Zyprexa Relprevv® injection PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Documented tolerance to the oral active ingredient; **AND**
- Clinically valid reason why the patient cannot use the preferred long-acting injectables

Zyprexa Zydis® ODT

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	Atypical Antipsychotics PA Form

Zyprexa Zydis® ODT PA Criteria:

See Abilify® tablets prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms or absorb medications through the GI tract; **OR**
- Non-response due to non-compliance

Antipsychotics: Typical

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antipsychotics prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** I/DD Prior Authorization Form

Chlorpromazine, fluphenazine, haloperidol, loxapine, perphenazine, pimozide, thioridazine, thiothixene, trifluoperazine

PDL Status	P
Quantity Limit	None
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Molindone, Orap®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Molindone, Orap® PA Criteria:
Standard NP Criteria

Atypical Antipsychotic and SSRI Combinations

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Antipsychotics prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** **I/DD Prior Authorization Form**

Fluoxetine/olanzapine capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Fluoxetine/olanzapine capsules PA Criteria:

- For diagnosis of depressive episodes associated with bipolar disorder; **AND**
  - Refractory to treatment with components taken separately
- For diagnosis of major depressive disorder:
  - Must have undergone an adequate trial of at least ONE agent in THREE of the following classes of antidepressants (unless contraindicated or intolerant to):
    - Selective serotonin reuptake inhibitors (SSRIs)
    - Serotonin-norepinephrine reuptake inhibitors (SNRIs)
    - New generation antidepressants (including bupropion, mirtazapine, etc.); **AND**
  - Refractory to treatment with components taken separately

Symbyax® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Symbyax® capsules PA Criteria:

See fluoxetine/olanzapine prior authorization criteria.

Cholinergic Muscle Stimulants

Pyridostigmine 30 mg tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Pyridostigmine 30 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Miscellaneous CNS Agents

Dextromethorphan/quinidine capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Dextromethorphan/quinidine capsules PA Criteria:

- Diagnosis of Pseudobulbar Affect (PBA); **AND**
- The following patient circumstances have been excluded:
  - Heart failure or high grade (second/third degree) atrioventricular block (AV) without an implanted pacemaker
  - Patient receiving drugs that prolong QT interval and are metabolized by CYP2D6 system
  - Prolonged QT interval (including congenital long QT syndrome) or a history of Torsades de Pointes
  - Concomitantly taking monoamine oxidase inhibitors (MAOIs) or have used a MAOI in the past 14 days; **AND**
- Clinically valid reason why brand Nuedexta® capsules cannot be used

Leqembi® IQLIK autoinjector

PDL Status	NP
Quantity Limit	4 syringes or autoinjectors/28 days
PA Criteria	See below
PA Form	General PA Form
Notes	The method of measuring amyloid beta plaque levels should be consistent

Leqembi® IQLIK autoinjector Initial PA Criteria:

- Diagnosis of Alzheimer's disease; **AND**
- Presence of amyloid beta pathology has been confirmed by documented by either ONE of the following:
  - Baseline positron emission tomography (PET); **OR**
  - Lumbar puncture results; **AND**
- If patient is less than 50 years old, patient has genetic testing results documenting a mutation in amyloid precursor protein (APP), presenilin-1 (PSEN1), or presenilin-2 (PSEN2); **AND**
- Patient has received at least 18 months of consistent Leqembi (lecanemab-irmb) intravenous formulation (i.e. chart notes, claims history); **AND**
- Prescriber has provided documentation of clinical improvement or stabilization while on Leqembi (lecanemab-irmb) intravenous formulation when compared to scores before starting therapy (i.e. Clinical Dementia Rating (CDR) global score, Memory Box score, Mini-Mental State Examination (MMSE) score, or Wechsler-Memory Scale-IV Logical Memory II (subscale) (WMS-IV LMII)); **AND**
- Prescriber has submitted clinical documentation (i.e. MRI) that demonstrates patient has not experienced any amyloid related imaging abnormalities -edema (ARIA-E) and -hemosiderin deposition (ARIA-H) prior to the 3rd, 5th, 7th, 14th, IV infusions and every 18 months thereafter; **AND**
- Prescriber attests that patient does NOT meet any of the following:
  - Vasogenic edema
  - Evidence of cerebral contusion
  - Aneurysm
  - Vascular malformation
  - Infective lesions
  - Multiple lacunar infarcts or stroke involving a major vascular territory
  - Severe small vessel or white matter disease
  - History of antithrombotic agent, other than aspirin or clopidogrel, with plans to use concurrently with Leqembi Iqlik
  - Not used in combination with other Aβ monoclonal antibodies (mAbs) for Alzheimer's Disease (e.g. Kisunla); **AND**
- Prescribed by, or in consultation with, a neurologist, psychiatrist, geriatrician, who specializes in dementia disorders

Leqembi® IQLIK autoinjector Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Provider has submitted documentation of decreases (or at a minimum, no increases) in amyloid beta plaque levels, as compared to baseline, confirmed by either an amyloid-positron emission tomography (PET) scan or amyloid protein concentrations in Cerebral Spinal Fluid (CSF) obtained from a lumbar puncture performed at 18month intervals.

Nuedexta®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Nuedexta® PA Criteria:

- Diagnosis of Pseudobulbar Affect (PBA); **AND**
- The following patient circumstances have been excluded:
  - Heart failure or high grade (second/third degree) atrioventricular block (AV) without an implanted pacemaker
  - Patient receiving drugs that prolong QT interval and are metabolized by CYP2D6 system
  - Prolonged QT interval (including congenital long QT syndrome) or a history of Torsades de Pointes
  - Concomitantly taking monoamine oxidase inhibitors (MAOIs) or have used a MAOI in the past 14 days

Mood Stabilizers

Carbamazepine suspension & tablets, Depakote® DR tablets, ER tablets, sprinkle capsules, Keppra® solution, tablets, Keppra XR® tablets, Lamictal® chewables, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carbamazepine suspension & tablets, Depakote® DR tablets, ER tablets, sprinkle capsules, Keppra® solution, tablets, Keppra XR® tablets, Lamictal® chewables, tablets

PA Criteria:

Standard NP Criteria

Lamictal® ODT, lamotrigine ODT

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamictal® ODT, lamotrigine ODT PA Criteria:

- Unable to swallow; **OR**
- Unable to absorb medications through the GI tract

Lamictal XR® tablets, lamotrigine tablets dose pack, Lithobid® tablets, Subvenite® starter kit

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lamictal XR® tablets, lamotrigine tablets dose pack, Lithobid® tablets, Subvenite® starter kit PA Criteria:

Standard NP Criteria

Subvenite® suspension

PDL Status	NP
Quantity Limit	40 mL/day
PA Criteria	See below
PA Form	General PA Form

Subvenite® suspension PA Criteria:

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why lamotrigine ODT cannot be used

Tegretol® chewables, Tegretol XR® tablets, Trileptal® suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tegretol® chewables, Tegretol XR® tablets, Trileptal® suspension, tablets PA Criteria:

Standard NP Criteria

Movement Disorders

Austedo® tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Austedo® tablets PA Criteria:

Diagnosis of tardive dyskinesia:

- Patient age ≥ 18 years; **AND**
- Documentation submitted of ONE of the following:
  - A completed AIMS assessment (e.g., score or copy of evaluation); **OR**
  - Prescriber documentation of symptoms consistent with Tardive Dyskinesia (e.g., chart notes or medical records); **AND**
- Prescribed by, or in consultation with, a neurologist or psychiatrist (or other mental health provider), provided patient has reasonable access; **AND**
- Documentation or claims history of current or former chronic patient use of a dopamine antagonist (e.g., antipsychotic, metoclopramide, prochlorperazine, promethazine)

Diagnosis of chorea related to Huntington's Disease:

- Physician is experienced in the treatment of Huntington's Disease or is in a Center of Excellence for Huntington's Disease; **AND**
- Patient does not have a history of untreated or inadequately treated depression or suicidal ideation

Austedo XR® tablets, titration packs

PDL Status	P
Quantity Limit	XR tablets: 1/day; Titration packs: 1/year
PA Criteria	See below
PA Form	General PA Form

Austedo XR® tablets PA Criteria:

See Austedo® prior authorization criteria.

Ingrezza® capsules, sprinkle capsules, initiation packs

PDL Status	P
Quantity Limit	Capsules/sprinkle capsules: 40 mg: 2/day; Capsules/sprinkle capsules: 60 & 80 mg: 1/day Titration pack: 1/ year
PA Criteria	See below
PA Form	General PA Form

Ingrezza® capsules, sprinkle capsules, initiation packs PA Criteria:

Diagnosis of tardive dyskinesia

- Patient age ≥ 18 years; **AND**
- Documentation submitted of ONE of the following:
- A completed AIMS assessment (e.g., score or copy of evaluation); **OR**
- Prescriber documentation of symptoms consistent with Tardive Dyskinesia (e.g., chart notes or medical records); **AND**
- Prescribed by, or in consultation with, a neurologist or psychiatrist (or other mental health provider), provided patient has reasonable access; **AND**
- Documentation or claims history of current or former chronic patient use of a dopamine antagonist (e.g., antipsychotic, metoclopramide, prochlorperazine, promethazine)

Diagnosis of chorea related to Huntington's Disease:

- Physician is experienced in the treatment of Huntington's Disease or is in a Center of Excellence for Huntington's Disease; **AND**
- Patient does not have a history of untreated or inadequately treated depression or suicidal ideation

Tetrabenazine tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tetrabenazine tablets PA Criteria:

- Diagnosis of chorea associated with Huntington's disease

Xenazine® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Xenazine® tablets PA Criteria:

- Diagnosis of chorea associated with Huntington's disease; **AND**
- Clinically valid reason why preferred generic tetrabenazine cannot be used

Sedative Hypnotic Agents

*For children (age < 21 years), larger quantities may be approved as medically necessary.

CLASS PRIOR AUTHORIZATION CRITERIA FOR PATIENTS WITH INTELLECTUAL AND DEVELOPMENTAL DISABILITIES (I/DD):

Sedative hypnotics prescribed for disorders related to Intellectual and Developmental Disabilities will be approved if **ONE** of the following is met:

- Prescribed by a Gold Card prescriber; **OR**
- Mental health assessment applicable to behavioral symptoms for which the medication is being prescribed; **AND**
  - Underlying physical condition such as pain, discomfort, and environmental issues have been evaluated, treated, and behavioral symptoms persist; **AND**
  - Non-pharmacological interventions have been tried, and behavioral symptoms persist (e.g., behavior plan, crisis intervention, and stabilization strategies or training) and training and support have been provided to family or other caregivers; **OR**
- Short-term therapy (less than 90 days) has been prescribed; **AND**
  - Behavioral symptoms are significant enough to place the person at potential risk or needing higher level of care or loss of community placement; **OR**
    - Continuation of existing therapy to address serious and ongoing behavioral symptoms while other prior authorization criteria are met; **AND**
  - Efficacy and potential side effects to be monitored; **AND**
  - Need for requested medication will be evaluated once other non-pharmacological interventions have been tried

Note the following:

- Drug specific step therapy, NP criteria and Brand Medically Necessary Criteria will still apply to this patient population.
- The I/DD Worksheet can be found at**:** I/DD Prior Authorization Form

Doxepin concentrate (10 mg/mL)

PDL Status	P
Quantity Limit	None*
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Eszopiclone tablets, ramelteon tablets, zaleplon capsules, zolpidem tablets

PDL Status	P
Quantity Limit	14/30 days*
PA Criteria	PA required for I/DD members only. See I/DD Criteria.

Ambien® tablets, Ambien CR® tablets, Belsomra® tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Ambien® tablets, Ambien CR® tablets, Belsomra® tablets PA Criteria:
Standard NP Criteria

Dayvigo® tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Dayvigo® tablets PA Criteria:

- Diagnosis of insomnia characterized by difficulties with sleep onset and/or sleep maintenance; **AND**
- Patient is 18 years of age or older; **AND**
- Narcolepsy and other insomnia-related disorders have been ruled out (e.g., movement, breathing, psychiatric disorders, medications); **AND**
- Trial (at least 3 weeks) of non-pharmacological therapies (e.g., stimulus control, sleep restriction, sleep hygiene measures and relaxation therapy); **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Patients who are pregnant should be registered in the Dayvigo® pregnancy registry

Doxepin tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Doxepin tablets PA Criteria:

- Documented trial and failure of (defined as ≥ 1 week) at an appropriate dose of preferred doxepin concentrate

Edluar® sublingual tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Edluar® sublingual tablets PA Criteria:

- Patient is unable to swallow solid dosage forms

Estazolam tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Estazolam tablets PA Criteria:

- Diagnosis of Insomnia; **AND**
- Medical documentation that rules out other insomnia related disorders (e.g., movement, breathing, psychiatric disorders and medication); **AND**
- Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., stimulus control, sleep restriction, sleep hygiene measures and relaxation therapy); **AND**
- Use of 2 preferred agents, unless patient has a contraindication or allergy; **AND**
- Due to increased risk of toxicity, patient should not be pregnant; **AND**
- Will not be taken concurrently with CNS stimulants, opiates, carisoprodol, meprobamate or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse

Flurazepam capsules

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Flurazepam capsules PA Criteria:

See estazolam prior authorization criteria

Halcion® capsules

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Halcion® capsules PA Criteria:

- Diagnosis of Insomnia; **AND**
- Medical documentation that rules out other insomnia related disorders (e.g., movement, breathing, psychiatric disorders & medication/substance use); **AND**
- Documented trial (at least 3 weeks) of non-pharmacological therapies (e.g., stimulus control, sleep restriction, sleep hygiene measures & relaxation therapy); **AND**
- Use of 2 preferred agents, unless patient has a contraindication or allergy; **AND**
- Clinical reason as to why patient cannot use generic equivalent; **AND**
- Due to increased risk of toxicity, patient should not be pregnant; **AND**
- Will not be taken concurrently with CNS stimulants, opiates, carisoprodol, meprobamate or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol **OR** drug dependence/abuse

Hetlioz® capsules

PDL Status	NP
Quantity Limit	30/60 days*
PA Criteria	See below
PA Form	General PA Form

Hetlioz® capsules PA Criteria:

- Treatment of non-24-hour sleep wake disorder (non-24 or N24) in members who are unable to distinguish between light and darkness in both eyes; **OR**
- Treatment of nighttime sleep disturbances in Smith-Magenis Syndrome (SMS) in patients 16 years of age and older; **AND**
- Trial and failure or contraindication to melatonin; **AND**
- Patient will not take any of the following:
  - Strong CYP1A2 inhibitors (e.g., fluvoxamine)
  - Strong CYP3A4 inducers (e.g., rifampin)

Hetlioz® suspension

PDL Status	NP
Quantity Limit	5 mL/day and 158 mL/60 days*
PA Criteria	See below
PA Form	General PA Form

Hetlioz® suspension PA Criteria:

- Treatment of nighttime sleep disturbances in Smith-Magenis Syndrome (SMS); **AND**
- Patient is at least 3 years of age but not greater than 15 years of age; **AND**
- Trial and failure or contraindication to melatonin; **AND**
- Patient is unable to swallow/absorb medications through the GI tract; **AND**
- Patient will not take any of the following:
  - Strong CYP1A2 inhibitors (e.g., fluvoxamine)
  - Strong CYP3A4 inducers (e.g., rifampin)

Quazepam tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Quazepam tablets PA Criteria:

See estazolam prior authorization criteria

Quviviq® tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Quviviq® tablets PA Criteria:

- Patient must be 18 years of age or older; **AND**
- Diagnosis of insomnia characterized by difficulties with sleep onset and/or sleep maintenance; **AND**
- Narcolepsy and other insomnia-related disorders have been ruled out (e.g., movement, breathing, psychiatric disorders, medications); **AND**
- Trial (at least 3 weeks) of non-pharmacological therapies (e.g., stimulus control, sleep restriction, sleep hygiene measures and relaxation therapy); **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Patient should not have any of the following diagnoses: Narcolepsy, COPD, or moderate to severe OSA; **AND**
- Concurrently not taking CNS stimulants, opiates, carisoprodol, meprobamate or barbiturates; **AND**
- Due to increased risk of dependency, patient does not have a history of alcohol or drug dependence/abuse; **AND**
- Patients who are pregnant should be registered in the Quviviq® pregnancy registry

Restoril® capsules

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Restoril® capsules PA Criteria:

See Halcion® prior authorization criteria.

Rozerem® tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Rozerem® tablets PA Criteria:
Standard NP Criteria

Tasimelteon capsules

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Tasimelteon capsules PA Criteria:
See Hetlioz® capsules prior authorization criteria; **AND**

- Clinically valid reason why Hetlioz® capsules cannot be used

Temazepam capsules, triazolam tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	Anti-Anxiety Agents PA Form

Temazepam capsules, triazolam tablets PA Criteria:
See estazolam prior authorization criteria.

Zolpidem capsules

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Zolpidem capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Zolpidem ER tablets, zolpidem tartrate sublingual tablets

PDL Status	NP
Quantity Limit	14/30 days*
PA Criteria	See below
PA Form	General PA Form

Zolpidem ER tablets, zolpidem tartrate sublingual tablets PA Criteria:
Standard NP Criteria

Skeletal Muscle Relaxants

Amrix® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Amrix® capsules PA Criteria:

- Diagnosis of an FDA-approved indication; **AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred cyclobenzaprine

Atmeksi® suspension

PDL Status	NP
Quantity Limit	30 mL/day
PA Criteria	See below
PA Form	General PA Form

Atmeksi® suspension PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Baclofen solution (1 mg/mL, 2 mg/mL)

PDL Status	NP
Quantity Limit	1 mg/mL: 80 mL/day; 2 mg/mL: 40 mL/day
PA Criteria	See below
PA Form	General PA Form

Baclofen solution PA Criteria:

- Diagnosis of spasticity with flexor spasms and concomitant pain, clonus, and/or muscular rigidity (e.g., multiple sclerosis, spinal cord injury, other spinal cord disease); **AND**
- Documented inability to swallow baclofen tablets; **AND**
- Clinically valid reason why baclofen suspension cannot be used (e.g., documented allergy or contraindication to an inactive ingredient that is NOT present in the suspension or tablets)

Baclofen suspension (5 mg/mL)

PDL Status	NP
Quantity Limit	16 mL/day
PA Criteria	See below
PA Form	General PA Form

Baclofen suspension PA Criteria:

- Diagnosis of spasticity with flexor spasms and concomitant pain, clonus, and/or muscular rigidity (e.g., multiple sclerosis, spinal cord injury, other spinal cord disease); **AND**
- Documented inability to swallow baclofen tablets

Baclofen 5 and 15 mg tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Baclofen 5 and 15 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Carisoprodol tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Carisoprodol tablets PA Criteria:

- Patient is 16 years of age or older; **AND**
- Contraindication, drug to drug interaction, or history of toxic side effects that will cause immediate or long-term damage with ALL preferred skeletal muscle relaxants; **AND**
- Patient does not have a history of, or received treatment for, drug dependency or drug abuse; **AND**
- Prescriber must have checked the Tennessee Controlled Substance Database for this patient within the last 30 days; **AND**
- Patient is not concurrently utilizing any other opioid therapy

Chlorzoxazone tablets (except 250 mg)

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Chlorzoxazone tablets PA Criteria:

Standard NP criteria

Chlorzoxazone 250 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Chlorzoxazone 250 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Cyclobenzaprine 7.5 mg tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Cyclobenzaprine 7.5 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Cyclobenzaprine ER capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Cyclobenzaprine ER capsules PA Criteria:

See Amrix® prior authorization criteria

Dantrolene capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dantrolene capsules PA Criteria:

Standard NP criteria

Fleqsuvy® suspension

PDL Status	NP
Quantity Limit	16 mL/day
PA Criteria	See below
PA Form	General PA Form

Fleqsuvy® suspension PA Criteria:

See baclofen suspension prior authorization criteria

Metaxalone 400 mg and 800 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Metaxalone 400 mg and 800 mg tablets PA Criteria:

Standard NP criteria

Metaxalone 640 mg tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Metaxalone 640 mg PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Methocarbamol 1000 mg tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Methocarbamol 1000 mg tablets PA Criteria:

- Clinically valid reason why preferred agents cannot be used

Norgesic® tablets, Norgesic Forte® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Norgesic® tablets, Norgesic Forte® tablets PA Criteria:

- Diagnosis of an FDA-approved indication**; AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Ontralfy® solution

PDL Status	NP
Quantity Limit	15 mL/day
PA Criteria	See below
PA Form	General PA Form

Ontralfy® solution PA Criteria:

- One of the following:
  - The patient has experienced therapeutic failure to 2 preferred medications; **OR**
  - Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications (please specify details and pend to RPh); **AND**
- Patient is unable to swallow solid dosage forms

Orphengesic Forte® tablets, orphenadrine/aspirin/caffeine tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Orphengesic Forte® tablets, orphenadrine/aspirin/caffeine tablets PA Criteria:

- Diagnosis of an FDA-approved indication**; AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Orphenadrine ER tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Orphenadrine ER tablets PA Criteria:
Standard NP criteria

Ozobax DS® suspension

PDL Status	NP
Quantity Limit	40 mL/day
PA Criteria	See below
PA Form	General PA Form

Ozobax DS® suspension PA Criteria:
See baclofen suspension prior authorization criteria

Soma®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Soma® PA Criteria:
See carisoprodol prior authorization criteria

Tanlor® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Tanlor® tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Tizanidine 2, 4, and 6 mg capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tizanidine 2, 4, and 6 mg capsules PA Criteria:
Standard NP criteria

Tizanidine 8 mg capsules

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Tizanidine capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Zanaflex® 2, 4, and 6 mg capsules and 4 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Zanaflex® 2, 4, and 6 mg capsules and 4 mg tablets PA Criteria:

Standard NP criteria

Zanaflex® 8 mg capsules

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Zanaflex® capsules and tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

V. DERMATOLOGICS

Acne Agents, Topical

Covered for recipients < 21 years old only.

Benzoyl peroxide 2.5%, 5%, 10% (excluding cleanser, gel, microspheres, & towelettes), clindamycin phosphate (excluding foam, lotion & 75 mL gel), clindamycin/benzoyl peroxide gel, erythromycin 2% solution, sodium sulfacetamide/sulfur

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

All branded single agent and combination products of benzoyl peroxide, clindamycin, erythromycin, and sodium sulfacetamide

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

All branded single agent and combination products of benzoyl peroxide, clindamycin, erythromycin, and sodium sulfacetamide PA Criteria:
Standard NP Criteria

Aczone®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Aczone® PA Criteria:

- Patient is at least 9 years of age and less than 21 years of age; **AND**
- Patient has a diagnosis of acne vulgaris; **AND**
- Clinically valid reason why generic dapsone gel cannot be used

Amzeeq®

PDL Status	NP
Quantity Limit	1 package/28 days
PA Criteria	See below
PA Form	General PA Form

Amzeeq® PA Criteria:

- Diagnosis of non-nodular moderate to severe acne vulgaris; **AND**
- Patient is at least 9 years of age and less than 21 years of age; **AND**
- Trial and failure, contraindication, or intolerance to ALL the following:
  - 2 preferred agents
  - Minocycline capsules; **AND**
- Prescriber must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Benzoyl peroxide cleansers, gel, microspheres, towelettes, Clindamycin phosphate foam, 75 mL gel, lotion

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Benzoyl peroxide cleansers, gel, microspheres, towelettes, Clindamycin phosphate foam, 75 mL gel, lotion PA Criteria:
Standard NP Criteria

Dapsone gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Dapsone gel PA Criteria:

- Patient is at least 9 years of age and less than 21 years of age; **AND**
- Patient has a diagnosis of acne vulgaris; **AND**
- Clinically valid reason why the preferred agents cannot be used

Dermatological kits

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Dermatological kits PA Criteria:

- Trial and failure of 3 preferred agents; **AND**
- Trial and failure of the individual components of the kit

Erythromycin/benzoyl peroxide gel, Ery® pad, erythromycin 2% gel, Ery® pad, erythromycin 2% gel, Sulfacetamide suspension

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Erythromycin/benzoyl peroxide gel, Ery® pad, erythromycin 2% gel, Ery® pad, erythromycin 2% gel, Sulfacetamide suspension PA Criteria:

Standard NP Criteria

Winlevi®

PDL Status	NP
Quantity Limit	1 tube/30 days
PA Criteria	See below
PA Form	General PA Form

Winlevi® PA Criteria:

- Diagnosis of acne vulgaris; **AND**
- Patient is at least 12 years of age and less than 21 years of age; **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents; **AND**
- Prescriber provides peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Agents for Burns, Topical

Silver sulfadiazine, SSD®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Mafenide packet, Silvadene®, Sulfamylon® cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Mafenide, Silvadene® Sulfamylon® cream PA Criteria:
Standard NP Criteria

Agents for Rosacea, Topical

Covered for recipients < 21 years old only.

Azelaic acid gel 15%, Finacea® foam

PDL Status	P
Quantity Limit	50 g/Rx
PA Criteria	None

Metronidazole cream, gel, lotion

PDL Status	P
Quantity Limit	60 g/Rx
PA Criteria	None

Brimonidine gel

PDL Status	NP
Quantity Limit	30 g/Rx
PA Criteria	See below
PA Form	General PA Form

Brimonidine gel PA Criteria:
Standard NP Criteria

Ivermectin cream, Miravaso® gel

PDL Status	NP
Quantity Limit	45 g/Rx
PA Criteria	See below
PA Form	General PA Form

Ivermectin cream, Miravaso® gel PA Criteria:
Standard NP Criteria

Oxymetazoline cream

PDL Status	NP
Quantity Limit	30 g/Rx
PA Criteria	See below
PA Form	General PA Form

Oxymetazoline cream PA Criteria:

- Patient age < 21 years of age; **AND**
- Patient has a diagnosis of persistent facial erythema associated with rosacea; **AND**
- Trial and failure, or contraindication, of 2 of the following: brimonidine, ivermectin, tetracycline, minocycline, doxycycline, erythromycin, clindamycin, benzoyl peroxide; **AND**
- Trial and failure of 2 preferred topical agents for rosacea; **AND**
- Clinically valid reason why brand Rhofade® cannot be used

Rhofade® cream

PDL Status	NP
Quantity Limit	30 g/Rx
PA Criteria	See below
PA Form	General PA Form

Rhofade® cream PA Criteria:

- Patient age < 21 years of age; **AND**
- Patient has a diagnosis of persistent facial erythema associated with rosacea; **AND**
- Trial and failure, or contraindication, of 2 of the following: brimonidine, ivermectin, tetracycline, minocycline, doxycycline, erythromycin, clindamycin, benzoyl peroxide; **AND**
- Trial and failure of 2 preferred topical agents for rosacea

Anesthetics, Topical

Lidocaine (excluding kits, lotion, solution)

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Lidocaine patch 5%

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Lidocaine patch 5% PA Criteria:

- Diagnosis of post-herpetic neuralgia

Lidocaine/prilocaine

PDL Status	P
Quantity Limit	30 g/Rx
PA Criteria	None

ZTLido®

PDL Status	P
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

ZTLido® PA Criteria:

- Diagnosis of post-herpetic neuralgia

Lidocaine/hydrocortisone

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Lidocaine/hydrocortisone PA Criteria:

- Diagnosis of FDA-approved indication; **AND**
- Clinically valid reason why the preferred topical anesthetics cannot be used

Lidocaine kits

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lidocaine kits PA Criteria:

- Diagnosis of FDA-approved indication; **AND**
- Clinically valid reason why the preferred topical anesthetics cannot be used; **AND**
- For combination kits, trial and failure of individual agents

Pliaglis® cream, Pramosone® 2.5-1% lotion

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Pliaglis® cream, Pramosone® 2.5-1% lotion PA Criteria:
Standard NP Criteria

Antibiotics, Topical

Mupirocin ointment

PDL Status	P
Quantity Limit	44 g/Rx
PA Criteria	None

Mupirocin cream

PDL Status	NP
Quantity Limit	30 g/Rx
PA Criteria	See below
PA Form	General PA Form

Mupirocin cream PA Criteria:
Standard NP Criteria

Silver nitrate solution 0.5%

PDL Status	NP
Quantity Limit	1 bottle/Rx
PA Criteria	See below
PA Form	General PA Form

Silver nitrate solution 0.5% PA Criteria:

- Will be used for granulation tissue removal; **AND**
- Patient cannot use a preferred PDL agent (e.g., topical antibiotics, topical steroids)

Antifungal Agents, Topical

Ciclopirox cream, 8% solution, Clotrimazole 1% cream, solution, Clotrimazole/betamethasone cream, lotion, Ketoconazole cream, shampoo, Nystatin/triamcinolone cream, ointment

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Nystatin topical powder

PDL Status	P
Quantity Limit	120 g/Rx
PA Criteria	None

Vusion®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Ciclopirox gel, suspension

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Ciclopirox gel, suspension PA Criteria:
Standard NP Criteria

Ciclopirox nail kit

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Ciclopirox nail kit PA Criteria:

- Clinically valid reason why the preferred ciclopirox 8% topical solution cannot be used

Clotrimazole 1% solution (Rx), Econazole, Ertaczo®, ketoconazole 2% foam, Ketodan® foam, Klayesta®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Clotrimazole 1% solution, Econazole, Ertaczo®, ketoconazole 2% foam, Ketodan® foam, Klayesta® PA Criteria:
Standard NP Criteria

Miconazole/zinc/petrolatum ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Miconazole/zinc/petrolatum ointment PA Criteria:

- Clinically valid reason why the preferred Vusion® ointment cannot be used

Naftifine cream, gel, Nyamyc®, Oxiconazole cream, lotion Oxistat®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Natifine cream, gel, Nyamyc®, Oxiconazole cream, lotion, Oxistat® PA Criteria:
Standard NP Criteria

Tavaborole solution

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Tavaborole solution PA Criteria:

- Diagnosis of mild to moderate onychomycosis of the toenails; **AND**
- Trial and failure, contraindication, or intolerance to the preferred topical ciclopirox 8% solution

Antineoplastics, Topical

Diclofenac 3% gel

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Diclofenac 3% gel PA Criteria:

- Diagnosis of actinic keratosis

Fluorouracil 5% cream, Imiquimod

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Bexarotene, Fluorouracil 0.5% cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Bexarotene, Fluorouracil 0.5% cream PA Criteria:

Standard NP Criteria

Hyftor®

PDL Status	NP
Quantity Limit	30 g/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	4 months

Hyftor® initial PA Criteria:

- Diagnosis of facial angiofibroma associated with tuberous sclerosis complex; **AND**
- Patient is 6 years of age or older; **AND**
- Prescribed by or in consultation with a dermatologist or neurologist; **AND**
- Patient is not a candidate for laser therapy or surgical treatments

Hyftor® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement in size or redness of facial angiofibroma)

Valchlor®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Valchlor® PA Criteria:

- Diagnosis of stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma; **AND**
- Patient has received skin directed therapy

Zyclara®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Zyclara® PA Criteria:

- Diagnosis of actinic keratosis; **OR**
- Diagnosis of basal cell carcinoma

Antipruritics/Antihistamines, Topical

Doxepin cream

PDL Status	NP
Quantity Limit	45 g/90 day
PA Criteria	See below
PA Form	General PA Form

Doxepin cream PA Criteria:

- Patient has moderate pruritus due to various forms of eczematous dermatitis, including atopic dermatitis and lichen simplex chronicus; **AND**
- Inadequate response, intolerance, or contraindication to BOTH of the following:
- A topical corticosteroid
- An oral antihistamine (first or second generation) or a topical antihistaminic agent

Prudoxin®, Zonalon®

PDL Status	NP
Quantity Limit	45 g/90 day
PA Criteria	See below
PA Form	General PA Form

Prudoxin®, Zonalon® PA Criteria:

See doxepin cream prior authorization criteria

Antipsoriatics, Oral

Acitretin

PDL Status	NP
Quantity Limit	10 mg: 3/day; 17.5, 22.5., & 25 mg: 2/day
PA Criteria	See below
PA Form	General PA Form
Notes	Will not be covered for the diagnosis of acne or rosacea for recipients ≥ 21 years of age.

Acitretin PA Criteria:

- Patient has a diagnosis of severe psoriasis; **AND**
- Minimum duration of a 4-week trial and failure, contraindication, or intolerance to at least two of the following:
  - Corticosteroids (e.g., betamethasone, clobetasol)
  - Vitamin D analogs (e.g., calcitriol, calcipotriene)
  - Tazarotene
  - Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Prescriber attests to each of the following:
  - Patient does-NOT have impaired liver or kidney function, or abnormally elevated lipid levels
  - Patient will NOT be receiving concomitant methotrexate (due to risk of hepatitis) or tetracyclines (due to risk of increased intracranial pressure)
  - If applicable, appropriate laboratory assessments and counseling have been conducted regarding risks associated with pregnancy

Methoxsalen

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	Will not be covered for the diagnosis of acne or rosacea for recipients ≥ 21 years of age.

Methoxsalen PA Criteria:

- Diagnosis of severe, recalcitrant, disabling psoriasis supported by biopsy; **AND**
- Minimum duration of a 4-week trial and failure, contraindication, or intolerance to at least two of the following:
  - Corticosteroids (e.g., betamethasone, clobetasol)
  - Vitamin D analogs (e.g., calcitriol, calcipotriene)
  - Tazarotene
  - Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

Antipsoriatics, Topical

Calcipotriene cream, scalp solution, Taclonex®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Calcipotriene cream, scalp solution, Taclonex® PA Criteria:

- Trial and failure, contraindication, or intolerance to ≥ 1 topical steroid

Tazarotene 0.05%cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Tazarotene 0.05% cream PA Criteria:

- Diagnosis of psoriasis

Tazarotene 0.1% cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Tazarotene 0.1% cream PA Criteria:

- Diagnosis of psoriasis; **OR**
- Diagnosis of acne in patients less than 21 years of age

Calcipotriene foam, ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Calcipotriene foam, ointment PA Criteria:

- Trial and failure, contraindication, or intolerance to ≥ 1 topical steroid; **AND**
- Trial and failure, contraindication, or intolerance to a preferred topical antipsoriatic agent

Calcitriol ointment, Calcipotriene/betamethasone, Calcitene® ointment, Enstilar®, Sorilux®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Calcitriol ointment, Calcipotriene/betamethasone, Calcitene® ointment, Enstilar®, Sorilux® PA Criteria:
See calcipotriene foam prior authorization criteria.

Tazarotene 1% gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Tazarotene 1% gel PA Criteria:

- Diagnosis of psoriasis **AND** Both of the following:
  - Trial and failure, contraindication, or intolerance to at least one topical steroid;
  - Trial and failure, contraindication, or intolerance to a preferred topical antipsoriatic agent**; OR**
- Diagnosis of acne in patients less than 21 years of age; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred topical retinoids

Vtama®

PDL Status	NP
Quantity Limit	60 g/28 days
PA Criteria	See below
PA Form	General PA Form

Vtama® Initial PA Criteria:

- Diagnosis of plaque psoriasis; **AND**
  - Minimum duration of a 4-week trial and failure, contraindication, or intolerance to at least two of the following:
    - Corticosteroids (e.g., betamethasone, clobetasol)
    - Vitamin D analogs (e.g., calcitriol, calcipotriene)
    - Tazarotene
    - Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus; **OR**
- Diagnosis of atopic dermatitis and Both of the following:
  - Trial and failure, contraindication, or intolerance to a topical corticosteroid **AND** a topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist

Vtama® Renewal PA Criteria:

- Documentation of positive clinical response to therapy as evidenced by one of the following:
  - Reduction in the body surface area (BSA) involvement from baseline
  - Improvement in symptoms (e.g., pruritus, inflammation) from baseline

Zoryve® 0.3% cream

PDL Status	NP
Quantity Limit	60 g/28 days
PA Criteria	See below
PA Form	General PA Form

Zoryve® 0.3% cream Initial PA Criteria:

- Diagnosis of plaque psoriasis; **AND**
- Patient is 2 years of age or older; **AND**
- Patient does not have moderate to severe liver impairment (Child-Pugh B or C); **AND**
- Request is for Zoryve 0.3% cream; **AND**
- Trial and failure, contraindication, or intolerance to TWO preferred topical antipsoriatic agents

Zoryve® 0.3% cream Renewal PA Criteria:

- Patient continues to be monitored for liver impairment; **AND**
- Documented clinical improvement in response to treatment (e.g. reduction in itch, rash, inflammation)

Antiseborrheic Agents

Selenium sulfide 2.5% lotion

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Ovace Plus®; Ovace Plus Wash®; selenium sulfide shampoo (except 2.3%); sodium sulfacetamide gel, liquid, shampoo (except 9.8%)

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Ovace Plus®; Ovace Plus Wash®; selenium sulfide shampoo (except 2.3%); sodium sulfacetamide gel, liquid, shampoo (except 9.8%) PA Criteria:
Standard NP Criteria

Zoryve® 0.3% topical foam

PDL Status	NP
Quantity Limit	1 can (60 g)/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Plaque psoriasis: 1 year Seborrheic dermatitis: Initial: 3 months; Renewal: 1 year

Zoryve® 0.3% topical foam Initial PA Criteria:
Seborrheic Dermatitis
- Diagnosis of seborrheic dermatitis; **AND**
- Patient is 9 years of age or older; **AND**
- Patient does not have moderate to severe liver impairment (Child-Pugh B or C); **AND**
- Trial and failure, contraindication, or intolerance to BOTH of the following agents:
  - Topical antifungals (ketoconazole, ciclopirox, miconazole, clotrimazole)
  - Topical corticosteroids

Plaque Psoriasis
- Diagnosis of mild to moderate plaque psoriasis; **AND**
- Patient is 12 years of age or older; **AND**
- Patient does not have moderate to severe liver impairment (Child-Pugh B or C); **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred topical antipsoriatic agents

Zoryve® 0.3% topical foam Renewal PA Criteria:
Seborrheic Dermatitis
- Patient continues to be monitored for liver impairment; **AND**
- Documented clinical improvement in response to treatment (e.g., decreased erythema, scaling, inflammation); **AND**
- Patient does not have any treatment limiting adverse effects

Plaque Psoriasis
- Patient continues to be monitored for liver impairment; **AND**
- Documented clinical improvement in response to treatment (e.g., decreased erythema, scaling, inflammation); **AND**
- Patient does not have any treatment limiting adverse effects

Antivirals, Topical

Acyclovir 5% ointment, penciclovir cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Acyclovir cream, Denavir® cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Acyclovir cream, Denavir® cream PA Criteria:
Standard NP Criteria

Zelsuvmi® 10.3% gel

PDL Status	NP
Quantity Limit	1 kit/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	3 months

Zelsuvmi® 10.3% gel PA Criteria:

- Diagnosis of Molluscum Contagiosum; **AND**
- Prescribed by, or in consultation with, a dermatologist; **AND**
- Patient meets ONE of the following:
  - Patient has a chronic skin condition;
  - Patient is Immunocompromised (e.g. auto-immune disease, HIV, etc.)
  - Patient has extensive involvement or experiences bleeds, secondary infections, or lesions in sensitive areas (e.g. facial, anogenital); **AND**
- Total treatment duration will not exceed 12 weeks

Atopic Dermatitis Agents, Topical

Eucrisa®

PDL Status	P
Quantity Limit	1 tube/month
PA Criteria	See below
PA Form	General PA Form

Eucrisa® PA Criteria:

- Diagnosis of atopic dermatitis; **AND**
- One of the following:
  - Patient is ≥ 2 years and meets BOTH of the following; **AND**
    - Trial and failure, contraindication, or intolerance to BOTH of the following:
      - A topical corticosteroid; **AND**
      - A topical calcineurin Inhibitor (e.g., Elidel or tacrolimus ointment)**; OR**
  - Patient is <2 years and greater than 3 months of age; **AND**
    - Trial and failure, contraindication, or intolerance to ONE topical corticosteroid

Pimecrolimus cream, tacrolimus ointment

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Anzupgo®

PDL Status	NP
Quantity Limit	60 g/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Anzupgo® Initial PA Criteria:

- Diagnosis of moderate to severe chronic hand eczema (CHE); **AND**
- Patient is 18 years of age or older; **AND**
- Prescribed by or in consultation with a specialist (i.e. dermatologist, allergist/immunologist); **AND**
- Trial and Failure, contraindication, or intolerance to BOTH of the following:
  - Preferred topical corticosteroid
  - Preferred Calcineurin inhibitor (e.g. tacrolimus ointment); **AND**
- Not used in combination with other Janus kinase (JAK) inhibitors or potent immunosuppressants (e.g., azathioprine or cyclosporine)

Anzupgo® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient demonstrates positive clinical response to therapy (e.g., reduction in erythema, scaling, hyperkeratosis, fissures)

Opzelura®

PDL Status	NP
Quantity Limit	240 g/month
PA Criteria	See below
PA Form	General PA Form

Opzelura® Initial PA Criteria:

- One of the following:
  - Diagnosis of mild to moderate atopic dermatitis (3-month approval duration) and ALL of the following:
    - Patient is not immunocompromised; **AND**
    - Opzelura will only be used for short-term and/or non-continuous chronic treatment; **AND**
    - Trial and failure, contraindication, or intolerance to Vtama®; **AND**
    - Patient is 2 years of age or older
  - Diagnosis of Nonsegmental Vitiligo (12-month approval duration); **AND**
    - Patient is 12 years of age or older; **AND**
- Patient is not breastfeeding; **AND**
- Trial and failure, contraindication, or intolerance to a topical corticosteroid; **AND**
- Trial and failure, contraindication, or intolerance to a topical calcineurin inhibitor; **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) **OR** potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the heart-related or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak Kinase inhibitor therapy is appropriate

Opzelura® Renewal PA Criteria:

- Positive response to therapy [e.g., reduction in symptoms (itch, rash, etc.), re-pigmentation, etc.]

Zoryve® 0.05% cream

PDL Status	NP
Quantity Limit	60 g/28 days
PA Criteria	See below
PA Form	General PA Form

Zoryve® 0.05% cream Initial PA Criteria:

- Diagnosis of mild to moderate atopic dermatitis; **AND**
- Patient is at least 2 yrs old and ≤ 6 years of age; **AND**
- Patient does not have moderate to severe liver impairment (Child-Pugh B or C); **AND**
- Request is for Zoryve 0.05% cream; **AND**
- Trial and failure, contraindication, or intolerance to ONE topical corticosteroid; **AND**
- Trial and failure, contraindication, or intolerance to ONE topical calcineurin inhibitor; **AND**
- Trial and failure, contraindication, or intolerance to Vtama®

Zoryve® 0.05% cream Renewal PA Criteria:

- Patient continues to be monitored for liver impairment; **AND**
- Documented clinical improvement in response to treatment (e.g. reduction in itch, rash, inflammation)

Zoryve® 0.15% cream

PDL Status	NP
Quantity Limit	60 g/28 days
PA Criteria	See below
PA Form	General PA Form

Zoryve® 0.15% cream Initial PA Criteria:

- Diagnosis of mild to moderate atopic dermatitis; **AND**
- Patient is 6 years of age or older; **AND**
- Patient does not have moderate to severe liver impairment (Child-Pugh B or C); **AND**
- Request is for Zoryve 0.15% cream; **AND**
- Trial and failure, contraindication, or intolerance to ONE topical corticosteroid; **AND**
- Trial and failure, contraindication, or intolerance to ONE topical calcineurin inhibitor; **AND**
- Trial and failure, contraindication, or intolerance to Vtama®

Zoryve® 0.15% cream Renewal PA Criteria:

- Patient continues to be monitored for liver impairment; **AND**
- Documented clinical improvement in response to treatment (e.g. reduction in itch, rash, inflammation)

Emollients

Ammonium lactate cream, lotion (Rx)

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Genital Wart Agents

Imiquimod cream, Condylox® gel

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Imiquimod pump, Veregen® ointment, Zyclara® pump

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Imiquimod pump, Veregen®, Zyclara® PA Criteria:
Standard NP Criteria

Keratolytic Agents

Generic urea products (Rx) (except 39, 39.5, and 41% cream), generic salicylic acid products (Rx)

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Brand urea products (Rx), brand salicylic acid products (Rx)

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Brand urea products, brand salicylic acid products PA Criteria:
Standard NP Criteria

Urea 39, 39.5, & 41% cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Urea 39, 39.5, & 41% cream PA Criteria:

- Clinically valid reason why urea 40% products cannot be used

Pediculocides/Scabicides

Please refer to the OTC Drug Formulary for coverage of OTC products.

Natroba® suspension, permethrin cream, spinosad suspension

PDL Status	P
Quantity Limit	2 packages/Rx
PA Criteria	None

Crotan® lotion

PDL Status	NP
Quantity Limit	1 bottle/Rx
PA Criteria	See below
PA Form	General PA Form

Crotan® lotion PA Criteria:

- Patient is being treated for scabies or pruritus; **AND**
- Patient has tried/failed permethrin (unless patient has a contraindication)

Ivermectin lotion

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Ivermectin lotion PA Criteria:
Standard NP Criteria

Malathion lotion, Ovide® lotion

PDL Status	NP
Quantity Limit	2 bottles/Rx
PA Criteria	See below
PA Form	General PA Form

Malathion lotion, Ovide® lotion PA Criteria:
Standard NP Criteria

Pruradik® lotion

PDL Status	NP
Quantity Limit	1 bottle/Rx
PA Criteria	See below
PA Form	General PA Form

Pruradik® lotion PA Criteria:

- Patient is being treated for scabies or pruritus; **AND**
- Patient has tried/failed permethrin (unless patient has a contraindication)

Sklice® lotion

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Sklice® lotion PA Criteria:
Standard NP Criteria

Retinoids, Oral

Absorica®, Absorica LD®, Accutane®, Amnesteem®, Claravis®, isotretinoin, Zenatane®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
Notes	Active registration and compliance with the iPLEDGE program is required by prescriber, patient, and pharmacy.

Absorica®, Absorica LD®, Accutane®, Amnesteem®, Claravis®, isotretinoin, Zenatane® PA Criteria:

- Diagnosis of chronic myelogenous leukemia, head or neck cancer, ichthyosis, keratosis follicularis, neuroblastoma, or pityriasis rubra pilaris will be reviewed on a case-by-case basis; **OR**
- Diagnosis of severe recalcitrant nodular acne; **AND**
  - Patient is < 21 years of age (will not be covered for acne or rosacea for recipients ≥ 21 years of age)

Retinoids, Topical

Adapalene 0.1% cream, 0.3% gel, 0.3% gel pump (Rx), Adapalene/benzoyl peroxide gel, Tretinoin cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Adapalene 0.1% cream, 0.3% gel, 0.3% gel pump Adapalene/benzoyl peroxide gel, Tretinoin cream PA Criteria:

- Patient is < 21 years old; **AND**
  - Diagnosis of acne, keratosis follicularis, verruca plana, or actinic keratosis; **OR**
- Patient is ≥ 21 years old: **AND**
  - Diagnosis of keratosis follicularis (1 year approval duration); **OR**
  - Diagnosis of verruca plana (2-month approval duration); **OR**
  - Diagnosis of actinic keratosis for the prevention of future lesions (1 year approval duration)

Tazarotene 0.1% cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year; Psoriasis: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Tazarotene 0.1% cream, PA Criteria:

- Patient is < 21 years old; **AND**
- One of the following:
  - Diagnosis of acne, keratosis follicularis, verruca plana, or actinic keratosis;
  - Diagnosis of psoriasis; **AND**
    - Patient has tried and failed at least one topical steroid or has a contraindication to the use of topical steroids; **OR**
- Patient is ≥ 21 years old: **AND**
  - Diagnosis of psoriasis; **AND**
    - Patient has tried and failed at least one topical steroid or has a contraindication to the use of topical steroids;

Clindamycin/tretinoin gel, Differin 0.1% cream, lotion, Differin 0.3% gel pump (Rx), tretinoin gel, tretinoin microsphere gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Clindamycin/tretinoin gel, Differin 0.1% cream, lotion, Differin 0.3% gel pump (Rx), tretinoin gel, tretinoin microsphere gel PA Criteria:

- Patient is < 21 years old; **AND**
  - Diagnosis of acne, keratosis follicularis, verruca plana, or actinic keratosis; **OR**
- Patient is ≥ 21 years old: **AND**
  - Diagnosis of keratosis follicularis (1 year approval duration); **OR**
  - Diagnosis of verruca plana (2-month approval duration); **OR**
  - Diagnosis of actinic keratosis for the prevention of future lesions (1 year approval duration); **AND**
- Patient has tried and failed 2 preferred products or has allergy, contraindication, drug-drug interaction or a history of unacceptable/toxic side effects with ALL preferred products

Fabior® foam

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year; Psoriasis: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Fabior® foam PA Criteria:

See tazarotene 0.1% cream prior authorization criteria; **AND**

- Patient has tried and failed 2 preferred products or has allergy, contraindication, drug-drug interaction or a history of unacceptable/toxic side effects with ALL preferred products

Tazarotene 0.1% foam

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year; Psoriasis: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Tazarotene 0.1% foam PA Criteria:

See Fabior® foam prior authorization criteria; **AND**

- Clinically valid reason why Fabior® cannot be used.

Tazarotene 0.1% gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form
PA Duration	Approval length dependent on indication Acne: 1 year; Verruca plana: 2 months; Actinic keratosis, keratosis follicularis, skin cancer: 1 year
Notes	Will not be covered for patients > 21 years old with a diagnosis of acne

Tazarotene 0.1% gel PA Criteria:

- Patient is < 21 years old; **AND**
  - Diagnosis of acne; **AND**
    - Trial and failure, contraindication, or intolerance to TWO preferred topical retinoids; **OR**
- Diagnosis of psoriasis; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred topical retinoids; **AND**
  - Trial and failure, contraindication, or intolerance to a preferred topical antipsoriatic agent

Topical Steroids: Least Potent

Hydrocortisone 0.5% cream & ointment (Rx), 1% cream, gel, lotion, & ointment (Rx), 2.5% cream, lotion & ointment

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Hydrocortisone 2.5% solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Hydrocortisone 2.5% solution PA Criteria:

Standard NP Criteria

Topical Steroids: Mild

Desonide 0.05% cream & ointment, fluocinolone 0.01% cream, oil, & solution

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Topical Steroids: Lower Mid-Strength

Betamethasone dipropionate 0.05% lotion, betamethasone valerate 0.1% cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Clocortolone 0.1% cream & pump, desonide 0.05% lotion, hydrocortisone butyrate 0.1% cream, lotion, ointment, & solution, hydrocortisone valerate 0.2% cream, Pandel® 0.1% cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Clocortolone 0.1% cream & pump, desonide 0.05% lotion, hydrocortisone butyrate 0.1% cream, lotion, ointment, & solution, hydrocortisone valerate 0.2% cream, Pandel® 0.1% cream PA Criteria:
Standard NP Criteria

Topical Steroids: Mid-Strength

Triamcinolone acetonide 0.1% cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Hydrocortisone valerate 0.2% ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Hydrocortisone valerate 0.2% ointment PA Criteria:
Standard NP Criteria

Topical Steroids: Upper Mid-Strength

Betamethasone valerate 0.1% cream, lotion, & ointment, fluticasone propionate 0.005% ointment, triamcinolone acetonide 0.025% cream, lotion, & ointment, triamcinolone acetonide 0.05% ointment, triamcinolone acetonide 0.1% lotion & ointment, triamcinolone acetonide 0.5% cream & ointment

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Amcinonide 0.1% cream,

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Amcinonide 0.1% cream PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Amcinonide 0.1% ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Amcinonide 0.1% ointment PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Betamethasone dipropionate 0.05% cream, betamethasone valerate 0.12% foam, desoximetasone 0.05% cream, fluocinonide 0.05% emulsified base cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Betamethasone dipropionate 0.05% cream, betamethasone valerate 0.12% foam, desoximetasone 0.05% cream, fluocinonide 0.05% emulsified base cream PA Criteria:
Standard NP Criteria

Topical Steroids: Potent

Betamethasone dipropionate, augmented 0.05% cream, fluocinonide 0.05% cream, ointment, & solution

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

ApexiCon E® 0.05% cream, betamethasone dipropionate, augmented 0.05% lotion, betamethasone dipropionate 0.05% ointment, desoximetasone 0.05% gel & ointment, desoximetasone 0.25% cream, ointment, & spray, diflorasone diacetate 0.05% cream &

ointment, fluocinonide 0.05% gel, halcinonide 0.1% cream, Topicort® 0.05% gel & ointment, Topicort®0.25% gel & ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

ApexiCon E® 0.05% cream, betamethasone dipropionate, augmented 0.05% lotion, betamethasone dipropionate 0.05% ointment, desoximetasone 0.05% gel & ointment, desoximetasone 0.25% cream, ointment, & spray, diflorasone diacetate 0.05% cream & ointment, fluocinonide 0.05% gel, halcinonide 0.1% cream, Topicort® 0.05% gel & ointment, Topicort®0.25% gel & ointment PA Criteria:
Standard NP Criteria

Halcinonide 0.1% solution

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Halcinonide 0.1% solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Halog® 0.1% cream & ointment

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Halog® 0.1% cream & ointment PA Criteria:
Standard NP Criteria

Halog® solution

PDL Status	NP
Quantity Limit	120 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Halog® solution PA Criteria:
Standard NP Criteria

Topical Steroids: Super Potent

Clobetasol 0.025% cream

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Clobetasol 0.025% cream PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Clobetasol propionate 0.05% cream, gel, lotion, ointment, & solution, clobetasol propionate emollient base 0.05% cream, fluocinonide 0.1% cream

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Betamethasone dipropionate, augmented 0.05% gel & ointment, clobetasol propionate 0.05% foam, shampoo, & spray, clobetasol propionate emollient base 0.05% foam, halobetasol propionate 0.05% cream & ointment, Ultravate® 0.05% lotion

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See below
PA Form	General PA Form

Clobetasol propionate 0.05% cream, gel, lotion, ointment, & solution, clobetasol propionate emollient base 0.05% cream, fluocinonide 0.1% cream, betamethasone dipropionate, augmented 0.05% gel & ointment, clobetasol propionate 0.05% foam, shampoo, & spray, clobetasol propionate emollient base 0.05% foam, halobetasol propionate 0.05% cream, & ointment, Ultravate® 0.05% lotion PA Criteria:

Standard NP Criteria

Halobetasol propionate 0.05% foam

PDL Status	NP
Quantity Limit	100 g/Rx
PA Criteria	See below
PA Form	General PA Form

Halobetasol propionate 0.05% foam PA Criteria:

- Diagnosis of an FDA-approved indication; **AND**
- Prescriber must submit peer-reviewed medical literature documenting why the requested agent is the only agent that will adequately treat the listed diagnosis as opposed to less costly alternatives.

Lexette® foam

PDL Status	NP
Quantity Limit	100 g/Rx
PA Criteria	See below
PA Form	General PA Form

Lexette® foam PA Criteria:

- Diagnosis of an FDA-approved indication; **AND**
- Prescriber must submit peer-reviewed medical literature documenting why the requested agent is the only agent that will adequately treat the listed diagnosis as opposed to less costly alternatives; **AND**
- Clinically valid reason why generic equivalent cannot be used

Wound Care, Topical

Filsuvez®

PDL Status	NP
Quantity Limit	15 tubes/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year
Notes	New wounds untreated with Filsuvez or recurrent reopened wounds are subject to initial criteria. Unhealed wounds > 6 months should rule out squamous cell and basal cell carcinoma.

Filsuvez® Initial PA Criteria:

- One of the following:
  - Diagnosis of Dystrophic Epidermolysis bullosa (EB);
  - Diagnosis of Junctional Epidermolysis bullosa (EB); **AND**
- Prescriber attests the target wound(s) meet ALL of the following:
  - Wound is clean in appearance and does not appear to be infected
  - Wound is 10 cm^2 to 50 cm^2; **AND**
- Patient will continue standard treatments for EB such as appropriate wound management and avoiding skin trauma; **AND**
- Prescribed by or in consultation with a dermatologist or wound management

Filsuvez® Renewal PA Criteria:

- Patient has a clinical response to therapy (e.g., decreased wound size, decreased frequency of wound dressing changes, reduction in pain)

VI. DIABETIC SUPPLIES

Blood Glucose Meters and Test Strips (OTC)

For agents in this class requiring PA, please submit the Diabetic Supply PA Form on faxed PA submissions.

Abbott Products

FreeStyle Meters: Freedom Lite, Lite, Precision Neo, Precision Xtra

PDL Status	P
Quantity Limit	1 meter/365 days
PA Criteria	None

FreeStyle Test Strips: Insulinx, Lite, Precision Neo, Precision Xtra

PDL Status	P
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	None

All other Abbott diabetic supplies:

PDL Status	P
Quantity Limit	Meters: 1 meter/365 days Test strips: Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	None

AgaMatrix Products

Various meters

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various AgaMatrix glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

Various test strips

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various AgaMatrix test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

Bayer Products

Bayer meters: Breeze 2, Contour

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Bayer glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

Bayer test strips: Breeze, Contour

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Bayer test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

All other Bayer diabetic supplies:

PDL Status	P
Quantity Limit	Meters: 1 meter/365 days Test strips: Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below

Various Bayer diabetic supplies PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

Home Diagnostics Products

Home Diagnostics: Various meters

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Home Diagnostics glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

Home Diagnostics: Various test strips

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Home Diagnostics test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

Johnson & Johnson Products

Johnson & Johnson meters: OneTouch Meters: UltraMini, Ping, Ultra2, UltraLink, UltraSmart

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Johnson & Johnson glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

Johnson & Johnson test strips

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Johnson & Johnson test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

All other Johnson & Johnson diabetic supplies:

PDL Status	P
Quantity Limit	Meters: 1 meter/365 days Test strips: Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below

Various Johnson & Johnson diabetic supplies PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

LifeScan Products

LifeScan: Various meters

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various LifeScan glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

LifeScan: Various test strips

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various LifeScan test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

Roche Products

Roche meters: Accu-Chek Aviva, Compact Plus

PDL Status	NP
Quantity Limit	1 meter/365 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Roche glucose meters PA Criteria

- Non-preferred meters will be approved for patients meeting ONE of the following criteria:
  - Patient is using an insulin pump that does not adequately communicate with a preferred meter
  - Patient requires a special meter due to visual impairment

Roche test strips

PDL Status	NP
Quantity Limit	Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Various Roche test strips PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

All other Roche diabetic supplies:

PDL Status	P
Quantity Limit	Meters: 1 meter/365 days Test strips: Age ≤ 5 years old: 306 strips/30 days; Age > 6 years old: 204 strips/30 days
PA Criteria	See below

Various Johnson & Johnson diabetic supplies PA Criteria

- Will be approved for individuals who meet prior authorization criteria and receive a prior authorization for the matching non-preferred glucometer

Continuous Glucose Monitors and Supplies

Dexcom®

Dexcom G6®, Dexcom G7®

PDL Status	P
Quantity Limit	Dexcom G6 Sensor: 3/30 days; Dexcom G6 Transmitter: 1/90 days; Dexcom G6 Receiver: 1/year Dexcom G7 10-day sensor: 3/30 days; Dexcom G7 15-day sensor: 2/30 days; Dexcom G7 Receiver: 1 year
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Dexcom G6®, Dexcom G7® Initial PA Criteria

- One of the following:
  - Patient is pregnant and has Type 1 Diabetes Mellitus (DM), Type 2 DM, or Gestational Diabetes**; OR**
  - Patient has a diagnosis of Type 1 DM Or Type 2 DM; **AND**
    - Patient is on insulin therapy or an insulin pump; **OR**
    - Patient is currently on glucose lowering therapy other than insulin therapy (e.g., SGLT2, metformin, SU, TZD, DPP-4) and meets ONE of the following:
      - History problematic hypoglycemia (e.g., frequent/severe or nocturnal hypoglycemia, hypoglycemia unawareness)
      - Dawn phenomenon with fasting blood sugars frequently exceeding 200 mg/dL
      - History of emergency room visit, or hospitalization related to ketoacidosis or hypoglycemia
      - Prescriber must submit medical records (e.g., chart notes) providing clinical rationale why the patient cannot use or continue to use traditional blood glucose monitoring devices (e.g. fingerstick); (Note: Patient convenience is NOT an approvable reason); **AND**
- Prescriber attests to BOTH the following:
  - Patient or caregiver will receive diabetes self-management education; **AND**
  - Patient or caregiver has received sufficient training using the requested CGM

Dexcom G6®, Dexcom G7® Renewal PA Criteria:

- Patient has received benefit from therapy which may include ONE or more of the following:
  - Decreased A1C from baseline
  - Decreased hypoglycemia episodes
  - Decreased percentage of time below therapeutic range (TBR)
  - Increased percentage of time in therapeutic range (TTR)

Senseonics® and Ascensia® Diabetes Care

Eversense® E3, E365, Mis

PDL Status	NP
Quantity Limit	Eversense E3: Sensor: 2/year; Transmitter: 1/year; Eversense E365: Sensor: 1/year; Transmitter: 1/year; Eversense Mis: Sensor: 1/90 days; Transmitter: 1/90 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Eversense® E3, E365, Mis PA Criteria

See Dexcom prior authorization criteria; **AND**

- Prescriber must submit clinical documentation (e.g., medical records) why the patient cannot use CGM Dexcom®

Abbott®

FreeStyle Libre 2®, FreeStyle Libre 3®

PDL Status	NP
Quantity Limit	Readers: 1/year; Sensors: 2/28 days; 2 kits/30 days
PA Criteria	See below
PA Form	Diabetic Supply PA Form

FreeStyle Libre 2®, FreeStyle Libre 3® PA Criteria

See Dexcom prior authorization criteria; **AND**

- Prescriber must submit clinical documentation (e.g., medical records) why the patient cannot use CGM Dexcom®

Medtronic®

Guardian 3®, Guardian 4®, Guardian Connect®, Guardian® CGM supplies

PDL Status	NP
Quantity Limit	Transmitter: 1/year; Sensors: 5/30 days; Charger: 1/year; Test plug: 1/year
PA Criteria	See below
PA Form	Diabetic Supply PA Form

Guardian 3®, Guardian 4®, Guardian Connect®, Guardian® CGM supplies PA Criteria

See Dexcom prior authorization criteria; **AND**

- Prescriber must submit clinical documentation (e.g., medical records) why the patient cannot use CGM Dexcom®

Insulin Management Systems

Omnipod 5®, Omnipod Dash®

PDL Status	P
Quantity Limit	Pods: 10/30 days; Device: 1/year
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Omnipod 5®, Omnipod Dash® Initial PA Criteria:

- Diagnosis of type 1 or type 2 diabetes mellitus; **AND**
- Submission of medical records (e.g., chart notes) documenting ONE of the following:
  - Patient is < 21 years of age and requires intensive insulin therapy and blood glucose monitoring (e.g.,3 or more injections, 3 or more blood glucose readings per day or utilizes a CGM)
  - Patient is ≥ 21 years of age and has difficulty maintaining stable blood glucose levels (hyperglycemia or hypoglycemia) or individual A1C/Time in Range (TIR) goals despite intensive insulin therapy and blood glucose monitoring (3 or more injections, 3 or more blood glucose readings per day, or utilizes a CGM)
  - Patient has physical impairments resulting in difficulty with self-injection of insulin (e.g., retinopathy, neuropathy)
  - Prescriber has provided clinical rationale for the use of disposable insulin delivery devices vs traditional insulin pump; **AND**
- Prescriber attests that the patient or caregiver will receive diabetes self-management education; **AND**
- Prescriber attests that the patient or caregiver has received sufficient training using the requested product

Omnipod 5®, Omnipod Dash® Renewal PA Criteria:

- Documentation of a positive clinical response (e.g. decrease HbA1C from baseline, decrease hypoglycemia episodes, decrease fasting and mealtime blood glucose levels)

Cequr Simplicity®

PDL Status	P
Quantity Limit	3-day patch: 10/30 days; 4-day patch: 8/32 days 7 day 1 unit patch: 4/28 days 7 day 2 unit patch: 5/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Cequr Simplicity® Initial PA Criteria:

- Diagnosis of type 1 or type 2 diabetes mellitus; **AND**
- Patient is ≥ 21 years old; **AND**
- Submission of medical records (e.g., chart notes) documenting ONE of the following:
  - Patient has difficulty maintaining stable blood glucose levels (hyperglycemia or hypoglycemia) or individual A1C/Time in Range (TIR) goals despite intensive insulin therapy and blood glucose monitoring (3 or more injections, 3 or more blood glucose readings per day, or utilizes a CGM)
  - Patient has physical impairments resulting in difficulty with self-injection of insulin (e.g., retinopathy, neuropathy)
  - Prescriber has provided clinical rationale for the use of disposable insulin delivery devices vs traditional insulin pump; **AND**
- Prescriber attests that the patient or caregiver will receive diabetes self-management education; **AND**
- Prescriber attests that the patient or caregiver has received sufficient training using the requested product

Cequr Simplicity® Renewal Criteria:

- Documentation of a positive clinical response (e.g. decrease HbA1C from baseline, decrease hypoglycemia episodes, decrease fasting and mealtime blood glucose levels)

iLet®, InPen®, Twiist®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

iLet®, InPen®, Twiist® PA Criteria:

- Prescriber must submit clinical documentation (e.g., medical records) why the patient cannot use the preferred products

V-Go® products

PDL Status	NP
Quantity Limit	30 patches/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

V-Go® products PA Criteria:

- Prescriber must submit clinical documentation (e.g., medical records) why the patient cannot use the preferred products

VII. ENDOCRINE AND METABOLIC

Adrenocorticotropins

Acthar® gel

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Acthar® PA Criteria:

- Appropriate FDA-approved diagnosis (e.g., diuresis in nephrotic syndrome, treatment of SLE or polymyositis, or acute MS exacerbation) for use **AND** has a contraindication, or intolerance to oral and injectable glucocorticoids; **OR**
- Diagnosis of infantile spasms

Cortrophin® gel

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Cortrophin® gel PA Criteria:

See Acthar® gel prior authorization criteria; **AND**

- Clinically valid reason why Acthar® gel cannot be used

Agents for Dyspareunia

Intrarosa® suppositories

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	This product is excluded from coverage in patients 21 years of age and older. Not a Covered Benefit.

Intrarosa® suppositories PA Criteria:

- Female younger than 21 years of age; **AND**
- Cessation of menses due to menopause; **AND**
- Painful intercourse

Osphena® tablets

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	This product is excluded from coverage in patients 21 years of age and older. Not a Covered Benefit.

Osphena® tablets PA Criteria:

See Intrarosa® prior authorization criteria

Agents for Gout

Allopurinol 200 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Allopurinol 200 mg tablets PA Criteria:
Standard NP Criteria

Colchicine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Colchicine capsules PA Criteria:

- Clinically valid reason why the preferred colchicine product cannot be used

Gloperba® solution

PDL Status	NP
Quantity Limit	300 mL/28 days
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	3 months

Gloperba® PA Criteria: (normal)

- Clinically valid reason why colchicine tablets or capsules cannot be used; **OR**
- Patient is unable to swallow sold dosage forms

Mitigare®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Mitigare® PA Criteria:

- Clinically valid reason why the preferred colchicine product cannot be used

Uloric®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Uloric® PA Criteria:

- Clinically valid reason as to why the preferred febuxostat cannot be used

Androgens

Gender dysphoria diagnosis is subject to PA review.

AndroGel® pump

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

AndroGel® pump Initial PA Criteria:

- Diagnosis of primary or hypogonadotropic hypogonadism; **AND**
- Documentation of at least 2 morning (before 10 am) total testosterone baseline levels done on separate days within the past 6 months that are considered low per laboratory testing

AndroGel® pump Renewal PA Criteria:

- Documentation of low or normal fasting testosterone level from previous 12 months

Danazol capsules

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Danazol capsules PA Criteria:

- Patient is male; **OR**
- Patient is female; **AND**
  - Danazol will be utilized for one of the following FDA approved indications:
    - HAE prophylaxis
    - Endometriosis; **AND**
  - Prescriber attests patient is not using for gender affirming care.

Testim® 1% gel (50 mg packet)

PDL Status	P
Quantity Limit	2 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testim® 1% gel PA Criteria:

See AndroGel® pump prior authorization criteria

Testosterone cypionate injection

PDL Status	P
Quantity Limit	4 mL/30 days
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone cypionate injectable PA Criteria:

See AndroGel® pump prior authorization criteria

Testosterone 1% gel (25 mg packet), testosterone 1.62% gel pump

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone 1% gel (25 mg packet), testosterone 1.62% gel packets PA Criteria:
See AndroGel® pump prior authorization criteria

Testosterone enanthate injection

PDL Status	P
Quantity Limit	4 mL/30 days
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone enanthate injection Initial PA Criteria:

- Diagnosis of primary or hypogonadotropic hypogonadism; **AND**
  - Documentation of at least 2 morning (before 10am) total testosterone baseline levels done on separate days within the past 6 months that are considered low per laboratory testing; **OR**
- Diagnosis of delayed puberty; **OR**
- Palliative treatment of androgen-responsive, advanced, inoperable, metastatic breast cancer in women who are 1-5 years postmenopausal and in premenopausal women who have benefited from oophorectomy

Testosterone enanthate injection Renewal PA Criteria:

- Documentation of low or normal fasting testosterone level from previous 12 months

Testosterone transdermal solution

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone transdermal solution PA Criteria:
See AndroGel® pump prior authorization criteria

Depo-Testosterone® injection

PDL Status	NP
Quantity Limit	4 mL/30 days
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Depo-Testosterone® injection PA Criteria:
See AndroGel® packets prior authorization criteria

Jatenzo® capsules

PDL Status	NP
Quantity Limit	2 caps/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Jatenzo® capsules PA Criteria:
See AndroGel® packets prior authorization criteria

Methitest® tablets

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Methitest® tablets Initial PA Criteria:

- One of the following:
  - Diagnosis of primary or hypogonadotropic hypogonadism; **AND**
    - Documentation of at least 2 morning (before 10am) total testosterone baseline levels done on separate days within the past 6 months that are considered low per laboratory testing; **OR**
  - Diagnosis of delayed puberty; **AND**
    - Clinically valid reason why preferred testosterone enanthate IM injection cannot be used; **OR**
  - Palliative treatment of androgen-responsive, advanced, inoperable, metastatic breast cancer in women who are 1-5 years postmenopausal and in premenopausal women who have benefited from oophorectomy

Methitest® tablets Renewal PA Criteria:

- Documentation of low or normal fasting testosterone level from previous 12 months

Methyltestosterone tablets

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Methyltestosterone tablets PA Criteria:
See Methitest® tablets prior authorization criteria

Natesto® nasal gel

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Natesto® nasal gel PA Criteria:
See AndroGel® packets prior authorization criteria

Testosterone 1% gel (50 mg packet), testosterone 1.62% gel packets

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone 1% gel (50 mg packet), testosterone 1.62% gel packets PA Criteria:
See AndroGel® pump prior authorization criteria; **AND**

- Trial and failure of TWO preferred products

Testosterone 2% gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Testosterone 2% gel PA Criteria:
See AndroGel® packets prior authorization criteria

Tlando® capsules

PDL Status	NP
Quantity Limit	2 capsules/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Tlando® capsules PA Criteria:
See AndroGel® packets prior authorization criteria

Undecatrex® capsules

PDL Status	NP
Quantity Limit	4 capsules/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Undecatrex® capsules PA Criteria:
See AndroGel® packets prior authorization criteria

Vogelxo® gel

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Vogelxo® gel PA Criteria:
See AndroGel® packets prior authorization criteria

Xyosted® injection

PDL Status	NP
Quantity Limit	2 mL/30 days
PA Criteria	See Below
PA Form	General PA Form
Notes	Requests for diagnosis of gender dysphoria will be reviewed on a case-by-case basis for determination.

Xyosted® injection PA Criteria:
See AndroGel® pump prior authorization criteria

Antidiuretics/Vasopressins

DDVAP tablets, DDAVP injection, desmopressin nasal spray

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

DDVAP tablets, DDAVP injection, desmopressin nasal spray PA Criteria:
Standard NP Criteria

Bone: Bisphosphonates

Alendronate tablets

PDL Status	P
Quantity Limit	5, 10, 40 mg: 1/day; 35, 70 mg: 4/28 days
PA Criteria	None

Alendronate solution

PDL Status	P
Quantity Limit	10 mL/day
PA Criteria	None

Atelvia® tablets

PDL Status	P
Quantity Limit	4/28 days
PA Criteria	None

Ibandronate tablets

PDL Status	P
Quantity Limit	1/28 days
PA Criteria	None

Actonel® tabs

PDL Status	NP
Quantity Limit	5, 30 mg: 1/day; 35 mg: 4/28 days; 150 mg: 1/28 days
PA Criteria	See Below
PA Form	General PA Form

Actonel® PA Criteria:

Standard NP Criteria

Binosto® effervescent tablets

PDL Status	NP
Quantity Limit	4/28 days
PA Criteria	See Below
PA Form	General PA Form

Binosto® effervescent tabletes PA Criteria:
Standard NP Criteria

Fosamax® tablets

PDL Status	NP
Quantity Limit	5, 10, 40 mg: 1/day; 35, 70 mg: 4/28 days
PA Criteria	See Below
PA Form	General PA Form

Fosamax® tablets PA Criteria:
Standard NP Criteria

Fosamax Plus D® tablets

PDL Status	NP
Quantity Limit	4/28 days
PA Criteria	See Below
PA Form	General PA Form

Fosamax Plus D® tablets PA Criteria:
Standard NP Criteria

Risedronate tablets

PDL Status	NP
Quantity Limit	5 & 30 mg: 1/day; 35 mg: 4/28 days; 150 mg: 1/28 days
PA Criteria	See Below
PA Form	General PA Form

Risedronate tablets PA Criteria:
Standard NP Criteria

Bone: Calcitonin

Calcitonin nasal spray

PDL Status	P
Quantity Limit	3.7 mL/30 days
PA Criteria	See Below
PA Form	General PA Form

Calcitonin Nasal Spray PA Criteria:

- Diagnosis of osteoporosis in postmenopausal women greater than five years post menopause; **AND**
- Trial and failure, contraindication, or intolerance to **BOTH** bisphosphonates **AND** raloxifene

Calcitonin injection

PDL Status	NP
Quantity Limit	1 mL/day
PA Criteria	See Below
PA Form	General PA Form

Calcitonin injection PA Criteria:

- Diagnosis of Paget's disease of the bone; **AND**
  - Trial and failure, contraindication, or intolerance to bisphosphonates; **OR**
- Treatment of hypercalcemia; **OR**
- Diagnosis of osteoporosis in postmenopausal women greater than five years post-menopause; **AND**
  - Trial and failure, contraindication, or intolerance to BOTH bisphosphonates **AND** raloxifene; **AND**
  - Trial and failure, contraindication, or intolerance to the preferred agent

Miacalcin® injection

PDL Status	NP
Quantity Limit	1 mL/day
PA Criteria	See Below
PA Form	General PA Form

Miacalcin® PA Criteria:

See calcitonin injection prior authorization criteria

Bone: SERMs

Raloxifene

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Evista®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Evista® PA Criteria:

Standard NP Criteria.

Bone: Parathyroid Hormone

Forteo® Injection

PDL Status	P
Quantity Limit	1 pen/28 days
PA Criteria	See Below
PA Form	General PA Form

Forteo® Injection Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Postmenopausal osteoporosis and patient is female; **OR**
  - Osteoporosis and patient is male; **OR**
  - Glucocorticoid-induced osteoporosis; **AND**
    - History of prednisone or its equivalent at a dose ≥ 5 mg/day; **AND**
- Patient meets ONE of the following:
  - Has a high risk for fracture (e.g., T-score of ≤ -3.0 even in the absence of fractures, T-score of -2.5 or below plus a fragility fracture, severe or multiple fractures, very high fracture probability by FRAX [e.g., major osteoporosis fracture > 30%, hip fracture > 4.5%]); **OR**
  - Unable to tolerate bisphosphonates or has a relative contraindication to oral bisphosphonates (achalasia, scleroderma esophagus, esophageal strictures); **OR**
  - Had an adequate trial of other osteoporosis therapies and had an insufficient response (fracture and/or loss of bone mineral density in spite of adherence to therapy); **AND**
- Total lifetime length of therapy with PTH analogs has not exceeded 2 years

Forteo® Injection Renewal PA Criteria:

- The prescriber attests that the patient remains at, or has returned to, having a high risk for fracture despite 24 months of cumulative use of parathyroid hormone analogs; **AND**
- The prescriber attests that the risk versus benefit of use greater than 24 months of cumulative use of parathyroid hormone analogs has been reviewed with the patient

Bonsity® Injection

PDL Status	NP
Quantity Limit	1 pen/28 days
PA Criteria	See Below
PA Form	General PA Form

Bonsity® Injection Initial PA Criteria:

See Forteo® prior authorization criteria; **AND**

- Clinically valid reason why preferred Forteo® cannot be used

Teriparatide Injection

PDL Status	NP
Quantity Limit	1 pen/28 days
PA Criteria	See Below
PA Form	General PA Form

Teriparatide Injection PA Criteria:

See Forteo® prior authorization criteria; **AND**

- Clinically valid reason why preferred Forteo® cannot be used

Tymlos® Injection

PDL Status	NP
Quantity Limit	1/28 days
PA Criteria	See Below.
PA Form	General PA Form

Tymlos® Injection Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Postmenopausal osteoporosis and patient is female
  - Osteoporosis and patient is male; **AND**
- Patient meets ONE of the following:
  - Has a high risk for fracture (e.g., T-score of ≤ -3.0 even in the absence of fractures, T-score of -2.5 or below plus a fragility fracture, severe or multiple fractures, very high fracture probability by FRAX [e.g., major osteoporosis fracture > 30%, hip fracture > 4.5%]); **OR**
  - Unable to tolerate bisphosphonates or has a relative contraindication to oral bisphosphonates (achalasia, scleroderma esophagus, esophageal strictures); **OR**
  - Had an adequate trial of other osteoporosis therapies and had an insufficient response (fracture and/or loss of bone mineral density in spite of adherence to therapy); **AND**
- Total lifetime length of therapy with PTH analogs has not exceeded 2 years

Tymlos® Injection Renewal PA Criteria:

- The prescriber attests that the patient remains at, or has returned to, having a high risk for fracture despite 24 months of cumulative use of parathyroid hormone analogs; **AND**
- The prescriber attests that the risk versus benefit of use greater than 24 months of cumulative use of parathyroid hormone analogs has been reviewed with the patient

Yorvipath® Injection

PDL Status	NP
Quantity Limit	2 pens/28 days
PA Criteria	See Below.
PA Form	General PA Form

Yorvipath® Injection Initial PA Criteria:

- Patient is 18 years or age or older; **AND**
- Diagnosis of hypoparathyroidism; **AND**
- Submission of medical records (e.g. chart notes or labs) demonstrating hypocalcemia in the setting of low serum parathyroid hormone (PTH) levels; **AND**
- Agent is prescribed by, or in consultations with, an endocrinologist

Yorvipath® Injection Renewal PA Criteria:

- Submission of medical records (e.g. chart notes) documenting a positive clinical response to therapy (e.g. albumin-corrected serum calcium in normal range, patient is independent from conventional therapy defined as no active vitamin D or ≤600 mg/day of calcium supplementation; **AND**
- Prescriber attest patient's serum calcium levels will be periodically monitored

Contraceptives, Non-Oral

Depo IM Provera®, Depo SubQ Provera®, medroxyprogesterone acetate injection

PDL Status	P
Quantity Limit	1 vial/ 90 days
PA Criteria	None

Etonogestrel-ethinyl estradiol vaginal ring

PDL Status	P
Quantity Limit	1/28 days
PA Criteria	None

Xulane®

PDL Status	P
Quantity Limit	3/28 days
PA Criteria	None

Annovera®

PDL Status	NP
Quantity Limit	1/year
PA Criteria	See Below
PA Form	General PA Form

Annovera® PA Criteria:

- Patient has tried and failed or had a contraindication to two preferred non-oral contraceptives; **AND**
- Clinically valid reason as to why preferred Etonogestrel-ethinyl estradiol vaginal ring cannot be used

Eluryng®, EnilloRing®, Haloette®

PDL Status	NP
Quantity Limit	1/28 days
PA Criteria	See Below
PA Form	General PA Form

Eluryng®, EnilloRing®, Haloette® PA Criteria:
Standard NP Criteria

Norelegestromin-ethinyl estradiol patch

PDL Status	NP
Quantity Limit	3/28 days
PA Criteria	See Below
PA Form	General PA Form

Norelegestromin-ethinyl estradiol patch PA Criteria:
Standard NP Criteria

Phexxi®

PDL Status	NP
Quantity Limit	12/month
PA Criteria	See Below
PA Form	General PA Form

Phexxi® PA Criteria:

- One of the following:
  - Patient has tried and failed or had a contraindication to two preferred non-oral contraceptives; **OR**
  - Patient has a contraindication or has experienced a clinically significant adverse event to hormonal agents; **AND**
- Provider attests that patient has been counseled regarding the higher rate of pregnancy prevention with the use of other contraception methods (e.g., injection, oral contraception, transdermal patch, vaginal ring) compared to Phexxi

Twirla®

PDL Status	NP
Quantity Limit	3/28 days
PA Criteria	See Below
PA Form	General PA Form

Twirla® PA Criteria:

- Trial and failure, or contraindication/intolerance of two preferred non-oral contraceptives; **AND**
- Avoid concomitant use of hepatitis C drug combinations containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir

Zafemy®

PDL Status	NP
Quantity Limit	3/28 days
PA Criteria	See Below
PA Form	General PA Form

Zafemy® PA Criteria:
Standard NP Criteria

Contraceptives, Oral

Please refer to OTC Drug Formulary for coverage of OTC products.

Preferred Oral Contraceptives on PDL

PDL Status	P
Quantity Limit	1.33 tablets/day
PA Criteria	None

Preferred Emergency Contraceptives on PDL

PDL Status	P
Quantity Limit	1/21 days
PA Criteria	None

Non-Preferred Contraceptives on PDL

PDL Status	NP
Quantity Limit	1.33 tablets/day
PA Criteria	See Below

Non-Preferred Contraceptives PA Criteria:
Standard NP Criteria

Corticosteroids, Oral

Alkindi Sprinkles®

PDL Status	NP
Quantity Limit	0.5, 1, 2 mg: 3/day; 5 mg: 4/day
PA Criteria	See Below.
PA Form	General PA Form

Alkindi Sprinkles® PA Criteria:

- Diagnosis of adrenocortical insufficiency; **AND**
- Patient is 18 years of age or younger

Eohilia®

PDL Status	NP
Quantity Limit	180 packs/ 365 days
PA Criteria	See Below.
PA Form	General PA Form
PA Approval Duration	3 months

Eohilia® PA Criteria:

- Patient is 11 years of age or older; **AND**
- Diagnosis of Eosinophilic esophagitis (EoE); **AND**
- Prescribed by, or in consultation with, a gastroenterologist, allergist, or immunologist; **AND**
- Prescriber attest patent meets both of the following:
  - Esophageal biopsy consists of ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf) following treatment course of a proton pump inhibitor; **OR**
  - Patient has symptoms of esophageal dysfunction (e.g., feeding difficulties, vomiting, pain, dysphagia); **AND**
- Trial and failure, or contraindication, to swallowed inhaled corticosteroids such as budesonide or fluticasone

Hemady®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below.
PA Form	General PA Form

Hemady® PA Criteria:

- Patient must be 18 years of age or older; **AND**
- Patient must have a diagnosis of Multiple Myeloma; **AND**
- Must be used in combination with other anti-myeloma agents; **AND**
- Patient does not have systemic fungal infection

Hydrocortisone sodium succinate injection

PDL Status	NP
PA Criteria	See Below.
PA Form	General PA Form

Hydrocortisone sodium succinate injection PA Criteria:

- Clinically valid reason why brand Solu-Cortef® injection cannot be used

Khindivi®

PDL Status	NP
PA Criteria	See Below.
PA Form	General PA Form

Khindivi® PA Criteria:

- Patient unable to swallow solid oral dosage forms

Prednisolone ODT

PDL Status	NP
PA Criteria	See Below.
PA Form	General PA Form

Prednisolone ODT PA Criteria:

- Unable to swallow; **OR**
- Unable to absorb medications through the GI tract

Prednisone DR tablets

PDL Status	NP
Quantity Limit	1 mg: 3/day; 2 mg: 2/day; 5 mg:12/day
PA Criteria	See Below.
PA Form	General PA Form

Prednisone DR tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Rayos® DR tablets

PDL Status	NP
Quantity Limit	1 mg: 3/day; 2 mg: 2/day; 5 mg:12/day
PA Criteria	See Below.
PA Form	General PA Form

Rayos® DR tablets PA Criteria:

Standard NP Criteria

Diabetes: Alpha-Glucosidase Inhibitors

Miglitol

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below

Miglitol PA Criteria:

- Trial and failure, or contraindication, or intolerance of the preferred acarbose

Diabetes: Amylin Analogs

Symlin Pen®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Symlin Pen® PA Criteria:

- Diagnosis of Type 1 or 2 diabetes; **AND**
- On insulin therapy; **AND**
- Failure to achieve adequate glycemic control (HbA1c ≤6.5%); **AND**
- Patients meeting any of the following will NOT be approved:
  - Recurrent, severe hypoglycemia requiring assistance during the past 6 months
  - Confirmed diagnosis of gastroparesis
  - Requiring the use of drugs that stimulate gastrointestinal motility

Diabetes: Biguanides

Metformin IR 500 mg, 850 mg, & 1000 mg tablets

PDL Status	P
Quantity Limit	500 mg: 4/day; 850 & 1000 mg: 2/day
PA Criteria	None

Metformin ER 500 mg, 750 mg tablets

PDL Status	P
Quantity Limit	500 mg: 4/day; 750 mg: 2/day
PA Criteria	None

Metformin ER Osmotic 500 mg, 1000 mg tablets

PDL Status	P
Quantity Limit	500 mg tablets: 4/day; 1000 mg tablets: 2/day
PA Criteria	None

Metformin IR 625 mg, 750 mg tablets

PDL Status	NP
Quantity Limit	625 mg: 3/day; 750 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Metformin IR Osmotic PA Criteria:

- Clinically valid reason why preferred metformin agents cannot be used

Metformin ER Modified Release 500 mg, 1000 mg tablets

PDL Status	NP
Quantity Limit	500 mg: 4/day 1000 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Metformin ER Modified Release PA Criteria:

Standard NP Criteria

Metformin Solution

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See Below
PA Form	General PA Form

Metformin Solution PA Criteria:

- Patient is ≤ 11 years of age; **OR**
- Patient is unable to swallow solid dosage forms

Riomet®

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See Below
PA Form	General PA Form

Riomet® PA Criteria:
See metformin solution prior authorization criteria

Diabetes: DPP-4 Inhibitors and Combinations

Janumet® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Janumet XR® tablets

PDL Status	P
Quantity Limit	50/500 mg, 100/1000 mg: 1/day; 50/1000 mg: 2/day
PA Criteria	None

Januvia® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Jentadueto® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Jentadueto® XR tablets

PDL Status	P
Quantity Limit	2.5/1000 mg: 2/day; 5/1000 mg: 1/day
PA Criteria	None

Saxagliptin tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Tradjenta® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Alogliptin tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Alogliptin PA Criteria:
Standard NP Criteria

Alogliptin/metformin tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Alogliptin/metformin PA Criteria:
Standard NP Criteria

Alogliptin/pioglitazone tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Alogliptin/pioglitazone PA Criteria:

- Clinically valid reason why a DPP-4 inhibitor and pioglitazone cannot be taken separately

Brynovin® solution

PDL Status	NP
Quantity Limit	4 mL/day
PA Criteria	See Below
PA Form	General PA Form

Brynovin® solution PA Criteria:

- Clinically valid reason why Januvia® cannot be used; **AND**
- Patient is unable to swallow solid dosage forms

Saxagliptin/metformin tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Saxagliptin/metformin PA Criteria:
Standard NP Criteria

Sitagliptin tablets (equivalent to Januvia®)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Sitagliptin PA Criteria:

- Clinically valid reason why brand Januvia® cannot be used

Sitagliptin tablets (equivalent to Zituvio®)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Sitagliptin PA Criteria:
Standard NP Criteria

Sitagliptin/metformin tablets (equivalent to Janumet®), sitagliptin/metformin XR tablets (equivalent to Januvia XR®)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Sitagliptin/metformin tablets, sitagliptin/metformin XR tablets PA Criteria:

- Clinically valid reason why brand Janumet® cannot be used

Sitagliptin/metformin tablets (equivalent to Zituvimet®), sitagliptin/metformin ER tablets (equivalent to Zituvimet XR®)

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Sitagliptin/metformin tablets, sitagliptin/metformin ER tablets PA Criteria:

- Clinically valid reason why Janumet or Janumet XR cannot be used

Zituvimet® tablets

PDL Status	NP
Quantity Limit	50/500 mg, 100/1000 mg: 1/day 50/1000 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Zituvimet® PA Criteria:

- Clinically valid reason why Janumet or Janumet XR cannot be used

Zituvimet XR® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Zituvimet XR® PA Criteria:

- Clinically valid reason why Janumet or Janumet XR cannot be used

Zituvio® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Zituvio® PA Criteria:

- Clinically valid reason why Januvia® cannot be used

Diabetes: Glucagon Agents

Baqsimi®, Gvoke

PDL Status	P
Quantity Limit	2/Rx
PA Criteria	None

Zegalogue®

PDL Status	NP
Quantity Limit	2/Rx
PA Criteria	See Below
PA Form	General PA Form

Zegalogue® PA Criteria:
Standard NP Criteria

Diabetes: Insulins, Rapid-Acting

Afrezza®

PDL Status	NP
Quantity Limit	Cartridges: 4-unit: 3/day; 8-unit: 6/day; 12-unit:6/day; Combo package: 1 box/month
PA Criteria	See Below
PA Form	General PA Form

Afrezza® PA Criteria:

- Patient is not a current smoker and does not have a history of smoking in the past 6 months; **AND**
- Prescriber attests that baseline spirometry has been performed prior to therapy and will be performed after 6 months of therapy, and every year thereafter; **AND**
- Patient does not have a history of chronic lung disease (e.g., asthma, COPD); **AND**
- Patient has ONE of the following diagnoses:
  - Type 2 Diabetes
  - Type 1 Diabetes while concurrently taking a long-acting insulin; **AND**
- One of the following:
  - Recipient or caregiver has problems with manual dexterity which may result in dosing errors (i.e., Parkinson's Disease, rheumatoid arthritis in the finger/hand joints, multiple sclerosis, lupus, osteoarthritis, stroke); **OR**
  - Recipient or caregiver has poor eyesight such that dosing errors may occur

Humalog U-200 KwikPen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Humalog U-200 KwikPen® PA Criteria:

- Patient requires > 100 insulin units/day

Humalog TempoPen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Humalog TempoPen® PA Criteria:

- Clinically valid reason why a preferred insulin pen (e.g. KwikPen®, FlexPen®) cannot be used

Kirsty® pen and vial

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Kirsty® pen and vial PA Criteria:

Standard NP Criteria

Lyumjev® KwikPen and vial

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Lyumjev® KwikPen and vial PA Criteria:

- Trial and failure or intolerance of 2 preferred, rapid-acting insulin agents; **AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Lyumjev TempoPen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Lyumjev TempoPen® PA Criteria:

- Clinically valid reason why a preferred insulin pen (e.g. KwikPen®, FlexPen®) cannot be used

Merilog® Solostar® and vial

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Merilog® Solostar® and vial PA Criteria:

Standard NP Criteria

Diabetes: Insulins, Intermediate-Acting

Humulin N KwikPen®

PDL Status	P
PA Criteria	See Below
PA Form	General PA Form

Humulin N KwikPen® PA Criteria:

- Prescriber must provide valid clinical rationale as to why patient is unable to utilize preferred Novolin N FlexPen®

Diabetes: Insulins, Long-Acting

Basaglar KwikPen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Basaglar KwikPen® PA Criteria:

- Contraindication or intolerance to a preferred insulin glargine pen that is not expected with the requested agent

Basaglar Tempo Pen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Basaglar Tempo Pen® PA Criteria:

- Clinically valid reason why a preferred insulin pen cannot be used (e.g. Solostar®)

Insulin degludec FlexTouch, Tresiba FlexTouch®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Insulin degludec FlexTouch, Tresiba FlexTouch® PA Criteria:

Standard NP Criteria

Rezvoglar®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Rezvoglar® PA Criteria:

- Contraindication or intolerance to a preferred insulin glargine pen that is not expected with the requested agent

Semglee®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Semglee® PA Criteria:

- Contraindication or intolerance to a preferred insulin glargine pen that is not expected with the requested agent

Diabetes: Meglitinides

Nateglinide

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Repaglinide

PDL Status	P
Quantity Limit	0.5 mg, 1 mg: 4/day 2 mg: 8/day
PA Criteria	None

Diabetes: SGLT2 Inhibitors and Combinations

Dapaglifozin tablets, Glyxambi® tablets, Invokana® tablets, Jardiance® tablets, Xigduo® XR tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Invokamet® tablets, Synjardy® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Synjardy XR® tablets

PDL Status	P
Quantity Limit	25/1000 mg: 1/day All other strengths: 2/day
PA Criteria	None

Dapaglifozin/metformin ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Dapaglifozin/metformin ER PA Criteria:

- Clinically valid reason why Xigduo XR® cannot be used

Dapagliflozin/saxagliptin 10/5 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Dapagliflozin/saxagliptin10/5 mg tablet PA Criteria:

- Trial and failure, contraindication, or intolerance to separate components (dapagliflozin and saxagliptin); **AND**
- Clinically valid reason why brand Qtern® cannot be used

Farxiga® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Farxiga® PA Criteria:

- Clinically valid reason why generic dapaglifozin cannot be used

Inpefa® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Inpefa® PA Criteria:

- Trial and failure of TWO preferred combination SGLT2/metformin ER agents (Synjardy XR®, Xigduo® XR)

Invokamet XR® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Invokamet XR® PA Criteria:

- Trial and failure of TWO preferred combination SGLT2/metformin ER agents (Synjardy XR®, Xigduo® XR)

Qtern® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Qtern® PA Criteria:

- Trial and failure, contraindication, or intolerance to separate components (dapagliflozin and saxagliptin)

Steglatro® tablets

PDL Status	NP
Quantity Limit	5 mg: 2/day 15 mg: 1/day
PA Criteria	See Below
PA Form	General PA Form

Steglatro® PA Criteria:

- Trial and failure or intolerance to TWO preferred single-entity SGLT2 agents (Farxiga®, Invokana®, Jardiance®)

Segluromet® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Segluromet® PA Criteria:

- Trial and failure of TWO preferred combination SGLT2/metformin immediate release agents (Invokamet®, Synjardy®)

Steglujan® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Steglujan® PA Criteria:

- Clinically valid reason why a preferred single-entity SGLT2 agent and DPP-4 inhibitor cannot be used as separate agents

Trijardy XR® tablets

PDL Status	NP
Quantity Limit	10/5/1000 mg, 2.5/5/1000 mg: 1/day; 5/2.5/1000 mg, 12.5/2.5/1000 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Trijardy XR® PA Criteria:

- Clinically valid reason as to why patient cannot use Glyxambi® and metformin ER as separate agents

Diabetes: Sulfonylureas

Glipizide 15 mg tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Glipizide 15 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Diabetes: Thiazolidinediones and Combinations

Pioglitazone tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Pioglitazone/metformin tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Actos® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Actos® PA Criteria:

Standard NP Criteria

ActoPlus Met® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

ActoPlus Met® PA Criteria:

Standard NP Criteria

Duetact® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Duetact® PA Criteria:

- Clinically valid reason why the patient cannot use pioglitazone and glimepiride as separate agents

Glimepiride 3 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Glimepiride 3 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Pioglitazone/glimepiride tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Pioglitazone/glimepiride PA Criteria:

- Clinically valid reason why the patient cannot use pioglitazone and glimepiride as separate agents

Estrogens, Oral

Gender dysphoria diagnosis is subject to PA review.

Conjugated estrogens tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Conjugated estrogens tablets PA Criteria:

- Clinically valid reason why brand Premarin® tablets cannot be used

Depo-Estradiol® injection, Delestrogen® injection, Estrace® tablets, estradiol cypionate injection, estradiol valerate injection, Menest® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Depo-Estradiol® injection, Delestrogen® injection, Estrace® tablets, estradiol cypionate injection, estradiol valerate injection, Menest® tablets PA Criteria:

Standard NP Criteria

Estrogens, Transdermal

Gender dysphoria diagnosis is subject to PA review.

Climara®, Dotti®, Lyllana®

PDL Status	P
Quantity Limit	4/28 days
PA Criteria	None

Estradiol biweekly patch

PDL Status	P
Quantity Limit	8/28 days
PA Criteria	None

Estradiol weekly patch

PDL Status	P
Quantity Limit	4/28 days
PA Criteria	None

Minivelle®, Vivelle Dot®

PDL Status	P
Quantity Limit	8/28 days
PA Criteria	None

Alora®

PDL Status	NP
Quantity Limit	8/28 days
PA Criteria	See Below

Alora® PA Criteria:
Standard NP Criteria

Divigel®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below

Divigel® PA Criteria:
Standard NP Criteria

Elestrin®

PDL Status	NP
Quantity Limit	1/28 days
PA Criteria	See Below

Elestrin® PA Criteria:
Standard NP Criteria

Estradiol gel

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below

Estradiol gel PA Criteria:
Standard NP Criteria

Menostar®

PDL Status	NP
Quantity Limit	4/28 days
PA Criteria	See Below

Menostar® PA Criteria:
Standard NP Criteria

Estrogens, Vaginal

Gender dysphoria diagnosis is subject to PA review.

Premarin® cream

PDL Status	P
Quantity Limit	2 grams/day
PA Criteria	None

Estrace®, Estradiol cream

PDL Status	NP
Quantity Limit	42.5 grams/Rx
PA Criteria	See Below

Estrace®, estradiol cream PA Criteria:
Standard NP Criteria

Estrogen/Progestins, Oral

Gender dysphoria diagnosis is subject to PA review.

Premphase®, Prempro®

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Estrogen/Progestins, Transdermal

Gender dysphoria diagnosis is subject to PA review.

CombiPatch®

PDL Status	P
Quantity Limit	8/28 days
PA Criteria	None

Climara Pro®

PDL Status	NP
Quantity Limit	4/28 days
PA Criteria	See Below

Climara Pro® PA Criteria:
Standard NP Criteria

Estrogen/SERM Agents

Gender dysphoria diagnosis is subject to PA review.

Duavee®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Duavee® PA Criteria:

- Patient has an intact uterus with a diagnosis of moderate to severe vasomotor symptoms associated with menopause; **OR**
- Patient has an intact uterus with a diagnosis of post-menopausal osteoporosis

GLP-1 Agonists

Please submit faxed PA requests using the **GLP-1 Agonist PA Form**.

Mounjaro® injection

PDL Status	P
Quantity Limit	2 mL/28 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Type 2 Diabetes:

Mounjaro® PA Criteria:

- Submission of medical records (e.g. chart notes) confirming diagnosis of type 2 diabetes as evidenced by ONE of the following laboratory values:
  - A1C greater than or equal to 6.5%
  - Fasting plasma glucose (FPG) greater than or equal to 126 mg/dL
  - 2-hour plasma glucose (PG) greater than or equal to 200 mg/dL during oral glucose tolerance test
  - Random plasma glucose greater than or equal to 200 mg/dL in patient with classic symptoms of hyperglycemia or hyperglycemic crisis**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patients does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)

Mounjaro® injection Renewal PA Criteria:

- Submission of recent medical records (e.g., chart notes and/or labs) documenting a positive clinical response to therapy (e.g. reduction in HbA1C, HbA1C goal achieved, improvement in fast blood glucose levels, improvement in ASCVD)**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)

Ozempic® injection

PDL Status	P
Quantity Limit	4 injections/month
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form
Notes	A1C level can be from early stages in treatment. If original HbA1c is unknown, or current HbA1c is controlled due to another current diabetic regimen, please include current regimen and current A1C.

Type 2 Diabetes:

Ozempic® injection Initial PA Criteria:

- Submission of medical records (e.g. chart notes) confirming diagnosis of type 2 diabetes as evidenced by ONE of the following laboratory values:
  - A1C greater than or equal to 6.5%
  - Fasting plasma glucose (FPG) greater than or equal to 126 mg/dL
  - 2-hour plasma glucose (PG) greater than or equal to 200 mg/dL during oral glucose tolerance test
  - Random plasma glucose greater than or equal to 200 mg/dL in patient with classic symptoms of hyperglycemia or hyperglycemic crisis**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® tabs [formerly Rybelsus®], Zepbound®)**; AND**
- Patients does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)

Ozempic® injection Renewal PA Criteria:

- Submission of recent medical records (e.g., chart notes and/or labs) documenting a positive clinical response to therapy (e.g. reduction in HbA1C, HbA1C goal achieved, improvement in fast blood glucose levels, improvement in ASCVD)**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® tabs [formerly Rybelsus®], Zepbound®)

Soliqua® injection

PDL Status	P
Quantity Limit	5 pens/ 30 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Soliqua® injection PA Criteria:

See Ozempic® injection prior authorization criteria; **AND**

- Patient is currently taking a long-acting insulin documented per TennCare paid claims

Wegovy® injection

PDL Status	P
Quantity Limit	4 injectors/28 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

CV Risk Reduction due to prior MI, prior stroke, or PD:

Wegovy® injection Initial PA Criteria:

- Treatment is being requested to reduce the risk of major adverse cardiovascular events**; AND**
- Patient is 18 years of age or older**; AND**
- Submitted medical documentation (e.g. chart notes) of initial BMI ≥ 27 kg/m^2**; AND**
- Submitted medical documentation (e.g. chart notes) of ONE of the following:
  - Prior myocardial infarction
  - Prior stroke (ischemic and hemorrhagic stroke)
  - Symptomatic peripheral arterial disease as evidenced by intermittent claudication with ankle–brachial index < 0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease**; AND**
- Patient is on optimized guideline-directed therapy including beta-blockers, RAS inhibitors, and lipid lowering agents to ensure reduced cardiovascular risk unless contraindicated (supported by medical documentation or claims history)**; AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g. dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management)**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)

Wegovy® injection Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., patient has not had a major cardiovascular event within the past 12 months, decreased body weight or waist circumference from baseline, decrease blood pressure, total cholesterol, LDL, or triglyceride levels from baseline)**; AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g. dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management)**; AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not being co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)**; AND**
- Patient is on optimized guideline-directed therapy including beta-blockers, RAS inhibitors, and lipid lowering agents to ensure reduced cardiovascular risk unless contraindicated (supported by medical documentation or claims history)

- Noncirrhotic nonalcoholic steatohepatitis (NASH) or Metabolic dysfunction associated steatohepatitis (MASH)**:**

Wegovy® injection Initial PA Criteria:

- Patient is greater than or equal to 18 years of age; **AND**
- Diagnosis of NASH or MASH; **AND**
- Submission of medical records (e.g. chart notes) confirming disease is fibrosis stage F2 or F3 as confirmed by ONE of the following:
  - FibroScan
  - Fibrosis-4 index (FIB-4) greater than 1.3
  - ELF score
  - Magnetic Resonance Elastography (MRE)
  - Liver Biopsy**; AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g. dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management); **AND**
- Prescribed by or in consultation with a gastroenterologist or hepatologist; **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patients does not have personal or immediate family history of medullary thyroid carcinoma or MEN2

Wegovy® injection Renewal PA Criteria:

- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Submission of recent medical records (e.g., chart notes and/or labs) documenting a positive clinical response to therapy (e.g. NASH/MASH disease symptoms)

Zepbound® single dose prefilled pen

PDL Status	P
Quantity Limit	4 injectors/28 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Obstructive Sleep Apnea:

Zepbound® single dose prefilled pen Initial PA Criteria:

- Patient is 18 years of age or older**; AND**
- Diagnosis of moderate to severe Obstructive Sleep Apnea (OSA)**; AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out**; AND**
- Submission of medical records (e.g. chart notes) documenting BOTH of the following:
  - Initial BMI ≥ 30 kg/m^2**; AND**
  - ≥ 15 respiratory events per hour of sleep confirmed by sleep study**; AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Trial and failure (minimum duration 3 months with documented compliance) of Continuous Positive Airway Pressure (CPAP) or BiPAP device, unless contraindicated**; AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not being co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)

Zepbound® single dose prefilled pen Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., decrease number of apneas and hypopneas during sleep from baseline); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not being co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)

Exenatide injection

PDL Status	NP
Quantity Limit	5 mcg: 1.2 mL/30 days 10 mcg: 2.4 mL/30 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form
Notes	A1C level can be from early stages in treatment. If original HbA1c is unknown, or current HbA1c is controlled due to another current diabetic regimen, please include current regimen and current A1C.

Type 2 Diabetes:

Exenatide injection Initial PA Criteria:

- Submission of medical records (e.g. chart notes) confirming diagnosis of type 2 diabetes as evidenced by ONE of the following laboratory values:
  - A1C greater than or equal to 6.5%
  - Fasting plasma glucose (FPG) greater than or equal to 126 mg/dL
  - 2-hour plasma glucose (PG) greater than or equal to 200 mg/dL during oral glucose tolerance test
  - Random plasma glucose greater than or equal to 200 mg/dL in patient with classic symptoms of hyperglycemia or hyperglycemic crisis; **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2); **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred agents

Exenatide injection Renewal PA Criteria:

- Submission of recent medical records (e.g., chart notes and/or labs) documenting a positive clinical response to therapy (e.g. reduction in HbA1C, HbA1C goal achieved, improvement in fast blood glucose levels, improvement in ASCVD); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)

Liraglutide injection (generic for Victoza®)

PDL Status	NP
Quantity Limit	3 pens/ 30 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Liraglutide injection (generic for Victoza®) PA Criteria:

See Ozempic® injection prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to 2 preferred GLP-1 agonist agents

Ozempic® tablets

PDL Status	NP
Quantity Limit	1 tab/day
PA Criteria	See below
PA Form	GLP-1 Agonist PA Form

Ozempic® tablets Initial PA Criteria:

- Submission of medical records (e.g. chart notes) confirming diagnosis of type 2 diabetes as evidenced by ONE of the following laboratory values:
  - A1C greater than or equal to 6.5%
  - Fasting plasma glucose (FPG) greater than or equal to 126 mg/dL
  - 2-hour plasma glucose (PG) greater than or equal to 200 mg/dL during oral glucose tolerance test
  - Random plasma glucose greater than or equal to 200 mg/dL in patient with classic symptoms of hyperglycemia or hyperglycemic crisis; **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, Zepbound®); **AND**
- Patients does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2); **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred agents;

Ozempic® tablets Renewal PA Criteria:

- Submission of recent medical records (e.g., chart notes and/or labs) documenting a positive clinical response to therapy (e.g. reduction in HbA1C, HbA1C goal achieved, improvement in fast blood glucose levels, improvement in ASCVD); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, Zepbound®);

Rybelsus® tablets

PDL Status	NP
Quantity Limit	1 tablet/day
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Rybelsus® tablets PA Criteria:

See Ozempic® tablets prior authorization criteria

Trulicity® injection

PDL Status	NP
Quantity Limit	2 mL/28 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Trulicity® PA Criteria:

See exenatide prior authorization criteria

Victoza® injection

PDL Status	NP
Quantity Limit	3 pens/30 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form
Notes	A1C level can be from early stages in treatment. If original HbA1c is unknown, or current HbA1c is controlled due to another current diabetic regimen, please include current regimen and current A1C.

Victoza® PA Criteria:

See Ozempic® injection prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to 2 preferred GLP-1 agonist agents

Wegovy® tablets

PDL Status	NP
Quantity Limit	1 tablet/day
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form
Notes	Patients not meeting the renewal criteria may be approved for up to one month to allow for titration off medication.

CV Risk Reduction due to prior MI, prior stroke, or PAD

Wegovy® tablets Initial PA Criteria:

- Treatment is being requested to reduce the risk of major adverse cardiovascular events; **AND**
- Patient must be 18 years of age or older meet labeled age minimum; **AND**
- Submitted medical documentation (e.g. chart notes) of initial BMI ≥ 27 kg/m^2; **AND**
- Submitted medical documentation (e.g. chart notes) of ONE of the following:
  - Prior myocardial infarction
  - Prior stroke (ischemic and hemorrhagic stroke)
  - Symptomatic peripheral arterial disease as evidenced by intermittent claudication with ankle–brachial index < 0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease; **AND**
- Patient is on optimized guideline-directed therapy including beta-blockers, RAS inhibitors, and lipid lowering agents to ensure reduced cardiovascular risk unless contraindicated (supported by medical documentation or claims history); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g. dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2); **AND**
- Submission of medical records (i.e. chart notes) providing a clinically valid reason why patient can't use Wegovy injectable (NOTE: Patient convenience, mild injection site reactions, or needle phobia is NOT an approvable reason. Patient must have contraindication to the excipients in the injectable Wegovy product.)

Wegovy® tablets Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., patient has not had a major cardiovascular event within the past 12 months, decreased body weight or waist circumference from baseline, decrease blood pressure, total cholesterol, LDL, or triglyceride levels from baseline); **AND**
- One of the following:
  - Patient is currently on the 25mg once daily maintenance dose; **OR**
  - Submission of medical records (i.e. chart notes) demonstrating a positive clinical response (e.g. decrease in body weight, blood pressure, heart rate, etc.) with no adverse effects (e.g. nausea, diarrhea, constipation, hypoglycemia, acute pancreatitis/gallbladder disease, etc.) at a lower maintenance dose; **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g. dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (e.g., liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient is on optimized guideline-directed therapy including beta-blockers, RAS inhibitors, and lipid lowering agents to ensure reduced cardiovascular risk unless contraindicated (supported by medical documentation or claims history)

Xultophy® injection

PDL Status	NP
Quantity Limit	5 pens/30 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Xultophy® PA Criteria:

See Ozempic® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to 2 preferred agents; **AND**
- Patient is currently taking, but inadequately controlled on, a long-acting insulin documented per TennCare paid claims

Zepbound KwikPen®

PDL Status	NP
Quantity Limit	4 injectors/28 days
PA Criteria	See Below
PA Form	GLP-1 Agonist PA Form

Obstructive Sleep Apnea:

Zepbound KwikPen® Initial PA Criteria:

- Patient is 18 years of age or older**; AND**
- Diagnosis of moderate to severe Obstructive Sleep Apnea (OSA)**; AND**
- Hypersomnolence secondary to another sleep disorder, neurologic disorder, medical condition, or by medicine or substance use has been ruled out**; AND**
- Submission of medical records (e.g. chart notes) documenting BOTH of the following:
  - Initial BMI ≥ 30 kg/m^2**; AND**
  - ≥ 15 respiratory events per hour of sleep confirmed by sleep study**; AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Trial and failure (minimum duration 3 months with documented compliance) of Continuous Positive Airway Pressure (CPAP) or BiPAP device, unless contraindicated**; AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not being co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2); **AND**
- Provider must submit medical documentation demonstrating a clinically valid reason why the patient is unable to use the preferred injectable formulation. Submission of medical records (e.g. chart notes) required.

Zepbound KwikPen® Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., decrease number of apneas and hypopneas during sleep from baseline); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Will not be used concomitantly with DPP-4 inhibitors; **AND**
- Will not being co-administered with another injectable or oral GLP-1 receptor agonists (e.g., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®)

GnRH/LHRH Agonists

Fensolvi®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Fensolvi® PA Criteria:

- Diagnosis of central precocious puberty in children (onset of secondary sexual development before 8 [girls] or 9 years of age [boys])

Lupron Ped-Depot®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Lupron Ped-Depot® PA Criteria:

- Diagnosis of central precocious puberty in children (onset of secondary sexual development before 8 [girls] or 9 years of age [boys])

Growth Hormone Agents

For agents in this class requiring PA, please submit the Growth Hormone PA Form on all faxed PA submissions.

Genotropin®

PDL Status	P
PA Criteria	See Below
PA Form	Growth Hormone PA Form
Notes	Growth hormone therapy will NOT be approved for idiopathic short stature (ISS)

Genotropin® PA Criteria:

- Agent is prescribed by, or in consultation with, an endocrinologist; **AND**
- Requested dose is within the approved dosage range for requested indication per clinical compendium; **AND**
- Dosage form resulting in least wastage of product is being used; **AND**
- ONE of the following:
  - Diagnosis of growth hormone (GH) deficiency in a child or adolescent confirmed by ONE of the following:
    - Lab results of TWO GH stimulation tests are both below the age adjusted normal range (< 10 mcg/mL); **OR**
    - Lab results to ONE GH stimulation test are the age adjusted below normal range (< 10 mcg/mL) and IGF-1 or IGFBP-3 levels are below the age and gender adjusted normal range; **OR**
  - Diagnosis of adult growth hormone deficiency; **AND**
    - Patient had a diagnosis of childhood-onset GHD; **AND**
    - Lab results to ONE GH stimulation test are below the age adjusted normal range; **OR**
  - Diagnosis of GH deficiency due to hypothalamic-pituitary disease or a known structural, genetic, or traumatic cause (e.g., pituitary tumor, pituitary surgical damage, head trauma, cranial irradiation) and ONE of the following:
    - Lab results to ONE GH stimulation test are below the age adjusted normal range
    - IGF-1 levels are below the age adjusted normal range normal for patient's age
    - The patient has deficiencies in 3 or more pituitary axes; **OR**
  - Patient is a newborn infant and has evidence of hypoglycemia **AND** either a low GH level (<20 ng/mL) or a low for age IGF-1/IGFBP-3 level; **OR**
  - Diagnosis of Small for Gestational Age (SGA) or fetal growth restriction (FGR) and patient is > 2 years old with a height at least 2 standard deviations below the population mean for age; **OR**
  - Diagnosis of short stature associated with Turner Syndrome or Noonan Syndrome; **OR**
  - Diagnosis of short stature associated with short stature homeobox (SHOX) gene deficiency; **OR**
  - Diagnosis of Prader-Willi Syndrome; **OR**
  - Patient has chronic kidney disease (CKD)

Egrifta SV®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Egrifta SV® PA Criteria:

- Diagnosis of Acquired Immunodeficiency Syndrome (AIDs) or Human Immunodeficiency Virus (HIV); **AND**
- Prescribed by, or in consultation with, an endocrinologist or provider with expertise in HIV; **AND**
- Waist circumference greater than or equal to 95 cm for males, or greater than or equal to 94 cm for females; **AND**
- Waist to hip ratio greater than or equal to 0.94 for males, or greater than or equal to 0.88 for females

Egrifta WR®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Egrifta WR® PA Criteria:

- Diagnosis of Acquired Immunodeficiency Syndrome (AIDs) or Human Immunodeficiency Virus (HIV); **AND**
- Prescribed by, or in consultation with, an endocrinologist or provider with expertise in HIV; **AND**
- Waist circumference greater than or equal to 95 cm for males, or greater than or equal to 94 cm for females; **AND**
- Waist to hip ratio greater than or equal to 0.94 for males, or greater than or equal to 0.88 for females

Humatrope®, Norditropin®, Nutropin AQ®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Humatrope® PA Criteria:

See Genotropin® prior authorization criteria

Ngenla®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Ngenla® Initial PA Criteria:

- Patient meets the minimum age requirement per label and is less than 18 years of age; **AND**
- Patient weighs at least 11.5kg; **AND**
- Agent is prescribed by, or in consultation with, an endocrinologist; **AND**
- Diagnosis of growth failure due to growth hormone deficiency has been confirmed by ONE of the following:
  - Lab results of TWO GH stimulation tests are both below the age adjusted normal range (< 10 mcg/mL); **OR**
  - Lab results to ONE GH stimulation test are the age adjusted below normal range (< 10 mcg/mL) and IGF-1 or IGFBP-3 levels are below the age and gender adjusted normal range; **OR**
- Epiphyses has not closed; **AND**
- Clinically valid reason why preferred Genotropin cannot be used

Ngenla® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests that patient has had a height increase of at least 2.5 cm/year over the previous year

Omnitrope®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Omnitrope® PA Criteria:

See Genotropin® prior authorization criteria

Serostim®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Serostim® Initial PA Criteria:

- Diagnosis of HIV-associated wasting syndrome or cachexia; **AND**
- One of the following:
  - Unintentional weight loss of >10% over the last 12 months
  - Unintentional weight loss of > 7.5% over the last 6-months
  - Loss of 5% body cell mass (BCM) within 6 months
  - Body mass index (BMI) < 20 kg/m2; **AND**
  - Body cell mass (BCM) below 40% total body weight in males or 35% total body weight in females; **AND**
- Nutritional evaluation since onset of wasting first occurred; **AND**
- Patient has not had weight loss due to other underlying treatable conditions (e.g., depression, mycobacterium avium complex, chronic infectious diarrhea, malignancy); **AND**
- Anti-retroviral therapy has been optimized to decrease the viral load and will be continued throughout the course of treatment; **AND**
- Trial and failure of megestrol

Serostim® Renewal PA Criteria:

- Evidence of positive response to therapy (i.e., > 2% increase in body weight and/or BCM); **AND**
- A target goal has not been achieved (i.e., weight, BCM, BMI)

Skytrofa®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Skytrofa® Initial PA Criteria:

- ONE of the following:
  - Patient is at least 1 year old and is less than 18 years of age and ALL of the following:
    - Patient weighs at least 11.5kg; **AND**
    - Diagnosis of growth failure due to growth hormone deficiency has been confirmed by ONE of the following:
      - Lab results of TWO GH stimulation tests are both below the age adjusted normal range (< 10 mcg/mL);
      - Lab results to ONE GH stimulation test are the age adjusted below normal range (< 10 mcg/mL) and IGF-1 or IGFBP-3 levels are below the age and gender adjusted normal range; **OR**
  - Patient is 18 years of age and older AND one of the following:
    - Diagnosis of adult growth hormone deficiency AND both of the following:
      - Submission of clinical records supporting a diagnosis of childhood-onset GHD; **AND**
      - Lab results to ONE GH stimulation test are below the age adjusted normal range; **OR**
  - Diagnosis of hormone deficiency due to hypothalamic-pituitary disease or structural damage (e.g., pituitary tumor, pituitary surgical damage, cranial irradiation, head trauma, subarachnoid hemorrhage) and ONE of the following:
    - Lab results of ONE GH stimulation test are both below the age adjusted normal range
    - Growth hormone levels of a GH stimulation test are below the age adjusted normal range
    - IGF-1 level is below the age adjusted normal range
    - The patient has deficiencies in 3 or more pituitary axes; **AND**
- Agent is prescribed by, or in consultation with, an endocrinologist; **AND**
- Clinically valid reason why preferred Genotropin cannot be used; **AND**
- Dosage form resulting in least wastage of product is being used

Skytrofa® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Evidence of positive response to therapy (i.e., increase in ht, change in truck percent fat from baseline, and/or normalization of IGF-1 levels);

Sogroya®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Sogroya® PA Criteria:

- Clinically valid reason as to why the patient cannot take the preferred Genotropin; **AND**
- Prescribed by, or in consultation with, an endocrinologist; **AND**
- One of the following:
  - Diagnosis of growth failure due to growth hormone (GH) deficiency confirmed by ONE of the following:
    - Lab results of 2 GH stimulation tests are below the age adjusted normal range (< 10 mcg/mL); **OR**
    - Lab results to ONE GH stimulation test are below the age adjusted norma (< 10 mcg/mL) and IGF-1 or IGFBP-3 levels are below the age and gender adjusted normal range; **OR**
  - Diagnosis of adult growth hormone deficiency **AND** both of the following:
    - Submission of clinical records supporting a diagnosis of childhood-onset GHD; **AND**
    - Lab results to ONE GH stimulation test are below the age adjusted normal range; **OR**
  - Diagnosis of hormone deficiency due to hypothalamic-pituitary disease or structural damage (e.g., pituitary tumor, pituitary surgical damage, cranial irradiation, head trauma, subarachnoid hemorrhage) and ONE of the following:
    - Lab results of ONE GH stimulation test are both below the age adjusted normal range
    - Growth hormone levels of a GH stimulation test are below the age adjusted normal range
    - IGF-1 level is below the age adjusted normal range
    - The patient has deficiencies in 3 or more pituitary axes; **OR**
  - Diagnosis of short stature born small for gestational age AND with no catch-up growth by two years of age; **AND**
    - Patient is > 2 years old with a height at least 2 standard deviations below the population mean for age; **OR**
  - Diagnosis of Growth Failure associated with Noonan Syndrome (NS)

Voxzogo®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Initial Voxzogo® PA Criteria:

- Diagnosis of achondroplasia; **AND**
- Prescribed by, or in consultation with, an endocrinologist; **AND**
- Patient has open epiphyses; **AND**
- Patient will not have limb-lengthening surgery during treatment with Voxzogo®; **AND**
- Provider attests that patient/caregiver has been properly trained on preparation and administration of Voxzogo

Renewal Voxzogo® PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Provider attests that patient has an annualized growth velocity $\geq$ 1.5 cm/year

Yuviwel®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Yuviwel® Initial PA Criteria:

- Diagnosis of achondroplasia; **AND**
- Submission of medical records (e.g. chart notes, medical records) confirming ONE of the following:
  - Presence of the FGFR3 variant; **OR**
  - Symptoms AND radiological findings consistent with achondroplasia; **AND**
- Documentation submitted shows patient has open epiphyses; **AND**
- Patient is 2 years of age and less than 18 years of age; **AND**
- Prescribed by, or in consultation with a geneticist or endocrinologist; **AND**
- Patient will not have limb-lengthening surgery during treatment with Yuviwel; **AND**
- Patient will not use in combination with Voxzogo; **AND**
- Provider attests that the patient/caregiver has been properly trained on preparation and administration of Yuviwel

Yuviwel® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Provider attests that patient has an annualized growth velocity of ≥ 1.5 cm/year

Zomacton®

PDL Status	NP
PA Criteria	See Below
PA Form	Growth Hormone PA Form

Zomacton® PA Criteria:

See Genotropin® prior authorization criteria

Hematopoietic Agents

Retacrit®

PDL Status	P
PA Criteria	See Below
PA Form	General PA Form

Retacrit® PA Criteria:

- Documentation that the patient has been assessed for iron deficiency and has adequate iron stores (defined as TSAT ≥ 20% or serum ferritin ≥ 100 mcg/L within the prior 3 months) or is receiving iron therapy before starting erythropoiesis stimulating agent; **AND**
- Other causes of anemia (e.g., hemolysis, bleeding, vitamin deficiency) have been ruled out; **AND**
- One of the following:
  - Anemia secondary to myelosuppressive chemotherapy; **AND**
    - Patient is at least 5 years of age and receiving concurrent myelosuppressive chemotherapy; **AND**
    - Upon initiation, there is at least 2 additional months of planned chemotherapy; **AND**
    - Hemoglobin ≤ 10 g/dL; **OR**
  - Anemia secondary to zidovudine treatment in HIV-infected patient; **AND**
    - Zidovudine dose is ≤ 4,200 mg/week; **AND**
    - Serum erythropoietin (EPO) levels ≤ 500 mU/mL; **AND**
    - ONE of the following:
      - For initial therapy, hemoglobin ≤ 10 g/dL; **OR**
      - For continuation of therapy, hemoglobin ≤ 12 g/dL; **OR**
  - Anemia secondary to hepatitis C virus (HCV) in patients receiving ribavirin and interferon-alfa therapy; **AND**

        - For initial therapy, hemoglobin ≤ 10 g/dL; **OR**
        - For continuation of therapy, hemoglobin ≤ 12 g/dL; **OR**
    - Anemia secondary to myelodysplastic syndrome (MDS), myeloproliferative neoplasms (MPN) – Myelofibrosis, or multiple myeloma; **AND**
        - Serum erythropoietin (EPO) levels ≤ 500 mU/mL; **AND**
        - ONE of the following:
            - For initial therapy, hemoglobin ≤ 10 g/dL; **OR**
            - For continuation of therapy, hemoglobin ≤ 12 g/dL; **OR**
- Anemia of prematurity, in combination with iron supplementation; **OR**
- Anemia secondary to rheumatoid arthritis; **OR**
- Anemia secondary to chronic kidney disease (CKD); **AND**
    - For initial therapy, hemoglobin ≤ 10 g/dL; **OR**
    - For continuation of therapy, hemoglobin ≤ 12 g/dL; **OR**
- Reduction of allogeneic blood transfusions in elective noncardiac, nonvascular surgery; **AND**
    - Patient is NOT willing to donate autologous blood pre-operatively
    - Hemoglobin is > 10 g/dL and ≤ 13 g/dL

Aranesp®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Aranesp® PA Criteria:
See Retacrit® prior authorization criteria

Epogen®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Epogen® PA Criteria:
See Retacrit® prior authorization criteria

Mircera®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Mircera® PA Criteria:

- Documentation that the patient has been assessed for iron deficiency and has adequate iron stores (defined as TSAT > 20% or serum ferritin ≥ 100 mcg/L within the prior 3 months) or is receiving iron therapy before starting erythropoiesis stimulating agent; **AND**
- Other causes of anemia (e.g., hemolysis, bleeding, vitamin deficiency, etc.) have been ruled out; **AND**
- Diagnosis of anemia secondary to chronic kidney disease (CKD) and ONE of the following:
    - Patient is 18 years of age or older; **AND**
        - For initial therapy, hemoglobin is < 10 g/dL; **OR**
        - For continuation of therapy, hemoglobin ≤ 12 g/dL; **OR**
    - Patient is between 3 months and 17 years of age; **AND**
        - Patient's hemoglobin stabilized following administration of another erythropoietin stimulating agent (ESA)

Procrit®

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Procrit® PA Criteria:
See Retacrit® prior authorization criteria

Vafseo®

PDL Status	NP
Quantity Limit	100 mg, 450 mg: 1/day 300 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Vafseo® PA Criteria:

- Documentation that the patient has been assessed for iron deficiency and has adequate iron stores (defined as TSAT > 20% or serum ferritin ≥ 100 mcg/L within the prior 3 months) or is receiving iron therapy before starting erythropoiesis stimulating agent; **AND**
- Diagnosis of anemia due to chronic kidney disease (CKD); **AND**
- Patient has been receiving dialysis for ≥ 3 months; **AND**
- Hemoglobin level <10 g/dL; **AND**
- Clinically valid reason why erythropoiesis stimulating agents (ESAs) cannot be used

Hyperparathyroid Agents

Cinacalcet

PDL Status	P
PA Criteria	See Below
PA Form	General PA Form

Cinacalcet PA Criteria:

- Secondary Hyperparathyroidism due to Chronic Kidney Disease (CKD), **AND** patient must be on dialysis; **OR**
- Parathyroid Carcinoma resulting in hypercalcemia; **OR**
- Severe Hypercalcemia in patients with primary HPT who are unable to undergo parathyroidectomy

Doxercalciferol capsules

PDL Status	NP
Quantity Limit	0.5, 2.5 mcg: 1/day; 1 mcg: 3/day
PA Criteria	See Below
PA Form	General PA Form

Doxercalciferol capsules PA Criteria:

- Recipients experiencing (or with a history of) hypercalcemia and/or hyperphosphatemia with calcitriol use; **AND**
- Trial and failure, contraindication, or intolerance to cinacalcet

Paricalcitol capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Paricalcitol capsules PA Criteria:

- Recipients experiencing (or with a history of) hypercalcemia and/or hyperphosphatemia with calcitriol use; **AND**
- Trial and failure, contraindication, or intolerance to cinacalcet

Rayaldee®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Rayaldee® PA Criteria:

- Secondary Hyperparathyroidism due to Stage 3 or Stage 4 Chronic Kidney Disease (CKD); **AND**
- Serum total 25-hydroxyvitamin D levels less than 30 ng/mL; **AND**
- Trial and failure, contraindication, or intolerance of cinacalcet

Sensipar

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Sensipar PA Criteria:

- See cinacalcet prior authorization criteria; **AND**
- Clinically valid reason why the preferred cinacalcet agent cannot be used

Zemplar® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Zemplar® capsules PA Criteria:

See doxercalciferol capsules prior authorization criteria

NK3 Antagonists

Lynkuet®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Lynkuet® PA Criteria:

- Diagnosis of moderate to severe vasomotor symptoms due to menopause; **AND**
- Trial and failure, contraindication, or intolerance to TWO of the following:
  - Gabapentin
  - Menopausal hormone therapy (e.g., estrogen monotherapy or estrogen + progesterone)
  - Oxybutynin
  - SSRI (e.g., paroxetine, escitalopram, citalopram)
  - SNRI (e.g., venlafaxine and desvenlafaxine)

Veozah®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Veozah® PA Criteria:

- Diagnosis of moderate to severe vasomotor symptoms due to menopause; **AND**
- Trial and failure, contraindication, or intolerance to TWO of the following:
  - Gabapentin
  - Menopausal hormone therapy (e.g., estrogen monotherapy or estrogen + progesterone)
  - Oxybutynin
  - SSRI (e.g., paroxetine, escitalopram, citalopram)
  - SNRI (e.g., venlafaxine and desvenlafaxine)

Pituitary Agents

Cabergoline tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Cabergoline tablets PA Criteria:

- Diagnosis of hyperprolactinemic disorder

Progestins, Oral

Gender dysphoria diagnosis is subject to PA review.

Megestrol suspension

PDL Status	P
Quantity Limit	40 mg: 20 mL/day 625 mg: 5 mL/day
PA Criteria	None

Norethindrone acetate

PDL Status	P
PA Criteria	See Below
PA Form	General PA Form

Norethindrone acetate PA Criteria:

- Diagnosis of endometriosis

Thyroid

Adthyza® tablets, Niva Thyroid®, NP Thyroid®, Thyquidity® solution, thyroid tablets, Tirosint-SOL® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Adthyza® tablets, Niva Thyroid®, NP Thyroid®, Thyquidity® solution, thyroid tablets, Tirosint-SOL® solution PA Criteria:
Standard NP Criteria

Uterine Disorder GnRH Modulators

Myfembree®

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Myfembree® Initial PA Criteria:

- Patient age is ≥ 18 years; **AND**
- One of the following:
  - Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas/fibroids; **AND**
    - Trial and failure of one of the following:
      - Estrogen/progestin contraceptive (including tablets, vaginal ring, or patch)
      - Progestin-only contraceptive (including tablets, IM injection, IUD); **OR**
  - Diagnosis of moderate to severe pain associated with endometriosis; **AND**
    - Trial and failure, or contraindication, to an analgesic (e.g., ibuprofen, meloxicam, naproxen); **AND**
    - Trial and failure of one of the following:
      - Estrogen/progestin contraceptive (including tablets, vaginal ring, or patch)
      - Progestin-only contraceptive (including tablets, IM injection, IUD); **AND**
- Patient must be premenopausal; **AND**
- Prescribed by, or in consultation with, an OB/GYN or reproductive specialist; **AND**
- Total treatment duration should not exceed 24 months due to risk of continued bone loss

Myfembree® Renewal PA Criteria:

- Patient has positive response to therapy (e.g., reduction in pain and discomfort from baseline, sustained reduction in menstrual blood loss per cycle); **AND**
- Total treatment duration should not exceed 24 months

Oriahnn®

PDL Status	P
Quantity Limit	1 box/28 days
PA Criteria	See Below
PA Form	General PA Form

Orihann® PA Criteria:

See Myfembree® prior authorization criteria

Orilissa®

PDL Status	P
Quantity Limit	150 mg: 1/day; 200 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form

Orilissa® Initial PA Criteria:

- Patient age is ≥ 18 years; **AND**
- Diagnosis of moderate to severe pain associated with endometriosis; **AND**
  - Trial and failure, or contraindication, to an analgesic (e.g., ibuprofen, meloxicam, naproxen); **OR**
  - Trial and failure of one of the following:
    - Estrogen/progestin contraceptive (including tablets, vaginal ring, or patch)
    - Progestin-only contraceptive (including tablets, IM injection, IUD); **AND**
- Patient must be premenopausal; **AND**
- Prescribed by, or in consultation with, an OB/GYN or reproductive specialist; **AND**
- Total treatment duration should not exceed 24 months due to risk of continued bone loss

Orilissa® Renewal PA Criteria (only for the 150 mg strength):

- Patient has positive response to therapy (e.g., reduction in pain and discomfort from baseline); **AND**
- Total treatment duration should not exceed 24 months

Vasopressor Antagonists

Jynarque® tablets, Jynarque Pak® tablets, tolvaptan tablets & tablet packs (generic for Jynarque®)

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Jynarque® tablets, Jynarque Pak® tablets, tolvaptan tablets & tablet packs Initial PA Criteria:

- Patient has a diagnosis of autosomal dominant polycystic kidney disease (ADPKD); **AND**
- Patient is 18 years of age or older; **AND**
- Prescribed by, or in consultation with, a nephrologist; **AND**
- Prescriber and patient are enrolled in the Jynarque REMS program; **AND**
- If the request is for brand Jynarque®, clinically valid reason why generic equivalents cannot be used

Samsca® tablets, tolvaptan tablets (generic for Samsca®)

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Samsca® tablets PA Criteria:

- Diagnosis of hyponatremia; **AND**
- Medication was initiated in a hospital setting

VIII. GASTROINTESTINAL

5-ASA Derivatives, Oral

Mesalamine DR 400 mg capsules

PDL Status	P
Quantity Limit	6/day
PA Criteria	None

Mesalamine DR 1.2 g tablets

PDL Status	P
Quantity Limit	1.2 g: 4/day
PA Criteria	None

Mesalamine ER 24-hour capsules 0.375 g

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Pentasa® capsules

PDL Status	P
Quantity Limit	250 mg: 16/day; 500 mg: 8/day
PA Criteria	See below
PA Form	General PA Form

Sulfasalazine tablets, sulfasalazine EC tablets

PDL Status	P
Quantity Limit	8/day
PA Criteria	None

Azulfidine® tablets, Azulfidine® EN tablets

PDL Status	NP
Quantity Limit	8/day
PA Criteria	See below
PA Form	General PA Form

Azulfidine® tablets, Azulfidine® EN tablets PA Criteria:
Standard NP Criteria

Balsalazide capsules

PDL Status	NP
Quantity Limit	9/day
PA Criteria	See below
PA Form	General PA Form

Balsalazide capsules PA Criteria:
Standard NP Criteria

Dipentum® capsules, Lialda® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Dipentum® capsules, Lialda® tablets PA Criteria:
Standard NP Criteria

Mesalamine DR tablets

PDL Status	NP
Quantity Limit	800 mg: 6/day;
PA Criteria	See below
PA Form	General PA Form

Mesalamine DR tablets PA Criteria:
Standard NP Criteria

Mesalamine ER capsules (equivalent to Pentasa®)

PDL Status	NP
Quantity Limit	250 mg: 16/day; 500 mg: 8/day
PA Criteria	None

Mesalamine ER capsules PA Criteria:

- Clinically valid reason why brand Pentasa® cannot be used

5-ASA Derivatives, Rectal

Canasa® suppository, mesalamine kit, SfRowasa® enema, Rowasa® kit

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Canasa® suppository, mesalamine kit, SfRowasa® enema, Rowasa® kit PA Criteria:
Standard NP Criteria

Agents for Chronic Constipation

Linzess® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Lubiprostone capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Symproic® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Symproic® tablets PA Criteria:

- Age ≥ 18 years; **AND**
- One of the following:
  - Diagnosis of opioid-induced constipation with chronic non-cancer pain
  - Diagnosis of opioid-induced constipation with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation; **AND**
- Submission of medical records (e.g., chart notes, control substance monitoring database) or confirmed pharmacy claims documenting at ≥1 of opioid therapy within the past 90 days; **AND**
- Prescriber attests that the requested drug will be discontinued when opioid treatment is discontinued; **AND**
- Patient does not have known or suspected gastrointestinal obstruction

Amitiza® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Amitiza® capsules PA Criteria:

Standard NP Criteria

Motegrity® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Motegrity® tablets PA Criteria:

- Age ≥ 18 years; **AND**
- Patient has diagnosis of chronic idiopathic constipation (CIC); **AND**
- Patient does not have intestinal perforation or obstruction due to structural or functional disorder of the gut wall, obstructive ileus, or severe inflammatory conditions of the intestinal tract (e.g., Crohn's disease, ulcerative colitis); **AND**
- Trial and failure of, or contraindication, or intolerance to lubiprostone AND Linzess®; **AND**
- If request is for brand Motegrity, clinically valid reason why generic prucalopride cannot be used

Movantik® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Movantik® tablets PA Criteria:

- Age ≥ 18 years; **AND**
- One of the following:
  - Diagnosis of opioid-induced constipation with chronic non-cancer pain
  - Diagnosis of opioid-induced constipation with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation; **AND**
- Submission of medical records (e.g., chart notes, control substance monitoring database) or confirmed pharmacy claims documenting at ≥ 1 month of opioid therapy within the past 90 days; **AND**
- Prescriber attests that Movantik® will be discontinued when opioid treatment is discontinued; **AND**
- Trial and failure of, or contraindication, or intolerance to Symproic®

Prucalopride tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Prucalopride tablets PA Criteria:

See Motegrity® prior authorization criteria

Agents for Irritable Bowel Syndrome (IBS)

Alosetron tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Alosetron tablets Initial PA Criteria:

- Patient is female and ≥ 18 years of age; **AND**
- Diagnosis of severe, diarrhea-predominant, irritable bowel syndrome (IBS); **AND**
- Chronic IBS symptoms lasting 6 months or more; **AND**
- Provider has ruled out anatomic or biochemical abnormalities of the GI tract; **AND**
- Patient is not concomitantly using fluvoxamine; **AND**
- Patient does not have a history of the following conditions:
  - Chronic or severe constipation or sequalae from constipation
  - Intestinal obstruction, stricture, toxic megacolon, gastrointestinal perforation, and/or adhesions
  - Ischemic colitis, impaired intestinal circulation, thrombophlebitis, or hypercoagulable state
  - Crohn's disease or ulcerative colitis
  - Diverticulitis
  - Severe hepatic impairment

Alosetron tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has not experienced any treatment-restricting adverse effects (e.g., severe constipation); **AND**
- Positive response to therapy (e.g., decrease stool frequency, frequent bowel urgency, and abdominal pain)

Linzess® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Lubiprostone capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Amitiza® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Amitiza® capsules PA Criteria:
Standard NP Criteria

Ibsrela® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Ibsrela® tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of irritable bowel syndrome with constipation (IBS-C); **AND**
- Patient does not have known or suspected mechanical gastrointestinal obstruction; **AND**
- Trial and failure, contraindication, or intolerance to lubiprostone AND Linzess®

Ibsrela® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has not experienced any treatment-restricting adverse effects (e.g., severe diarrhea); **AND**
- Positive response to therapy (e.g., decrease stool frequency, frequent bowel urgency, and abdominal pain)

Lotronex® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Lotronex® tablets PA Criteria:
See alosetron prior authorization criteria; **AND**

- Clinically valid reason why the preferred generic alosetron cannot be used

Viberzi® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Viberzi® tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of severe, diarrhea-predominant, irritable bowel syndrome (IBS); **AND**
- Patient does not have history of the following:
  - Alcohol abuse/addiction or drink more than 3 alcoholic drinks per day
  - Pancreatitis or structural diseases of the pancreas
  - Severe hepatic impairment (Child Pugh Class-C)
  - Severe constipation
  - Absence of gallbladder
  - Biliary duct (gallbladder) obstruction or Sphincter of Oddi disease/dysfunction

Viberzi® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has not experienced any treatment-restricting adverse effects (e.g., severe diarrhea); **AND**
- Positive response to therapy (e.g., decrease stool frequency, frequent bowel urgency, and abdominal pain)

Antidiarrheals

Please refer to OTC Drug Formulary for coverage of OTC products.

Lomotil® tablets, loperamide capsules, Motofen® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lomotil® tablets, loperamide capsules, Motofen® tablets PA Criteria:

Standard NP Criteria

Mytesi®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Mytesi® PA Criteria:

- Patient has non-infectious diarrhea of at least one month duration; **AND**
- Patient has a diagnosis of HIV or AIDS; **AND**
- Patiently is currently receiving anti-retroviral therapy

Opium tincture

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Opium tincture PA Criteria:

Standard NP Criteria

Antiemetics: 5-HT3 Antagonists

Ondansetron ODT, tablets (except 16 mg ODT)

PDL Status	P
Quantity Limit	30/90 days
PA Criteria	See below
PA Form	General PA Form
Notes	Prior authorization is not required for quantities up to 30 tablets per 90 days. For requests that exceed the quantity limit, the criteria below must be met.

Ondansetron ODT, tablets (except 16 mg ODT) Quantity Limit Criteria:

- For requests that exceed the quantity limit, ONE of the following must be met:
  - Receiving highly or moderately emetogenic chemotherapy
  - Receiving radiation therapy
  - Treatment is for post-operative nausea and vomiting (PONV)
  - Nausea or vomiting associated with pregnancy and trial and failure of TWO conventional antiemetics (i.e., metoclopramide, prochlorperazine, dexamethasone, Diclegis®)

Anzemet® tablets

PDL Status	NP
Quantity Limit	2/30 day
PA Criteria	See below
PA Form	General PA Form

Anzemet® tablets PA Criteria:

- ONE of the following:
  - Receiving highly or moderately emetogenic chemotherapy
  - Receiving radiation therapy
  - Treated for post-operative nausea and vomiting (PONV); **AND**
- Trial and failure, contraindication, or intolerance to a preferred 5-HT3 antagonist

Granisetron tablets

PDL Status	NP
Quantity Limit	Tabs: 60/30 days; Injection: 2 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Granisetron PA Criteria:

See Anzemet® prior authorization criteria.

Ondansetron 16 mg ODT

PDL Status	NP
Quantity Limit	30/90 days
PA Criteria	See below
PA Form	General PA Form

Ondansetron 16 mg ODT PA Criteria:

- Clinically valid reason why the patient cannot use the preferred agents

Ondansetron solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Notes	Prior authorization is not required for patients < 6 years of age

Ondansetron solution PA Criteria:

- Patient < 6 years of age; **OR**
- The requested dose is not achievable with ondansetron ODT; **OR**
- Allergy or intolerance to inactive ingredient in ODT tab (e.g., dye, filler, excipient)

Sancuso® patch

PDL Status	NP
Quantity Limit	1/30 days
PA Criteria	See below
PA Form	General PA Form

Sancuso® patch PA Criteria:

See Anzemet® prior authorization criteria.

Antiemetics: Anticholinergics

Promethazine solution, tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	Promethazine PA Form
Notes	Prior authorization is not required for patients ≥ 2 years old

Promethazine solution, tablets PA Criteria:

- Patient is < 2 years of age; **AND**
- Prescriber documents medical necessity; **AND**
- Prescriber is aware of contraindication and agrees to accept risk

Antivert® chewables, tablets, Compro® suppositories

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Antivert® chewables, tablets, Compro® suppositories capsules PA Criteria:

Standard NP criteria

Promethazine suppositories

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	Promethazine PA Form

Promethazine suppositories PA Criteria:

- One of the following:
  - Patient is > 2 years old; **OR**
  - Patients < 2 years of age AND both of the following:
    - Prescriber documents medical necessity; **AND**
    - Prescriber is aware of contraindication and agrees to accept risk; **AND**
- Patient has tried and failed 2 preferred antiemetics from any of the following classes:
  - Antiemetics: 5-HT3 Antagonists
  - Antiemetics: Anticholinergics
  - Antiemetics: Miscellaneous
  - Antiemetics: NK-1 Antagonists

Scopolamine patches

PDL Status	NP
Quantity Limit	10 patches/30 days
PA Criteria	See below
PA Form	General PA Form

Scopolamine patches PA Criteria:

- One of the following:
  - Recipient has tried and failed, or is intolerant to TWO of the following agents: meclizine, promethazine, dimenhydrinate, diphenhydramine or metoclopramide
  - Unable to take oral medications
  - Therapy is needed for an extended period of time where taking short acting agents would not be feasible
  - Has a tracheotomy or is ventilator dependent

Tigan® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tigan® injection capsules PA Criteria:

Standard NP criteria

Transderm Scop® patches

PDL Status	NP
Quantity Limit	10 patches/30 days
PA Criteria	See below
PA Form	General PA Form

Transderm Scop® patches PA Criteria:

- One of the following:
  - Recipient has tried and failed, or is intolerant to TWO of the following agents: meclizine, promethazine, dimenhydrinate, diphenhydramine or metoclopramide
  - Unable to take oral medications
  - Therapy is needed for an extended period of time where taking short acting agents would not be feasible
  - Has a tracheotomy or is ventilator dependent; **AND**
- If request is for brand Transderm Scop® patches, clinically valid reason why generic scopolamine patches cannot be used

Antiemetics: Delta-9-THC Derivatives

Dronabinol capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dronabinol capsules PA Criteria:

- Request is for the treatment of severe nausea/vomiting associated with cancer chemotherapy for patients actively being treated for cancer; **AND**
  - Trial and failure, intolerance, intolerance, medical reason, or contraindication that prohibits taking aprepitant + 5-HT3 receptor antagonist + corticosteroid; **OR**
- Request is for the treatment of anorexia associated with weight loss in patients with AIDS**; AND**
  - Trial and failure, intolerance, or contraindication to megestrol acetate oral suspension

Dronabinol solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dronabinol solution PA Criteria:

See dronabinol capsules prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why brand Syndros cannot be used

Marinol® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Marinol® capsules PA Criteria:

See dronabinol prior authorization criteria.

Syndros® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Syndros® solution PA Criteria:

See dronabinol prior authorization criteria; **AND**

- Patient is unable to swallow solid dosage forms

Antiemetics: Miscellaneous

Diclegis®

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Bonjesta®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Bonjesta® PA Criteria:

- Patient has a diagnosis of pregnancy-induced nausea or vomiting; **AND**
- Patient has failed documented conservative measures (e.g., dietary changes, trigger avoidance); **AND**
- Clinically valid reason as to why preferred Diclegis® cannot be used

Doxylamine/pyridoxine

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Doxylamine/pyridoxine PA Criteria:

- Clinically valid reason as to why preferred Diclegis® cannot be used

Antiemetics: NK-1 Antagonists

Aprepitant capsules, capsule packs

PDL Status	P
Quantity Limit	40 mg: 1/30 days; 80 mg: 4/30 days: 125 mg: 2/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	See below
Notes	Will be approved for 3 days/treatment course. If chemotherapy is more frequent than weekly, may approve a quantity sufficient for THREE days beyond the chemotherapy duration.

Aprepitant capsules, capsule packs PA Criteria:

- ONE of the following:
  - Patient is receiving a highly emetogenic chemotherapy regimen; **OR**
  - Patient is receiving a moderately emetogenic chemotherapy regimen and has failed two other antiemetic regimens; **OR**
  - Prevention of Postoperative Nausea and Vomiting (PONV) with trial and failure or contraindication to a 5HT3-receptor antagonist (PA Duration: 1 month); **OR**
  - Patient has refractory nausea that would require hospitalization (PA Duration: 1 month)

Akynzeo® capsules

PDL Status	NP
Quantity Limit	2/30 days
PA Criteria	See below
PA Form	General PA Form

Akynzeo® capsules PA Criteria:

- ONE of the following:
  - Patient is receiving a highly emetogenic chemotherapy regimen
  - Patient is receiving a moderately emetogenic chemotherapy regimen and has failed other previous antiemetic regimens; **AND**
- Trial and failure, contraindication, or intolerance to aprepitant

Emend® capsules, capsule packs

PDL Status	NP
Quantity Limit	80 mg: 4/30 days; Tri-Pack: 2 packs/30 days
PA Criteria	See below
PA Form	General PA Form

Emend® capsule, capsule packs PA Criteria:

See aprepitant prior authorization criteria; **AND**

- Clinically valid reason why preferred aprepitant cannot be used

Emend® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Emend® suspension PA Criteria:
See aprepitant prior authorization criteria; **AND**

- Patient is 6 months to 11 years old; **OR**
- Patient is unable to swallow solid dosage forms

Antispasmodics/Anticholinergics

Glycopyrrolate solution

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Glycopyrrolate solution PA Criteria:

- Patients unable to swallow tablets; **OR**
- Patient is < 8 years of age

Chlordiazepoxide/clidinium capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Chlordiazepoxide/clidinium capsules PA Criteria:
Standard NP Criteria

Dicyclomine 40 mg tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Dicyclomine 40 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Cuvposa® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Cuvposa® solution PA Criteria:

- Patients unable to swallow tablets; **OR**
- Patient is < 8 years of age

Glycate® tablets, glycopyrrolate 1.5 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Glycate® tablets, glycopyrrolate 1.5 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Levsin® SL tablets, tablets, methscopolamine tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Glycate® tablets, Levsin® SL tablets, tablets, methscopolamine tablets PA Criteria:
Standard NP Criteria

Combination Products for H. pylori

Pylera® capsules, Talicia® capsules

PDL Status	P
Quantity Limit	1 box/Rx; 2 courses of therapy/year
PA Criteria	See below
PA Form	General PA Form
Notes	See Appropriate Diagnosis PA bypass list who need treatment for *H. pylori.*

Pylera® capsules, Talicia® capsules PA Criteria:

- Documentation of recent positive *H. pylori* test

Bismuth subcitrate/metronidazole/tetracycline capsules

PDL Status	P
Quantity Limit	1 box/Rx; 2 courses of therapy/year
PA Criteria	See below
PA Form	General PA Form

Bismuth subcitrate/metronidazole/tetracycline capsules PA Criteria:

- Documentation of recent positive *H. pylori* test; **AND**
- Trial and failure, contraindication, or intolerance to a preferred combination agent; **OR**
- Trial and failure, contraindication, or intolerance to H. pylori treatment prescribed as separate agents (e.g., amoxicillin + clarithromycin + omeprazole or bismuth subsalicylate + metronidazole + tetracycline)

Lansoprazole/amoxicillin/clarithromycin combination pack

PDL Status	P
Quantity Limit	1 box/Rx; 2 courses of therapy/year
PA Criteria	See below
PA Form	General PA Form

Lansoprazole/amoxicillin/clarithromycin combination pack PA Criteria:

- Documentation of recent positive *H. pylori* test; **AND**
- Documentation of clarithromycin-sensitive strain of H. pylori is present; **AND**
- Trial and failure, contraindication, or intolerance to a preferred combination agent; **OR**
- Trial and failure, contraindication, or intolerance to H. pylori treatment prescribed as separate agents (e.g., amoxicillin + clarithromycin + omeprazole or bismuth subsalicylate + metronidazole + tetracycline)

Voquezna Dual Pak®

PDL Status	P
Quantity Limit	1 box/Rx; 2 courses of therapy/year
PA Criteria	See below
PA Form	General PA Form

Voquezna Dual Pak® PA Criteria:

- Documentation of recent positive *H. pylori* test; **AND**
- Trial and failure, contraindication, or intolerance to a preferred combination agent; **OR**
- Trial and failure, contraindication, or intolerance to H. pylori treatment prescribed as separate agents (e.g., amoxicillin + clarithromycin + omeprazole or bismuth subsalicylate + metronidazole + tetracycline)

Voquezna Triple Pak®

PDL Status	P
Quantity Limit	1 box/Rx; 2 courses of therapy/year
PA Criteria	See below
PA Form	General PA Form

Voquezna Triple Pak® PA Criteria:

- Documentation of recent positive *H. pylori* test; **AND**
- Documentation of clarithromycin-sensitive strain of H. pylori is present; **AND**
- Trial and failure, contraindication, or intolerance to a preferred combination agent; **OR**
- Trial and failure, contraindication, or intolerance to H. pylori treatment prescribed as separate agents (e.g., amoxicillin + clarithromycin + omeprazole or bismuth subsalicylate + metronidazole + tetracycline)

Fecal Microbiota

Vowst® capsules

PDL Status	NP
Quantity Limit	12 caps/year
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	2 months

Vowst® capsules PA Criteria:

- Patient is ≥ 18 years old; **AND**
- Treatment is to prevent the recurrence of *Clostridioides difficile* infection (CDI); **AND**
- Patient has had three or more episodes of CDI within the past year; **AND**
- Submission of medical records (e.g. chart notes, lab test) of a positive C. difficile stool test with toxin A/B results within the previous 30 days; **AND**
- Patient has completed a full treatment course with ONE of the following antibiotic therapies with the course ending 2 to 4 days prior to initiating Vowst:
  - Fidaxomicin
  - Vancomycin; **AND**
- Prescribed by, or in consultation with, an infectious disease specialist or gastroenterologist; **AND**
- The agent will not to be used in combination with other products for prevention of CDI, such as Zinplava or Rebyota

Gallstone Solubilizing Agents

Ursodiol capsules (300 mg), tablets (250 and 500 mg)

PDL Status	P
Quantity Limit	300 mg caps: 3/day; 250 mg tabs: 3/day 500 mg tabs: 2/day
PA Criteria	None

Chenodal® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Chenodal® tablets PA Criteria:

Standard NP Criteria

Ctexli® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Ctexli® tablets Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of Cerebrotendinous Xanthomatosis (CTX); **AND**
- Diagnosis has been confirmed by the presence of pathogenic variants in CYP27A1 gene; **AND**
- Prescribed by, or in consultation with, a medical geneticist, neurologist, metabolic disorder sub-specialist, or physician who specializes in the treatment of CTX

Ctexli® tablets Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., stabilization or improvement in symptoms of CTX or reduction in cholesterol metabolites (e.g., plasma cholestanol and urine 23S-pentol)

Cholbam® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Cholbam® capsules PA Criteria:

- One of the following:
  - Diagnosis of Bile Acid Synthesis Disorders due to Single Enzyme Defects (SED); **OR**
  - Agent will be used as adjunctive treatment for manifestations of Peroxisomal Disorders (PDs); **AND**
- Prescribed by a hepatologist or gastroenterologist

Iqirvo® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Iqirvo® tablets PA Criteria:

- Patient has a diagnosis of primary biliary cholangitis (PBC) **AND**
- Prescribed by, or in consultation with, a hepatologist or gastroenterologist **AND**
- ONE of the following:
  - Both of the following:
    - Will be taken in combination with ursodiol; **AND**
    - Submitted lab documentation indicates the patient had an inadequate response (no reduction in ALP or total bilirubin after 1-year trial) to ursodiol; **OR**
  - Patient has a contraindication, or intolerance to ursodiol

Livdelzi® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Livdelzi® capsules PA Criteria:

See Iqirvo® prior authorization criteria

Reltone® capsules

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Reltone® capsules PA Criteria:

Standard NP Criteria

Urso-Forte® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Urso-Forte® tablets PA Criteria:

Standard NP Criteria

Hepatotropics

Rezdiffra® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Rezdiffra® tablets Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) or metabolic dysfunction associated steatohepatitis (MASH); **AND**
- Submission of medical records (e.g. chart notes) confirming disease is fibrosis stage F2 or F3 as confirmed by ONE of the following:
  - FibroScan
  - Fibrosis-4 index (FIB-4) greater than 1.3
  - ELF score
  - Magnetic Resonance Elastography (MRE)
  - Liver Biopsy; **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Patient has had an inadequate response, intolerance, or contraindication to a minimum 1 year trial of a stable subcutaneous GLP-1 (i.e., semaglutide, Wegovy® injection) maintenance dose (confirmed by claims history); **AND**
- Patient does not have untreated hyperthyroidism or hypothyroidism; **AND**
- Patient does not have decompensated cirrhosis (Child-Pugh Class B or C); **AND**
- Prescribed by, or in consultation with, a gastroenterologist or hepatologist; **AND**
- Prescriber attests patient is not consuming alcohol above recommended daily limits (20 g/day for women or 30 g/day for men)

Rezdiffra® tablets Renewal PA Criteria

- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g. NASH resolution, fibrosis stage improvements); **AND**
- Prescriber attests patient is not consuming alcohol above recommended daily limits (20 g/day for women or 30 g/day for men); **AND**
- Patient does not have untreated hyperthyroidism or hypothyroidism

H2 Receptor Antagonists

Nizatidine capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Nizatidine capsules PA Criteria:

Standard NP Criteria

Ranitidine tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Ranitidine tablets PA Criteria:
Standard NP Criteria

Inflammatory Bowel Disease, Miscellaneous Agents

Budesonide ER tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Budesonide foam

PDL Status	P
Quantity Limit	66.8 g/day
PA Criteria	None

Hydrocortisone AC 25 mg suppositories

PDL Status	P
Quantity Limit	12/30 days
PA Criteria	None

Hydrocortisone AC 30 mg suppositories

PDL Status	NP
Quantity Limit	12/30 days
PA Criteria	None

Hydrocortisone AC 30 mg suppositories PA Criteria:
Standard NP Criteria

Laxatives

Clenpiq® oral solution, GoLYTELY® solution, Kristalose® packets,

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Clenpiq® oral solution, GoLYTELY® solution, Kristaolse® packets PA Criteria:
Standard NP Criteria

Lactulose packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lactulose packets PA Criteria:

- Trial and failure, or contraindication to one preferred agent; **AND**
- Clinically valid reason why brand Kristalose packets cannot be used

PEG-3350 with flavor packs (generic of Gavilyte®), PEG-3350 electrolyte powder with ascorbic acid (generic of Moviprep®), Suflave® powder for solution, Suprep® bowel prep kit

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

PEG-3350 with flavor packs (generic of Gavilyte®), PEG-3350 electrolyte powder with ascorbic acid (generic of Moviprep®), Suflave® powder for solution, Suprep® bowel prep kit PA Criteria:
Standard NP Criteria

Sutab® tablets

PDL Status	NP
Quantity Limit	24 tablets/colonoscopy
PA Criteria	See below
PA Form	General PA Form

Sutab® tablets PA Criteria:
Standard NP Criteria

Motility Agents

Metoclopramide solution, tablets

PDL Status	P
Quantity Limit	12-week duration limit
PA Criteria	None

Gimoti® nasal spray

PDL Status	NP
Quantity Limit	1 bottle/Rx; 12-week duration limit
PA Criteria	See below
PA Form	General PA Form

Gimoti® nasal spray PA Criteria:

- Patient must have acute and recurrent diabetic gastroparesis; **AND**
- Patient is ≥ 18 years of age; **AND**
- Patient does not have a history of tardive dyskinesia (TD) or dystonic reaction to metoclopramide; **AND**
- Clinically valid reason why metoclopramide tablets or solution cannot be used

Metoclopramide ODT

PDL Status	NP
Quantity Limit	12-week duration limit
PA Criteria	See below
PA Form	General PA Form

Metoclopramide ODT PA Criteria:

- Unable to swallow solid dosage forms; **OR**
- Unable to absorb medications through the GI tract

Reglan® tablets

PDL Status	NP
Quantity Limit	12-week duration limit
PA Criteria	See below
PA Form	General PA Form

Reglan® tablets PA Criteria:
Standard NP Criteria

Mucosal Protectants

Carafate® tablets, Cytotec® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carafate® tablets, Cytotec® tablets PA Criteria:
Standard NP Criteria

Sucralfate suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sucralfate suspension PA Criteria:

- Patient is < 13 years of age; **OR**
- Trial and failure, or intolerance to, sucralfate tablets, **OR**
- Has documented difficulty swallowing/dysphagia

Pancreatic Enzymes

Pertzye® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pertzye® capsules PA Criteria:
Standard NP Criteria

Proton Pump Inhibitors and Related Agents

Please refer to the OTC Drug Formulary for coverage of OTC products.

The quantity limit for proton pump inhibitors is 1 per day.

Proton Pump Inhibitors Quantity Limit Criteria:

- If request is for twice daily dosing, one of the following must be met:
  - Treatment of *H. Pylori* (1-month duration)
  - Treatment of GI Bleed/Hemorrhagic Gastritis (12-month duration)
  - Patient has a diagnosis of Barrett's Esophagus with documentation of uncontrolled reflux symptoms or esophagitis (following a trial of once daily PPI therapy)
  - Patient is being treated for Zollinger-Ellison Syndrome or gastrinoma (12-month duration)
  - Uncontrolled symptoms following a 30-day trial of once daily PPI therapy (1-month duration); renewals will require member to attempt step down to once daily PPI therapy. If patient fails step down to once daily dosing, they will not be asked to step down again.

Esomeprazole capsules, lansoprazole capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Lansoprazole ODT

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Lansoprazole ODT PA Criteria:

- Patient is 12 years of age or older; **AND**
- Unable to swallow solid dosage forms; **AND**
- Clinically valid reason why Protonix® packets cannot be used

Nexium® packets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
Notes	PA not required for patients < 6 years old.

Nexium® packets PA Criteria:

- Unable to swallow solid dosage forms

Omeprazole capsules (Rx), omeprazole/sodium bicarbonate capsules (Rx), pantoprazole packets & tablets, Protonix® packets, rabeprazole tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Dexilant® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Dexilant® capsules PA Criteria:
Standard NP Criteria

Dexlansoprazole capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Dexlansoprazole capsules PA Criteria:

- Clinically valid reason why brand Dexilant® cannot be used

Esomeprazole packets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Esomeprazole packets PA Criteria:

- Unable to swallow solid dosage forms; **AND**
- Trial, failure, contraindication, or intolerance to Protonix® suspension and Nexium® granules; **AND**
- Clinically valid reason why patient cannot use brand Nexium® granules

Konvomep® suspension

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See below
PA Form	General PA Form
Notes	PA not required for patients < 6 years old

Konvomep® suspension PA Criteria:

- Unable to swallow solid dosage forms; **AND**
- Trial, failure, contraindication, or intolerance to two preferred non-solid oral dosage agents

Nexium® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Nexium® capsules PA Criteria:
Standard NP Criteria

Omeprazole/sodium bicarbonate packets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Omeprazole/sodium bicarbonate packets PA Criteria:

- Trial and failure, contraindication, or intolerance of two (2) preferred non-solid oral agents; **OR**
- Patient is unable to swallow solid dosage forms

Prevacid® capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Prevacid® capsules PA Criteria:

Standard NP Criteria

Prevacid SoluTab®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Prevacid SoluTab® PA Criteria:

- Unable to swallow solid dosage forms; **AND**
- Trial, failure, contraindication, or intolerance to two preferred non-solid oral agents; **AND**
- Clinically valid reason why generic lansoprazole ODT cannot be used

Prilosec® packets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Prilosec® packets PA Criteria:

- Trial and failure, contraindication, or intolerance of two (2) preferred non-solid oral agents; **OR**
- Patient is unable to swallow solid dosage forms

Protonix® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Protonix® tablets PA Criteria:

Standard NP Criteria

Voquezna® tabs

PDL Status	NP
Quantity Limit	1/day AND 10 mg: 180 days/year or 20 mg: 60 days/year
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Active treatment of erosive esophagitis: 2 months; Maintenance treatment of healed erosive esophagitis: 6 months; Non-erosive gastroesophageal reflux disease (GERD): 1 month

Voquezna® PA Criteria:

- Patient is 18 years of age or older; **AND**
- Trial and failure, contraindication, or intolerance to TWO preferred proton pump inhibitors
- One of the following diagnoses:
  - Active treatment of erosive esophagitis; **AND**
    - Request is for Voquezna 20 mg tablets; **OR**
  - Maintenance treatment of healed erosive esophagitis; **AND**
    - Request is for Voquezna 10 mg tablets; **OR**
  - Non-erosive gastroesophageal reflux disease (GERD); **AND**
    - Request is for Voquezna 10 mg tablets

IX. IMMUNOLOGIC

Allergen Specific Immunotherapy

Grastek®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Prior authorizations may be processed for Grastek® between January 1 and March 31; with PA requests being accepted 2 weeks prior to this period. Requests received after March 31 will not be processed.

Grastek® PA Criteria:

- Diagnosis of grass pollen-induced allergic rhinitis with or without conjunctivitis; **AND**
- Documentation initial dose was administered in the physician office or medical facility; **AND**
- Must be prescribed by an allergy/immunology specialist; **AND**
- Patient's diagnosis is confirmed with documentation of **ONE** of the following:
  - A positive skin test to **ONE** of the pollen extracts contained in the requested agent
  - Pollen specific IgE antibodies to **ONE** of the pollen extracts contained in the requested agent; **AND**
- Trial and failure, contraindication, or intolerance to **ONE** agent from **TWO** of the following classes:
  - Oral antihistamine
  - Intranasal antihistamine
  - Intranasal corticosteroid
  - Leukotriene receptor antagonist; **AND**
- Documented trial/failure or drug-drug interaction of subcutaneous allergen immunotherapy (SCIT, or allergy shots) [Note: Failure defined as lack of efficacy, allergic reaction, documented intolerable side effects; agent will not be approved for needle phobia]; **AND**
- Patient has been prescribed and trained to administer epinephrine in case of severe allergic reaction; **AND**
- Oral Anti-allergens will NOT be approved if patient meets ANY of the following:
  - Patient experienced a severe reaction post initial dose administered in the physician's office
  - Patient has concomitant allergen immunotherapy
  - Patient has a history of severe, unstable, or uncontrolled asthma
  - Patient has a history of eosinophilic esophagitis; **AND**
- Treatment is requested within 12 weeks prior to season of allergen being treated (Grass season: April-September)

Odactra®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Odactra® PA Criteria:

- Diagnosis of house dust mite (HDM) induced allergic rhinitis with or without conjunctivitis; **AND**
- Patient's diagnosis confirmed with documentation of ONE of the following:
  - Confirmed in vitro IgE antibodies to Dermatophagoides farinae or Dermatophagoides pteronyssinus HDMs
  - Confirmed skin testing to licensed HDM allergen extracts; **AND**
- Prescribed by or in consultation with an allergy/immunology specialist; **AND**
- Documentation initial dose was administered in the physician office or medical facility; **AND**
- Trial and failure, contraindication, or intolerance to ONE agent from TWO of the following classes:
  - Oral antihistamine
  - Intranasal antihistamine
  - Intranasal corticosteroid
  - Leukotriene receptor antagonist; **AND**
- Patient has been prescribed and trained to administer epinephrine in case of severe allergic reaction; **AND**
- Oral Anti-allergens will NOT be approved if patient meets ANY of the following:
  - Patient experienced a severe reaction post initial dose administered in the physician's office
  - Patient has concomitant allergen immunotherapy
  - Patient has a history of severe, unstable, or uncontrolled asthma
  - Patient has a history of eosinophilic esophagitis

Oralair®

PDL Status	NP
Quantity Limit	Tablets: 1/day; Dose pack: total max limit 100 mg IR/300 mg IR
PA Criteria	See Below
PA Form	General PA Form
Notes	Prior authorizations may be processed for Oralair® between December 1 and March 31; with PA requests being accepted 2 weeks prior to this period. Requests received after March 31 will not be processed.

Oralair® PA Criteria:

- Diagnosis of grass pollen-induced allergic rhinitis with or without conjunctivitis; **AND**
- Documentation initial dose was administered in the physician office or medical facility; **AND**
- Must be prescribed by an allergy/immunology specialist; **AND**
- Patient's diagnosis is confirmed with documentation of **ONE** of the following:
  - A positive skin test to **ONE** of the pollen extracts contained in the requested agent
  - Pollen specific IgE antibodies to **ONE** of the pollen extracts contained in the requested agent; **AND**
- Trial and failure, contraindication, or intolerance to **ONE** agent from **TWO** of the following classes:
  - Oral antihistamine
  - Intranasal antihistamine
  - Intranasal corticosteroid
  - Leukotriene receptor antagonist; **AND**
- Documented trial/failure or drug-drug interaction of subcutaneous allergen immunotherapy (SCIT, or allergy shots) [Note: Failure defined as lack of efficacy, allergic reaction, documented intolerable side effects; agent will not be approved for needle phobia]; **AND**
- Patient has been prescribed and trained to administer epinephrine in case of severe allergic reaction; **AND**
- Oral Anti-allergens will NOT be approved if patient meets ANY of the following:
  - Patient experienced a severe reaction post initial dose administered in the physician's office
  - Patient has concomitant allergen immunotherapy
  - Patient has a history of severe, unstable, or uncontrolled asthma
  - Patient has a history of eosinophilic esophagitis; **AND**
- Treatment is requested within 4 months prior to season of allergen being treated (Grass season: April-September)

Ragwitek®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Prior authorizations may be processed for Ragwitek® between May 1st through July 31st; with PA requests being accepted 2 weeks prior to this period. Requests received after July 31st will not be processed.

Ragwitek® PA Criteria:

- Diagnosis of grass pollen-induced allergic rhinitis with or without conjunctivitis; **AND**
- Documentation initial dose was administered in the physician office or medical facility; **AND**
- Must be prescribed by an allergy/immunology specialist; **AND**
- Patient's diagnosis is confirmed with documentation of **ONE** of the following:
  - A positive skin test to **ONE** of the pollen extracts contained in the requested agent
  - Pollen specific IgE antibodies to **ONE** of the pollen extracts contained in the requested agent; **AND**
- Trial and failure, contraindication, or intolerance to **ONE** agent from **TWO** of the following classes:
  - Oral antihistamine
  - Intranasal antihistamine
  - Intranasal corticosteroid
  - Leukotriene receptor antagonist; **AND**
- Documented trial/failure or drug-drug interaction of subcutaneous allergen immunotherapy (SCIT, or allergy shots) [Note: Failure defined as lack of efficacy, allergic reaction, documented intolerable side effects; agent will not be approved for needle phobia]; **AND**
- Patient has been prescribed and trained to administer epinephrine in case of severe allergic reaction; **AND**
- Oral Anti-allergens will NOT be approved if patient meets ANY of the following:
  - Patient experienced a severe reaction post initial dose administered in the physician's office
  - Patient has concomitant allergen immunotherapy
  - Patient has a history of severe, unstable, or uncontrolled asthma
  - Patient has a history of eosinophilic esophagitis; **AND**
- Treatment is requested within 12 weeks prior to season of allergen being treated (Ragweed season: August-December)

Anti-Rheumatic: Kinase Inhibitors

Rinvoq® ER tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Ankylosing Spondylitis (AS), non-radiographic axial Spondylarthritis (nr-axSpA), or Psoriatic Arthritis (PsA)

Rinvoq® ER tablets Initial PA Criteria:

- Diagnosis of active ankylosing spondylitis; **OR**
- Diagnosis of active psoriatic arthritis (PSA); **OR**
- Diagnosis of active non-radiographic axial spondylarthritis (nr-axSpA) with objective signs of inflammation; **AND**
- Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel); **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the risks of heart-related events (heart attack, stroke, blood clots, etc.) or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate

Rinvoq® ER tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, lower UC disease activity index)

Atopic Dermatitis (AD)

Rinvoq® ER tablets Initial PA Criteria:

- Diagnosis of moderate to severe Atopic Dermatitis; **AND**
- Trial and failure, contraindication, or intolerance to 1 topical corticosteroid of medium-to-high potency; **AND**
- Trial and failure, contraindication, or intolerance to a topical calcineurin inhibitor; **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the risks of heart-related events (heart attack, stroke, blood clots, etc.) or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate

Rinvoq® ER tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, lower UC disease activity index)

Crohn's Disease (CD) or Ulcerative Colitis (UC)

Rinvoq® ER tablets Initial PA Criteria:

- Diagnosis of moderately to severely active Crohn's Disease; **OR**
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
- Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g., Humira); **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the risks of heart-related events (heart attack, stroke, blood clots, etc.) or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate

Rinvoq® ER tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, lower UC disease activity index)

Giant Cell Arteritis (GCA)

Rinvoq® ER tablets Initial PA Criteria:

- Diagnosis of giant cell arteritis (GCA); **AND**
- Trial and failure, contraindication, or intolerance to systemic glucocorticoids, azathioprine, or methotrexate; **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the risks of heart-related events (heart attack, stroke, blood clots, etc.) or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate

Rinvoq® ER tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, lower UC disease activity index)

Polyarticular Juvenile Idiopathic Arthritis (pJIA) or Rheumatoid Arthritis (RA)

Rinvoq® ER tablets Initial PA Criteria:

- Diagnosis of active polyarticular juvenile idiopathic arthritis (pJIA); **OR**
- Diagnosis of moderately to severely active rheumatoid arthritis (RA); **AND**
- Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, OR sulfasalazine; **AND**
- Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel); **AND**
- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh the risks of heart-related events (heart attack, stroke, blood clots, etc.) or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate

Rinvoq® ER tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, lower UC disease activity index)

Rinvoq LQ® solution

PDL Status	P
Quantity Limit	30 mL/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Rinvoq LQ® solution Initial PA Criteria:

- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab) **OR** potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh heart-related events or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered, and it has been determined that Jak inhibitor therapy is appropriate; **AND**
- One of the following:
  - Diagnosis of active Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
    - Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, **OR** sulfasalazine; **OR**
  - Diagnosis of active psoriatic arthritis (PSA); **AND**
    - Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel); **AND**
- One of the following:
  - Patient weighs <30 kg
  - Patient is unable to swallow solid dosage forms

Rinvoq LQ® solution Renewal PA Criteria:

- One of the following:
  - Patient weighs <30 kg
  - Patient is unable to swallow oral dosage forms; **AND**
- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts)

Xeljanz® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Xeljanz® tablets Initial PA Criteria:

- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab) **OR** potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh heart-related events or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate; **AND**
- One of the following:
  - Diagnosis of moderately to severely active Rheumatoid Arthritis (RA), or active Polyarticular Course Juvenile Idiopathic Arthritis (pcJIA); **AND**
    - Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, **OR** sulfasalazine; **AND**
    - Trial and failure or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel)
  - Diagnosis of active Psoriatic Arthritis (PsA); **AND**
    - Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel); **OR**
  - Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
    - Trial and failure, contraindication, or intolerance to Humira
  - Diagnosis of Ankylosing spondylitis; **AND**
    - Trial and failure, contraindication, or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel)

Xeljanz® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, lower UC disease activity index)

Olumiant® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year
Notes	Will not be covered for COVID-19 treatment in post hospitalized patients

Rheumatoid Arthritis

Olumiant® tablets Initial PA Criteria:

- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) **OR** potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh heart-related events or cardiovascular risk factors
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate; **AND**
- Diagnosis of moderately to severely active Rheumatoid Arthritis; **AND**
  - Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, **OR** sulfasalazine; **AND**
  - Trial and failure, contraindication, or intolerance a preferred TNF-inhibitors (e.g., Enbrel, Humira); **AND**
  - Trial and failure, contraindication, or intolerance to ONE preferred agent; **OR**

Olumiant® tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, lower UC disease activity index)

Alopecia Areata

Refer to Criteria for Non-PDL Agents for this indication.

Tofacitinib solution

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tofacitinib solution Initial PA Criteria:

See Xeljanz® solution PA criteria.

Tofacitinib tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tofacitinib tablets Initial PA Criteria:

See Xeljanz® tablets PA criteria.

Tofacitinib ER tablets (11 mg)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tofacitinib ER tablets (11 mg) Initial PA Criteria:
See Xeljanz XR® 11 mg tablets PA criteria.

Tofacitinib ER tablets (22 mg)

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tofacitinib ER tablets (2 mg) Initial PA Criteria:
See Xeljanz XR®22 mg tablets PA criteria.

Xeljanz® solution

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Xeljanz® solution Initial PA Criteria:

- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) **OR** potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh heart-related events or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate; **AND**
- Diagnosis of active Polyarticular Juvenile Idiopathic Arthritis (pcJIA); **AND**
- Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, **OR** sulfasalazine; **AND**
- Trial and failure or intolerance to a TNF-inhibitor (e.g. Humira, Enbrel); **AND**
- One of the following:
  - Patient weighs <40 kg
  - Patient is unable to swallow oral dosage forms

Xeljanz® solution Renewal PA Criteria

- One of the following:
  - Patient weighs <40 kg
  - Patient is unable to swallow oral dosage forms; **AND**
- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, lower UC disease activity index)

Xeljanz XR® 11 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Xeljanz XR® 11 mg tablets Initial PA Criteria:

See Xeljanz® tablet prior authorization criteria; **AND**

- Clinically valid reason why the preferred Xeljanz immediate release product cannot be used

Xeljanz XR® 22 mg tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Xeljanz XR® 22 mg tablets Initial PA Criteria:

- Prescriber attests to each of the following:
  - Patient is not concurrently taking biologic agents (i.e., adalimumab, anakinra, etanercept, rituximab, tocilizumab, infliximab, abatacept) OR potent immunosuppressants (i.e., azathioprine, cyclosporine); **AND**
  - Benefits of using this agent outweigh heart-related events or cardiovascular risk factors; **AND**
  - Risk of malignancy has been considered and it has been determined that Jak inhibitor therapy is appropriate; **AND**
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
- Trial and failure, contraindication, or intolerance to the preferred adalimumab product; **AND**
- Clinically valid reason why the preferred Xeljanz® immediate release product cannot be used

Xeljanz XR® 22 mg Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, lower UC disease activity index)

Disease Modifying Anti-Rheumatic Drugs (DMARDs)

Sulfasalazine tablets, sulfasalazine EC tablets

PDL Status	P
Quantity Limit	8/day
PA Criteria	None

Arava® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Arava® tablets PA Criteria:

Standard NP Criteria

Azulfidine® tablets, Azulfidine® EN tablets

PDL Status	NP
Quantity Limit	8/day
PA Criteria	See below
PA Form	General PA Form

Azulfidine®, Azulfidine® EN PA Criteria:
Standard NP Criteria

Jylamvo® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Jylamvo® solution PA Criteria:

- Dosing that will not allow the use of preferred methotrexate tablets; **OR**
- Patient unable to swallow methotrexate tablets

Otrexup® autoinjector

PDL Status	NP
Quantity Limit	4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form

Otrexup® autoinjector PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial/failure of TWO preferred DMARD agents; **AND**
  - Must have an allergy or contraindication to benzoyl alcohol or other preservative contained in injectable methotrexate that is not in requested agent; **OR**
  - Patient is experiencing dexterity issues without assistance to a caregiver who can administer the requested agent; **OR**
- Diagnosis of psoriasis; **AND**
- Trial and failure of TWO topical antipsoriatic agents; **AND**
  - Clinically valid reason why oral methotrexate cannot be used; **AND**
  - One of the following:
    - Patient has an allergy or contraindication to benzoyl alcohol or other preservative in injectable methotrexate that is not in requested agent
    - Patient is experiencing dexterity issues without assistance of a caregiver who can administer the requested agent

Plaquenil® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Plaquenil® tablets PA Criteria:
Standard NP Criteria

Rasuvo® autoinjector

PDL Status	NP
Quantity Limit	4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form

Rasuvo® autoinjector PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial/failure of TWO preferred DMARD agents; **AND**
  - Must have an allergy or contraindication to benzoyl alcohol or other preservative contained in injectable methotrexate that is not in requested agent; **OR**
  - Patient is experiencing dexterity issues without assistance to a caregiver who can administer the requested agent; **OR**
- Diagnosis of psoriasis; **AND**
- Trial and failure of TWO topical antipsoriatic agents; **AND**
  - Clinically valid reason why oral methotrexate cannot be used; **AND**
  - One of the following:
    - Patient has an allergy or contraindication to benzoyl alcohol or other preservative in injectable methotrexate that is not in requested agent
    - Patient is experiencing dexterity issues without assistance of a caregiver who can administer the requested agent

Sovuna® tablets, Trexall® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sovuna® tablets, Trexall® tablets PA Criteria:
Standard NP Criteria

Xatmep® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Xatmep® solution PA Criteria:

- Age ≤ 12 years; **AND**
- One of the following:
  - Dosing that will not allow the use of preferred methotrexate tablets
  - Patient unable to swallow methotrexate tablet

Immunomodulators: TNF Inhibitors

Adalimumab-adbm (NDCs 00597), adalimumab-adaz, adalimumab-fkjp, Amjevita®, Cyltezo®

PDL Status	P
Quantity Limit	2 injectors/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Adalimumab-adbm (NDCs 00597), adalimumab-adaz, adalimumab-fkjp, Amjevita®, Cyltezo® Initial PA Criteria:

- Diagnosis of Ankylosing Spondylitis (AS); **OR**
- Diagnosis of chronic, moderate to severe Plaque Psoriasis (PsO) and both the following:
  - Trial and failure of ONE topical treatment of a corticosteroid, calcipotriene, **OR** tazarotene; **AND**
  - Trial and failure, or contraindication, of ONE oral treatment with acitretin, methotrexate, **OR** cyclosporine; **OR**
- Diagnosis of Generalized pustular psoriasis (GPP) confirmed by ONE of the following:
  - Presence of sterile, macroscopically visible pustules on non-acral skin and pustulation is NOT restricted to psoriatic plaques (i.e., occurs outside of psoriatic plaques)
  - Skin biopsy confirming presence of Kogoj's spongiform pustules
  - Genetic confirmation of IL36RN, CARD14, or AP1S3 mutation; **OR**
- Diagnosis of active Psoriatic Arthritis (PsA); **OR**
- Diagnosis of active Juvenile Psoriatic arthritis (JPsA); **OR**
- Diagnosis of Rheumatoid Arthritis (RA) or Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, **OR** sulfasalazine; **OR**
- Diagnosis of moderate*ly* to severely active Ulcerative Colitis (UC); **OR**
- Diagnosis of moderately to severely active Crohn's Disease (CD) and ONE of the following:
  - Previous trial and failure of infliximab in the past 365 days
  - Crohn's disease is classified as moderate, severe, or fistulizing
  - ≥ 90 days of drug therapy with one of the following: azathioprine, mercaptopurine, mesalamine, methotrexate, or systemic glucocorticoid; **OR**
- Diagnosis of moderate to severe Hidradenitis Suppurativa (HS); **OR**
- Diagnosis of non-infectious intermediate, posterior or panuveitis (UV); **AND**
  - Diagnosis of Uveitis must be by, or in consultation with, an ophthalmologist; **AND**
  - ≥ 90 days of drug therapy with one of the following: oral/injectable steroid therapy, methotrexate, mycophenolate, azathioprine, cyclosporine, tacrolimus, cyclophosphamide

Adalimumab-adbm (NDCs 00597), adalimumab-adaz, adalimumab-fkjp, Amjevita®, Cyltezo® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score, <UC disease activity index, reduction in inflammatory bumps/abscesses, decreases in flares)

Enbrel®, Enbrel Mini-Cartridge®, Enbrel SureClick®

PDL Status	P
Quantity Limit	25 mg dose: 8 syringes/28 days; 50 mg dose: 4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Enbrel®, Enbrel Mini-Cartridge®, Enbrel SureClick® Initial PA Criteria:

- Diagnosis of Ankylosing Spondylitis; **OR**
- Diagnosis of chronic, moderate to severe Plaque Psoriasis; **AND**
  - Trial and failure of ONE topical treatment of a corticosteroid, calcipotriene, **OR** tazarotene; **AND**
  - Trial and failure, or contraindication, of ONE oral treatment with acitretin, methotrexate, **OR** cyclosporine; **OR**
- Diagnosis of active Psoriatic Arthritis (PsA); **OR**
- Diagnosis of active Juvenile Psoriatic arthritis (JPsA); **OR**
- Diagnosis of Rheumatoid Arthritis (RA) or Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, **OR** sulfasalazine

Enbrel®, Enbrel Mini-Cartridge®, Enbrel SureClick® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, a 50% reduction of total Psoriasis Area Severity Index (PASI) score)

Hadlima®, Humira®, Simlandi®

PDL Status	P
Quantity Limit	2 syringes/28 days; Starter packs: 1 kit/28 days Hidradenitis Suppurativa (HS) diagnosis only: 4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Hadlima®, Humira®, Simlandi® Initial PA Criteria:

- Diagnosis of Ankylosing Spondylitis (AS); **OR**
- Diagnosis of chronic, moderate to severe Plaque Psoriasis (PsO) and both the following:
  - Trial and failure of ONE topical treatment of a corticosteroid, calcipotriene, **OR** tazarotene; **AND**
  - Trial and failure, or contraindication, of ONE oral treatment with acitretin, methotrexate, **OR** cyclosporine; **OR**
- Diagnosis of Generalized pustular psoriasis (GPP) confirmed by ONE of the following:
  - Presence of sterile, macroscopically visible pustules on non-acral skin and pustulation is NOT restricted to psoriatic plaques (i.e., occurs outside of psoriatic plaques)
  - Skin biopsy confirming presence of Kogoj's spongiform pustules
  - Genetic confirmation of IL36RN, CARD14, or AP1S3 mutation; **OR**
- Diagnosis of active Psoriatic Arthritis (PsA); **OR**
- Diagnosis of active Juvenile Psoriatic arthritis (JPsA); **OR**
- Diagnosis of Rheumatoid Arthritis (RA) or Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, **OR** sulfasalazine; **OR**
- Diagnosis of moderate*ly* to severely active Ulcerative Colitis (UC); **OR**
- Diagnosis of moderately to severely active Crohn's Disease (CD) and ONE of the following:
  - Previous trial and failure of infliximab in the past 365 days
  - Crohn's disease is classified as moderate, severe, or fistulizing
  - ≥ 90 days of drug therapy with one of the following: azathioprine, mercaptopurine, mesalamine, methotrexate, or systemic glucocorticoid; **OR**
- Diagnosis of moderate to severe Hidradenitis Suppurativa (HS); **OR**
- Diagnosis of non-infectious intermediate, posterior or panuveitis (UV); **AND**
  - Diagnosis of Uveitis must be by, or in consultation with, an ophthalmologist; **AND**
  - ≥ 90 days of drug therapy with one of the following: oral/injectable steroid therapy, methotrexate, mycophenolate, azathioprine, cyclosporine, tacrolimus, cyclophosphamide

Hadlima®, Humira®, Simlandi® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score, <UC disease activity index, reduction in inflammatory bumps/abscesses, decreases in flares)

Abrilada®, adalimumab-aacf, adalimumab-aaty, adalimumab-adbm (NDCs 82009), adalimumab-ryvk

PDL Status	NP
Quantity Limit	2 injectors/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Abrilada®, adalimumab-aacf, adalimumab-aaty, adalimumab-adbm (NDCs 82009), adalimumab-ryvk Initial PA Criteria:

- Diagnosis of one of the following:
  - Ankylosing Spondylitis (AS)
  - Active Psoriatic Arthritis (PsA)
  - Rheumatoid Arthritis (RA)
  - Polyarticular Juvenile Idiopathic Arthritis (pJIA)
  - Plaque Psoriasis (PsO)
  - Moderately to severely active Crohn's Disease (CD)
  - Hidradenitis Suppurativa (HS)
  - Moderately to severely active Ulcerative Colitis (UC)
  - Non-infectious intermediate, posterior or panuveitis (UV); **AND**
- Clinically valid reason why ALL the preferred adalimumab products cannot be used

Abrilada®, adalimumab-aacf, adalimumab-aaty, adalimumab-adbm (NDCs 82009), adalimumab-ryvk Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score, endoscopic remission)

Cimzia®

PDL Status	NP
Quantity Limit	2 kits/28 days (4 syringes)
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Cimzia® Initial PA Criteria:

- One of the following:
  - Diagnosis of one of the following:
    - Ankylosing spondylitis (AS)
    - Axial spondyloarthritis, non-radiographic (nr-axSpA)
    - Active psoriatic arthritis (PsA)
    - Rheumatoid arthritis (RA)
    - Plaque psoriasis (PsO);
    - Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred immunomodulators with same indication; **OR**
  - Diagnosis of moderately to severely active Crohn's disease; **AND**
    - Trial and failure, contraindication, or intolerance to a preferred adalimumab product Entyvio, or infliximab

Cimzia® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score, endoscopic remission)

Hulio®, Hyrimoz®, Idacio®

PDL Status	NP
Quantity Limit	2 injectors/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Hulio®, Hyrimoz®, Idacio® PA Criteria:
See Abrilada® prior authorization criteria.

Simponi®

PDL Status	NP
Quantity Limit	1 syringe/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Simponi® Initial PA Criteria:

- Diagnosis of Ankylosing Spondylitis, active Psoriatic Arthritis (PsA), or Rheumatoid Arthritis (RA):
  - Trial and failure, contraindication, or intolerance to 2 preferred immunomodulators with same indication; **OR**
- Diagnosis of moderately to severely active Ulcerative Colitis (UC):
  - Trial and failure to two of the following (or have an intolerance or contraindication to all agents):
  - A preferred adalimumab product
  - Entyvio
  - Infliximab
  - A preferred JAK inhibitor (e.g., Xeljanz and Rinvoq)

Simponi® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score, endoscopic remission)

Yuflyma®, Yusimry®

PDL Status	NP
Quantity Limit	2 injectors/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Yuflyma®, Yusimry® PA Criteria:
See Abrilada® prior authorization criteria.

Zymfentra®

PDL Status	NP
Quantity Limit	2/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Zymfentra® Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of ONE of the following:
  - Moderately to severely active Crohn's disease
  - Moderately to severely active Ulcerative Colitis; **AND**
- Submission of medical records demonstrating a positive clinical response following a treatment minimum of 10 weeks of infliximab IV; **AND**
- Must provide last administration date for the IV infliximab doses; **AND**
- Provider attests to ALL of the following:
  - Patient has completed the full IV regimen as directed per prescribing
  - Patient has not experienced any severe adverse reactions
  - Patient has demonstrated a positive response to therapy
  - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Zymfentra® Renewal PA Criteria

- Diagnosis of ONE of the following:
  - Moderately to severely active Crohn's disease
  - Moderately to severely active Ulcerative Colitis; **AND**
- Disease response to therapy and tolerability compared to baseline (e.g., decreased UC disease activity index, endoscopic remission, decreased number of soft stools, decreased abdominal pain)

Immunomodulators: Interleukin Inhibitors

Kineret® prefilled syringe

PDL Status	P
Quantity Limit	1 syringe/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Kineret® prefilled syringe Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis; **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, **OR** sulfasalazine; **OR**
- Diagnosis of Neonatal-Onset Multisystem Inflammatory Disease (NOMID)
- Diagnosis of Deficiency of Interleukin-1 Receptor Antagonist (DIRA)

Kineret® prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts)

Pyzchiva® prefilled syringe, Selarsdi® prefilled syringe and 45 mg/0.5 mL vials

PDL Status	P
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Pyzchiva® prefilled syringe, Selarsdi® prefilled syringe and 45 mg/0.5 mL vials Initial PA Criteria:

- Diagnosis of Plaque Psoriasis (PsO) or active Psoriatic Arthritis (PsA); **AND**
  - Trial and failure, contraindication, or intolerance to ONE preferred immunomodulator with same indication; **OR**
- Diagnosis of moderately to severely active Crohn's disease (CD); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Pyzchiva® prefilled syringe, Selarsdi® prefilled syringe and 45 mg/0.5 mL vials Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Taltz® autoinjector, prefilled syringe

PDL Status	P
Quantity Limit	1 syringe/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Taltz® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of chronic, moderate to severe Plaque Psoriasis; **AND**
  - Trial and failure to ONE topical treatment of a corticosteroid, calcipotriene, **OR** tazarotene; **AND**
  - Trial and failure, or contraindication, to ONE oral treatment with acitretin, methotrexate, **OR** cyclosporine; **OR**
- Diagnosis of active Psoriatic Arthritis (PsA); **OR**
- Diagnosis of Axial spondylarthritis (axSpA), Active Ankylosing spondylitis (AS), or Active non-radiographic axial spondyloarthritis (nr-axSpA)

Taltz® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Tremfya® autoinjector, pen, prefilled syringe

PDL Status	P
Quantity Limit	1 syringe (1 mL) or autoinjector/56 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tremfya® autoinjector, pen, prefilled syringe Initial PA Criteria:

- Diagnosis of Plaque Psoriasis (PsO) or active psoriatic arthritis (PsA); **AND**
  - Trial and failure of ONE preferred immunomodulator agents with the same indication; **OR**
- Diagnosis of moderately to severely active Crohn's disease; **AND**
  - One of the following:
    - Patient is receiving or has received IV induction dosing; **AND**
      - Provider must provide last administration date for the IV formulation; **AND**
      - Provider attests to ALL of the following:
        - Patient has completed the full IV regimen as directed per prescribing
        - Patient has not experienced any severe adverse reactions
        - Patient has demonstrated a positive response to therapy
        - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug); **OR**
    - Patient is receiving SQ induction dosing; **AND**
      - Patient will begin induction dosing with 400 mg (2-200 mg) subcutaneous (SQ) doses at weeks 0, 4, and 8; **AND**
      - Trial and failure, contraindication, or intolerance to a preferred adalimumab product, Entyvio, or infliximab; **OR**
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
  - One of the following:
    - Patient is receiving or has received IV induction dosing; **AND**
      - Provider must provide last administration date for the IV formulation; **AND**
      - Provider attests to ALL of the following:
        - Patient has completed the full IV regimen as directed per prescribing
        - Patient has not experienced any severe adverse reactions
        - Patient has demonstrated a positive response to therapy
        - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug); **OR**
    - Patient is receiving SQ induction dosing; **AND**
      - Patient will begin induction dosing with 400 mg (2-200 mg) subcutaneous (SQ) doses at weeks 0, 4, and 8; **AND**
      - Trial and failure of ONE of the following (or have an intolerance or contraindication to all agents):
        - A preferred adalimumab product
        - Entyvio
        - Infliximab
        - A preferred JAK inhibitor (e.g. Xeljanz or Rinvoq);

Tremfya® autoinjector, pen, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Tyenne® autoinjector, prefilled syringe

PDL Status	P
Quantity Limit	4 injections (3.6 mL)/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tyenne® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis; **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, **OR** sulfasalazine; **OR**
- Diagnosis of active Systemic Juvenile Idiopathic Arthritis (SJIA); **OR**
- Diagnosis of Polyarticular Juvenile Idiopathic Arthritis (pJIA); **OR**
- Diagnosis of Giant Cell Arteritis (GCA) and ONE of the following:
  - Trial and failure, contraindication, or intolerance to systemic glucocorticoids, azathioprine, or methotrexate; **OR**
  - Patient will be utilizing systemic glucocorticoids with tocilizumab

Tyenne® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease number of tender/swollen joint course)

Actemra® autoinjector, prefilled syringe, Actemra ACTPen® autoinjector

PDL Status	NP
Quantity Limit	3.6 mL/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Actemra® autoinjector, prefilled syringe, Actemra ACTPen® autoinjector Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or active Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, OR sulfasalazine; **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used*;* **OR**
- Diagnosis of active Systemic Juvenile Idiopathic Arthritis (SIJA); **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used; **OR**
- Diagnosis of Giant Cell Arteritis (CGA); **AND**
  - One of the following:
    - Trial and failure, contraindication, or intolerance to systemic glucocorticoids, azathioprine, or methotrexate; **OR**
    - Patient will be utilizing systemic glucocorticoid with tocilizumab; **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used; **OR**
- Diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD); **AND**
  - Patient is 18 years of age or older; **AND**
  - Patient's onset of disease was 5 years ago or less; **AND**
  - Patient has active disease with elevated inflammatory markers or platelets

Actemra® autoinjector, prefilled syringe, Actemra ACTPen® autoinjector Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts)

Arcalyst® injection

PDL Status	NP
Quantity Limit	8 vials/month
PA Criteria	See below
PA Form	General PA Form

Arcalyst® injection PA Criteria:

- Diagnosis of Cryopyrin-Associated Periodic Syndromes (CAPS), including Familial Cold Auto-inflammatory Syndrome (FCAS), and Muckle-Wells Syndrome (MWS); **OR**
- Diagnosis of Deficiency of Interleukin-1 Receptor Antagonist (DIRA); **AND**
- Patient has tried and failed or have contraindication or intolerance to preferred agent Kineret; **OR**
- Diagnosis of recurrent pericarditis (RP) and meets all of the following; **AND**
- Trial and failure, contraindication, or intolerance to ONE of the following:
  - Colchicine
  - Corticosteroids
  - NSAIDS

Avtozma® autoinjector

PDL Status	NP
Quantity Limit	3.6 mL/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Avtozma® autoinjector Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or active Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to methotrexate, leflunomide, OR sulfasalazine; **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used*;* **OR**
- Diagnosis of active Systemic Juvenile Idiopathic Arthritis (SIJA); **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used; **OR**
- Diagnosis of Giant Cell Arteritis (CGA); **AND**
  - One of the following:
    - Trial and failure, contraindication, or intolerance to systemic glucocorticoids, azathioprine, or methotrexate; **OR**
    - Patient will be utilizing systemic glucocorticoid with tocilizumab; **AND**
  - Clinically valid reason why the preferred product Tyenne cannot be used;

Avtozma® autoinjector Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts)

Bimzelx® autoinjector, prefilled syringe

PDL Status	NP
Quantity Limit	1 injection/28 days
PA Criteria	See below
PA Form	General PA Form

Bimzelx® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of moderate to severe Plaque Psoriasis (PsO), Active Psoriatic Arthritis (PsA), or Ankylosing Spondylitis (AS); **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred immunomodulators with the same indication; **OR**
- Diagnosis of Non-Radiographic Axial spondyloarthritis (nr-axSpA); **AND**
  - Trial and failure, contraindication, or intolerance to Taltz®; **OR**
- Diagnosis of moderate to severe Hidradenitis Suppurativa (HS); **AND**
  - Trial and failure, contraindication, or intolerance of a preferred adalimumab product

Bimzelx® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., reduction of total PASI score, reduction in inflammatory bumps/abscesses)

Cosentyx® autoinjector, prefilled syringe

PDL Status	NP
Quantity Limit	300 mg dose: 2 pens/28 days; 150 mg dose: 1 pen/28 days; Hidradenitis Suppurativa (HS) diagnosis only: 300 mg dose, 4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Cosentyx® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of chronic, moderate to severe Plaque Psoriasis (PsO), Ankylosing Spondylitis (AS), active Psoriatic Arthritis (PsA); **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred immunomodulators with same indication; **OR**
- Diagnosis of active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation; **AND**
  - Trial and failure, contraindication, or intolerance of Taltz®; **OR**
- Diagnosis of Active Enthesitis-related arthritis (ERA); **AND**
  - Failed an adequate trial of ONE NSAID (unless contraindicated); **AND**
- Diagnosis of moderate to severe Hidradenitis Suppurativa (HS); **AND**
  - Trial and failure, contraindication, or intolerance of a preferred adalimumab product

Cosentyx® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of tender and swollen joint counts, a 50% reduction of total PASI score, reduction in inflammatory bumps/abscesses, decreases in flares)

Icotyde® tablets

PDL Status	NP
Quantity Limit	1 tablet/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Icotyde® tablets PA Criteria:

- Diagnosis of chronic moderate to severe Plaque Psoriasis (PsO); **AND**
- Trial and failure to TWO preferred immunomodulator agents with the same indication, one of which must be Tremfya; **AND**
- Submission of medical records (i.e., chart notes) providing a clinically valid reason why patient can't use Tremfya injectable (NOTE: Patient convenience, mild injection site reactions, or needle phobia is NOT an approvable reason. Patient must have contraindication to the excipients in the injectable Tremfya product.)

Icotyde® tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g. a 50% reduction of total PASI score)

Imuldosa® prefilled syringe

PDL Status	NP
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Imuldosa® prefilled syringe Initial PA Criteria:

- Diagnosis of Plaque Psoriasis (PsO) or active Psoriatic Arthritis (PsA); **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred immunomodulators with same indication; **AND**
  - Clinically valid reason why a preferred ustekinumab agent cannot be used; **OR**
- Diagnosis of moderately to severely active Crohn's disease (CD); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Imuldosa® prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Kevzara® autoinjector, prefilled syringe

PDL Status	NP
Quantity Limit	2 pens or syringes/30 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Kevzara® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or active Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - Trial and failure, contraindication, or intolerance to 2 preferred immunomodulators with same indication: **OR**
- Diagnosis of Polymyalgia Rheumatic (PR); **AND**
  - Trial and failure, contraindication, or intolerance to systemic corticosteroids; **AND**

Kevzara® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts)

Omvoh® autoinjector, prefilled syringe

PDL Status	NP
Quantity Limit	2 autoinjectors/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Omvoh® autoinjector, prefilled syringe Initial PA Criteria:

- One of the following:
  - Diagnosis of Ulcerative Colitis; **OR**
  - Diagnosis of moderately to severely active Crohn's disease (CD); **AND**
- Patient is completing or has received required IV induction dosing; **AND**
- Provider must provide last administration date for the IV formulation; **AND**
- Provider attests to ALL of the following:
  - Patient has completed the full IV regimen as directed per prescribing
  - Patient has not experienced any severe adverse reactions
  - Patient has demonstrated a positive response to therapy
  - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Omvoh® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., endoscopic remission)

Otulfi® prefilled syringe and 45 mg/0.5 mL vials

PDL Status	NP
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Otulfi® prefilled syringe and 45 mg/0.5 mL vials Initial PA Criteria:

See Imuldosa® prior authorization criteria.

Skyrizi® autoinjector, cartridge, prefilled syringe

PDL Status	NP
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Skyrizi® autoinjector, cartridge, prefilled syringe Initial PA Criteria:

- Diagnosis of Plaque Psoriasis (PsO) or active psoriatic arthritis (PsA); **AND**
  - Trial and failure to TWO preferred immunomodulator agents with the same indication, one of which must be Tremfya; **OR**
- Diagnosis of moderately to severely active Crohn's disease (CD); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug); **OR**
- Diagnosis of moderately to severely active Ulcerative Colitis (UC); **AND**
  - Patient is completing or has received required IV induction dosing; **AND**
  - Provider must provide last administration date for the IV formulation; **AND**
  - Provider attests to ALL of the following:
    - Patient has completed the full IV regimen as directed per prescribing
    - Patient has not experienced any severe adverse reactions
    - Patient has demonstrated a positive response to therapy
    - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Skyrizi® autoinjector, cartridge, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Steqeyma® prefilled syringe and 45 mg/0.5 mL vials

PDL Status	NP
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Steqeyma® prefilled syringe and 45 mg/0.5 mL vials Initial PA Criteria:

See Imuldosa® prior authorization criteria.

Spevigo® prefilled syringe

PDL Status	NP
Quantity Limit	2/28 days
PA Criteria	See below
PA Form	General PA Form
Notes	The Spevigo subcutaneous formulation is not FDA approved for the treatment of GPP flare and will not be approved for that diagnosis. A SQ loading dose is not required following treatment of a GPP flare with IV Spevigo

Spevigo® prefilled syringe Initial PA Criteria:

- Diagnosis of Generalized pustular psoriasis (GPP) confirmed by ONE of the following:
  - Presence of sterile, macroscopically visible pustules on non-acral skin and pustulation is NOT restricted to psoriatic plaques (i.e. occurs outside of psoriatic plaques)
  - Skin biopsy confirming presence of Kogoj's spongiform pustules;
  - Genetic confirmation of IL36RN, CARD14, or AP1S3 mutation; **AND**
- Patient is 12 years of age and older and weights at least 40 kg; **AND**
- Prescriber attest to ALL of the following:
  - Treatment is NOT for an active GPP flare
  - Patient has been evaluated and screened for the presence of latent TB infection prior to initiating treatment
  - Patient will not receive live vaccines during therapy and 16 weeks after treatment; **AND**
- Trial and failure to BOTH of the following (or have an intolerance or contraindication to all agents):
  - A TNF inhibitor (e.g. adalimumab, infliximab, and etanercept)
  - Taltz; **AND**
  - Prescribed by, or in consultation with, a dermatologist

Spevigo® prefilled syringe Renewal PA Criteria:

- Submission of medical records (e.g. chart notes) documenting disease response to therapy and tolerability compared to baseline (e.g., decreased number of GPP flares)

Stelara® prefilled syringe and 45 mg/0.5 mL vials, ustekinumab prefilled syringe and 45 mg/0.5 mL vials, Yesintek® prefilled syringe and 45 mg/0.5 mL vials

PDL Status	NP
Quantity Limit	Crohn's Disease, Ulcerative Colitis: 1 injection/56 days; Plaque Psoriasis, Psoriatic Arthritis: 1 injection/84 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Stelara® prefilled syringe and 45 mg/0.5 mL vials, ustekinumab prefilled syringe and 45 mg/0.5 mL vials, Yesintek® prefilled syringe and 45 mg/0.5 mL vials Initial PA Criteria:

See Imuldosa® prior authorization criteria.

Immunomodulators: Miscellaneous

Orencia® autoinjector, prefilled syringe

PDL Status	P
Quantity Limit	4 mL/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Orencia® autoinjector, prefilled syringe Initial PA Criteria:

- Diagnosis of Rheumatoid Arthritis (RA) or Polyarticular Juvenile Idiopathic Arthritis (pJIA); **AND**
  - One of the following:
    - Patient is NOT receiving any IV infusion loading dose or IV induction dosing; **AND**
      - Trial and failure, contraindication, or intolerance to leflunomide, methotrexate, or sulfasalazine; **OR**
    - Patient is receiving or has received an IV infusion loading dose or induction dosing; **AND**
      - Provider must provide last administration date for the IV formulation; **AND**
      - Provider attests to ALL of the following:
        - Patient has completed the full IV regimen as directed per prescribing
        - Patient has not experienced any severe adverse reactions
        - Patient has demonstrated a positive response to therapy
        - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug); **OR**
- Diagnosis of active Psoriatic Arthritis PsA)

Orencia® autoinjector, prefilled syringe Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Otezla® tablets

PDL Status	P
Quantity Limit	30 mg: 2/day Starter pack: 1/Rx
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Otezla® tablets Initial PA Criteria:

- Diagnosis of Plaque Psoriasis (PsO); **AND**
  - Trial and failure to a topical treatment of a corticosteroid, calcipotriene, **OR** tazarotene; **AND**
  - Trial and failure, or contraindication, to at least one of the following oral agents: Soriatane®, methotrexate, cyclosporine; **OR**
- Diagnosis of active Psoriatic Arthritis (PsA); **OR**
- Diagnosis of oral lesions associated with Behçet's Disease; **AND**
  - Patient has active oral ulcers; **AND**
  - Trial and failure, contraindication, or intolerance to colchicine

Otezla® tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decrease in number of swollen joint counts, a 50% reduction of total PASI score)

Entyvio® pen

PDL Status	NP
Quantity Limit	2 injections/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 4 months; Renewal: 1 year

Entyvio® pen Initial PA Criteria:

- One of the following:
  - Diagnosis of moderate to severe Crohn's disease
  - Diagnosis of moderate to severe ulcerative colitis (UC); **AND**
- Patient is completing or has received required IV induction dosing; **AND**
- Provider must provide last administration date for the IV formulation; **AND**
- Provider attests to ALL of the following:
  - Patient has completed the full IV regimen as directed per prescribing
  - Patient has not experienced any severe adverse reactions
  - Patient has demonstrated a positive response to therapy
  - Provider has provided administration education to patient and/or caregiver (NOTE: provider accepts all professional responsibilities and risks associated with administration of drug);

Entyvio® pen Renewal PA Criteria:

- Patient is established on Entyvio therapy for ≥ 14 weeks (supported by paid claims or chart notes); **AND**
- Documentation of positive disease response to therapy and tolerability compared to baseline (e.g., decreased UC disease activity index, endoscopic remission, decreased stool frequency)

Otezla XR® tablets

PDL Status	NP
Quantity Limit	1/day; Starter pack: 1/Rx
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Otezla XR® tablets Initial PA Criteria:

See Otezla® prior authorization criteria; **AND**

- Patient has been titrated to maintenance dose with Otezla IR product; **AND**
- Clinically valid reason as to why the preferred Otezla formulation cannot be used

Sotyktu® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Sotyktu® tablets Initial PA Criteria:

- One of the following:
  - Diagnosis of moderate to severe Plaque Psoriasis (PsO); **OR**
  - Diagnosis of active psoriatic arthritis (PsA); **AND**
- Trial and failure, contraindication, or intolerance to 2 preferred immunomodulators with same indication

Sotyktu® tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., a 50% reduction of total PASI score, decrease in the number of swollen joint counts)

Velsipity® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 3 months; Renewal: 1 year

Velsipity® tablets Initial PA Criteria:

- Diagnosis of moderately to severely active ulcerative colitis (UC); **AND**
- Patient is ≥ 18 years old; **AND**
- Trial and failure to two of the following (or have an intolerance or contraindication to all agents):
- A preferred adalimumab product
    - Entyvio
    - Infliximab
    - A preferred JAK inhibitor (e.g. Xeljanz or Rinvoq); **AND**
- Patient does not have a recent (within the previous 6 months) myocardial infarction, unstable angina, stroke, transient ischemic attack (TIA), decompensated heart failure with hospitalization, or Class III/IV heart failure; **AND**
- Patient does not have a history or presence of Mobitz Type II second-degree, or third-degree atrioventricular block, sick sinus syndrome, or sinoatrial block (unless treated with a functioning pacemaker)

Velsipity® tablets Renewal PA Criteria

- Disease response to therapy and tolerability compared to baseline (e.g., endoscopic remission, decreased stool frequency, decreased rectal bleeding)

Immunosuppressants

Astagraf XL® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Astagraf XL® capsules PA Criteria:

- Trial and failure, contraindication, or intolerance to ONE preferred agent

Azasan® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Azasan® tablets PA Criteria:

- Diagnosis of rheumatoid arthritis; **AND**
    - Trial and failure of ONE preferred agent with the same indication (e.g., azathioprine); **OR**
- All transplant recipients will be allowed a prior authorization for any drug

CellCept® capsules, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

CellCept® capsules, tablets PA Criteria:

- All transplant recipients will be allowed a prior authorization for any drug

Envarsus XR® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Envarsus XR® tablets PA Criteria:

- Trial and failure, contraindication, or intolerance to ONE preferred agent

Imuran® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Imuran® tablets PA Criteria:

See Azasan® prior authorization criteria.

Myfortic® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Myfortic® tablets PA Criteria:

See CellCept® capsules, tablets prior authorization criteria.

Neoral® capsules, solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Neoral® capsules, solution PA Criteria:

- Diagnosis of rheumatoid arthritis or plaque psoriasis; **AND**
  - Trial and failure of ONE preferred agent with the same indication (e.g., cyclosporine, GENGRAF); **OR**
- All transplant recipients will be allowed a prior authorization for any drug

Prograf® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Prograf® capsules PA Criteria:

See CellCept® capsules, tablets prior authorization criteria.

Prograf® packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Prograf® packets PA Criteria:

- Trial and failure, contraindication, or intolerance to ONE preferred agent

Sandimmune® capsules, solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Sandimmune® capsules, solution PA Criteria:

See CellCept® capsules, tablets prior authorization criteria.

Tacrolimus ER capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tacrolimus ER capsules PA Criteria:

- Trial and failure, contraindication, or intolerance to ONE preferred agent; **AND**
- Clinically valid reason why brand Astagraf XL® cannot be used

Zortress® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
Note	The PA requirement may be overridden at POS via an ICD-10 code override.

Zortress® tablets PA Criteria:
See CellCept® capsules, tablets prior authorization criteria.

Lupus and Lupus Nephritis Agents

Benlysta®

PDL Status	NP
Quantity Limit	4 syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Benlysta® Initial PA Criteria:
- Patient is ≥ 5 years of age; **AND**
- Diagnosis of one of the following:
  - Active systemic lupus erythematosus (SLE); **OR**
  - Active lupus nephritis; **AND**
    - Diagnosis has been confirmed by biopsy or biopsy is contraindicated in the patient***; AND***
- Prescribed by, or consultation with, a rheumatologist or nephrologist; **AND**
- Must be used in combination with standard treatment regimens (e.g., corticosteroids, mycophenolate, azathioprine); **AND**
- Patient does not have severe active central nervous system lupus

Benlysta® Renewal PA Criteria:
- Patient meets the initial criteria; **AND**
- Positive clinical response to therapy (e.g., reduction in corticosteroid dose, reduction in flares, improvement in organ dysfunction)

Lupkynis®

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Lupkynis® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of active lupus nephritis; **AND**
- Diagnosis has been confirmed by biopsy or biopsy is contraindicated in the patient; **AND**
- Must take in combination with mycophenolate mofetil and corticosteroids; **AND**
- Will NOT take in combination with cyclophosphamide or Benlysta; **AND**
- Prescribed by, or in consultation with, a rheumatologist or nephrologist; **AND**
- Will not be used concomitantly with strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin); **AND**
- Patient is not pregnant

Lupkynis® Renewal PA Criteria:

- Must take in combination with mycophenolate mofetil and corticosteroids; **AND**
- Will NOT take in combination with cyclophosphamide or Benlysta; **AND**
- Prescribed by, or in consultation with, a rheumatologist or nephrologist; **AND**
- Patient has experienced a positive response to therapy (evidence of long-term preservation of kidney function, prevention of disease flares, prevention of organ damage); **AND**
- Patient has not experienced treatment-limiting adverse effects (eGFR decline, blood pressure increase, hypertensive crisis)

Multiple Sclerosis Agents, Injectable

Avonex®, Avonex Pack®

PDL Status	P
Quantity Limit	4/28 days
PA Criteria	None

Copaxone®

PDL Status	P
Quantity Limit	20 mg: 1 mL/day; 40 mg: 12 mL/30 days
PA Criteria	None

Betaseron®

PDL Status	NP
Quantity Limit	14/28 days
PA Criteria	See below
PA Form	General PA Form

Betaseron® PA Criteria:

Standard NP Criteria

Glatiramer

PDL Status	NP
Quantity Limit	20 mg: 1 mL/day; 40 mg: 12 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Glatiramer PA Criteria:
Standard NP Criteria

Glatopa®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Glatopa® PA Criteria:
Standard NP Criteria

Kesimpta®

PDL Status	NP
Quantity Limit	Initiation: 3 pens for the first month; Maintenance: 1 pen/month
PA Criteria	See below
PA Form	General PA Form

Kesimpta® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting Multiple Sclerosis [RRMS], active secondary progressive disease [SPMS]); **AND**
- Prescriber attests that initial dose was administered under the guidance of a healthcare professional; **AND**
- Patient meets ONE of the following:
  - Patient has active secondary progressive disease (SPMS); **OR**
  - Trial and failure, contraindication, or intolerance to one disease modifying therapy for MS; **OR**
  - Submission of medical records (e.g., chart notes) documenting clinical features of highly active MS such as high radiological burden of disease, high relapse frequency, severe relapse(s) requiring corticosteroids and/or hospitalization, severe physical or cognitive impairment; **AND**
- Patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment; **AND**
- For patients of reproductive potential, the following has been addressed:
  - Patient has been counseled to use effective contraception during treatment and for 6 months after the last dose; **AND**
  - Lactating women will be counseled to discontinue breastfeeding during treatment; **AND**
  - Provider has confirmed (via pregnancy test) that the patient is not pregnant prior to receiving treatment

Kesimpta® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Plegridy®

PDL Status	NP
Quantity Limit	2 pens/28 days
PA Criteria	See below
PA Form	General PA Form

Plegridy® PA Criteria:

- Patient is ≥ 18 years old; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Diagnosis of relapsing forms of Multiple Sclerosis which include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease; **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred injectable MS agents

Rebif®

PDL Status	NP
Quantity Limit	6 mL/28 days
PA Criteria	See below
PA Form	General PA Form

Rebif® PA Criteria:

Standard NP Criteria

Multiple Sclerosis Agents, Oral

Dalfampridine ER

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Dimethyl fumarate

PDL Status	P
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Dimethyl fumarate Initial PA Criteria:

- Patient must be the labeled age minimum; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e., clinically isolated syndrome, relapsing-remitting disease, active secondary progressive disease)

Dimethyl fumarate Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Fingolimod

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Fingolimod PA Criteria:

See dimethyl fumarate prior authorization criteria

Teriflunomide

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Teriflunomide PA Criteria:
See dimethyl fumarate prior authorization criteria

Ampyra®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Ampyra® PA Criteria:
- Clinically valid reason why preferred dalfampridine cannot be used

Aubagio®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Aubagio® PA Criteria:
See dimethyl fumarate prior authorization criteria; **AND**
- Clinically valid reason why preferred dalfampridine cannot be used

Bafiertam®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Bafiertam® Initial PA Criteria:
- Patient is ≥ 18 years old; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Trial and failure, contraindication, or intolerance of teriflunomide or fingolimod; **AND**
- Trial and failure, contraindication, or intolerance of dimethyl fumarate; **AND**

Bafiertam® Renewal PA Criteria:
- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Cladribine tablets

PDL Status	NP
Quantity Limit	40 tabs/2 years
PA Criteria	See below
PA Form	General PA Form

Cladribine tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- ONE of the following:
  - Diagnosis of relapsing-remitting multiple sclerosis [RRMS]; AND
    - Trial and failure, contraindication, or intolerance to teriflunomide, dimethyl fumarate, or fingolimod; **OR**
  - Diagnosis of active secondary progressive disease [SPMS]; AND
- For patients of reproductive potential:
  - Provider has counseled patient to use contraception during treatment and for 6 months after the last dose
  - Lactating women will be counseled to discontinue breastfeeding during treatment
  - Provider has confirmed (via pregnancy test) that the patient is not pregnant prior to receiving treatment; AND
- Prescriber attests to ALL of the following:
  - Patient does not have a current malignancy
  - Patient does not have serious active chronic infections such as HIV, tuberculosis, and active hepatitis
  - Patient has been screened for the presence of tuberculosis; **AND**
- Clinically valid reason why brand Mavenclad® tablets cannot be used

Cladribine tablets Renewal PA Criteria

- At least 43 weeks (approx. 10 months) has/will have elapsed since the end of the first treatment course; **AND**
- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Gilenya®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Gilenya® Initial PA Criteria:

- Patient is ≥ 10 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Diagnosis of a relapsing form of multiple sclerosis (i.e., clinically isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Clinically valid reason why preferred fingolimod cannot be used

Gilenya® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Mavenclad® tablets

PDL Status	NP
Quantity Limit	40 tabs/2 years
PA Criteria	See below
PA Form	General PA Form

Mavenclad® tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- ONE of the following:
  - Diagnosis of relapsing-remitting multiple sclerosis [RRMS]; AND
    - Trial and failure, contraindication, or intolerance to teriflunomide, dimethyl fumarate, or fingolimod; **OR**
  - Diagnosis of active secondary progressive disease [SPMS]; AND
- For patients of reproductive potential:
  - Provider has counseled patient to use contraception during treatment and for 6 months after the last dose
  - Lactating women will be counseled to discontinue breastfeeding during treatment
  - Provider has confirmed (via pregnancy test) that the patient is not pregnant prior to receiving treatment; AND
- Prescriber attests to ALL of the following:
  - Patient does not have a current malignancy
  - Patient does not have serious active chronic infections such as HIV, tuberculosis, and active hepatitis
  - Patient has been screened for the presence of tuberculosis

Mavenclad® tablets Renewal PA Criteria

- At least 43 weeks (approx. 10 months) has/will have elapsed since the end of the first treatment course; **AND**
- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Mayzent®

PDL Status	NP
Quantity Limit	Starter pack: 1 pack/Rx; 0.25 mg tabs: 4/day; 2 mg tabs: 1/day
PA Criteria	See below
PA Form	General PA Form

Mayzent® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Patient does NOT have any of the following:
    - Recent (within 6-months) myocardial infarction, unstable angina, stroke, transient ischemic attack (TIA), decompensated heart failure with hospitalization, or Class III/IV heart failure; **OR**
    - Prolonged QTc interval at baseline (> 500 msec); **OR**
    - History of Mobitz Type II second- or third-degree atrioventricular block or sick sinus syndrome (unless treated with a functioning pacemaker); **OR**
        - CYP2C9*3/*3 genotype; **OR**
        - Active infection (including clinically important localized infections); **AND**
- Patient will not be initiating therapy after previous treatment with alemtuzumab (Lemtrada); **AND**
- For female patients of reproductive potential, the following has been addressed:
    - Provider has counseled patient to use effective contraception during treatment with therapy
    - Lactating patient has been counseled on the risks versus benefits of breastfeeding while on treatment

Mayzent® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Ponvory®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Ponvory® Initial PA Criteria:

- Patient ≥ 18 years old; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- ONE of the following:
  - Diagnosis of relapsing-remitting multiple sclerosis [RRMS]; **AND**
    - Trial and failure, contraindication, or intolerance to teriflunomide, dimethyl fumarate, or fingolimod; **OR**
  - Diagnosis of active secondary progressive disease [SPMS]; **AND**
- Patient must NOT meet any of the following:
  - Recent (within 6-months) myocardial infarction, unstable angina, stroke, transient ischemic attack (TIA), decompensated heart failure with hospitalization, or Class III/IV heart failure; **OR**
  - Presence of Mobitz type II second-degree, third degree atrioventricular (AV) block, sick sinus syndrome unless the patient has a functioning pacemaker; **AND**
- For female patients of reproductive potential, all the following has been addressed:
  - Provider has counseled patient to use effective contraception during treatment
  - Lactating patients have been counseled on the risks versus benefits of breastfeeding while on treatment

Ponvory® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Tascenso ODT®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Tascenso ODT® Initial PA Criteria:

- Patient is ≥ 10 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Patient is unable to swallow sold dosage forms

Tascenso ODT® Renewal PA Criteria:

- Patient is unable to swallow sold dosage forms; **AND**
- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Tecfidera®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Tecfidera® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Clinically valid reason why preferred dimethyl fumarate cannot be used

Tecfidera® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Vumerity®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Vumerity® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has a relapsing form of multiple sclerosis (i.e. clinically Isolated syndrome, relapsing-remitting disease, active secondary progressive disease); **AND**
- Trial and failure, contraindication, or intolerance of teriflunomide or fingolimod; **AND**
- Trial and failure, contraindication, or intolerance of dimethyl fumarate

Vumerity® Renewal Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression)

Zeposia®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Zeposia® Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- ONE of the following:
  - Diagnosis of relapsing forms of multiple sclerosis, including clinical isolated syndrome, relapsing-remitting disease, and active secondary progressive disease: **AND**
    - Prescribed by, or in consultation with, a neurologist; **AND**
    - Trial and failure, contraindication, or intolerance to teriflunomide, dimethyl fumarate, **OR** fingolimod (not required for SPMS); **OR**
  - Diagnosis of moderately to severely active ulcerative colitis (UC) in adults; **AND**
    - Trial and failure to two of the following (or have an intolerance or contraindication to all agents):
      - A preferred adalimumab product
      - Entyvio
      - Infliximab
      - A preferred JAK inhibitor (e.g. Xeljanz and Rinvoq); **AND**
- Patient does NOT have any of the following:
  - Recent (within the previous 6-months) myocardial infarction, unstable angina, stroke, transient ischemic attack (TIA), decompensated heart failure with hospitalization, or Class III/IV heart failure
  - Severe untreated sleep apnea
  - History or presence of Mobitz Type II second-degree, or third-degree atrioventricular block, sick sinus syndrome, or sinoatrial block (unless treated with a functioning pacemaker)
  - Concomitantly taking a monoamine oxidase (MAO) inhibitor; **AND**
- For female patients of reproductive potential, all the following has been addressed:
  - Patient has been counseled to use effective contraception during treatment
  - Lactating patients have been counseled on the risks versus benefits of breastfeeding while on treatment

Zeposia® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., improvement or stabilization in radiologic disease activity, clinical relapses, disease progression, endoscopic remission)

Respiratory and Allergy Biologics

Adbry® injection

PDL Status	P
Quantity Limit	Initial month: 6 pens or syringes; Maintenance: 4 pens or syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Adbry® injection Initial PA Criteria:

- Patient is ≥ 12 years of age; **AND**
- Diagnosis of moderate to severe atopic dermatitis with ≥ 1 of the following:
  - Involvement of at least 10% of body surface area (BSA)
  - Scoring Atopic Dermatitis (SCORAD) score of 20 or more
  - Investigator's Global Assessment (IGA) with a score ≥ 3
  - Eczema Area and Severity Index (EASI) score of ≥ 16
  - Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (documented by claims) or contraindication to both of the following:
  - A topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone)
  - A topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Adbry® injectionRenewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SCORAD)

Dupixent® injection

PDL Status	P
Quantity Limit	2 syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Allergic Fungal Rhinosinusitis (AFRS)

Dupixent® injection Initial PA Criteria:

- Diagnosis of Allergic Fungal Rhinosinusitis (AFRS); **AND**
- Patient has previously had sino-nasal surgery; **AND**
- Patient is ≥ 6 years of age; **AND**
- Symptoms persist after trial of TWO of the following classes of agents
  - Nasal saline irrigations
  - Intranasal corticosteroids
  - Oral antifungals
  - Oral corticosteroids; **AND**
- Prescribed by, or in consultation with, an allergist, immunologist, otolaryngologist, or pulmonologist

Dupixent® injection Renewal PA Criteria:

- Patient continues to exhibit signs or symptoms of ongoing allergic fungal rhinosinusitis (AFRS); **AND**
- Documentation of positive clinical response to therapy (i.e. decrease in mucus discharge, decrease in loss of smell, decrease in recurrence of nasal polyps, etc.)

Asthma

Dupixent® injection Initial PA Criteria:

- Diagnosis of moderate to severe asthma; **AND**
- Patient is ≥ 6 years old; **AND**
- One of the following:
  - Asthma is an eosinophilic phenotype as defined by a baseline (pre- treatment) peripheral blood eosinophil level ≥ 150 cells/µL or peripheral blood eosinophil levels > 300 cells/mcL; **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Asthma is inadequately controlled as shown by one of the following:
  - Two more exacerbations requiring systemic corticosteroids within the past 12 months; **OR**
  - Asthma-related emergency treatment (e.g., hospitalization, intensive care unit stay, or mechanical ventilation in the past 12 months; **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Patient is currently being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including BOTH of the following:
    - One medium or high dose inhaled corticosteroid (ICS); **AND**
    - One additional asthma controller medication [e.g., LABA, leukotriene antagonist, theophylline]; **OR**
  - One maximally dosed combination inhaled ICS/ LABA or ICS/LAMA/LABA product; **AND**
- Will be used as adjunct therapy along with above asthma treatment; **AND**
- Prescribed by, or in consultation with, a pulmonologist, allergist, or immunologist

Dupixent® injection Renewal Criteria:

- Documentation of a positive clinical response (e.g., reduction in exacerbations, improvement of FEV1, decreased use of rescue medications); **AND**
- Patient is being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both a high-dose ICS and an additional asthma controller medication
  - One maximally dosed combination inhaled ICS/ LABA or ICS/LAMA/LABA product

Atopic Dermatitis

Dupixent® injection Initial PA Criteria:

- Patient is ≥ 6-months of age; **AND**
- Diagnosis of moderate to severe atopic dermatitis with ≥ 1 of the following:
  - Involvement of at least 10% of body surface area (BSA)
  - Scoring Atopic Dermatitis (SC**OR**AD) score of 20 or more
  - Investigator's Global Assessment (IGA) with a score ≥ 3
  - Eczema Area and Severity Index (EASI) score of ≥ 16
  - Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (documented by claims) or contraindication to both of the following:
  - A topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone)
  - A topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Dupixent® injection Renewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SCORAD)

Bullous Pemphigoid (BP)

Dupixent® injection Initial PA Criteria:

- Diagnosis of Bullous Pemphigoid (BP); **AND**
- Patient age ≥ 18 years; **AND**
- Inadequate response, intolerance, or contraindication to one of the following:
- A topical corticosteroid,
- Oral corticosteroid,
- Doxycycline; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Dupixent® injection Renewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SCORAD)

Chronic Spontaneous Urticaria (CSU) or Chronic Idiopathic Urticaria (CIU)

Dupixent® injection Initial PA Criteria:

- Patient is ≥ 2 years of age; **AND**
- Diagnosis of chronic spontaneous idiopathic urticaria (CSU) or chronic idiopathic urticaria (CIU); **AND**
- Patient remains symptomatic despite a 2-week trial to BOTH the following taken in combination:
    - A second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine, loratadine); **AND**
    - One of the following:
        - Different second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine)
        - First generation H1-antihistamine (e.g., diphenhydramine, chlorpheniramine, hydroxyzine)
        - H2-receptor antihistamine (e.g., famotidine, cimetidine, ranitidine)
        - Leukotriene modifier (e.g., montelukast); **AND**
- Prescribed by, or in consultation with, an allergist, dermatologist, or immunologist

Dupixent® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, itch severity, hives)

Chronic Rhinosinusitis with Nasal Polyposis (CRSwNP)

Dupixent® injection Initial PA Criteria:

- Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) confirmed by ONE of the following:
    - Presence of bilateral nasal polyps
    - Patient has previously required surgical removal of bilateral nasal polyps; **AND**
- Patient is ≥ 12 years of age; **AND**
- One of the following:
    - Patient has required prior sinus surgery; **OR**
    - Patient has required systemic corticosteroids for CRSwNP
    - Symptoms persist after trial of TWO of the following classes of agents:
        - Nasal saline irrigations
        - Intranasal corticosteroids
        - Antileukotriene agents; **AND**
- Must be used in combination with intranasal corticosteroid, unless contraindication or intolerance; **AND**
- Prescribed by, or in consultation with, an allergist, immunologist, otolaryngologist, or pulmonologist

Dupixent® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy; **AND**
- Will continue to use in combination with intranasal corticosteroids

Chronic Obstructive Pulmonary Disease (COPD)

Dupixent® injection Initial PA Criteria:

- Diagnosis of COPD; **AND**
- Patient is ≥ 18 years of age; **AND**
- Eosinophilic phenotype confirmed by peripheral blood eosinophil levels > 300 cells/mcL; **AND**
- COPD is inadequately controlled as shown by one of the following:
  - Two or more exacerbations requiring systemic corticosteroids and/or antibiotics within the past 12 months; **OR**
  - COPD-related emergency treatment (e.g., hospitalization in the past 12 months, mechanical ventilation); **AND**
- Patient is currently receiving standard of care COPD treatment, unless contraindicated (i.e., ICS/LAMA/LABA); **AND**
- Post-bronchodilator FEV1/FVC ratio <0.7 and FEV1 ≤ 79%; **AND**
- Medication will be used as maintenance add-on therapy

Dupixent® injection Renewal PA Criteria:

- Positive clinical response to treatment (e.g., improved FEV1 from baseline, reduction in COPD exacerbations); **AND**
- Patient is currently receiving standard of care COPD treatment, unless contraindicated (i.e., ICS/LAMA/LABA); **AND**
- Medication will be used as maintenance add-on therapy

Eosinophilic Esophagitis (EoE)

Dupixent® injection Initial PA Criteria:

- Diagnosis of eosinophilic esophagitis (EOE) as confirmed by an esophageal biopsy demonstrating ≥ 15 intraepithelial eosinophils per high power field (documentation required); **AND**
- Patient is ≥ 1 years of age and weighs at least 15 kg; **AND**
- Patient is experiencing symptoms related to esophageal dysfunction (e.g., dysphagia, food impaction, chest pain that may not respond to antacids, gastroesophageal reflux disease-like symptoms); **AND**
- Symptoms are inadequately controlled after a trial of ONE of the following:
  - Proton pump inhibitor (e.g., pantoprazole, omeprazole); **OR**
  - Topical esophageal corticosteroids (e.g., budesonide, fluticasone); **AND**
- Prescribed by, or in consultation with, a gastroenterologist, allergist, or immunologist

Dupixent® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., reduction of eosinophil count, improvement of symptoms)

Prurigo Nodularis (PN)

Dupixent® injection Initial PA Criteria:

- Diagnosis of Prurigo Nodularis (PN); **AND**
- Patient age ≥ 18 years; **AND**
- Patient has 20 or more nodular lesions (IGA PN-S ≥ 3); **AND**
- Inadequate response, intolerance, or contraindication to a topical steroid; **AND**
  - Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Dupixent® injection Renewal PA Criteria:

- Documentation of positive clinical response (e.g., improved symptoms or IGA PN-S)

Ebglyss® injection

PDL Status	P
Quantity Limit	Initial 6 months: 4 pens or syringes/28 days; Maintenance: 1 pen or syringe/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Ebglyss® injection Initial PA Criteria:

- Patient is ≥ 12 years of age; **AND**
- Diagnosis of moderate to severe atopic dermatitis with ≥ 1 of the following:
  - Involvement of at least 10% of body surface area (BSA)
  - Scoring Atopic Dermatitis (SCORAD) score of 20 or more
  - Investigator's Global Assessment (IGA) with a score ≥ 3
  - Eczema Area and Severity Index (EASI) score of ≥ 16
  - Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (documented by claims) or contraindication to both of the following:
  - A topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone)
  - A topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Ebglyss® injection Renewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SCORAD)

Fasenra® autoinjector

PDL Status	P
Quantity Limit	Initial (first 3 doses): 1/30 days; Maintenance: 1/56 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Asthma

Fasenra® autoinjector Initial PA Criteria:

- Diagnosis of moderate to severe asthma; **AND**
- Patient is ≥ 6 years old; **AND**
- One of the following:
  - Asthma is an eosinophilic phenotype as defined by a baseline (pre- treatment) peripheral blood eosinophil level ≥ 150 cells/µL or peripheral blood eosinophil levels > 300 cells/mcL; **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Asthma is inadequately controlled as shown by one of the following:
  - Two more exacerbations requiring systemic corticosteroids within the past 12 months; **OR**
  - Asthma-related emergency treatment (e.g., hospitalization, intensive care unit stay, or mechanical ventilation); **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Patient is currently being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both of the following:
    - One high dose inhaled corticosteroid (ICS)
    - One additional asthma controller medication [e.g., LABA, leukotriene receptor antagonist, theophylline]; **OR**
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product; **AND**
- Will be used as adjunct therapy along with above asthma treatment; **AND**
- Prescribed by, or in consultation with, a pulmonologist, allergist, or immunologist

Fasenra® autoinjector Renewal PA Criteria:

- Documentation of a positive clinical response (e.g., reduction in exacerbations, improvement of FEV1, decreased use of rescue medications); **AND**
- Patient is being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both a high-dose ICS and an additional asthma controller medication
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product

Eosinophilic Granulomatosis with Polyangiitis (EGPA)

Fasenra® autoinjector Initial PA Criteria:

- Diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA); **AND**
- Patient is ≥ 18 years of age; **AND**
- Patient is currently taking standard therapy [i.e., systemic glucocorticoids (e.g., prednisolone, prednisone) with or without immunosuppressive therapy (e.g., cyclophosphamide, rituximab)]; **AND**
- One of the following:
- Patient has relapsing disease (defined as a relapse requiring additional or dose escalation of corticosteroids or immunosuppressant, or hospitalization); **OR**
- Refractory disease as defined as failure to attain remission within the prior 6 months following induction treatment with standard therapy regimens; **AND**
- Patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy); **AND**
- Prescribed by, or in consultation with, a pulmonologist, rheumatologist, allergist, or immunologist

Fasenra® autoinjector Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., reduced frequency or severity of relapses, reduction or discontinuation of corticosteroids or immunosuppressants)

Hypereosinophilic Syndrome (HES)

Refer to Interim Criteria for Agents Awaiting PAC Review for this indication.

Nemluvio® injection

PDL Status	P
Quantity Limit	2 injections/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Atopic Dermatitis Initial: 6 months; Renewal: 1 year Prurigo Nodularis: 1 year

Atopic Dermatitis

Nemluvio® injection Initial PA Criteria:

- Patient is ≥ 12 years of age; **AND**
- Diagnosis of moderate to severe atopic dermatitis with ≥ 1 of the following:
  - Involvement of at least 10% of body surface area (BSA)
  - Scoring Atopic Dermatitis (SC**OR**AD) score of 20 or more
  - Investigator's Global Assessment (IGA) with a score ≥ 3
  - Eczema Area and Severity Index (EASI) score of ≥ 16
  - Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (documented by claims) or contraindication to both of the following:
  - A topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone)
  - A topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist

Nemluvio® injection Renewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SC**OR**AD)

Prurigo Nodularis (PN)

Nemluvio® injection Initial PA Criteria:

- Diagnosis of Prurigo Nodularis (PN); **AND**
- Patient age ≥ 18 years; **AND**
- Patient has 20 or more nodular lesions (IGA PN-S ≥ 3); **AND**
- Inadequate response, intolerance, or contraindication to a topical steroid; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist; **AND**
- Trial and failure, contraindication, or intolerance to Dupixent

Nemluvio® injection Renewal PA Criteria:

- Documentation of positive clinical response (e.g., improved symptoms or IGA PN-S)

Nucala® injection

PDL Status	P
Quantity Limit	3 pens or syringes/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Asthma

Nucala® injection Initial PA Criteria:

- Diagnosis of moderate to severe asthma; **AND**
- Patient is ≥ 6 years old; **AND**
- One of the following:
  - Asthma is an eosinophilic phenotype as defined by a baseline (pre- treatment) peripheral blood eosinophil level ≥ 150 cells/µL or peripheral blood eosinophil levels > 300 cells/mcL; **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Asthma is inadequately controlled as shown by one of the following:
  - Two more exacerbations requiring systemic corticosteroids within the past 12 months
  - Asthma-related emergency treatment (e.g., hospitalization, intensive care unit stay, or mechanical ventilation)
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Patient is currently being treated with ONE of the following, unless there is a contraindication:
- Combination therapy including both of the following:
    - One high dose inhaled corticosteroid (ICS)
    - One additional asthma controller medication [e.g., LABA, leukotriene receptor antagonist, theophylline]; **OR**
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product; **AND**
- Will be used as adjunct therapy along with above asthma treatment; **AND**
- Prescribed by, or in consultation with, a pulmonologist, allergist, or immunologist

Nucala® injection Renewal PA Criteria:

- Documentation of a positive clinical response (e.g., reduction in exacerbations, improvement of FEV1, decreased use of rescue medications); **AND**
- Patient is being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both a high-dose ICS and an additional asthma controller medication
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product

Chronic Obstructive Pulmonary Disease (COPD)

Nucala® injection Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Eosinophilic phenotype confirmed by peripheral blood eosinophil levels > 150 cells/mcL; **AND**
- COPD is inadequately controlled as shown by one of the following:
  - Two or more exacerbations requiring systemic corticosteroids and/or antibiotics within the past 12 months; **OR**
  - COPD-related emergency treatment (e.g., hospitalization in the past 12 months, mechanical ventilation); **AND**
- Patient is currently receiving standard of care COPD treatment, unless contraindicated (i.e., ICS/LAMA/LABA); **AND**
- Post-bronchodilator FEV1/FVC ratio <0.7 and FEV1 ≤ 79%; **AND**
- Medication will be used as maintenance add-on therapy

Nucala® injection Renewal PA Criteria:

- Positive clinical response to treatment (e.g., improved FEV1 from baseline, reduction in COPD exacerbations); **AND**
- Patient is currently receiving standard of care COPD treatment, unless contraindicated (i.e., ICS/LAMA/LABA); **AND**
- Medication will be used as maintenance add-on therapy

Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)

Nucala® injection Initial PA Criteria:

- Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) confirmed by ONE of the following:
  - Presence of bilateral nasal polyps
  - Patient has previously required surgical removal of bilateral nasal polyps; **AND**
- Patient is ≥ 18 years of age; **AND**
- One of the following:
  - Patient has required prior sinus surgery; **OR**
  - Patient has required systemic corticosteroids for CRSwNP; **OR**
  - Symptoms persist after trial of TWO of the following classes of agents:
    - Nasal saline irrigations
    - Intranasal corticosteroids
    - Antileukotriene agents; **AND**
- Must be used in combination with intranasal corticosteroid, unless contraindication or intolerance; **AND**
- Prescribed by, or in consultation with, an allergist, immunologist, otolaryngologist, or pulmonologist

Nucala® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy; **AND**
- Will continue to use in combination with intranasal corticosteroids

Eosinophilic Granulomatosis with Polyangiitis (EGPA)

Nucala® injection Initial PA Criteria:

- Diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA); **AND**
- Patient is ≥ 18 years of age; **AND**
- Patient is currently taking standard therapy [i.e., systemic glucocorticoids (e.g., prednisolone, prednisone) with or without immunosuppressive therapy (e.g., cyclophosphamide, rituximab)]; **AND**
- One of the following:
  - Patient has relapsing disease (defined as a relapse requiring additional or dose escalation of corticosteroids or immunosuppressant, or hospitalization); **OR**
  - Refractory disease as defined as failure to attain remission within the prior 6 months following induction treatment with standard therapy regimens; **AND**
- Patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy); **AND**
- Prescribed by, or in consultation with, a pulmonologist, rheumatologist, allergist, or immunologist

Nucala® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., reduced frequency or severity of relapses, reduction or discontinuation of corticosteroids or immunosuppressants)

Hypereosinophilic Syndrome (HES)

Nucala® injection Initial PA Criteria:

- Patient is > 12 years of age; **AND**
- Patient has had HES for > 6-months without an identifiable non-hematologic secondary cause (e.g., drug hypersensitivity, parasitic helminth infection, HIV infection, non-hematologic malignancy); **AND**
- Patient does not have FIP1L1-PDGFRα kinase-positive HES; **AND**
- Baseline (pre-Nucala treatment) blood eosinophil level ≥1000 cells/µL (documentation required); **AND**
- Patient is currently receiving a stable dose of background HES therapy (e.g., oral corticosteroid, immunosuppressor); **AND**
- Prescribed by, or in consultation with a pulmonologist, rheumatologist, allergist, or immunologist

Nucala® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy

Tezspire® injection

PDL Status	P
Quantity Limit	1 pen or syringe/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Asthma

Tezspire® injection Initial PA Criteria:

- Diagnosis of severe asthma; **AND**
- Patient is ≥ 12 years old; **AND**
- Patient has inadequately controlled asthma as shown by one of the following:
  - Two more exacerbations requiring systemic corticosteroids within the past 12 months; **OR**
  - Asthma-related emergency treatment (e.g., hospitalization, intensive care unit stay, or mechanical ventilation); **OR**
  - Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Patient is currently being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both of the following:
    - One medium or high dose inhaled corticosteroid (ICS); **AND**
    - One additional asthma controller medication [e.g., LABA, leukotriene antagonist, theophylline]; **OR**
  - One maximally dosed combination inhaled ICS/ LABA or ICS/LAMA/LABA product; **AND**
- Will be used as adjunct therapy along with above asthma treatment; **AND**
- Prescribed by, or in consultation with, a pulmonologist, allergist, or immunologist

Tezspire® injection Renewal PA Criteria:

- Documentation of a positive clinical response (e.g., reduction in exacerbations, improvement of FEV1, decreased use of rescue medications); **AND**
- Patient is being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both a high-dose ICS and an additional asthma controller medication
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product

Chronic Rhinosinusitis with Nasal Polyps:

Tezspire® injection Initial PA Criteria:

- Diagnosis of chronic rhinosinusitis with nasal polyps (CRSwNP) confirmed by ONE of the following:
  - Presence of bilateral nasal polyps;
  - Patient has previously required surgical removal of bilateral nasal polyps; **AND**
- Patient is ≥ 12 years old; **AND**
  - One of the following:
    - Patient has required prior sinus surgery; **OR**
    - Patient has required systemic corticosteroids for CRSwNP; **OR**
    - Symptoms persist after trial of TWO of the following classes of agents:
      - Nasal saline irrigations
      - Intranasal corticosteroids
      - Antileukotriene agents; **AND**
- Must be used in combination with a intranasal corticosteroid, unless contraindicated or intolerance; **AND**
- Prescribed by, or in consultation with, an allergist, immunologist, otolaryngologist, or pulmonologist

Tezspire® injection Renewal PA Criteria:

- Documentation of a positive clinical response to therapy; **AND**
- Will continue to use in combination with intranasal corticosteroids

Xolair® injection

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year for all indications except IgE-Mediated Food Allergy IgE-Mediated Food Allergy: 1 year

Asthma

Xolair® injection Initial PA Criteria:

- Diagnosis of moderate to severe persistent asthma; **AND**
- Patient is ≥ 6 years old; **AND**
- Dose requested is consistent with corresponding weight and IgE level per manufacturer's dosing chart; **AND**
- Baseline (pre-treatment) serum total IgE level greater than or equal to 30 IU/mL and less than or equal to 1300 IU/mL; **AND**
- Positive skin test or in vitro reactivity to a perennial aeroallergen; **AND**
- Patient has inadequately controlled asthma as shown by one of the following:
  - Two more exacerbations requiring systemic corticosteroids within the past 12 months; **OR**
  - Asthma-related emergency treatment (e.g., hospitalization, intensive care unit stay, or mechanical ventilation; **OR**
- Patient is currently dependent on oral corticosteroids for the treatment of asthma; **AND**
- Patient is currently being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both of the following:
    - One medium or high dose inhaled corticosteroid (ICS); **AND**
    - One additional asthma controller medication [e.g., LABA, leukotriene antagonist, theophylline]; **OR**
  - One maximally dosed combination inhaled ICS/ LABA or ICS/LAMA/LABA product; **AND**
- Will be used as adjunct therapy along with above asthma treatment; **AND**
- Prescribed by, or in consultation with, a pulmonologist, allergist, or immunologist

Xolair® injection Renewal PA Criteria:

- Documentation of a positive clinical response (e.g., reduction in exacerbations, improvement of FEV1, decreased use of rescue medications); **AND**
- Patient is being treated with ONE of the following, unless there is a contraindication:
  - Combination therapy including both a high-dose ICS and an additional asthma controller medication; **OR**
  - One maximally dosed combination ICS/ LABA or ICS/LAMA/LABA product

Chronic Idiopathic Urticaria (CIU)

Xolair® injection Initial PA Criteria:

- Patient is ≥ 12 years of age; **AND**
- Diagnosis of chronic spontaneous idiopathic urticaria (CSU) or chronic idiopathic urticaria (CIU); **AND**
- Patient remains symptomatic despite a 2-week trial to BOTH the following taken in combination:
  - A second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine, loratadine); **AND**
  - One of the following:
    - Different second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine)
    - First generation H1-antihistamine (e.g., diphenhydramine, chlorpheniramine, hydroxyzine)
    - H2-receptor antihistamine (e.g., famotidine, cimetidine, ranitidine)
    - Leukotriene modifier (e.g., montelukast); **AND**
- Prescribed by, or in consultation with, an allergist, dermatologist, or immunologist

Xolair® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, itch severity, hives)

IgE-Mediated Food Allergy

Xolair® injection Initial PA Criteria:

- Diagnosis of IgE-mediated food allergy confirmed by BOTH of the following:
  - History of type I allergic reactions; **AND**
  - Food specific skin prick testing (SPT) or IgE antibody in vitro testing; **AND**
- Xolair is to be used in combination with food allergen avoidance; **AND**
- Dose requested is consistent with corresponding weight and IgE level per manufacturer's dosing chart; **AND**
- Prescribed by, or in consultation with allergist or immunologist

Chronic rhinosinusitis with nasal polyps (CRSwNP)

Xolair® injection Initial PA Criteria:

- Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) confirmed by ONE of the following:
  - Presence of bilateral nasal polyps
  - Patient has previously required surgical removal of bilateral nasal polyps; **AND**
- Patient is ≥ 18 years of age; **AND**
- One of the following:
  - Patient has required prior sinus surgery; **OR**
  - Patient has required systemic corticosteroids for CRSwNP; **OR**
  - Symptoms persist after trial of TWO of the following classes of agents:
    - Nasal saline irrigations
    - Intranasal corticosteroids
    - Antileukotriene agents; **AND**
- Must be used in combination with intranasal corticosteroid, unless contraindication or intolerance; **AND**
- Prescribed by, or in consultation with, an allergist, immunologist, otolaryngologist, or pulmonologist

Xolair® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy; **AND**
- Will continue to use in combination with intranasal corticosteroids

Cibinqo® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Cibinqo® tablets Initial PA Criteria:

- Patient is ≥ 12 years of age; **AND**
- Diagnosis of moderate to severe atopic dermatitis with ≥ 1 of the following:
  - Involvement of at least 10% of body surface area (BSA)
  - Scoring Atopic Dermatitis (SCORAD) score of 20 or more
  - Investigator's Global Assessment (IGA) with a score ≥ 3
  - Eczema Area and Severity Index (EASI) score of ≥ 16
  - Incapacitation due to AD lesion location (e.g., head and neck, palms, soles, or genitalia); **AND**
- Trial and failure (documented by claims) or contraindication to both of the following:
  - A topical corticosteroid of medium to high potency (e.g., mometasone, fluocinolone)
  - A topical calcineurin inhibitor; **AND**
- Prescribed by, or in consultation with, a dermatologist, allergist, or immunologist
- Trial and failure, contraindication, or intolerance of a preferred agent indicated for atopic dermatitis (e.g., Adbry, Dupixent, Ebglyss, Nemluvio)

Cibinqo® tablets Renewal PA Criteria:

- Documented positive response to therapy (e.g., improved pruritus, BSA involvement, EASI, IGA, or SCORAD)

Rhapsido® tablets

PDL Status	NP
Quantity Limit	2 tablets/day
PA Criteria	See below
PA Form	General PA Form
PA Duration	Initial: 6 months; Renewal: 1 year

Rhapsido® tablets Initial PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of chronic spontaneous urticaria (CSU); **AND**
- Patient remains symptomatic despite a 2-week trial of BOTH the following taken in combination;
  - A second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine, loratadine); **AND**
  - One of the following:
    - Different second generation H1-antihistamine (e.g., cetirizine, fexofenadine, levocetirizine)
    - First generation H1-antihistamine (e.g., diphenhydramine, chlorpheniramine, hydroxyzine)
    - H2-receptor antihistamine (e.g., famotidine, cimetidine, ranitidine)
    - Leukotriene modifier (e.g., montelukast); **AND**
- Trial and failure, contraindication or intolerance to Dupixent or Xolair; **AND**
- Prescribed by, or in consultation with, an allergist, dermatologist, or immunologist

Rhapsido® tablets Renewal PA Criteria:

- Prescribed by, or in consultation with, an allergist, dermatologist, or immunologist; **AND**
- Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, itch severity, hives)

X. MISCELLANEOUS

Oral Iron Chelators

Deferasirox tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Deferasirox Initial PA Criteria:

- Patient has a diagnosis of ONE of the following:
  - Chronic iron overload due to blood transfusions in patients 2 years of age and older; **AND**
  - Non-transfusion-dependent thalassemia (NTDT) in patients aged 10 and older; **AND**
    - ONE of the following:
      - Serum ferritin > 300 mcg/L; **OR**
      - Liver iron concentration is >5 mgFe/g dw; **AND**
- Patient's platelet count is greater than $50x10^9$/L; **AND**
- Patient's creatinine clearance is greater than 40 mL/min; **AND**
- Patient does not have severe (Child-Pugh C) hepatic impairment

Deferasirox Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g. reduction or normalization of serum ferritin)

Deferiprone tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Deferiprone Initial PA Criteria:

- Patient has a diagnosis of ONE of the following:
  - Transfusional iron overload due to thalassemia syndromes regardless of prior chelation exposure
  - Transfusional iron overload in patients with sickle cell disease or other anemias; **AND**
- Patient is 8 years of age and up (tablets); **OR** 3 years of age and up (solution); **AND**
- ONE of the following:
  - Serum ferritin > 1,000 mcg/L
  - Liver iron concentration is > 3.2 Fe/g dw L; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects to all preferred medications

Deferasirox granule packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Deferasirox granule packets PA Criteria:

See deferasirox tablets prior authorization criteria; **AND**

- Clinically valid reason as to why patient cannot use deferasirox tabs

Deferasirox tablets for suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Deferasirox tablets for suspension Initial PA Criteria:

- Patient has a diagnosis of ONE of the following:
  - Chronic iron overload due to blood transfusions in patients 2 years of age and older; **AND**
  - Non-transfusion-dependent thalassemia (NTDT) in patients aged 10 and older; **AND**
    - ONE of the following:
      - Serum ferritin > 300 mcg/L; **OR**
      - Liver iron concentration is >5 mgFe/g dw; **AND**
- Patient's platelet count is greater than $50x10^9$/L; **AND**
- Patient's creatinine clearance is greater than 40 mL/min; **AND**
- Patient does not have severe (Child-Pugh C) hepatic impairment; **AND**
- Patient is unable to swallow solid dosage forms

Deferasirox tablets for suspension Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g. reduction or normalization of serum ferritin)

Exjade® tablets for suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Exjade® tablets for suspension Initial PA Criteria:

- Patient has a diagnosis of ONE of the following:
  - Chronic iron overload due to blood transfusions in patients 2 years of age and older; **AND**
  - Non-transfusion-dependent thalassemia (NTDT) in patients aged 10 and older; **AND**
    - ONE of the following:
      - Serum ferritin > 300 mcg/L; **OR**
      - Liver iron concentration is >5 mgFe/g dw; **AND**
- Patient's platelet count is greater than $50x10^9$/L; **AND**
- Patient's Creatinine clearance is greater than 40 mL/min; **AND**
- Patient does not have severe (Child-Pugh C) hepatic impairment
- Trial and failure, contraindication, or intolerance to one of the preferred products

Exjade® tablets for suspension Renewal PA Criteria:

- Patient has a positive clinical response to therapy (e.g. reduction or normalization of serum ferritin)

Ferriprox® solution, tablets, Ferriprox Twice-a-Day®

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Ferriprox® solution, tablets, Ferriprox Twice-a-Day® Initial PA Criteria:

- Patient has a diagnosis of ONE of the following:
  - Transfusional iron overload due to thalassemia syndromes regardless of prior chelation exposure
  - Transfusional iron overload in patients with sickle cell disease or other anemias; **AND**
- Patient is 8 years of age and up (tablets); **OR** 3 years of age and up (solution); **AND**
- ONE of the following:
  - Serum ferritin > 1,000 mcg/L
  - Liver iron concentration is > 3.2 Fe/g dw L; **AND**
- Absolute Neutrophil Count (ANC) has been measured prior to starting therapy; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects to all preferred medications

Ferriprox® solution, tablets, Ferriprox Twice-a-Day® Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g. reduction or normalization of serum ferritin)

Jadenu® sprinkle granules

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Jadenu® sprinkle granules Initial PA Criteria:

See deferasirox tablets prior authorization criteria; **AND**

- Clinically valid reason as to why patient cannot use deferasirox tabs

Jadenu® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Jadenu® tablets Initial PA Criteria:

See Exjade® tablets for suspension prior authorization criteria.

Oral Iron Supplements

Accrufer®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Accrufer® PA Criteria:

- Patient has iron deficiency; **AND**
- Patient is 10 years of age or older; **AND**
- Patient must NOT meet any of the following:
  - Hemochromatosis and other iron overload syndromes
  - Receiving repeated blood transfusions or intravenous iron supplementation
  - Irritable bowel disease (IBD) flare
  - Concomitant use of dimercaprol

Saliva Stimulating Agents

Pilocarpine

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Cevimeline, Evoxac®

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Cevimeline, Evoxac® PA Criteria:

- Trial and failure, contraindication, or intolerance of pilocarpine

XI. ONCOLOGY AGENTS

Acute Myeloid Leukemia (AML) Agents

Venclexta® tablets

PDL Status	P
Quantity Limit	Ramp-Up Phase Dosing: Dispense 7-day supply of 10 mg tabs (for 20 mg dose); followed by 7-day supply of 50 mg tabs 100 mg tabs: 4/day; Starter Pack: 1 pack/year
PA Criteria	None

Daurismo® tablets

PDL Status	NP
Quantity Limit	25 mg: 84/28 days; 100 mg: 28/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Daurismo® tablets Initial PA Criteria:

- Newly diagnosed acute myeloid leukemia (AML); **AND**
- ONE of the following:
  - Patient ≥ 75 years of age
  - Patient has comorbidities that preclude the use of intensive induction chemotherapy (i.e., Severe Cardiac Disease, Baseline serum creatinine > 1.3 mg/dL, or Baseline Eastern Cooperative Oncology Group (ECOG) performance status of ≥ 2); **AND**
- Women of child-bearing potential must have a negative pregnancy test; **AND**
- Female patients of reproductive potential and males undergoing treatment with female partners of reproductive should use effective contraception during treatment and for at least 30 days after treatment; **AND**
- Daurismo® will be used in conjunction with low-dose subcutaneous cytarabine

Daurismo® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., CPK elevation, QTc-interval prolongation, low platelet or neutrophil counts, grade 4 nonhematologic toxicity)

Onureg® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Onureg® tablets Initial PA Criteria:

- Diagnosis of acute myeloid leukemia; **AND**
- Patient has achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy; **AND**
- Female patients of reproductive potential have a negative pregnancy test and have been advised that:
  - Female patients should use effective contraception during treatment and for at least 6-months after treatment
  - Males undergoing treatment with female partners of reproductive age should use effective contraception during treatment and for at least 3 months after treatment due to male mediated teratogenicity

Onureg® tablets Renewal PA Criteria:

- Patient must continue to meet the initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., myelosuppression, renal impairment, hepatic impairment)

Vanflyta® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Criteria

Vanflyta® tablets Initial PA Criteria:

- Patient has newly diagnosed acute myeloid leukemia (AML); **AND**
- AML is FLT3 internal tandem duplication (ITD)-positive as detected by an FDA-approved test; **AND**
- Vanflyta will be used in combination with cytarabine and anthracycline induction and high dose cytarabine consolidation therapy followed by maintenance therapy; **AND**
- Vanflyta will not be used as maintenance monotherapy following allogeneic hematopoietic stem cell transplantation (HSCT); **AND**
- Patient and prescriber are enrolled in the Vanflyta REMS program; **AND**
- One of the following:
  - Female patients of reproductive potential should use effective contraception during treatment and for 7 months after last dose.
  - Males with female partners of reproductive potential should use effective contraception during treatment and for 4 months after the last dose

Vanflyta® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., hypokalemia, hypomagnesemia, long QT syndrome)

Xospata® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Xospata® tablets Initial PA Criteria:

- Patient has a diagnosis of acute myeloid leukemia (AML); **AND**
- AML is positive for FLT3 mutation as detected by an FDA-approved; **AND**
- Females of child-bearing potential had a negative pregnancy test within 7 days before starting Xospata®; **AND**
- Female and male patients of reproductive potential have been advised to use effective contraception during treatment and for at least 6 and 4 months, respectively, after the last dose

Xospata® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., QT prolongation, differentiation syndrome, encephalopathy, pancreatitis)

Alkylating Agents

Lomustine capsules

PDL Status	P
Quantity Limit	1 capsule/Rx
PA Criteria	None

Antimetabolites

Inqovi® tablets

PDL Status	NP
Quantity Limit	5/28-day cycle
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 3 months; Renewal:3 months

Inqovi® tablets Initial PA Criteria:

- One of the following:
  - Patient has tried and failed or is not a candidate for Allogenic stem cell transplantation AND Diagnosis of myelodysplastic syndromes (MDS), patients previously treated and untreated, de novo and secondary MDS with the following French American-British subtypes:
    - Refractory anemia
    - Refractory anemia with ringed sideroblasts
    - Refractory anemia with excess blasts
    - Chronic myelomonocytic leukemia [CMML])
    - Intermediate-1, intermediate-2, and high-risk international prognostic IPSS groups; **OR**
  - Patient is newly diagnosed with Acute Myeloid Leukemia (AML); **AND**
    - Patient is using in combination with venetoclax; **AND**
    - ONE of the following:
      - Patient is 75yrs of age or older;
      - Patient has co-morbidities that prevent use of intensive induction chemotherapy; **AND**
- Prescriber will obtain baseline CBC, creatinine clearance (CrCl), and liver enzymes prior to therapy and prior to each cycle; **AND**
- Patient must not be pregnant or breastfeeding; **AND**
- Female patients should use effective contraception during treatment and for at least 6-months after treatment; **AND**
- Males undergoing treatment with female partners of reproductive age should use effective contraception during treatment and 3 months after treatment due to male mediated teratogenicity; **AND**
- Will not be used concomitantly with drugs metabolized by cytidine deaminase enzyme (i.e., gemcitabine, capecitabine, cytarabine, azacytidine)

Inqovi® tablets Renewal PA Criteria:

- Continues to meet initial criteria; **AND**
- Patient has positive disease response, defined as disease stabilization; **AND**
- Prescriber attests to delay next cycle and reduce dose if patient experiences elevated liver enzymes or renal impairment **OR** if patient's absolute neutrophil count (ANC) is less than 1,000 cells/microL and platelet count is less than 50,000 cell/microL

Purixan® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Purixan® suspension PA Criteria:

- Diagnosis of acute lymphocytic leukemia (ALL); **AND**
- ONE of the following:
  - For patients ≤ 11 years of age, no prior authorization required
  - For patients > 11 years of age, Purixan will be approved for patients unable to swallow tablets

Colorectal Cancer Agents, Miscellaneous

Lonsurf® tablets

PDL Status	P
Quantity Limit	8/day
PA Criteria	None

Fruzaqla® capsules

PDL Status	NP
Quantity Limit	5 mg: 21/28 days; 1 mg: 84/28 days
PA Criteria	See below
PA Form	General PA Form

Fruzaqla® capsules Initial PA Criteria:

- Diagnosis of metastatic colorectal cancer; **AND**
- Patient has tried and failed, contraindication, or intolerance to ALL of the following chemotherapy-based regimens:
    - Fluoropyrimidine,
    - Oxaliplatin
    - Irinotecan
    - Anti- vascular endothelial growth factor (VEGF) therapy (e.g., bevacizumab); **AND**
- If RAS wild-type, patient has tried and failed, contraindication, or intolerance to anti-epidermal growth factor receptor (EGFR) therapy (e.g., cetuximab, panitumumab); **AND**
- Prescribed by or in consultation with an oncologist

Fruzaqla® capsulesRenewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hypertension, hemorrhagic events)

EGFR Inhibitors

Gefitinib tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Gefitinib tablets PA Criteria:

- Clinically valid reason why brand Iressa® cannot be used

Lazcluze® tablets

PDL Status	NP
Quantity Limit	80 mg: 2/day; 240 mg: 1/day
PA Criteria	See below
PA Form	General PA Form

Lazcluze® tablets Initial PA Criteria

- Diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC); **AND**
- Disease is positive for epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations as detected by an FDA-approved test; **AND**
- Lazcluze will be used in combination with Rybrevant; **AND**
- Prescribed by, or in consultation with, an oncologist

Lazcluze® tablets Renewal PA Criteria

- Patient continues to meet the initial criteria; **AND**
- Patient does not have unacceptable toxicity (e.g., interstitial lung disease, keratitis, venous thromboembolic events)

Tarceva® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tarceva® PA Criteria:
Standard NP Criteria

Vizimpro® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Vizimpro® tablets PA Criteria:

- Patient has a diagnosis of metastatic non-small cell lung cancer (NSCLC); **AND**
- Disease is positive for EGFR mutations as confirmed by FDA approved Test (e.g., cobas® EGFR Mutation Test v2); **AND**
- Requested agent will be prescribed by, or in consultation with, an oncologist

Enzyme Inhibitors: ALK

Ensacove® capsules

PDL Status	P
Quantity Limit	25 mg caps: 2/day; 100 mg caps: 2/day
PA Criteria	None

Lorbrena® tablets

PDL Status	P
Quantity Limit	25 mg: 3/day; 100 mg: 1/day
PA Criteria	None

Xalkori Sprinkles®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Xalkori Sprinkles® PA Criteria:

- Patient is unable to swallow solid dosage forms.

Enzyme Inhibitors: BCR-ABL Kinase

Bosulif® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Bosutinib tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Bosutinib tablets PA Criteria:

- Clinically valid reason why preferred Bosulif® cannot be used

Danziten® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Danziten® tablets Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Adult patient with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase; **OR**
  - Adult patient with chronic phase and accelerated phase Ph+ CML; **AND**
    - Patient must be resistant to or have contraindication or intolerant to imatinib; **AND**
- Prescribed by, or in consultation with, an oncologist;

Danziten® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., QTc prolongation, edema, hepatic or renal toxicity)

Gleevec® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Gleevec® tablets PA Criteria:

- Clinically valid reason why generic imatinib cannot be used

Imkeldi® solution

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See below
PA Form	General PA Form

Imkeldi® solution PA Criteria:

- Patient is <8 years old; **OR**
- Patient is unable to swallow oral dosage forms

Nilotinib capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Nilotinib capsules PA Criteria:

- Clinically valid reason why preferred Tasigna® cannot be used

Phyrago® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Phyrago® tablets PA Criteria:

- Clinically valid reason why preferred dasatinib cannot be used

Scemblix®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Scemblix® PA Criteria:

- Patient has ONE of the following:
- Newly diagnosed Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase; **OR**
- Philadelphia chromosome-positive CML in chronic phase (Ph+ CML-CP) previously treated with two or more tyrosine kinase inhibitors (TKIs); **OR**
- Ph+ CML-CP with the T315I mutation; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Females of reproductive potential will use effective contraception during treatment and for 1 week after the last dose; **AND**
- Patient will not breastfeed during treatment and for 1 week after the last dose

Sprycel® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sprycel® tablets PA Criteria:

- Clinically valid reason why preferred dasatinib cannot be used

Enzyme Inhibitors: BRAF Kinase & MEK

Braftovi® capsules

PDL Status	P
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Braftovi® capsules Initial PA Criteria:

- Prescribed by, or in consultation with, an oncologist; **AND**
- One of the following:
  - Diagnosis of unresectable or metastatic melanoma; **AND**
    - Patient is positive for BRAF V600E or V600K mutation as confirmed by an FDA-approved test; **AND**
    - Prescribed in combination with Mektovi®
  - Diagnosis of metastatic colorectal cancer (CRC); **AND**
    - Cancer is positive for BRAF V600E mutation as confirmed by an FDA-approved test after prior therapy
  - Diagnosis of metastatic non-small cell lung cancer (NSCLC);
    - Cancer is positive for BRAF V600E mutation, as detected by an FDA-approved test; **AND**
    - Prescribed in combination with Mektovi®

Braftovi® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- No unacceptable disease progression or unacceptable toxicity

Mektovi® tablets

PDL Status	P
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Mektovi® tablets Interim PA Criteria:

- Prescribed by, or in consultation with, an oncologist; **AND**
- Prescribed in combination with Braftovi®; **AND**
- One of the following:
  - Diagnosis of unresectable or metastatic melanoma; **AND**
    - Patient is positive for BRAF V600E or V600K mutation as confirmed by an FDA-approved test; **AND**
  - Diagnosis of metastatic non-small cell lung cancer (NSCLC); **AND**
    - Cancer is positive for BRAF V600E mutation as detected by an FDA-approved test; **AND**

Mektovi® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- No unacceptable disease progression or unacceptable toxicity

Gomekli® capsules, tablets for oral suspension

PDL Status	NP
Quantity Limit	1 mg cap: 6/day; 2 mg cap: 4/day; 1 mg susp tab: 8/day
PA Criteria	See below
PA Form	General PA Form

Gomekli® capsules, tablets for oral suspension Initial PA Criteria:

- Diagnosis of neurofibromatosis type 1 (NF1); **AND**
- Patient has plexiform neurofibromas that are BOTH of the following:
  - Not amenable to complete resection
  - Causing significant morbidity (e.g., disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment); **AND**
- Prescribed by, or in consultation with, an oncologist or neurologist; **AND**
- If the request is for the tablets for oral suspension, patient is unable to swallow solid dosage forms

Gomekli® capsules, tablets for oral suspension Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests that patient has experienced improvement in disease severity and/or symptoms; **AND**
- Patient does not have unacceptable toxicity (e.g., retinal pigment epithelial detachment, ventricular dysfunction, rash)

Koselugo® capsules

PDL Status	NP
Quantity Limit	10 mg: 10/day; 25 mg: 4/day
PA Criteria	See below
PA Form	General PA Form

Koselugo® capsules Initial PA Criteria:

- Diagnosis of neurofibromatosis type 1 (NF1); **AND**
- Patient has symptomatic, inoperable plexiform neurofibromas (PN); **AND**
- Prescribed by, or in consultation with, an oncologist or neurologist

Koselugo® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests that patient has experienced improvement in disease severity and/or symptoms; **AND**
- Patient does not have unacceptable toxicity (e.g., retinal pigment epithelial detachment, severe diarrhea, rash, increased bleeding, myalgia)

Koselugo® Sprinkle Capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Koselugo® Sprinkle Capsules Initial PA Criteria:

- Diagnosis of neurofibromatosis type 1 (NF1); **AND**
- Patient has symptomatic, inoperable plexiform neurofibromas (PN); **AND**
- Prescribed by, or in consultation with, an oncologist or neurologist; **AND**
- Patient is unable to swallow solid dosage forms

Koselugo® Sprinkle Capsules Initial PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests that patient has experienced improvement in disease severity and/or symptoms; **AND**
- Patient does not have unacceptable toxicity (e.g., retinal pigment epithelial detachment, severe diarrhea, rash, increased bleeding, myalgia); **OR**
- Patient is NOT experiencing an adverse reaction or hepatic impairment that requires a dose modification or reduction

Mekinist® solution

PDL Status	NP
Quantity Limit	
PA Criteria	See below
PA Form	General PA Form

Mekinist® solution PA Criteria:

- Patient is <8 years old; **OR**
- Patient is unable to swallow solid dosage forms

Ojemda® suspension, tablets

PDL Status	NP
Quantity Limit	24/28 days
PA Criteria	See below
PA Form	General PA Form

Ojemda® suspension, tablets Initial PA Criteria:

- Diagnosis of relapsed or refractory pediatric low-grade glioma (PLGG); **AND**
- Patient has BRAF fusion or rearrangement or BRAF V600 mutation; **AND**
- Prescribed by, or in consultation with, an oncologist

Ojemda® suspension, tablets Renewal PA Criteria:

- Patient demonstrates disease stabilization or clinical response to therapy (e.g., stabilized or decrease tumor size, decreased pain, improved vision, increased quality of life)

Tafinlar® tablets for oral suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Tafinlar® tablets for oral suspension PA Criteria:

- Patient is <8 years old; **OR**
- Patient is unable to swallow solid dosage forms

Enzyme Inhibitors: BTK

Brukinsa® capsules, tablets

PDL Status	P
Quantity Limit	Capsules: 4/day; Tablets: 2/day
PA Criteria	None

Calquence® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Imbruvica® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Imbruvica® suspension PA Criteria:

- Patient is unable to swallow capsules or tablets

Jaypirca® tablets

PDL Status	NP
Quantity Limit	50 mg: 1/day; 100 mg: 2/day
PA Criteria	See below
PA Form	General PA Form

Jaypirca® tablets Initial PA Criteria:

- One of the following:
  - Diagnosis of relapsed or refractory mantle cell lymphoma (MCL); AND
    - Patient has received two prior therapies including a BTK inhibitor (e.g., Ibrutinib, acalabrutinib, zanubrutinib); OR
  - Diagnosis of refractory or relapsed chronic lymphocytic leukemia (CLL) or small cell lymphocytic lymphoma (SLL); AND
    - Patient has received ONE prior therapy including a BTK inhibitor (e.g., Ibrutinib, acalabrutinib, zanubrutinib)

Jaypirca® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Absence of unacceptable toxicity from Jaypirca (e.g., hemorrhage, severe infections, myelosuppression (neutropenia, thrombocytopenia, anemia), atrial fibrillation/flutter, second primary malignancies); **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread

Enzyme Inhibitors: CDK

Kisqali® tablets

PDL Status	P
Quantity Limit	63 tabs/28 days
PA Criteria	None

Enzyme Inhibitors: FGFR

Balversa® tablets

PDL Status	NP
Quantity Limit	3 mg: 3/day; 4 mg: 2/day; 5 mg: 1/day
PA Criteria	See below
PA Form	General PA Form

Balversa® tablets Initial PA Criteria:

- Patient has a diagnosis of locally advanced or metastatic urothelial carcinoma; **AND**
- Patient has a susceptible FGFR3 or FGFR2 genetic alteration as confirmed by an FDA-approved diagnostic; **AND**
- Patient has progressed during or following ≥ 1 prior line of platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Provider attests to ALL the following:
    - Patient has received a baseline ophthalmological examination (e.g., assessment of visual acuity, slit lamp examination, fundoscopy, and optical coherence tomography)
    - Patient has had a baseline serum phosphate level measurement and it is within normal limits
    - Patient phosphate intake is restricted to < 800 mg per day
    - Patient will not concomitantly take the requested agent with a strong CYP2C9 or CYP3A4 inhibitors (e.g., fluconazole, itraconazole) or with strong CYP2C9 or CYP3A4 inducers (e.g., rifampicin) or, if therapy is unavoidable, prescriber attestation that the patient will be monitored for adverse reactions

Balversa® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., central serous retinopathy/retinal pigment epithelial detachment (CSR/RPED), severe hyperphosphatemia)

Lytgobi® tablets

PDL Status	NP
Quantity Limit	12 mg: 84/month; 16 mg: 112/month; 20 mg: 140/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Lytgobi® tablets Initial PA Criteria:

- Patient has diagnosis of unresectable, locally advanced, or metastatic intrahepatic cholangiocarcinoma; **AND**
- Patient has fibroblast growth factor receptor 2 (FGFR2) gene fusions or other rearrangements as detected by
- FDA approved test; **AND**
- The patient has progressed on at least one systemic therapy; **AND**
- The prescriber attest to ALL of the following:
  - Patient will have an ophthalmological examination including optical coherence tomography (OCT)
  - performed prior to initiation of therapy, every 2 months for the first 6 months of treatment and every 3
  - months thereafter, and urgently at any time for visual symptoms
  - Prescriber will obtain baseline phosphate levels and monitor for hyperphosphatemia throughout treatment
  - Patient is not pregnant
  - Female patients of reproductive potential and males with female partners of reproductive age have been advised to use effective contraception during treatment and for at least 1 week after the last dose
  - Patient is not concomitantly taking strong dual P-gp and CYP3A Inducers (e.g. rifampin)

Lytgobi® tablets Renewal PA Criteria:

- Positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., retinal pigment epithelial detachment, severe hyperphosphatemia)

Pemazyre® tablets

PDL Status	NP
Quantity Limit	14 tabs/21 days
PA Criteria	See below
PA Form	General PA Form

Pemazyre® tablets Initial PA Criteria:

- One of the following:
  - Diagnosis of previously treated unresectable locally advanced or metastatic cholangiocarcinoma with a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test
  - Diagnosis of relapsed or refractory myeloid/lymphoid neoplasms (MLNs) with FGFR1 rearrangement; **AND**
- Prescriber attests to ALL the following:
  - Patient will have an ophthalmological examination including optical coherence tomography (OCT) performed prior to initiation of therapy, every 2 months for the first 6 months of treatment and every 3 months thereafter, and urgently at any time for visual symptoms
  - Prescriber will obtain baseline phosphate levels and monitoring for hyperphosphatemia
  - Females and males with female partners will be advised to use effective contraception during treatment and for 1 week after the final dose due to embryo-fetal toxicity
  - Patient is not concomitantly taking strong and moderate CYP3A Inducers

Pemazyre® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., retinal pigment epithelial detachment, severe hyperphosphatemia)

Enzyme Inhibitors: HER2 Targeted Therapies

Hernexeos® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Hernexeos® tablets Initial PA Criteria:

- Diagnosis of unresectable or metastatic non-squamous non-small cell lung cancer (NCSLC); **AND**
- Patient is positive for HER2 (ERBB2) tyrosine kinase domain activating mutations as detected by an FDA approved test; **AND**
- Patient is 18 years of age or older; **AND**
- Prescribed by, or in consultation with an oncologist

Hernexeos® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hepatotoxicity, left ventricular dysfunction, interstitial lung disease/pneumonitis)

Hyrnuo® tablets

PDL Status	NP
Quantity Limit	4 tablets/day
PA Criteria	See below
PA Form	General PA Form

Hyrnuo ® tablets PA Criteria:

- Diagnosis of unresectable or metastatic non-squamous non-small cell lung cancer (NCSLC); **AND**
- Patient is positive for HER2 (ERBB2) tyrosine kinase domain activating mutations as detected by an FDA approved test; **AND**
- Patient is 18 years of age or older; **AND**
- Patient has documented disease progression on at least one prior systemic therapy (e.g., platinum and Carboplatin based chemotherapy); **AND**
- Prescribed by, or in consultation with, an oncologist

Hyrnuo® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hepatotoxicity, left ventricular dysfunction, interstitial lung disease/pneumonitis)

Lapatinib tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lapatinib tablets Initial PA Criteria:

Standard NP criteria

Tukysa® tablets

PDL Status	NP
Quantity Limit	50 mg: 10/day; 150 mg: 4/day
PA Criteria	See below
PA Form	General PA Form

Tukysa® Initial PA Criteria:

- ONE of the following:
  - Diagnosis of advanced unresectable or metastatic HER2-positive breast cancer and both of the following:
    - Patient has received at least one or more prior anti-HER2 based regimen;
    - Must be used in combination with trastuzumab and capecitabine; **OR**
  - Diagnosis of RAS wild-type, HER2-positive unresectable or metastatic colorectal cancer and both of the following:
    - Patient has documented disease progression on at least one prior systemic therapy (e.g., fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy)
    - Must be used in combination with trastuzumab; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Prescriber attests to ALL of the following:
  - Patient has baseline ALT, AST, and bilirubin measured and within normal limits
  - Patient continues to receive ALT/AST and bilirubin monitoring every 3 weeks during treatment

Tukysa® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., diarrhea, hepatotoxicity)

Enzyme Inhibitors: Isocitrate Dehydrogenase (IDH)

Tibsovo®

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Rezlidhia®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Rezlidhia® Initial PA Criteria:

- Patient has diagnosis of relapsed or refractory acute myeloid leukemia (AML); **AND**
- Patient has an isocitrate dehydrogenase-1 (IDH1) mutation, as detected by an FDA-approved test

Rezlidhia® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., hepatoxicity, differentiation syndrome)

Voranigo®

PDL Status	NP
Quantity Limit	10 mg: 2/day; 40 mg: 1/day
PA Criteria	See below
PA Form	General PA Form

Voranigo® Initial PA Criteria:

- Diagnosis of grade 2 astrocytoma or oligodendroglioma; **AND**
- Tumor has susceptible isocitrate dehydrogenase-1 (IDH1) or isocitrate dehydrogenase-2 (IDH2) mutation; **AND**
- Patient has had prior surgery including biopsy, sub-total resection, or gross total resection; **AND**
- Prescribed by, or in consultation with, an oncologist

Voranigo® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., hepatoxicity)

Enzyme Inhibitors: KRAS

Krazati® tablets

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Krazati® tablets PA Criteria:

- Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC) as confirmed by an FDA-approved test; **AND**
  - Patient has received at least one prior systemic therapy (e.g., immune checkpoint inhibitors [anti- PD-1, PD-L1 immunotherapy], platinum-based chemotherapy); **AND**
  - Patient has at least one measurable lesion as defined by Response Evaluation Criteria in Solid Tumors (RECIST v1.1); **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Prescriber attests that patient is not pregnant or breastfeeding during treatment and for 1 week after the final dose; **AND**
- Prescriber attests that patient will be monitored for the following:
  - Hepatotoxicity: Monitor liver function tests (ALT, AST, and total bilirubin) prior to the start of Krazati, every 3 weeks for the first 3 months of treatment then once monthly as clinically indicated
  - Interstitial Lung Disease (ILD)/Pneumonitis: Monitor for new or worsening pulmonary symptoms; **AND**
- Prescriber attests that Patient will not take Krazati with:
  - Acid-reducing agents (e.g., proton pump inhibitors, H_2 receptor antagonists, antacids, etc.)
  - Strong CYP3A4 inducers (e.g., rifampin, carbamazepine, etc.)

Lumakras® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lumakras® tablets PA Criteria:

- ONE of the following:
  - Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC) as confirmed by an FDA-approved test for detection of KRAS G12C; **AND**
    - Patient has at least one measurable lesion as defined by Response Evaluation Criteria in Solid Tumors (RECIST v1.1); **AND**
  - Diagnosis of KRAS G12C-mutated locally advanced or metastatic colorectal cancer (CRC) as confirmed by an FDA-approved test; **AND**
    - Trial and failure, contraindication, or intolerance to ALL the following chemotherapy-based regimens:
      - Fluoropyrimidine
      - Oxaliplatin
      - Irinotecan; **AND**
    - Agent will be used in combination with panitumumab; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient has received at least one prior systemic therapy (e.g., immune checkpoint inhibitors [anti- PD-1, PD-L1 immunotherapy], platinum-based chemotherapy); **AND**
- Prescriber attests that patient is not pregnant or breastfeeding during treatment with Lumakras and for 1 week after the final dose; **AND**
- Prescriber attests that Patient will be monitored for the following:
  - Hepatotoxicity: Liver function tests ((ALT, AST, and total bilirubin) prior to the start of Lumakras, every 3 weeks for the first 3 months of treatment then once monthly as clinically indicated
  - Interstitial Lung Disease (ILD)/Pneumonitis: New or worsening pulmonary symptoms; **AND**
- Prescriber attests that the patient will not take Lumakras with:
  - Acid-reducing agents (e.g., proton pump inhibitors, H_2 receptor antagonists, antacids.)
  - Strong CYP3A4 inducers (e.g., rifampin, carbamazepine)

Enzyme Inhibitors: MEK/FAK Kinase Inhibitors

AVMAPKI/FAKZYNJA Co-Pack®

PDL Status	NP
Quantity Limit	1 co-pack (66 tablets)/28 days
PA Criteria	See below
PA Form	General PA Form

AVMAPKI/FAKZYNJA Co-Pack® Initial PA Criteria:

- Diagnosis of recurrent low-grade serous ovarian cancer; **AND**
- Tumor is KRAS-mutated; **AND**
- Patient has received at least one prior *guideline-recommended* systemic therapy (e.g., *Paclitaxel/carboplatin plus maintenance letrozole or other hormonal therapy; paclitaxel/carboplatin/bevacizumab plus maintenance bevacizumab; or hormone therapy)*; **AND**
- Prescribed by, or in consultation with, an oncologist

AVMAPKI/FAKZYNJA Co-Pack® Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. ocular toxicities, serious skin toxicities, hepatotoxicity, rhabdomyolysis, and embryo-fetal toxicity)

Enzyme Inhibitors: Menin

Komzifti® capsules

PDL Status	NP
Quantity Limit	3 capsules/day
PA Criteria	See below
PA Form	General PA Form
PA Duration	Initial: 6 months; Renewal: 1 year

Komzifti® capsules Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of relapsed or refractory acute myeloid leukemia (AML); **AND**
- Disease is positive for Nucleophosmin 1 (NPM1) mutation; **AND**
- Patient is refractory to standard treatment option (cytarabine plus daunorubicin or idarubicin, venetoclax plus azacitidine or decitabine, etc.); **AND**
- Prescribed by, or in consultation with, an oncologist or hematologist

Komzifti® capsules Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., differentiation syndrome, QTc-interval prolongation)

Revuforj® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Revuforj® tablets Initial PA Criteria:

- Diagnosis of relapsed or refractory acute leukemia (e.g. acute myeloid leukemia (AML), Acute lymphoblastic leukemia (ALL), or Mixed phenotype acute leukemia (MPAL)); **AND**
- One of the following:
  - Disease is positive for lysine methyltransferase 2A gene (KMT2A) translocation; **OR**
  - Disease is positive for nucleophosmin 1 (NPM1) mutation; **AND**
- Patient is refractory to standard treatment option (cytarabine plus daunorubicin or idarubicin, venetoclax plus azacitidine or decitabine, etc.); **AND**
- Prescribed by, or in consultation with, an oncologist or hematologist

Revuforj® tablets Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., differentiation syndrome, QTc-interval prolongation)

Enzyme Inhibitors: MET

Tabrecta®

PDL Status	P
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Tabrecta® Initial PA Criteria:

- Patient must have metastatic non-small cell lung cancer (NSCLC); **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient must have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping in tumor specimens as confirmed by an FDA-approved test; **AND**

Tabrecta® Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., interstitial lung disease, liver enzymes outside of normal limits)

Tepmetko®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Tepmetko® Initial PA Criteria:

- Diagnosis of metastatic non-small cell lung cancer (NSCLC); **AND**
- Disease is harboring mesenchymal-epithelial transition (MET) exon 14 skipping alterations; **AND**
- Prescribed by, or in consultation with, an oncologist

Tepmetko® Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. interstitial lung disease, liver enzymes outside of normal limits)

Enzyme Inhibitors: MTOR

Afinitor Disperz®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Afinitor Disperz® PA Criteria:

- Patient is unable to swallow solid dosage forms

Afinitor® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Afinitor® tablets PA Criteria:

Standard NP Criteria

Everolimus dispersible tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Everolimus dispersible tablets PA Criteria:

- Patient is unable to swallow solid dosage forms
-

Enzyme Inhibitors: PARP

Lynparza® tablets, Rubraca® tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Talzenna® capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Talzenna® capsules Initial PA Criteria:

- One of the following:
  - Diagnosis of HER2-negative locally advanced or metastatic breast cancer; **AND**
    - Patient has a BRCA-positive mutated germline confirmed by an FDA-approved test (e.g., BRACAnalysis CDx); **AND**
    - Patient must have received treatment with an anthracycline and/or a taxane (unless contraindicated) as neoadjuvant, adjuvant, and/or metastatic treatment; **AND**
    - If patient received prior platinum-based chemotherapy, disease progression nor relapse were experienced within 6-months of receiving neoadjuvant or adjuvant platinum therapy; **OR**
  - Diagnosis of metastatic castration-resistant prostate cancer; **AND**
    - Patient has homologous recombination repair (HRR) gene mutation; **AND**
    - Patient must use in combination with Xtandi; **AND**
    - Patient has had a bilateral orchiectomy **OR** will receive a gonadotropin-releasing hormone (GnRH)-analog (e.g., leuprolide, goserelin, triptorelin); **AND**
- Provider will monitor complete blood counts at baseline and monthly thereafter; **AND**
- Patient does not have untreated CNS metastases (patient has completed definitive local therapy and may have stable CNS lesions on repeat brain imaging); **AND**
- Patient will not use requested agent in combination with any other PARP inhibitors; **AND**
- Patient has not received prior therapy with a PARP-inhibitor (e.g., Lynparza)

Talzenna® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Tumor response has been demonstrated with either stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from; **AND**
- Patient has not developed myelodysplastic syndrome (MDS)/acute myeloid leukemia (AML)

Zejula® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Enzyme Inhibitors: RET

Retevmo® capsules, tablets

PDL Status	P
Quantity Limit	40 mg capsules and tablets: 3/day; All other strengths capsules and tablets: 2/day
PA Criteria	See below
PA Form	General PA Form

Retevmo® capsules, tablets Initial PA Criteria:

- Patient must have ONE of the following diagnoses:
- Locally advanced or metastatic RET fusion-positive non-small cell lung cancer (NSCLC)
- Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) who require systemic therapy
- Advanced or metastatic RET fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory
- Locally advanced or metastatic solid tumors with a RET gene fusion that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options; **AND**
- Prescribed by, or in consultation with, an oncologist

Retevmo® capsules, tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., severe or life-threatening hemorrhaging, uncontrolled blood pressure, interstitial lung disease, elevated liver enzymes, QT interval prolongation)

Gavreto® capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Gavreto® capsules Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Metastatic RET fusion-positive non-small cell lung cancer (NSCLC) that is detected by an FDA approved test
  - Advanced or metastatic RET fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate); **AND**
- Requested agent will be prescribed by, or in consultation with, an oncologist

Gavreto® capsules Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., interstitial lung disease, elevated liver enzymes, severe or life-threatening hemorrhaging, uncontrolled blood pressure)

Enzyme Inhibitors: Tropomyosin Receptor Kinase (TRK)

Augtyro® capsules

PDL Status	NP
Quantity Limit	40 mg caps: 8/day; 160 mg caps: 2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Augtyro® capsules Initial PA Criteria:

- Patient has diagnosis of ONE of the following:
  - Diagnosis of locally advanced or metastatic Non-Small Cell Lung Cancer (NSCLC); **AND**
    - Tumor is ROS1 rearrangement positive; **OR**
  - NTRK Gene Fusion-Positive Solid Tumor and BOTH of the following:
    - Disease is locally advanced or metastatic or where surgical resection is likely to result in severe morbidity
    - Disease has progressed following treatment or there is no satisfactory alternative treatment; **OR**
  - Diagnosis of secretory breast cancer or mammary analogue secretory cancer; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- For patients with reproductive potential, prescriber attest to all of the following:
  - Patient is not pregnant prior to initiation of therapy
  - Female patients have been advised to use effective contraception during treatment and for 2 months after the final dose
  - Female patients have been advised to not breastfeed during treatment and 10 days after the final dose
  - Male patients with female partners of reproductive potential have been advised to use effective contraception during treatment and for 4 months after the final dose

Augtyro® capsules Renewal PA Criteria:

- Patient must continue to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hepatotoxicity, central nervous system effects, hyperuricemia, skeletal fractures, creatine phosphokinase elevation, interstitial lung disease/pneumonitis)

Ibtrozi® capsules

PDL Status	NP
Quantity Limit	3 capsules/day
PA Criteria	See below
PA Form	General PA Form

Ibtrozi® capsules Initial PA Criteria:

- Diagnosis of locally advanced or metastatic non-small cell lung cancer; **AND**
- ROS1-positive mutation present; **AND**
- Prescribed by, or in consultation with, an oncologist

Ibtrozi® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient does not show evidence of progressive disease or unacceptable toxicity (e.g., hepatotoxicity, interstitial lung disease, QTc interval prolongation, etc.)

Rozlytrek® capsules

PDL Status	NP
Quantity Limit	100 mg capsules: 4/day; 200 mg capsules: 3/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Rozlytrek® capsules Initial PA Criteria:

- Patient meets ONE of the following disease specific criteria:
  - Diagnosis of recurrent, advanced, or metastatic Non-Small Cell Lung Cancer (NSCLC); **AND**
    - Disease is ROS1 positive as detected by an FDA-approved test
  - NTRK Gene Fusion-Positive Solid Tumor; **AND**
    - Presence of a neurotrophic tyrosine receptor kinase (NTRK) gene fusion as detected by an FDA-approved test, without a known acquired resistance mutation; **AND**
    - Disease is metastatic or where surgical resection is likely to result in severe morbidity; **AND**
    - Disease has progressed following treatment or there is no satisfactory alternative treatment; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient does not have a history of prolonged QTc interval (e.g., QTc interval > 450 milliseconds); **AND**
- Patient will not use therapy in combination with drugs which prolong QT-interval; **AND**
- Patient will not use therapy with other NTRK-inhibitor therapy or ROS1-directed therapy; **AND**
- Patient does not have signs and symptoms of hyperuricemia as evidenced by a baseline serum; **AND**
- Patient will avoid concomitant use with moderate or strong CYP3A inducers or inhibitors; **AND**
- Provider attests to perform ALL the following:
  - Assess left ventricular ejection fraction (LVEF) prior to initiation of Rozlytrek in patients with symptoms or known risk factors for CHF
  - Monitor liver tests, including ALT and AST, every 2 weeks during the first month of the patient's treatment, then monthly thereafter, and as clinically indicated
  - Assess serum uric acid levels prior to initiation and periodically during treatment with Rozlytrek
  - Assess QT interval and electrolytes at baseline and periodically during treatment patients who have or who are at risk for QTc interval prolongation
  - Advise females of reproductive potential of the potential risk to a fetus and use of effective contraception during treatment and for 5 weeks following the final dose
  - Advise males with female partners of reproductive potential to use effective contraception during treatment and for 3 months after the final dose

Rozlytrek® capsules Renewal PA Criteria:

- Patient must continue to meet the initial criteria; **AND**
- Documented disease response with treatment, as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- There is absence of unacceptable toxicity from the drug (e.g., congestive heart failure, hepatotoxicity, central nervous system effects, hyperuricemia, QT-interval prolongation, visual disturbances)

Rozlytrek® packets

PDL Status	NP
Quantity Limit	600 mg/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Rozlytrek® packets PA Criteria:

See Rozlytrek® capsules prior authorization criteria; **AND**

- Clinically valid reason why Rozlytrek capsules cannot be used

Vitrakvi® capsules, solution

PDL Status	NP
Quantity Limit	25 mg capsules: 3/day; 100 mg capsules: 2/day; 20 mg/mL solution: 10 mL/day
PA Criteria	See below
PA Form	General PA Form

Vitrakvi® capsules, solution Initial PA Criteria:

- Patient has a solid tumor (e.g., soft tissue sarcoma, salivary gland, infantile fibrosarcoma, thyroid, lung, or gastrointestinal stromal tumors); **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient meets ALL the following:
  - Presence of a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation
  - Disease is metastatic or surgical resection is likely to result in severe morbidity
  - Disease has progressed following treatment or there is no satisfactory alternative treatment; **AND**
- Provider attests to ALL the following:
  - Monitor liver tests including ALT and AST every 2 weeks during the first month of treatment, then monthly thereafter and as clinically indicated
  - Advise females with reproductive potential and males with female partners of reproductive potential to use effective contraception during treatment and for 1 week after the final dose

Vitrakvi® capsules, solution Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity such as severe neurotoxicity, hepatotoxicity; (adverse effects resolve following dose recommendations/no permanent discontinuation required)

Hormonal Agents: Aromatase Inhibitors

Arimidex® tablets, Aromasin® tablets, Femara® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Arimidex® tablets, Aromasin® tablets, Femara® tablets PA Criteria:
Standard NP Criteria

Hormonal Agents: Anti-Androgens, First Generation

Casodex® tablets, Nilandron® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Casodex® tablets, Nilandron® tablets PA Criteria:
Standard NP Criteria

Hormonal Agents: Anti-Androgens Second Generation

Abiraterone tablets, Abirtega® tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Abiraterone tablets, Abirtega® tablets Initial PA Criteria:

- Patient has diagnosis of ONE of the following:
  - Metastatic castration-resistant prostate cancer (mCRPC)
  - Metastatic castration-sensitive prostate cancer (mCSPC); **AND**
- Will be used in combination with prednisone; **AND**
- ONE of the following:
  - Medication will be used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard, Lupron), goserelin (Zoladex), triptorelin (Trelstar))
  - Patient has had a bilateral orchiectomy; **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least three weeks after the last dose of abiraterone or Abirtega

Abiraterone tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug (e.g., hypokalemia, fluid retention, cardiovascular adverse reactions due to mineralocorticoid excess, adrenocortical insufficiency, hepatotoxicity, etc.)

Xtandi® capsules

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Xtandi® capsules Initial PA Criteria:

- Patient has diagnosis of ONE of the following:
  - Castration-resistant prostate cancer (CRPC); **AND**
    - ONE of the following:
      - Medication will be used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard, Lupron), goserelin (Zoladex), triptorelin (Trelstar))
      - Patient has had a bilateral orchiectomy; OR
  - Metastatic castration-sensitive prostate cancer (mCSPC); **AND**
    - ONE of the following:
      - Medication will be used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard, Lupron), goserelin (Zoladex), triptorelin (Trelstar))
      - Patient has had a bilateral orchiectomy; OR
  - Non-metastatic castration-sensitive prostate cancer (nmCSPC) with biochemical recurrence at high risk for metastasis (high-risk BCR); **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least three months after the last dose of Xtandi;

Xtandi® capsules Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug (e.g., seizure, posterior reversible encephalopathy syndrome (PRES), hypersensitivity, ischemic heart disease, falls and fractures, dysphagia or choking, etc.)

Akeega® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Akeega® tablets Initial PA Criteria:

- Diagnosis of one of the following:
  - Diagnosis of metastatic castration-resistant prostate cancer (mCRPC) with a deleterious or suspected deleterious BRCA-mutated (BRCAm) germline confirmed by an FDA approved test; **OR**
  - Diagnosis of metastatic castration-sensitive prostate cancer (mCSPC) with a deleterious or suspected deleterious BRCA2-mutated (BRCA2m) germline confirmed by an FDA approved test; **AND**
- Will be taken in combination with prednisone; **AND**
- ONE of the following:
  - Medication will used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard®, Lupron), goserelin (Zoladex®), triptorelin (Trelstar®))
  - Patient has had a bilateral orchiectomy; **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least four months after the last dose of Akeega

Akeega® tablets Renewal PA Criteria

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug (e.g., hepatotoxicity, fractures, hypertension)

Erleada® tablets

PDL Status	NP
Quantity Limit	60 mg tablets: 4/day; 240 mg tablets: 1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Erleada® tablets Initial PA Criteria:

- Patient has diagnosis of ONE of the following:
  - Non-metastatic castration-resistant prostate cancer (nmCRPC)
  - Metastatic castration-sensitive prostate cancer (mCSPC); **AND**
- ONE of the following:
  - Medication will used in combination with gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard®, Lupron), goserelin (Zoladex®), triptorelin (Trelstar®))
  - Patient has had a bilateral orchiectomy; **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least three months after the last dose of Erleada

Erleada® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. (e.g., seizures, excessive falls and/or fractures and any other Grade 3 or above side effects that are intolerable to patient, etc.)

Nubeqa® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Nubeqa® tablets Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Non-metastatic castration-resistant prostate cancer (nmCRPC); **OR**
  - Metastatic castration-sensitive prostate cancer (mCSPC); **AND**
  - Metastatic castration-sensitive prostate cancer (mCSPC) and will be used in combination with docetaxel; **AND**
- ONE of the following:
  - Medication will used in combination with with gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard®, Lupron), goserelin (Zoladex®), triptorelin (Trelstar®))
  - Patient has had a bilateral orchiectomy; **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least one week after the last dose of Nubeqa

Nubeqa® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug.(e.g., elevated hepatic enzymes, hyperbilirubinemia, neutropenia, or any other Grade 3 or above side effects that are intolerable to patient, etc.)

Xtandi® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Xtandi® tablets PA Criteria:

See Xtandi capsules PA Criteria; **AND**

- If request is for Xtandi tablets, clinically valid reason why Xtandi capsules cannot be used

Yonsa® tablets

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Yonsa® tablets Initial PA Criteria:

- Patient has metastatic castration-resistant prostate cancer (mCRPC); **AND**
- Will be taken in combination with methylprednisolone; **AND**
- ONE of the following:
  - Medication will used in combination with with with gonadotropin-releasing hormone (GnRH) analog (e.g., leuprolide (Eligard®, Lupron), goserelin (Zoladex®), triptorelin (Trelstar®))
  - Patient has a bilateral orchiectomy; **AND**
- Patients with female partners of reproductive potential should use effective contraception during treatment and for at least three weeks after the last dose of Yonsa

Yonsa® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Tumor response with stabilization of disease or decrease in size of tumor or tumor spread
- Absence of unacceptable toxicity from the drug (e.g., elevated hepatic enzymes, hypokalemia, fluid retention, hypertension)

Zytiga® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Zytiga® tablets PA Criteria:

See abiraterone PA Criteria; **AND**

- If request is for Zytiga, clinically valid reason why generic equivalents cannot be used

Hormonal Agents: GnRH Agonists & LNRH Analogs

Eligard® subcutaneous injection

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Eligard® subcutaneous injection PA Criteria:

- Diagnosis of prostate cancer in male patient

Leuprolide injection kit

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Leuprolide injection kit PA Criteria:

- Leuprolide will be approved for patients meeting **ONE** of the following criteria:
  - Diagnosis of prostate cancer in male patient
  - Diagnosis of central precocious puberty in children (onset of secondary sexual development before 8 [girls] or 9 years of age [boys])

Lupron Depot® injection kit

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lupron Depot® injection kit PA Criteria:

- Will be approved for self-administering patients with ONE of the following:
  - Diagnosis of prostate cancer in male patient
  - Diagnosis of endometriosis in female patient
  - Diagnosis of uterine leiomyomas in female patient
  - Diagnosis of recurrent ovarian carcinoma

Orgovyx® tablets

PDL Status	NP
Quantity Limit	30/month (32 tablets for initial month of therapy)
PA Criteria	See below
PA Form	General PA Form

Orgovyx® tablets PA Criteria:

- Diagnosis of advanced prostate cancer in male patient; **AND**
- Male patients with female partners of reproductive potential have been advised to use effective contraception during treatment and for two weeks after the last dose; **AND**
- Patient will not take requested medication with ANY of the following:
  - P-GP Inhibitors
  - Strong CYP3A Inducers
  - Cisapride
  - Pimozide
  - Thioridazine

Vabrinty® injection kit

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vabrinty® injection kit PA Criteria:

See Eligard® prior authorization criteria; **AND**

- Clinically valid reason why brand Eligard® cannot be used

Hormonal Agents: SERM/SERD

Inluriyo® tablets

PDL Status	NP
Quantity Limit	2 tablets/day
PA Criteria	See below
PA Form	General PA Form

Inluriyo ® tablets PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of advanced or metastatic breast cancer that meets ALL of the following:
  - Estrogen receptor (ER)-positive;
  - Human epidermal growth factor receptor 2 (HER2)-negative;
  - Estrogen receptor-1 (ESR1)-mutated; **AND**
- Presence of an estrogen receptor-1 (ESR1) mutation as confirmed by an FDA-approved test; **AND**
- Patient has documented disease progression on at least one prior treatment of endocrine therapy (e.g. tamoxifen, anastrozole, exemestane, letrozole, etc.); **AND**
- ONE of the following:
  - Request is for a male patient who will be using in combination with a gonadotropin-releasing hormone (GnRH) agent (e.g., leuprolide, goserelin, triptorelin); **OR**
  - Request is for a pre- or peri-menopausal female patient who will be using in combination with a gonadotropin-releasing hormone (GnRH) agent (e.g., leuprolide, goserelin, triptorelin); **OR**
  - Request is for a post-menopausal female; **AND**
- Prescribed by, or in consultation with, an oncologist or hematologist.

Inluriyo® Renewal Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hepatotoxicity, left ventricular dysfunction, interstitial lung disease/pneumonitis)

Orserdu® tablets

PDL Status	NP
Quantity Limit	86 mg: 3/day; 345 mg: 1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Orserdu® tablets Initial PA Criteria:

- Diagnosis of advanced or metastatic breast cancer that meets ALL of the following:
  - Estrogen receptor (ER)-positive;
  - Human epidermal growth factor receptor 2 (HER2)-negative;
  - Estrogen receptor-1 (ESR1)-mutated; **AND**
- Patient has received at least one endocrine based regimen; **AND**
- Patient has ESR1 mutation confirmed by FDA-approved test; **AND**
- If female, patient is postmenopausal; **AND**
- Orserdu will be used as monotherapy; **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient must not be breastfeeding; **AND**
- Males undergoing treatment with female partners of reproductive age should be advised to use effective contraception during treatment and for 1 week after the final dose

Orserdu® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., dyslipidemia, musculoskeletal pain)

Toremifene tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Toremifene tablets PA Criteria:

Standard NP Criteria

Veppanu® tablets

PDL Status	NP
Quantity Limit	1 tab/day
PA Criteria	See below
PA Form	General PA Form

Veppanu® tablets Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of advanced or metastatic breast cancer that meets ALL of the following:
  - Estrogen receptor (ER)-positive;
  - Human epidermal growth factor receptor 2 (HER2)-negative;
  - Estrogen receptor-1 (ESR1)-mutated; **AND**
- Presence of an estrogen receptor-1 (ESR1) mutation as confirmed by an FDA-approved test; **AND**
- Patient has documented disease progression on at least one prior treatment of endocrine therapy (e.g., tamoxifen, anastrozole, exemestane, letrozole, etc.); **AND**
- Patient must not be pregnant or breastfeeding; **AND**
- Females of reproductive potential and males undergoing treatment with female partners of reproductive age should be advised to use effective contraception during treatment and for 1 week after the final dose
- Prescribed by, or in consultation with, an oncologist or hematologist.

Veppanu® tablets Renewal Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. QTc interval prolongation, musculoskeletal pain, hepatic injury, lab abnormalities)

Multikinase Inhibitors: Renal and Thyroid Cancers

Fotivda® capsules

PDL Status	NP
Quantity Limit	21/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Fotivda® capsules Initial PA Criteria:

- Patient has diagnosis of relapsed or refractory advanced renal cell carcinoma (RCC); **AND**
- Patient has had two or more prior systemic therapies [two kinase inhibitors (KIs), a KI plus an immune checkpoint inhibitor, or a KI plus other systemic agents]; **AND**
- Prescriber attests to ALL the following:
  - Patient's blood pressure will be assessed prior to and during therapy
  - Patient will be closely monitored due to increased risk of Arterial and venous Thromboembolic Events, Hemorrhagic Events, Proteinuria, and Thyroid Dysfunction
  - Fotivda will be withheld for at least 24 days before elective surgery and will not administer for at least 2 weeks following major surgery and adequate wound healing
  - Patient's baseline liver function tests will be assessed
  - Female and male patients of reproductive potential have been advised to use effective contraception during treatment and for one month after the last dose
  - Agent will not be co-administered with strong CYP3A inducers
  - Patient does not have a history of allergic reactions to tartrazine (only applies to requests for Fotivda 0.89 mg)
  - Female patients are not pregnant or breastfeeding; **AND**
- Will not use in patients with any of the following:
  - Strong CYP3A inducers
  - History of allergic reactions to tartrazine

Fotivda® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests to positive response to therapy indicated by tumor response with stabilization of disease **OR** decrease in size of tumor or tumor spread; **AND**
- Patient has absence of unacceptable toxicity from the drug (e.g., uncontrolled hypertension, onset of cardiac failure, arterial and venous Thromboembolic Events, hemorrhagic events, proteinuria, thyroid dysfunction, onset of Reversible Posterior Leukoencephalopathy Syndrome (RPLS), or increased LFT's)

Nexavar® tablets, sunitinib capsules, Votrient® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Nexavar® tablets, sunitinib capsules, Votrient® tablets PA Criteria:

Standard NP Criteria

Multiple Myeloma Agents

Lenalidomide capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Lenalidomide capsules PA Criteria:

- Clinically valid reason why Revlimid® cannot be used

Pomalyst® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Pomalyst® capsules PA Criteria:

- Clinically valid reason why generic pomalidomide cannot be used

Xpovio® tablets

PDL Status	NP
Quantity Limit	4 packs/month
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Xpovio® tablets Initial PA Criteria:

- Diagnosis of multiple myeloma; **AND**
- Patient has received at least one prior therapy; **AND**
- Prescribed by, or in consultation with, an oncologist

Xpovio® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests that the patient has experienced lack of disease progression, and/or improvement in symptoms; **AND**
- Patient has absence of unacceptable toxicity from the drug (e.g., thrombocytopenia, neutropenia, gastrointestinal toxicity, hyponatremia, neurological toxicity)

Myelofibrosis

Jakafi® tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Besremi® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Besremi® injection PA Criteria:

- Diagnosis of polycythemia vera; **AND**
- Prescribed by, or in consultation with, an oncologist or hematologist; **AND**
- Patient does not have ANY of the following:
  - Severe, acute, or unstable cardiovascular disease
  - Existence of, or history of severe psychiatric disorders, particularly severe depression, suicidal ideation, or suicide attempt
  - Hypersensitivity to interferon or to any component of BESREMI
  - Hepatic impairment (Child-Pugh B or C)
  - History or presence of active serious or untreated autoimmune disease; **AND**
- Patient is not an immunosuppressed transplant recipient; **AND**
- For women of childbearing age, provider has confirmed that the patient is not pregnant prior to receiving treatment; **AND**
- Patients of reproductive potential will be counseled to use effective contraception during treatment and for at least 8 weeks after the final dose

Inrebic® capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Inrebic® capsules Initial PA Criteria:

- Patient has a diagnosis of primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis; **AND**
- Patient is considered intermediate-2 risk or high-risk; **AND**
- Patient's platelet count ≥ 50 x 109/L; **AND**
- Provider attests patient is not thiamine deficient (vitamin B1) and will monitor thiamine level during treatment

Inrebic® capsules Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., reduction in symptoms, decreased spleen size); **AND**
- Absence of unacceptable toxicity (e.g., encephalopathy, anemia, thrombocytopenia, hepatoxicity, major adverse cardiovascular events, thrombosis, and malignancies); **AND**
- Prescriber agrees to continue monitoring thiamine (vitamin B1) levels

Jakafi XR® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Jakafi XR® tablets PA Criteria:

- Clinically valid reason why Jakafi® immediate release tablets cannot be used

Ojjaara® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Ojjaara® tablets Initial PA Criteria:

- Patient has a diagnosis of primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis; **AND**
- Patient is considered intermediate-1, intermediate-2, or high-risk; **AND**
- Patient is anemic (e.g., hemoglobin (Hb) < 10 g/dL and/or hematocrit (Hct) < 30%)

Ojjaara® tablets Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., reduction in symptoms, decreased spleen size, decreased number of transfusion); **AND**
- Absence of unacceptable toxicity (e.g., thrombocytopenia, neutropenia, hepatotoxicity, major adverse cardiovascular events, thrombosis, and malignancies)

Vonjo® capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Vonjo® capsules Initial PA Criteria:

- Patient has a diagnosis of primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis; **AND**
- Patient is considered intermediate risk or high-risk; **AND**
- Platelet count is below 50 x 10^9/L

Vonjo® capsules Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., reduction in symptoms, decreased spleen size, decreased number of transfusions); **AND**
- Absence of unacceptable toxicity (e.g., thrombocytopenia, major adverse cardiovascular events, thrombosis, malignancies)

Phosphoinositide-3-Kinase Inhibitors

Copiktra®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Copiktra® Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Relapsed or refractory chronic lymphocytic leukemia (CLL)
  - Small lymphocytic lymphoma (SLL); **AND**
- Patient has not received therapy with ANY of the following:
  - Phosphoinositide 3-kinase (PI3K) inhibitors (e.g., Zydelig)
  - Bruton's tyrosine kinase (BTK) inhibitor (e.g., Imbruvica and Calquence); **AND**
- Used as a single agent; **AND**
- Patient has received at least 2 prior therapies which included treatment with ofatumumab (Arzerra); **AND**
- Patient must not have undergone prior autologous hematopoietic stem cell transplant (HSCT) within 180 days of the first dose OR prior allogeneic transplant; **AND**
- Patient does not have an active infection, including clinically important localized infections; **AND**
- Patient must not be pregnant or breastfeeding; **AND**
- Prescriber attests to counsel and/or monitor regarding the following:
  - Signs and symptoms of infection
  - Diarrhea and colitis
  - Cutaneous reactions
  - Pneumonitis
  - Hepatotoxicity
  - Neutropenia
  - Female patients of reproductive potential should use effective contraception during treatment and for at least 1 month after treatment
  - Males undergoing treatment with female partners of reproductive age should use effective contraception during treatment and 1 month after treatment due to male mediated teratogenicity

Copiktra® Renewal PA Criteria:

- Patient continues to meet criteria identified in initial criteria; **AND**
- Disease response as defined by lack of disease progression, improvement in tumor size and/or improvement in patient symptoms; **AND**
- Absence of unacceptable toxicity from the drug (e.g., active/severe infections, hematologic toxicity [neutropenia], severe diarrhea or colitis, hepatotoxicity, pneumonitis, severe cutaneous reactions [Stevens-Johnson syndrome and toxic epidermal necrolysis], anaphylaxis)

PI3K Inhibitors

Itovebi® tablets

PDL Status	NP
Quantity Limit	3 mg: 2/day; 9 mg: 1/day
PA Criteria	See below
PA Form	General PA Form

Itovebi® tablets Initial PA Criteria:

- Patient has hormone endocrine-resistant[SB1] , receptor (HR)-positive, HER2-negative locally advanced or metastatic breast cancer; **AND**
- Patient has a PIK3CA mutation as detected by an FDA-approved test; **AND**
- Patient has experienced disease recurrence on or after completing adjuvant endocrine therapy; **AND**
- Agent is being given in combination with Ibrance and fulvestrant; **AND**
- Agent is prescribed by, or in consultation with, an oncologist

Itovebi® tablets Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has clinical response defined as disease stabilization or decrease in size or spread of tumor; **AND**
- Patient does not have unacceptable toxicity (e.g., hyperglycemia, diarrhea, stomatitis)

Piqray® tablets, tablet packs

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Piqray® tablets, tablet packs Initial PA Criteria:

- Patient has hormone receptor-(HR) positive, HER2-negative advanced or metastatic breast cancer; **AND**
- Agent is prescribed by, or in consultation with, an oncologist; **AND**
- Patient has experienced disease progression on or after an endocrine based regimen for advanced disease or has relapsed disease within 12 months after completion of adjuvant endocrine therapy; **AND**
- Patient has a PIK3CA-mutation as detected by an FDA-approved test; **AND**
- Piqray® will be given in combination with fulvestrant; **AND**

Piqray® tablets, tablet packs Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has tumor response with stabilization of disease or decrease in the size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity such as severe cutaneous reaction or pneumonitis

Truqap® tablets, tablet packs

PDL Status	NP
Quantity Limit	64/28 days
PA Criteria	See below
PA Form	General PA Form

Truqap® tablets, tablet packs Initial PA Criteria:

- One of the following:
  - All of the following:
    - Patient has hormone receptor-positive, HER2-negative locally advanced or metastatic breast cancer; **AND**
    - Patient has one or more PIK3CA/AKT1/PTEN-alterations as detected by an FDA-approved test; **AND**
    - Patient has experienced disease progression or on after an endocrine based regimen for advanced disease OR has relapsed disease within 12 months after completion of adjuvant endocrine therapy; **AND**
    - Agent is being given in combination with fulvestrant; **OR**
  - All of the following:
    - Patient has PTEN-deficient metastatic androgen pathway modulation-naïve or -sensitive (mAPMN/S) prostate cancer; **AND**
    - PTEN-deficiency has been detected by an FDA-authorized test; **AND**
    - Agent is being given in combination with abiraterone and prednisone; **AND**
- Agent is prescribed by, or in consultation with, an oncologist

Truqap® tablets, tablet packs Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has clinical response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., hyperglycemia, diarrhea, cutaneous adverse reactions)

Rare/Miscellaneous Oncology Conditions

Ayvakit® injection

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Ayvakit® injection Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Unresectable or metastatic gastrointestinal stromal tumors (GIST) with platelet-derived growth factor-alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations
  - Indolent systemic mastocytosis (ISM)
  - Advanced systemic mastocytosis (AdvSM) Note: Includes aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SM-AHN), and mast cell leukemia (MCL); **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**

Ayvakit® injection Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. intracranial hemorrhage, edema, fatigue/asthenia, cognitive dysfunction)

Hydrea® capsules

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Hydrea® capsules PA Criteria:
Standard NP Criteria

Iwilfin® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Iwilfin® tablets Initial PA Criteria:

- Diagnosis of high-risk neuroblastoma (HRNB); **AND**
- Patient had a partial response to prior multiagent, multimodality therapy; **AND**
- Patient has received anti-GD2 immunotherapy (e.g., dinutuximab); **AND**
- Prescribed by, or in consultation with, an oncologist

Iwilfin® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. hepatotoxicity, hearing loss, myelosuppression, pneumonia)

Lifyorli® capsules

PDL Status	NP
Quantity Limit	27 capsules/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Lifyorli® capsules Initial PA Criteria:

- Diagnosis of platinum-resistant epithelial ovarian, fallopian tube, or primary peritoneal cancer; **AND**
- Patient has failed at least one prior systemic treatment regimen which has included bevacizumab; **AND**
- Patient is not receiving systemic glucocorticoids for lifesaving purposes (e.g., immunosuppression after organ transplantation); **AND**
- Prescribed by, or in consultation with, an oncologist

Lifyorli® Capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g. febrile neutropenia, adrenal insufficiency)

Modeyso® capsules

PDL Status	NP
Quantity Limit	20 capsules/28 days
PA Criteria	See below
PA Form	General PA Form

Modeyso® capsules Initial PA Criteria:

- Diagnosis of diffuse midline glioma; **AND**
- Patient is positive for H3 K27M mutation; **AND**
- Patient has progressed on at least one prior therapy (e.g., radiation therapy)
- Prescribed by, or in consultation with, an oncologist

Modeyso® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., QT prolongation, Torsades de pointes, polymorphic ventricular tachycardia, confusional state)

Ogsiveo® tablets

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form

Ogsiveo® tablets Initial PA Criteria:

- Diagnosis of progressing desmoid tumor (also known as aggressive fibromatosis); **AND**
- Prescriber has reviewed and evaluated appropriate treatment options and attests that the patient requires systemic therapy; **AND**
- Prescribed by, or in consultation with, an oncology, hematology, or gastroenterology specialist

Ogsiveo® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., diarrhea, ovarian toxicity, hepatotoxicity, electrolyte abnormalities)

Qinlock® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Qinlock® tablets Initial PA Criteria:

- Diagnosis of unresectable, locally advanced, or metastatic gastrointestinal stromal tumor (GIST); **AND**
- Prescribed by, or in consultation with, an oncologist; **AND**
- Patient has been previously treated with at least THREE kinase systemic therapies (e.g., imatinib, avapritinib, sunitinib, regorafenib)

Qinlock® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., Palmar-Plantar Erythrodysesthesia Syndrome, impaired wound healing, new cutaneous malignancies, uncontrolled hypertension, cardiac dysfunction)

Rezurock® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Rezurock® tablets Initial PA Criteria:

- Patient has diagnosis of Chronic Graft-Versus-Host Disease; **AND**
- Patient is 12 years of age or older; **AND**
- Patient has a history of allogenic hematopoietic cell transplant (HCT); **AND**
- Agent is prescribed by, or in consultation with, an oncologist, hematologist, or bone marrow transplant specialist; **AND**
- Patient has had a previous failure of at least one systemic corticosteroid therapy (i.e., methylprednisolone, prednisone, etc.); **AND**
- Patient has had a previous failure of at least one non-steroidal systemic immunosuppressant therapy (e.g., abatacept, alemtuzumab, calcineurin inhibitor, etanercept, hydroxychloroquine, ibrutinib, imatinib, interleukin-2, low-dose methotrexate, mT**OR** inhibitor, mycophenolate mofetil, pentostatin, rituximab, ruxolitinib, etc.); **AND**
- Prescriber attests, if applicable, that patient will be advised that effective contraception should be used during treatment and for at least one week after last dose

Rezurock® tablets Renewal PA Criteria:

- Patient continues to meet the initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., laboratory abnormalities, infections, asthenia, nausea, diarrhea, edema, hemorrhage)

Romvimza® capsules

PDL Status	NP
Quantity Limit	8/28 days
PA Criteria	See below
PA Form	General PA Form

Romvimza® capsules Initial PA Criteria:

- Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT); **AND**
- Patient has severe morbidity or functional limitations (e.g., worst stiffness numeric rating scale [NRS] of > 4); **AND**
- Patient is not a candidate for surgical resection associated with potential worsening, functional limitation, or severe morbidity; **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist

Romvimza® capsules Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not show evidence of progressive disease and unacceptable toxicity (e.g., laboratory abnormalities, hepatotoxicity, edema, rash, hypertension)

Turalio® capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Turalio® capsules Initial PA Criteria:

- Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT); **AND**
- Patient has severe morbidity or functional limitations (e.g., worst stiffness numeric rating scale [NRS] of ≥ 4); **AND**
- Patient is not a candidate for surgical resection associated with potential worsening, functional limitation, or severe morbidity; **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist; **AND**
- Prescriber is enrolled in the Turalio REMS Program

Turalio® capsules Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., laboratory abnormalities, hepatotoxicity, nausea, abdominal pain)

Welireg® tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Welireg® tablets Initial PA Criteria:

- ONE of the following:
  - Diagnosis of Von Hippel-Lindau (VHL) disease and require therapy for ONE of the following VHL-associated cancers, not requiring immediate surgery:
    - Renal cell carcinoma (RCC)
    - Central nervous system (CNS) hemangioblastomas
    - Pancreatic neuroendocrine tumors (pNET); **OR**
  - Diagnosis of advanced renal cell carcinoma (RCC) with a clear cell component (ccRCC) and ONE of the following:
    - Patient is using as adjuvant treatment in combination with pembrolizumab or pembrolizumab and berahyaluronidase alfa-pmph, for intermediate-high or high risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions; **OR**
    - Patient has tried and failed, has a contraindication, or intolerance to ALL of the following:
      - Programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor (e.g., nivolumab, avelumab, pembrolizumab)
      - Vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) (e.g., Cabometyx, Inlyta, Lenvima, Nexavar, Sutent); **OR**
  - Diagnosis of locally advanced, unresectable, or metastatic pheochromocytoma or paraganglioma (PPGL); **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist; **AND**
- Patient is not pregnant or breastfeeding; **AND**
- Females of reproductive potential and males with female partners of reproductive potential should be advised to use effective non-hormonal contraception during treatment and for 1 week after the last dose

Welireg® tablets Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has positive disease response defined as disease stabilization or decrease in size of tumor or tumor spread; **AND**
- Patient does not have unacceptable toxicity (e.g., anemia, hypoxia)

XII. OPHTHALMICS

Dry Eye Disease Agents

Restasis® emulsion (single dose vials)

PDL Status	P
Quantity Limit	60 vials/30 days
PA Criteria	See Below
PA Form	General PA Form

Restasis® emulsion PA Criteria:

- Treatment of vernal keratoconjunctivitis (VKC) (i.e., severe atopic keratoconjunctivitis); **OR**
- Diagnosis of dry eye disease [i.e., dry eye syndrome, keratoconjunctivitis sicca (KCS)]

Xiidra® drops

PDL Status	P
Quantity Limit	2 vials/day
PA Criteria	See Below
PA Form	General PA Form

Xiidra® drops PA Criteria:

- Diagnosis of dry eye disease [i.e., dry eye syndrome, keratoconjunctivitis sicca (KCS)]; **AND**
- Trial and failure or contraindication to Restasis® (trial duration ≥ 12 weeks confirmed by paid claims)

Cequa® solution

PDL Status	NP
Quantity Limit	2 vials/day
PA Criteria	See Below
PA Form	General PA Form

Cequa® solution PA Criteria:

- Diagnosis of dry eye disease [i.e., dry eye syndrome, keratoconjunctivitis sicca (KCS)]; **AND**
- Trial and failure, or contraindication, to both the following:
  - Restasis® (trial duration > 12 weeks confirmed by paid claims)
  - Xiidra® (trial duration > 12 weeks confirmed by paid claims)

Cyclosporine emulsion 0.05% (single dose vials)

PDL Status	NP
Quantity Limit	60 vials/30 days
PA Criteria	See Below
PA Form	General PA Form

Cyclosporine emulsion 0.05%® PA Criteria:

- One of the following:
  - Treatment of vernal keratoconjunctivitis (VKC) (i.e., severe atopic keratoconjunctivitis)
  - Diagnosis of dry eye disease [i.e., dry eye syndrome, keratoconjunctivitis sicca (KCS)]; **AND**
- Clinically valid reason why the preferred Restasis® cannot be used

Miebo® drops

PDL Status	NP
Quantity Limit	3 bottles/30 days
PA Criteria	See Below
PA Form	General PA Form

Miebo® drops PA Criteria:
See Cequa® prior authorization criteria

Restasis Multidose® emulsion

PDL Status	NP
Quantity Limit	1 bottle/30 days
PA Criteria	See Below
PA Form	General PA Form

Restasis Multidose® emulsion PA Criteria:
See cyclosporine emulsion 0.05% prior authorization criteria

Tryptyr® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tryptyr® solution PA Criteria:

- Diagnosis of dry eye disease [i.e., dry eye syndrome, keratoconjunctivitis sicca (KCS)]; **AND**
- Trial and failure, or contraindication, to both the following:
  - Restasis® (trial duration > 12 weeks confirmed by paid claims); **AND**
  - Xiidra® (trial duration > 12 weeks confirmed by paid claims)

Tyrvaya® nasal solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tyrvaya® nasal solution PA Criteria:
See Cequa® prior authorization criteria

Vevye® solution

PDL Status	NP
Quantity Limit	3 bottles/30 days
PA Criteria	See Below
PA Form	General PA Form

Vevye® solution PA Criteria:
See Cequa® prior authorization criteria

Ophthalmic Alpha-2 Agonists

Apraclonidine, Brimonidine 0.2%, & Alphagan P®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Brimonidine 0.1%, Brimonidine 0.15%, & Iopdine®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Brimonidine 0.1%, Brimonidine 0.15%, & Iopidine® PA Criteria:
Standard NP Criteria

Ophthalmic Antibiotics

Ciprofloxacin, Erythromycin, Moxifloxacin, Neomycin/bac/poly B, Neomycin/poly/B/gramicidin, Polymyxin B/TMP, Sulfacetamide Soln, Tobramycin

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

AzaSite® & Besivance®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

AzaSite® & Besivance® PA Criteria:
Standard NP Criteria

Ciloxan®

PDL Status	NP
Quantity Limit	10 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Ciloxan® PA Criteria:
Standard NP Criteria

Gentamicin

PDL Status	NP
Quantity Limit	15 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Gentamicin PA Criteria:
Standard NP Criteria

Gatifloxacin 0.5% soln, Levofloxacin 0.5% soln, Moxifloxacin (3xday), Sulfacetamide ointment, Tobrex®, & Vigamox®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Gatifloxacin 0.5% soln, Levofloxacin 0.5% soln, Moxifloxacin (3xday), Sulfacetamide ointment, Tobrex®, Vigamox® PA Criteria:
Standard NP Criteria

Ophthalmic Antibiotic/Steroid Combinations

Neomycin/BAC/Poly B/HC, Sulfacetamide/Prednisolone, Tobramycin/Dexamethasone, Zylet®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Loteprednol/tobramycin ophthalmic suspension

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Loteprednol/tobramycin ophthalmic suspension PA Criteria:

- Clinically valid reason why brand Zylet® cannot be used

Maxitrol®, Neomycin/Poly B/HC, TobraDex®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Maxitrol®, Neomycin/Poly B/HC, TobraDex® PA Criteria:

Standard NP Criteria

Ophthalmic Antifungals

Natacyn®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Natacyn® PA Criteria:

- Diagnosis of ophthalmic fungal infection

Ophthalmic Antivirals

Trifluridine & Zirgan®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Ophthalmic Anti-Allergic

Please refer to OTC Drug Formulary for coverage of OTC products.

Azelastine 0.05% ophthalmic solution

PDL Status	P
Quantity Limit	6 mL/Rx
PA Criteria	None

Cromolyn Sodium 4% ophthalmic solution

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Alocril® 2% ophthalmic solution, Alomide 0.1% ophthalmic solution

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Alocril®, Alomide® PA Criteria:

- Trial and failure, contraindication, or intolerance to TWO preferred Rx or Covered OTC Ophthalmic Anti-Allergics

Bepotastine 1.5% ophthalmic solution

PDL Status	NP
Quantity Limit	10 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Bepotastine PA Criteria:

- Trial and failure, contraindication, or intolerance to TWO preferred Rx or Covered OTC Ophthalmic Anti-Allergics

Epinastine 0.05% ophthalmic solution

PDL Status	NP
Quantity Limit	5 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Epinastine PA Criteria:

- Trial and failure, contraindication, or intolerance to TWO preferred Rx or Covered OTC Ophthalmic Anti-Allergics

Verkazia® 0.1% ophthalmic emulsion

PDL Status	NP
Quantity Limit	120/30 days
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Verkazia® Initial PA Criteria:

- Diagnosis of moderate to severe vernal keratoconjunctivitis; **AND**
- Trial and failure, contraindication, or intolerance of one agent in ALL the following categories:
  - Ophthalmic antihistamines (e.g., azelastine, olopatadine)
  - Ophthalmic mast cell stabilizers (e.g., cromolyn sodium)
  - Ophthalmic corticosteroids (e.g., dexamethasone, prednisolone, fluorometholone)

Verkazia® Renewal PA Criteria:

- Patient demonstrates positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms (e.g., itching, photophobia, papillary hypertrophy, mucus discharge, conjunctival hyperaemia)

Zerviate® 0.24% ophthalmic solution

PDL Status	NP
Quantity Limit	30 vials/Rx
PA Criteria	See Below
PA Form	General PA Form

Zerviate® PA Criteria:

- Trial and failure, contraindication, or intolerance to TWO preferred Rx or Covered OTC Ophthalmic Anti-Allergics

Ophthalmic Beta Blockers

Carteolol & Timolol maleate

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Betaxolol, Betoptic-S®, Istalol®, Levobunolol, Timolol gel soln, Timoptic Ocudose®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Betaxolol, Betoptic-S®, Istalol®, Levobunolol, Timolol gel soln, Timoptic Ocudose® PA Criteria:
Standard NP Criteria

Ophthalmic Carbonic Anhydrase Inhibitors

Azopt®

PDL Status	P
Quantity Limit	15 mL/30 days
PA Criteria	None

Dorzolamide & Dorzolamide/Timolol

PDL Status	P
Quantity Limit	10 mL/30 days
PA Criteria	None

Brinzolamide

PDL Status	NP
Quantity Limit	15 mL/30 days
PA Criteria	See Below
PA Form	General PA Form

Brinzolamide® PA Criteria:
Standard NP Criteria

Cosopt®

PDL Status	NP
Quantity Limit	10 mL/30 days
PA Criteria	See Below
PA Form	General PA Form

Cosopt® PA Criteria:
Standard NP Criteria

Cosopt PF®

PDL Status	NP
Quantity Limit	2 vials/day
PA Criteria	See Below
PA Form	General PA Form

Cosopt PF® PA Criteria:
Standard NP Criteria

Ophthalmic Kinase Inhibitors

Rhopressa®

PDL Status	P
Quantity Limit	5 mL/30 days
PA Criteria	None
PA Form	General PA Form

Rhopressa® PA Criteria:

- Patient has a diagnosis of ocular hypertension or open-angle glaucoma; **AND**
- Patient has tried/failed or is intolerant to either a prostaglandin inhibitor or beta-adrenergic antagonist

Rocklatan®

PDL Status	P
Quantity Limit	5 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Rocklatan® PA Criteria:
See Rhopressa® prior authorization criteria

Glaucoma Combinations

Combigan® & Simbrinza®

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Brimonidine/Timolol

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Brimonidine/Timolol PA Criteria:
Standard NP Criteria

Miotics

Pilocarpine 12.5%

PDL Status	NP
Quantity Limit	5 mL/30 days
PA Criteria	See Below
PA Form	General PA Form

Pilocarpine 12.5% PA Criteria:

- Diagnosis of presbyopia; **AND**
- Patient is 18 years of age or older; **AND**
- Patient is not a candidate for surgery or surgery was non-curative; **AND**
- Clinically valid reason as to why the preferred pilocarpine strengths cannot be used

Vuity®

PDL Status	NP
Quantity Limit	5 mL/30 days
PA Criteria	See Below
PA Form	General PA Form

Vuity® PA Criteria:

See pilocarpine 1.25% prior authorization criteria

Miscellaneous Ophthalmics

Cystaran®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Cystaran® PA Criteria:

- Diagnosis of cystinosis

Cystadrops®

PDL Status	NP
Quantity Limit	4 packages/28 days
PA Criteria	See Below
PA Form	General PA Form

Cystadrops® PA Criteria:

- Diagnosis of cystinosis; **AND**
- Prescriber must provide a clinically valid reason as to why Cystaran cannot be used

Oxervate®

PDL Status	NP
Quantity Limit	2 mL/day (lifetime therapy QL=112 mL for 8 weeks of therapy)
PA Criteria	See Below
PA Form	General PA Form

Oxervate® PA Criteria:

- Patient must be ≥ 2 years of age; **AND**
- Patient must have a diagnosis of moderate to severe (stage 2 or stage 3) neurotrophic keratitis (NK); **AND**
- Prescribed by, or in consultation with, an ophthalmologist; **AND**
- Prescriber attests that patient or caregiver has been counseled on proper administration technique

Xdemvy®

PDL Status	NP
Quantity Limit	1 bottle (10 mL)/50 days
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	2 months

Xdemvy® PA Criteria:

- Diagnosis of Demodex blepharitis; **AND**
- Patient has collarettes, cylindrical deposits at the base of eyelashes, confirmed by slit lamp examination; **AND**
- Prescribed by or in consultation with an ophthalmologist or optometrist

Mydriatics

Atropine & Cyclopentolate

PDL Status	P
Quantity Limit	15 mL/Rx
PA Criteria	None

Phenylephrine

PDL Status	P
Quantity Limit	2.5%: 15 mL/Rx 10%: 5 mL/Rx
PA Criteria	None

Tropicamide

PDL Status	P
Quantity Limit	15 mL/Rx
PA Criteria	None

Cyclogyl®

PDL Status	NP
Quantity Limit	15 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Cyclogyl® PA Criteria:
Standard NP Criteria

Cyclomydril®

PDL Status	NP
Quantity Limit	5 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Cyclomydril® PA Criteria:
Standard NP Criteria

Mydriacyl®

PDL Status	NP
Quantity Limit	15 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Mydriacyl® PA Criteria:
Standard NP Criteria

Ophthalmic NSAIDs

Diclofenac ophthalmic solution, flurbiprofen ophthalmic solution, ketorolac ophthalmic solution

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Acular LS® ophthalmic solution, Acuvail® ophthalmic solution, BromSite® ophthalmic solution, bromfenac ophthalmic solution, Ilevro® ophthalmic suspension, Nevanac® ophthalmic suspension, & Prolensa® ophthalmic solution

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Acular LS®, Acuvail®, BromSite®, Bromfenac, Ilevro®, Nevanac®, & Prolensa® PA Criteria:
Standard NP Criteria

Ophthalmic Prostaglandin Agonists

Latanoprost ophthalmic solution 0.005%, Lumigan® 0.01% ophthalmic solution, & Travatan Z® ophthalmic solution 0.004%

PDL Status	P
Quantity Limit	5 mL/Rx
PA Criteria	None

Zioptan® PF ophthalmic solution 0.0015%

PDL Status	P
Quantity Limit	1 container/day
PA Criteria	None

Bimatoprost 0.01% ophthalmic solution (generic for Lumigan®)

PDL Status	NP
Quantity Limit	5 mL/ Rx
PA Criteria	See Below
PA Form	General PA Form

Bimatoprost 0.01% ophthalmic solution PA Criteria:

- Clinically valid reason why brand Lumigan 0.01% cannot be used

Bimatoprost 0.03% ophthalmic solution (generic for Lumigan®)

PDL Status	NP
Quantity Limit	5 mL/ Rx
PA Criteria	See Below
PA Form	General PA Form

Bimatoprost 0.03% solution PA Criteria:
Standard NP Criteria

Iyuzeh® PF solution

PDL Status	NP
Quantity Limit	1 container/day
PA Criteria	See Below
PA Form	General PA Form

Iyuzeh® PF solution PA Criteria:

- Clinically valid reason why preferred Travatan Z® cannot be used

Tafluprost PF ophthalmic solution

PDL Status	NP
Quantity Limit	1 container/day
PA Criteria	See Below
PA Form	General PA Form

Tafluprost PF ophthalmic solution PA Criteria:
Standard NP Criteria

Travoprost BAK free ophthalmic solution

PDL Status	NP
Quantity Limit	5 mL/ Rx
PA Criteria	See Below
PA Form	General PA Form

Travoprost ophthalmic solution PA Criteria:

- Clinically valid reason why preferred Travatan Z® cannot be used

Vyzulta® ophthalmic solution, Xalatan® ophthalmic solution, & Xelpros® ophthalmic emulsion

PDL Status	NP
Quantity Limit	5 mL/ Rx
PA Criteria	See Below
PA Form	General PA Form

Vyzulta® ophthalmic solution, Xalatan® ophthalmic solution, & Xelpros® ophthalmic emulsion PA Criteria:
Standard NP Criteria

Ophthalmic Steroids

Alrex®, Durezol®, & Fluorometholone

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Loteprednol 0.5% susp

PDL Status	P
Quantity Limit	15 mL/Rx
PA Criteria	None

Pred Mild® & Prednisolone acetate

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Difluprednate

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Difluprednate PA Criteria:

- Clinically valid reason why the preferred Durezol® cannot be used

Eysuvis®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Eysuvis PA Criteria:

- Patient is being treated for symptoms of Dry Eye disease; **AND**
- Patient has had a trial and failure of Restasis; **AND**
- Patient has had a trial and failure of a preferred loteprednol product (e.g., Alrex, Lotemax suspension)

Flarex®, FML Forte®, FML Liquifilm®, Lotemax SM® gel, Lotemax® ointment, Loteprednol 0.5% gel

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Flarex®, FML Forte®, FML Liquifilm®, Lotemax SM® gel, Lotemax® ointment, Loteprednol 0.5% gel PA Criteria:
Standard NP Criteria

Loteprednol 0.2% susp

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Loteprednol 0.2% susp PA Criteria:

- Clinically valid reason why the preferred Alrex® cannot be used

Maxidex®, Prednisolone sodium phosphate, & Pred Forte®

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	General PA Form

Maxidex®, Prednisolone sodium phosphate, & Pred Forte® PA Criteria:
Standard NP Criteria

XIII. OTICS

Otic Quinolones

Ciprofloxacin Otic 0.2% solution

PDL Status	P
Quantity Limit	14/Rx
PA Criteria	None

Ofloxacin otic solution

PDL Status	P
Quantity Limit	10 mL/Rx
PA Criteria	None

Otic Steroid/Antibiotic Combinations

Hydrocortisone/neomycin/polymyxin B otic solution, otic suspension

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Ciprofloxacin/dexamethasone otic suspension

PDL Status	P
Quantity Limit	7.5 mL/Rx
PA Criteria	None

Cipro HC® otic suspension

PDL Status	P
Quantity Limit	10 mL/Rx
PA Criteria	None

Ciprofloxacin/fluocinolone preservative free otic solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Ciprofloxacin/fluocinolone preservative free otic solution PA Criteria:
Standard NP Criteria

Ciprofloxacin/hydrocortisone otic suspension

PDL Status	NP
Quantity Limit	10 mL/Rx
PA Criteria	See Below
PA Form	General PA Form

Ciprofloxacin/hydrocortisone otic suspension PA Criteria:
Standard NP Criteria

Miscellaneous Otics

Please refer to OTC Drug Formulary for coverage of OTC products.

Hydrocortisone/Acetic Acid Otic Solution

PDL Status	P
Quantity Limit	10 mL/Rx
PA Criteria	None

Fluocinolone acetonide (otic) oil 0.01%

PDL Status	P
Quantity Limit	20 mL/Rx
PA Criteria	None

Clotic® solution

PDL Status	NP
Quantity Limit	28/Rx
PA Criteria	See below
PA Form	General PA Form

Clotic® solution l PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

DermOtic® otic oil

PDL Status	NP
Quantity Limit	20 mL/Rx
PA Criteria	See below
PA Form	General PA Form

DermOtic® otic oil PA Criteria:
Standard NP Criteria

XIV. RARE CONDITIONS

Activated PI3K Delta Syndrome (APDS)

Joenja®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Joenja® Initial PA Criteria:

- Patient is ≥ 4 years of age and weighs ≥ 27 kg; **AND**
- Diagnosis of activated phosphoinositide 3-kinase delta syndrome (APDS); **AND**
- Diagnosis has been confirmed by the presence of an APDS-associated genetic variant in either PIK3CD or PIK3R1; **AND**
- Documentation of clinical findings and manifestations consistent with APDS (e.g., recurrent respiratory tract infections and viral infections, lymphadenopathy, hepatosplenomegaly, autoimmune cytopenia); **AND**
- Prescribed by, or in consultation with, hematologist, allergist, or immunologist; **AND**
- For patients with reproductive potential, the prescriber attests to all of the following:
  - Patient is not pregnant prior to initiation of therapy
  - Patient has been counseled on potential risk during pregnancy
  - Patient has been advised to use effective contraception during treatment and for 1 week after the last dose
  - Patient has been advised to not breastfeed during treatment and for 1 week after the last dose

Joenja® Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decreased lymph node size, increased functional B cell counts, decreased infections/hospitalizations, and decreased utilization of immunoglobulin replacement therapy)

Alkaptonuria (AKU)

Harliku® tablets

PDL Status	Awaiting PAC Review
Quantity Limit	1 tablet/day
PA Criteria	See below
PA Form	General PA Form

Harliku® tablets Initial PA Criteria:

- Diagnosis of alkaptonuria (AKU); **AND**
- Used for the reduction of urine homogentisic acid; **AND**
- Disease is confirmed by ONE of the following:
  - Presence of a mutation in the homogentisate 1,2-dioxygenase (HGD) gene as detected by an FDA-approved test
  - Presence of elevated homogentisic acid (HGA) levels in urine greater than 0.4 g/24h; **AND**
- Prescribed by or in consultation with a specialist (i.e., someone experienced with treatment of inborn errors of metabolism or disorder manifestations such as rheumatologist, cardiologist, orthopedist, neurologist, hepatologist, dermatologist, urologist)

Harliku® tablets Renewal PA Criteria:

- Patient demonstrates positive clinical response to therapy (e.g., reduced levels of urinary HGA)

Amyotrophic Lateral Sclerosis (ALS)

Radicava ORS®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Radicava ORS® Initial PA Criteria:

- Submission of medical records (e.g., chart notes, diagnostic tests, nerve conduction studies, lab values) to support a diagnosis of "definite" or "probable" ALS per the revised EL Escorial diagnostic criteria; **AND**
- Prescribed by, or in consultation with, a neurologist; **AND**
- Patient has scores ≥ 2 in all items of the ALS Functional Rating Scale-Revised (ALSFRS-R) criteria at the start of treatment; **AND**
- Patient has a forced vital capacity (%FVC) greater than or equal to 80% at the start of treatment
- Patient must not be pregnant

Radicava ORS® Renewal PA Criteria

- Prescribed by, or in consultation with, a neurologist; **AND**
- Documentation of positive clinical response to therapy (e.g., slowing in the decline of functional abilities); **AND**
- Patient is not dependent on invasive ventilation or tracheostomy

Teglutik®

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 6 months

Teglutik® PA Criteria:

- Diagnosis of Amyotrophic Lateral Sclerosis (ALS); **AND**
- Patient is unable to swallow tablets

Antineutrophil Cytoplasmic Autoantibody (ANCA)

Tavneos®

PDL Status	NP
Quantity Limit	6 caps/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Tavneos® Initial PA Criteria:

- Diagnosis of severe active anti-neutrophil cytoplasmic autoantibody ANCA-associated vasculitis (granulomatosis with polyangiitis [GPA] or microscopic polyangiitis [MPA]) confirmed by ONE of the following:
  - ANCA test positive for proteinase 3 (PR3) antigen
  - ANCA test positive for myeloperoxidase (MPO) antigen
  - Tissue biopsy; **AND**
- Prescribed by, or in consultation with, a rheumatologist, nephrologist, pulmonologist, or a provider with expertise in vascular medicine; **AND**
- Will be used as adjunctive therapy with standard therapy (e.g., cyclophosphamide, azathioprine, mycophenolate, rituximab); **AND**
- Patient is concurrently on glucocorticoids or has an intolerance or contraindication to glucocorticoids

Tavneos® Renewal PA Criteria:

- Patient continues to meet initial approval criteria; **AND**
- Disease response to therapy and tolerability compared to baseline

CHAPLE Disease

Veopoz®

PDL Status	NP
Quantity Limit	8 vials/28 days
PA Criteria	See below
PA Form	General PA Form

Veopoz® Initial PA Criteria:

- Diagnosis of CD55-deficient protein-losing enteropathy (CHAPLE disease); **AND**
- Patient has documentation of genetic testing confirming biallelic CD55 loss-of-function mutation; **AND**
- Prescriber attests to ALL of the following:
  - Patient has received *or will receive* Veopoz IV loading dose;
  - Patient has completed or updated meningococcal vaccination at least 2 weeks prior to administering the first dose of Veopoz unless the risk of delaying therapy outweighs the risk; **AND**
- Prescribed by or in consultation with a clinical specialist knowledgeable in appropriate disease management (e.g., geneticist, gastroenterologist, hematologist)

Veopoz® Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., normalization of serum albumin, decreased abdominal pain, diarrhea, facial edema, and peripheral edema)

Congenital Adrenal Hyperplasia (CAH)

Crenessity®

PDL Status	NP
Quantity Limit	50 mg: 2/day; 100 mg: 4/day; Solution: 4 mL/day
PA Criteria	See below
PA Form	General PA Form

Crenessity® Initial PA Criteria:

- Patient is 4 years of age and older; **AND**
- Diagnosis of classic congenital adrenal hyperplasia (CAH); **AND**
- Patient is currently receiving glucocorticoid replacement therapy for adrenal insufficiency; **AND**
- Crenessity will be used as adjunctive treatment with glucocorticoid replacement to control androgens; **AND**
- Prescribed by, or in consultation with, an endocrinologist

Crenessity® Renewal PA Criteria:

- Crenessity will be used as adjunctive treatment with glucocorticoid replacement to control androgens; **AND**
- Positive response to therapy (e.g., improvement or stabilization in steroid regimen or androgen levels)

Duchenne Muscular Dystrophy (DMD)

Emflaza® suspension, tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Emflaza® suspension, tablets Initial PA Criteria:

- Documentation of a confirmed diagnosis of Duchenne muscular dystrophy (DMD); **AND**
- Age ≥ 2 years; **AND**
- Patient has experienced ≥ ONE of the following adverse reactions directly attributable to therapy with prednisone:
- Significant weight gain (e.g., crossing 2 percentile lines and/or reaching 98th percentile for age and sex)
- Significant behavioral changes negatively impacting function at school, home, day care, etc.

Emflaza® Renewal PA Criteria:

- Patient has received benefit from therapy [e.g., stability or slowing in the decline of symptoms (motor function, respiratory function, musculature strength), quality of life]

Agamree® suspension

PDL Status	NP
Quantity Limit	3 bottles (300 mL)/month
PA Criteria	See below
PA Form	General PA Form

Agamree® suspension PA Criteria:

See Emflaza® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to Emflaza®

Deflazacort suspension, tablets, Kymbee® suspension, tablets, Jaythari® suspension, tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Deflazacort suspension, tablets PA Criteria:

See Emflaza® prior authorization criteria; **AND**

- Clinically valid reason why preferred Emflaza® cannot be used

Duvyzat® suspension

PDL Status	NP
Quantity Limit	12 mL/day
PA Criteria	See below
PA Form	General PA Form

Duvyzat® suspension Initial PA Criteria:

- Documentation of a confirmed diagnosis of Duchenne muscular dystrophy (DMD); **AND**
- Age ≥ 6 years; **AND**
- Patient retains meaningful voluntary motor function (e.g., patient can speak, manipulate objects using upper extremities, ambulate); **AND**
- Will be used in combination with a stable dose of corticosteroids (e.g. prednisone, Emflaza, Agamree); **AND**
- Prescribed by, or in consultation, with a neuromuscular specialist or medical geneticist

Duvyzat® Renewal PA Criteria:

- Patient retains meaningful voluntary motor function (e.g., patient can speak, manipulate objects using upper extremities, ambulate); **AND**
- Will be used in combination with a stable dose of corticosteroids (e.g. prednisone, Emflaza, Agamree); **AND**
- Patient has received benefit from therapy [e.g., stability or slowing in the decline of symptoms (motor function, respiratory function, musculature strength), quality of life]

Fatty Acid Oxidation Disorder (FAOD)

Dojolvi®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Dojolvi® Initial PA Criteria:

- Diagnosis of long-chain fatty acid oxidation disorders (LC-FAOD) as confirmed by two of the following:
  - Acylcarnitine profile
  - Molecular/genetic test
  - Fibroblast test; **AND**
- Patient does not have pancreatic insufficiency; **AND**
- Prescribed by, or in consultation with, a clinical specialist knowledgeable in appropriate disease-related dietary management (e.g., geneticist, cardiologist, gastroenterologist, etc.); **AND**
- For patients receiving another medium-chain triglyceride product, discontinue prior to the first dose of Dojolvi®

Dojolvi® Renewal PA Criteria:

- Evidence of positive clinical response from baseline (e.g., reduction in signs/symptoms such as hypoglycemia, hepatopathy, skeletal myopathy, rhabdomyolysis, cardiomyopathy, etc.)

Familial Chylomicronemia Syndrome (FCS)

Tryngolza®

PDL Status	NP
Quantity Limit	1/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Familial Chylomicronemia Syndrome

Tryngolza® Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of Familial Chylomicronemia Syndrome (FCS) confirmed by ONE of the following:
- Genetic variants in the FCS-causing genes (e.g. GPIHBP1, APOA5I, APOC2I and LMF1)
- A North American FCS (NAFCS) Score of greater than or equal to 45
- Fasting triglyceride levels > 880 mg/dL with lack of response to triglyceride lowering therapy (e.g. statins, fibrates, omega-3 fatty acids); **AND**
- Prescriber attests patient will follow a low-fat diet (≤ 20 g/day) while on Tryngolza; **AND**
- Prescribed by, or in consultation with, a cardiologist, endocrinologist, gastroenterologist, or lipidologist

Tryngolza® Renewal PA Criteria

- Documentation of positive clinical response to therapy (e.g., triglyceride reduction from baseline, decreased incidence of acute pancreatitis)

Severe Hypertriglyceridemia

See Interim Criteria for Agents Awaiting PAC Review

Fibrodysplasia ossificans progressive (FOP)

Sohonos®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sohonos® PA Criteria:

- Diagnosis of fibrodysplasia ossificans progressive (FOP); **AND**
- One of the following:
  - Female aged ≥ 8 years of age
  - Male aged ≥ 10 years of age; **AND**
- Diagnosis of FOP confirmed by one of the following:
  - Mutation in the ALK2/ACVR1 gene
  - Classic FOP clinical features such as malformation of big toe and progressive heterotopic endochondral ossification in ribbons, sheets, and plates
  - Radiographic bone scans detecting heterotopic ossification (HO); **AND**
- Prescriber attests to all of the following:
  - Patient is not pregnant
  - Female patients of reproductive potential will be counseled to use effective contraception during treatment with therapy and for at least 1 month after last dose
  - For pediatric patients, premature epiphyseal closure has not occurred

Friedreich's Ataxia

Skyclarys®

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See below
PA Form	General PA Form

Skyclarys® Initial PA Criteria:

- Patient is ≥ 16 years old; **AND**
- Patient has diagnosis of Friedreich's ataxia (FA); **AND**
- Patient has documentation of genetic testing confirming frataxin (FXN) gene mutation; **AND**
- Prescribed by, or in consultation with, a neurologist, geneticist, or cardiologist

Skyclarys® Renewal PA Criteria

- Patient has disease stabilization or clinical response to therapy

Glucagon-Like Peptide-2 (GLP-2) Analog

Gattex®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Gattex® Initial PA Criteria:

- Diagnosis of short bowel syndrome, **AND**
- Patient is dependent on parenteral nutrition and/or fluids/electrolytes; **AND**
- Submission of medical records (e.g. chart notes) documenting that patient has been unable to significantly reduce PN/IV support

Gattex® Renewal PA Criteria:

- Submission of medical records (e.g. chart notes) demonstrating a positive response to therapy (e.g. decreased frequency or volume of parenteral nutrition and/or fluids/electrolytes from baseline)

Hereditary Angioedema (HAE) Agents

Sajazir® prefilled syringe

PDL Status	P
Quantity Limit	6 injections/28 days
PA Criteria	See below
PA Form	General PA Form

Sajazir® prefilled syringe Initial PA Criteria:

- Prescribed by, or in consultation with, a specialist in allergy, immunology, hematology, pulmonology, or genetics; **AND**
- Patient is ≥ 18 years of age; **AND**
- Submission of medical records (e.g. chart notes and labs) documenting a diagnosis consistent with 1 of the following HAE subtypes:
  - Type I:
    - Low C1 inhibitor (C1-INH) antigenic level (below the limit of normal defined by the lab performing the test); **AND**
    - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **OR**
  - Type II:
    - Normal to elevated C1-INH antigenic level; **AND**
    - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **OR**
  - Type III:
    - Normal C1-INH antigenic level; **AND**
    - One of the following:
      - Confirmed presence of a FXII, angiopoietin-1, plasminogen, KNG1, MYOF, or HS3ST6 gene mutation; **OR**
      - Recurring angioedema attacks that are refractory to high-dose antihistamines with confirmed family history of angioedema; **AND**
- Medication will be using to treat acute HAE attacks; **AND**
- Medication will not be used in combination with other approved treatments for acute HAE attacks; **AND**
- Prescriber attests patient is avoiding all possible triggers for HAE attacks

Sajazir® prefilled syringe Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., disappearance of swelling from an acute attack, relief from pain); **AND**
- Absence of unacceptable toxicity from drug (e.g., severe hypersensitivity reactions)

Takhzyro® injection

PDL Status	P
Quantity Limit	2 injections/28 days
PA Criteria	See below
PA Form	General PA Form

Takhzyro® injection Initial PA Criteria:

- Prescribed by, or in consultation with, a specialist in allergy, immunology, hematology, pulmonology, or genetics; **AND**
- Patient must be ≥ the labeled age minimum (Andembry ≥ 12 years; Dawnzera ≥ 12 years; Haegarda ≥6 years; Orladeyo ≥12 years; Takhzyro ≥2 years); **AND**
- Submission of medical records (e.g. chart notes and labs) documenting a diagnosis consistent with 1 of the following HAE subtypes:
  - Type I:
    - Low C1 inhibitor (C1-INH) antigenic level (below the limit of normal defined by the lab performing the test); **AND**
    - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **OR**
  - Type II:
    - Normal to elevated C1-INH antigenic level; **AND**
    - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **AND**
  - Type III:
    - Normal C1-INH antigenic level; **AND**
    - One of the following:
      - Confirmed presence of a FXII, angiopoietin-1, plasminogen, KNG1, MYOF, or HS3ST6 gene mutation; **OR**
      - Recurring angioedema attacks that are refractory to high-dose antihistamines with confirmed family history of angioedema; **AND**
- Will not be used in combination with other routine prophylaxis HAE agents (e.g., Haegarda, Takhzyro, Orladeyo); **AND**
- Prescriber attests patient is avoiding all possible triggers for HAE attacks

Takhzyro® injection Renewal PA Criteria:

- Improvement in severity and duration of attacks have been achieved and sustained; **AND**
- Absence of unacceptable toxicity from the drug (e.g., severe hypersensitivity reactions, thromboembolic events)

Andembry® autoinjector

PDL Status	NP
Quantity Limit	Loading dose: 2 syringes Maintenance dose: 1 syringe/month
PA Criteria	See below
PA Form	General PA Form

Andembry® autoinjector PA Criteria:

See Takhzyro® prior authorization criteria

Dawnzera® autoinjector

PDL Status	NP
Quantity Limit	1 autoinjector/month
PA Criteria	See below
PA Form	General PA Form

Dawnzera® autoinjector PA Criteria:

See Takhzyro® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to all preferred HAE prophylaxis agents;

Ekterly® tablets

PDL Status	NP
Quantity Limit	8 tablets/month
PA Criteria	See below
PA Form	General PA Form

Ekterly® tablets PA Criteria:

See Sajazir® prior authorization criteria; **AND**

- For patients 18 years of age or older, clinically valid reason why preferred injectable (e.g., Sajazir®) cannot be used.

Firazyr® prefilled syringe

PDL Status	NP
Quantity Limit	6 injections/month
PA Criteria	See below
PA Form	General PA Form

Firazyr® prefilled syringe PA Criteria:

See Sajazir® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to 2 preferred agents

Haegarda® injection

PDL Status	NP
Quantity Limit	2 injections/28 days
PA Criteria	See below
PA Form	General PA Form

Haegarda® injection PA Criteria:

See Takhzyro® prior authorization criteria

Icatibant prefilled syringe

PDL Status	NP
Quantity Limit	6 injections/28 days
PA Criteria	See below
PA Form	General PA Form

Icatibant prefilled syringe PA Criteria:

See Sajazir® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to 2 preferred agents

Orladeyo® capsules, Orladeyo® packets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

- **Orladeyo® capsules, Orladeyo® packets Initial PA Criteria:**
  - Prescribed by, or in consultation with, a specialist in allergy, immunology, hematology, pulmonology, or genetics; **AND**
  - Patient must be ≥ the labeled age minimum (Andembry ≥ 12 years; Dawnzera ≥ 12 years; Haegarda ≥6 years; Orladeyo ≥ 2 years, Takhzyro ≥ 2 years); **OR**
  - One of the following:
    - If the request is for Orladeyo packets, the patient is 2-11 years old; **OR**
    - If the request is for Orladeyo capsules, the patient is ≥ 12 years old; **AND**
  - Submission of medical records (e.g. chart notes and labs) documenting a diagnosis consistent with 1 of the following HAE subtypes:
    - Type I:
      - Low C1 inhibitor (C1-INH) antigenic level (below the limit of normal defined by the lab performing the test); **AND**
      - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **OR**
    - Type II:
      - Normal to elevated C1-INH antigenic level; **AND**
      - Low C1-INH functional level (below the limit of normal defined by the lab performing the test); **OR**
    - One of the following:
      - Confirmed presence of a FXII, angiopoietin-1, plasminogen, KNG1, MYOF, or HS3ST6 gene mutation; **OR**
      - Recurring angioedema attacks that are refractory to high-dose antihistamines with confirmed family history of angioedema; **AND**
  - Will not be used in combination with other routine prophylaxis HAE agents (e.g., Haegarda, Takhzyro, Orladeyo); **AND**
  - Prescriber attests patient is avoiding all possible triggers for HAE attacks;

Orladeyo® capsules, Orladeyo® packets Renewal PA Criteria:

- Improvement in severity and duration of attacks have been achieved and sustained; **AND**
- Absence of unacceptable toxicity from the drug (e.g., severe hypersensitivity reactions, thromboembolic events)

Homocystinuria Agents

Cystadane®

PDL Status	P
Quantity Limit	6 g/day
PA Criteria	See below
PA Form	General PA Form

Cystadane® PA Criteria:

- Diagnosis of moderate to severe hyperhomocysteinemia
- Genetic test confirming ONE of the following:
  - Cystathionine beta-synthase (CBS) deficiency
  - 5,10-methylenetetrahydrofolate reductase (MTHRF) deficiency
  - Cobalamin cofactor metabolism (cbl) defect; **AND**
- Prescribed by, or in consultation with, a physician who specializes in the treatment of inherited metabolic disorders; **AND**
- Patient had an inadequate response or is unable to be managed by diet and vitamin supplementation with folic acid, vitamin B12, and vitamin B6

Betaine anhydrous powder

PDL Status	NP
Quantity Limit	6 g/day
PA Criteria	See below
PA Form	General PA Form

Betaine anhydrous powder PA Criteria:

See Cystadane® prior authorization criteria; **AND**

- Clinically valid reason why preferred Cystadane® cannot be used

Hutchinson-Gilford Progeria Syndrome

Zokinvy®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Zokinvy® Initial PA Criteria:

- Patient has a diagnosis of Hutchinson-Gilford Progeria Syndrome; **OR**
- Patient has processing deficient Progeroid Laminopathies with either:
  - Heterozygous LMNA mutation with progerin-like protein accumulation
  - Homozygous or compound heterozygous ZMPSTE24 mutations; **AND**
  - Patient must be 12 months of age or older; **AND**
- Patient must have a body surface area (BSA) of 0.39 m2 and above; **AND**
- Females must use effective contraception due to embryo-fetal toxicity; **AND**
- Patient must not meet any of the following:
  - Other Progeroid Syndromes or processing proficient Progeroid Laminopathies
  - Concomitant use of strong or moderate CYP3A inhibitors or inducers
  - Concomitant use of midazolam
  - Concomitant use of lovastatin, simvastatin, and atorvastatin
  - Patient is pregnant

Zokinvy® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has experienced a positive response to therapy, as documented by provider; **AND**
- Patient has not experienced treatment-limiting adverse effects (e.g., laboratory Abnormalities: changes in electrolytes, complete blood counts, and liver enzymes, decrease in renal function, retinal toxicity)

Hyperoxaluria Agents

Rivfloza®

PDL Status	NP
Quantity Limit	1/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Rivfloza® PA Criteria:

- Patient is 2 years of age or older; **AND**
- Diagnosis of primary hyperoxaluria type 1 (PH1) confirmed by ONE of the following:
  - Genetic testing demonstrating mutation in the alanine-glyoxylate aminotransferase (AGXT) gene
  - Liver biopsy demonstrating absent or reduced alanine-glyoxylate aminotransferase (AGT) activity; **AND**
- Patient has ONE of the following*:*
  - Elevated urinary oxalate excretion
  - Elevated plasma oxalate levels
  - Urinary oxalate creatinine ratio above the age-specific upper limit of normal
- Patient has relatively preserved kidney function (e.g., eGFR ≥ 30 mL/min/1.73 m^2); **AND**
- Prescribed by, or in consultation with, a hematologist, nephrologist, urologist, or geneticist

Rivfloza® Renewal PA Criteria:

- Patient has positive clinical response to therapy (e.g., decreased urinary oxalate excretion or plasma concentration, decreased number or size of kidney stones, improved kidney function)

Hypophosphatasia (HPP) Agents

Strensiq®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year
Notes	80 mg/0.8 mL vial will not be approved for pediatric patients weighing < 40 kg

Strensiq® Initial PA Criteria:

- Diagnosis of perinatal, infantile, or juvenile-onset hypophosphatasia (HPP); **AND**
- Onset of clinical signs and symptoms of HPP prior to age 19 years (e.g., rickets, skeletal deformities, fractures, respiratory compromise, vitamin B6 dependent seizure, craniosynostosis, dental abnormalities, severe osteopenia); **AND**
- Clinical diagnosis of HPP evidenced by one of the following:
  - Serum alkaline phosphatase (ALP) below age-adjusted normal range
  - Genetic confirmation of ALPL mutation;
  - Elevated plasma pyridoxal 5'-phosphate (PLP) levels; **AND**
- Prescribed by, or in consultation with, a physician who specializes in the treatment of inherited metabolic disorders

Strensiq® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., healing of the skeletal manifestations, improved respiratory, motor function, and linear growth); **AND**
- Prescribed by, or in consultation with, a physician who specializes in the treatment of inherited metabolic disorders

IBAT (Ileal Bile Acid Transporter) Inhibitors

Bylvay®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Bylvay® PA Criteria:

- One of the following:
  - Diagnosis of progressive familial intrahepatic cholestasis (PFIC); **AND**
    - Patient does not have ABCB11 variant resulting in non-functional or complete absence of bile salt export pump protein (BSEP-3)
  - Diagnosis of Alagille syndrome (ALGS) confirmed by presence of the JAG1 or Notch2 gene mutation; **AND**
- Prescribed by, or in consultation with, hepatologist or gastroenterologist; **AND**
- Patient is experiencing moderate to severe pruritus confirmed by ONE of the following:
  - Total serum bile acid > 3x the upper limit of normal
  - Conjugated bilirubin > 1 mg/dL.
  - Fat soluble vitamin deficiency otherwise unexplainable.
  - GGT > 3x the upper limit of normal
  - Intractable pruritus explainable only by liver disease; **AND**
- Trial and failure to at TWO other conventional treatments for the symptomatic relief of pruritus (e.g., bile acid-binding agents, naltrexone, phenobarbital, rifampin, ursodiol); **AND**
- Provider attests to monitor the following:
  - Liver-function tests at baseline and during treatment
  - Fat-soluble vitamin (FSV) levels at baseline and during treatment

Livmarli® solution, tablets

PDL Status	NP
Quantity Limit	Solution: None Tablets: 1/day
PA Criteria	See below
PA Form	General PA Form

Livmarli® solution, tablets PA Criteria:

- One of the following:
  - Diagnosis of progressive familial intrahepatic cholestasis (PFIC); **AND**
    - Patient does not have ABCB11 variant resulting in non-functional or complete absence of bile salt export pump protein (BSEP-3)
  - Diagnosis of Alagille syndrome (ALGS) confirmed by presence of the JAG1 or Notch2 gene mutation; **AND**
- Prescribed by, or in consultation with, hepatologist or gastroenterologist; **AND**
- Patient is experiencing moderate to severe pruritus confirmed by ONE of the following:
  - Total serum bile acid > 3x the upper limit of normal
  - Conjugated bilirubin > 1 mg/dL.
  - Fat soluble vitamin deficiency otherwise unexplainable.
  - GGT > 3x the upper limit of normal
  - Intractable pruritus explainable only by liver disease; **AND**
- Trial and failure to at TWO other conventional treatments for the symptomatic relief of pruritus (e.g., bile acid-binding agents, naltrexone, phenobarbital, rifampin, ursodiol); **AND**
- Provider attests to monitor the following:
  - Liver-function tests at baseline and during treatment
  - Fat-soluble vitamin (FSV) levels at baseline and during treatment

IgA Nephropathy (IgAN)

Fabhalta® capsules

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Fabhalta® capsules Initial PA Criteria:

- One of the following:
  - Diagnosis of primary IgA immunoglobulin A nephropathy (IgAN) confirmed by kidney biopsy; **AND**
    - Patient is at risk of rapid disease progression (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
    - Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor; **AND**
    - Patient has an eGFR > 20 mL/min/1.73 m2; **OR**
  - Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) confirmed by flow cytometry; **AND**
    - Patient has symptoms attributed to PNH (e.g., anemia, extreme fatigue, difficulty swallowing, recurrent abdominal pain, smooth muscle dystonia, thrombosis, CKD, organ damage secondary to chronic hemolysis); **AND**
  - Diagnosis of complement 3 glomerulopathy (C3G) confirmed by kidney biopsy; **AND**
    - Medication is being used to reduce proteinuria (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
    - Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor; **AND**
    - Patient has an eGFR > 30 mL/min/1.73 m2; **AND**
- Prescribed by, or in consultation with, a hematologist, nephrologist, or oncologist; **AND**
- Prescriber is enrolled in the Fabhalta REMS Program; **AND**
- One of the following:
  - Will not be used concurrently with another complement inhibitor; **OR**
  - Patient is currently receiving another complement inhibitor (e.g., Empaveli, Soliris, Ultomiris) which will be discontinued and Fabhalta will be initiated in accordance with the FDA approved labeling

Fabhalta® capsules Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., increased or stabilization of hemoglobin levels, reduction in transfusions, improvement in hemolysis, decrease in LDH); **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist; **AND**
- Patient is not receiving Fabhalta® in combination with another complement inhibitor

Filspari® tablets

PDL Status	NP
Quantity Limit	200 mg tablets: 1/day; 400 mg tablets: 2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Filspari® tablets Initial PA Criteria:

- One of the following:
  - All of the following:
    - Diagnosis of primary IgA immunoglobulin A nephropathy (IgAN) confirmed by kidney biopsy; **AND**
    - Patient is 18 years of age or older; **AND**
    - Patient is at risk of rapid disease progression (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
    - Patient has an eGFR > 30 mL/min/1.73 m2; **AND**
    - Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor; **OR**
  - All of the following:
    - Diagnosis of proteinuria associated with focal segmental glomerulosclerosis (FSGS) without nephrotic syndrome confirmed by one of the following:
      - Kidney biopsy
      - Documentation of a genetic mutation in a podocyte protein associated with FSGS (e.g., CD2AP, INF2, NPHS1, NPHS2); **AND**
    - Patient has a UPCR ≥ 1.5 g/g (170 mg/mmol); **AND**
    - Patient has an eGFR ≥30 mL/min/1.73 m2; **AND**
    - Patient is 8 years of age or older; **AND**
- Use of renin-angiotensin-aldosterone system (RAAS) inhibitors (e.g., ACE inhibitors, ARBs), endothelin receptor antagonists (e.g., Letairis, Opsumit, Tracleer), and aliskiren will be discontinued prior to initiating treatment; **AND**
- Prescriber and patient are enrolled in the Filspari REMS program; **AND**
- Prescribed by, or in consultation with, a nephrologist

Filspari® tablets Renewal PA Criteria:

- Patient has positive clinical response to Filspari therapy (e.g., reduction of proteinuria from baseline, decreased UPCR)

Tarpeyo® capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Tarpeyo® capsules PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of primary IgA immunoglobulin A nephropathy (IgAN) confirmed by kidney biopsy; **AND**
- Patient is at risk of rapid disease progression (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
- Patient has an eGFR > 30 mL/min/1.73 m2; **AND**
- Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor, unless contraindicated; **AND**
- Prescribed by, or in consultation with, a nephrologist

Vanrafia® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Vanrafia® tablets PA Criteria:

See Filspari® prior authorization criteria

Voyxact® prefilled syringe

PDL Status	Awaiting PAC Review
Quantity Limit	2 mL/28 days
PA Criteria	See below
PA Form	General PA Form
PA Duration	Initial: 6 months; Renewal: 1 year

Voyxact® prefilled syringe Initial PA Criteria:

- Patient is 18 years or older; **AND**
- Diagnosis of primary IgA immunoglobulin A nephropathy (IgAN) confirmed by kidney biopsy; **AND**
- Patient is at risk of rapid disease progression (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
- Prescriber attests patient is being monitored for active infections during therapy and that no live vaccines have been received 30 days prior to starting Voyxact® or while receiving treatment with Voyxact®; **AND**
- Inadequate response to a stable maximally tolerated doses of an ARB or ACE inhibitor for a trial of 3 months or greater; **AND**
- Submission of recent medical records (e.g., chart notes and/or labs) documenting that patient has an EGFR greater than or equal to 30 mL/min/1.73 m^2 AND proteinuria greater than or equal to 0.75g/day; **AND**
- Patient is not concurrently taking other treatments for IgAN (e.g., Fabhalta®, Tarpeyo®, etc.); **AND**
- Prescribed by, or in consultation with, a nephrologist

Voyxact® prefilled syringe Renewal PA Criteria:

- Patient has a positive clinical response to Voyxact® therapy (e.g., reduction of proteinuria from baseline, decreased UPCR)

IGF-1 Deficiency

Increlex®

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year
Notes	Will not be approved for patients with active or secondary neoplasms, secondary forms of IGF-1 deficiency, weight loss management, nor as a substitute for growth hormone.

Increlex® Initial PA Criteria:

- Patient is < 21 years old; **AND**
- Epiphyses is open (therapy will not be approved once epiphyseal fusion occurs); **AND**
- One of the following:
  - Diagnosis of growth failure due to severe primary IGF-1 deficiency defined by the following (documentation required):
    - Height standard deviation score ≤ -3
    - Basal IGF-1 standard deviation score ≤ -3
    - Normal or elevated growth hormone
  - Diagnosis of growth hormone (GH) gene deletion in a patient who has developed neutralizing antibodies to GH; **AND**
    - Secondary causes of IGF-1 deficiency have been ruled out (e.g., hypothyroidism, malnutrition, hepatic disease, GHD, chronic corticosteroid treatment); **AND**
- Secondary causes of IGF-1 deficiency have been ruled out (e.g., hypothyroidism, malnutrition, hepatic disease, GHD, chronic corticosteroid treatment); **AND**
- Patient will not be treated with concurrent growth hormone therapy

Increlex® Renewal PA Criteria:

- Patient is < 21 years old; **AND**
- Prescriber attests patient has had a height increase of ≥ 2 cm/year over the previous year of treatment; **AND**
- Epiphyses is open; **AND**
- Patient is not treated with concurrent growth hormone therapy

Lambert-Eaton Myasthenic Syndrome (LEMS)

Firdapse®

PDL Status	NP
Quantity Limit	10/day
PA Criteria	See below
PA Form	General PA Form

Firdapse® Initial PA Criteria:

- Diagnosis of Lambert-Eaton Myasthenic Syndrome (LEMS) confirmed by a positive anti-P/Q type voltage-gated calcium channel antibody test; **AND**
- Patient is ≥ 6 years old; **AND**
- Patient does not have a history of seizures; **AND**
- Patient does not have a hypersensitivity to amifampridine or another aminopyridine (such as dalfampridine [Ampyra®])

Firdapse® Renewal PA Criteria:

- Patient has not experienced any treatment-restricting adverse effects; **AND**
- Patient must demonstrate disease improvement, stabilization, and/or slowing in the rate of decline due to the medication

Leptin Deficiency

Myalept®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Myalept® Initial PA Criteria:

- Diagnosis of congenital or acquired lipodystrophy**; AND**
- Leptin deficiency confirmed by laboratory testing; **AND**
- Patient has one of the following complications of lipodystrophy:
  - Diabetes mellitus
  - Hypertriglyceridemia
  - Hepatic steatosis
  - Polycystic ovarian syndrome
  - Acanthosis nigricans; **AND**
- Requested agent will be used as adjunct to dietary management of lipodystrophy; **AND**
- Documented baseline HbA1C, fasting glucose, triglycerides, and liver enzymes provided; **AND**
- Patient does NOT have HIV-related or partial lipodystrophy or metabolic disease without concurrent evidence of generalized lipodystrophy; **AND**
- Prescriber is enrolled in the Myalept REMS program

Myalept® Renewal PA Criteria:

- Documented positive clinical response to therapy (e.g., improved glycemic control, decrease in triglycerides)

Lysosomal Storage Disease

Aqneursa®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Aqneursa® Initial PA Criteria:

- Patient weighs at least 15 kg; **AND**
- Diagnosis of Niemann-Pick disease type C confirmed by one of the following:
- Genetically confirmed mutations in both alleles of NPC1 or NPC2; **OR**
- Mutation in only one allele of NPC1 or NPC2 plus either positive filipin staining or elevated cholestane triol/oxysterols (>2 x upper limit of normal); **AND**
- Patient has at least one neurological symptom of the disease (e.g., hearing loss, vertical supranuclear gaze palsy, ataxia, dementia, dystonia, seizures, dysarthria, or dysphagia); **AND**
- Prescribed by, or in consultation with, a neurologist, a medical geneticist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of lysosomal storage disorders

Aqneursa® Renewal PA Criteria

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., improvement or stabilization in neurological symptoms of disease)

Cerdelga®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form

Cerdelga® PA Criteria:

Standard NP Criteria

Galafold®

PDL Status	NP
Quantity Limit	14/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Galafold® Initial PA Criteria:

- Patient is ≥ 18 years old; **AND**
- Documented diagnosis of Fabry disease with biochemical/genetic confirmation by 1 of the following:
  - Males only: α-galactosidase A (α-Gal A) activity in plasma, isolated leukocytes, and/or cultured cells
  - Plasma or urinary globotriaosylceramide (Gb3/GL-3) or globotriaosylsphingosine (lyso-Gb3)
  - Detection of pathogenic mutations in the GALA/GLA gene by molecular genetic testing; **AND**
- Patient has an amenable GLA mutation (as defined in the migalastat labeling or determined by a clinical genetics professional); **AND**
- Will NOT be used in combination with agalsidase beta or Elfabrio (pegunigalsidase alfa); **AND**
- Prescribed by, or in consultation with, clinical genetics professional with knowledge in management of Fabry disease

Galafold® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests to patient compliance with therapy; **AND**
- Disease response to treatment as defined by a reduction in urine GL-3 and/or GL-3 inclusions compared to pre-treatment baseline or there has been improvement in clinical symptoms (e.g. stabilization of kidney function, slow or prevention of organ function decline); **AND**
- Absence of unacceptable toxicity (e.g., kidney infections);

Miplyffa®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Miplyffa® Initial PA Criteria:

- Patient is 2 years of age or older; **AND**
- Diagnosis of Niemann-Pick disease type C confirmed by ONE of the following:
  - Genetically confirmed mutations in both alleles of NPC1 or NPC2; **OR**
  - Mutation in only one allele of NPC1 or NPC2 plus either positive filipin staining or elevated cholestane triol/oxysterols (>2 x upper limit of normal); **AND**
- Patient has at least one neurological symptom of the disease (e.g., hearing loss, vertical supranuclear gaze palsy, ataxia, dementia, dystonia, seizures, dysarthria, dysphagia); **AND**
- Miplyffa will be used in combination with miglustat; **AND**
- Prescribed by, or in consultation with, a neurologist, a medical geneticist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of lysosomal storage disorders

Miplyffa® Renewal PA Criteria:

- Patient continued to meet initial criteria; **AND**
- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., improvement or stabilization in neurological symptoms of disease)

Opfolda®

PDL Status	NP
Quantity Limit	8/28 days
PA Criteria	See below
PA Form	General PA Form

Opfolda® PA Criteria:

- Patient is ≥ 18 years old and weighs at least 40 kg; **AND**
- Diagnosis of late-onset Pompe disease confirmed by ONE of the following:
  - Documentation demonstrating deficiency of acid alpha-glucosidase (GAA) enzyme activity
  - Molecular genetic test demonstrating pathogenic variants in GAA; **AND**
- Prescriber attest patient did not have clinical improvement on enzyme replacement therapy (e.g., Lumizyme, Nexviazyme, Elfabrio); **AND**
- Must be used in combination with Pombiliti (cipaglucosidase alfa); **AND**
- Prescribed by, or in consultation with, a neurologist, a medical geneticist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of lysosomal storage disorders

Procysbi®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Procysbi® Initial PA Criteria:

- Diagnosis of nephropathic cystinosis; **AND**
- Patient is ≥ 1 year old; **AND**
- Trial and failure, contraindication, or intolerance to Cystagon®; **AND**
- WBC cystine levels or plasma cysteamine concentration will be monitored

Procysbi® Renewal PA Criteria:

- Documentation of positive clinical response to therapy; **AND**
- WBC cystine levels or plasma cysteamine concentration will be monitored

Neuromyelitis Optica Spectrum Disorder (NMOSD)

Enspryng®

PDL Status	NP
Quantity Limit	Loading dose: 3 syringes/28 days; Maintenance: 1/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Enspryng® Initial PA Criteria:

- Diagnosis of neuromyelitis optica spectrum disorder (NMOSD); **AND**
- Patient > 12 years old of age; **AND**
- Patient is anti-aquaporin-4 (AQP4) antibody positive; **AND**
- Patient has been screened, and does not have any of the following:
  - Active Hepatitis B infection
  - Active or untreated latent tuberculosis
  - Active infection; **AND**
- Prescribed by or in consultation with a neurologist or ophthalmologist
- **Ensprying® Renewal PA Criteria:**
- Patient has demonstrated positive response to therapy

Myasthenia Gravis

Zilbrysq®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Zilbrysq® Initial PA Criteria:

- Diagnosis of generalized myasthenia gravis (gMG); **AND**
- Documented positive serology for acetylcholine receptor (AChR) autoantibodies; **AND**
- Patient has a baseline MG-Activities of Daily Living (MG-ADL) total score of ≥6; **AND**
- Patient has tried and failed, or has contraindication, or intolerance to TWO of the following:
  - Corticosteroids
  - Azathioprine
  - Cyclosporine
  - Mycophenolate mofetil
  - Methotrexate
  - Tacrolimus; **AND**
- Prescribed by, or in consultation with, a neurologist or neuromuscular specialist; **AND**
- Prescriber is enrolled in the Zilbrysq REMS Program; **AND**
- Patient has not failed a previous course of Zilbrysq, Ultomiris, or Soliris therapy; **AND**
- Patient is not receiving Zilbrysq in combination with another complement inhibitor (e.g., Soliris, Ultomiris)

Renewal Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., reduction in MG-ADL score or improvement in talking, chewing, swallowing, breathing, double vision, eyelid drop, movement)

Paroxysmal Nocturnal Hemoglobinuria (PNH)

Empaveli® injection

PDL Status	NP
Quantity Limit	200 mL/30 days
PA Criteria	See below
PA Form	General PA Form

Empaveli® injection Initial PA Criteria:

- Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) confirmed by flow cytometry; **AND**
- One of the following:
  - Patient has symptoms attributed to PNH (e.g., anemia, extreme fatigue, difficulty swallowing, recurrent abdominal pain, smooth muscle dystonia; thrombosis, CKD, organ damage secondary to chronic hemolysis); **OR**
  - Diagnosis of C3 glomerulopathy (C3G) or primary immune-complex membranoproliferative glomerulonephritis (IC-MPGN); **AND**
- Prescribed by, or in consultation with, a nephrologist, hematologist or oncologist; **AND**
- One of the following:
  - Will not be used concurrently combination with another complement inhibitor (e.g., Soliris, Ultomiris); **OR**
  - Patient is currently receiving another complement inhibitor (e.g., Fabhalta, Soliris, Ultomiris) which will be discontinued and Empaveli will be initiated in accordance with the FDA approved labeling; **AND**
- One of the following:
  - The requested quantity does not exceed 1,080 mg twice weekly ; **OR**
  - The requested quantity is for 1,080 mg every 3 days and lactate dehydrogenase (LDH) is >2 the upper limit of the normal range (LDH level documentation is required)

Empaveli® injection Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., increased or stabilization of hemoglobin levels, reduction in transfusions, improvement in hemolysis, decrease in LDH); **AND**
- Prescribed by, or in consultation with, a hematologist, hematologist, or oncologist; **AND**
- Patient is not receiving Empaveli in combination with another complement inhibitor; **AND**
- One of the following:
  - The requested quantity does not exceed 1,080 mg twice weekly
  - The requested quantity is for 1,080 mg every 3 days and lactate dehydrogenase (LDH) is >2 the upper limit of the normal range (LDH level documentation is required)

Fabhalta®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Fabhalta® Initial PA Criteria:

- One of the following:
  - Diagnosis of primary IgA immunoglobulin A nephropathy (IgAN) confirmed by kidney biopsy; **AND**
    - Patient is at risk of rapid disease progression (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
    - Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor; **AND**
    - Patient has an eGFR > 20 mL/min/1.73 m2; **OR**
  - Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) confirmed by flow cytometry; **AND**
    - Patient has symptoms attributed to PNH (e.g., anemia, extreme fatigue, difficulty swallowing, recurrent abdominal pain, smooth muscle dystonia, thrombosis, CKD, organ damage secondary to chronic hemolysis); **AND**
  - Diagnosis of complement 3 glomerulopathy (C3G) confirmed by kidney biopsy; **AND**
    - Medication is being used to reduce proteinuria (e.g., proteinuria > 0.5 g/day or UPCR > 0.8 g/g); **AND**
    - Inadequate response to a stable max tolerated doses of an ARB or an ACE inhibitor; **AND**
    - Patient has an eGFR > 30 mL/min/1.73 m2; **AND**
- Prescribed by, or in consultation with, a hematologist, nephrologist, or oncologist; **AND**
- Prescriber is enrolled in the Fabhalta REMS Program; **AND**
- One of the following:
  - Will not be used concurrently with another complement inhibitor; **OR**
  - Patient is currently receiving another complement inhibitor (e.g., Empaveli, Soliris, Ultomiris) which will be discontinued and Fabhalta will be initiated in accordance with the FDA approved labeling

Fabhalta® Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g., increased or stabilization of hemoglobin levels, reduction in transfusions, improvement in hemolysis, decrease in LDH); **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist; **AND**
- Patient is not receiving Fabhalta® in combination with another complement inhibitor

Voydeya®

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Voydeya® Initial PA Criteria:

- Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) confirmed by flow cytometry; **AND**
- Patient has symptoms attributed to PNH (e.g., anemia, extreme fatigue, difficulty swallowing, recurrent abdominal pain, smooth muscle dystonia, thrombosis, CKD, organ damage secondary to chronic hemolysis); **AND**
- Patient is experiencing extravascular hemolysis (EVH) while on complement C5 inhibitor Ultomiris or Soliris; **AND**
- Prescriber attests Voydeya will be used in combination with Ultomiris or Soliris; **AND**
- Prescriber is enrolled in the Voydeya REMS Program; **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist

Voydeya® Renewal PA Criteria:

- Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy (e.g. hemoglobin stabilization, decreased number of blood transfusions, improvement in signs and symptoms of the disease); **AND**
- Prescribed by, or in consultation with, a hematologist or oncologist; **AND**
- Patient will continue to use Voydeya in combination with Ultomiris or Soliris

Phenylketonuria (PKU)

Palynziq® prefilled syringe

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Palynziq® prefilled syringe PA Criteria:

- Diagnosis of Phenylketonuria (PKU); **AND**
- Prescribed by, or in consultation with, a physician who specializes in the treatment of inherited metabolic disorders; **AND**
- Patient will receive first dose of Palynziq® in prescribing MD's office; **AND**
- Patient meets ONE of the following:
  - Patient has blood phenylalanine (Phe) concentrations > 600 µmol/L; **OR**
  - Prescriber attests patient cannot maintain a healthy diet with Phe restriction; **OR**
  - Patient has neurocognitive deficits; **OR**
  - Trial and failure, contraindication, or intolerance of sapropterin

Sapropterin packets, tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sapropterin PA Criteria:

- Patient has diagnosis of Phenylketonuria (PKU); **AND**
- Prescribed by, or in consultation with, a metabolic specialist; **AND**
- Prescriber attests that Phenylalanine (Phe) levels cannot be maintained within recommended range (120-360 umol/L) with dietary intervention alone; **AND**
- Medication will be used in conjunction with a phenylalanine restricted diet

Javygtor® packets, tablets, Kuvan® packets, tablets, Sephience® packets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Javygtor® packets, tablets, Kuvan® packets, tablets, Sephience® packets PA Criteria:

See sapropterin prior authorization criteria; **AND**

- Clinically valid reason why a preferred PKU agent cannot be used

PIK3CA-Related Overgrowth Spectrum (PROS)

Vijoice® packets, tablets

PDL Status	NP
Quantity Limit	6/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Vijoice® Initial PA Criteria:

- Diagnosis of PIK3CA-Related Overgrowth Spectrum (PROS); **AND**
- Patient has a mutation of the PIK3CA gene; **AND**
- Patient is 2 years of age or older; **AND**
- Patient has severe manifestations of PROS and requires systemic therapy; **AND**
- Vijoice will NOT be used for an oncology diagnosis; **AND**
- Prescriber attests to monitor, and potentially discontinue Vijoice treatment, if patient shows any of the following:
  - Signs or symptoms of severe cutaneous adverse reactions (SCARs)
  - New or worsening respiratory symptoms or is suspected to have developed pneumonitis
  - Severe diarrhea
  - Severe hyperglycemia
  - Severe hypersensitivity; **AND**
- Female and male patients of reproductive potential have been advised to use effective contraception during treatment and for one week after the last dose

Vijoice® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Prescriber attests patient has had ≥ 20% reduction from baseline in the measurable target lesion volume confirmed by at least one subsequent imaging assessment

Prader Willi Syndrome (PWS)

Vykat XR®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Vykat XR® Initial PA Criteria:

- Patient is greater than or equal to 4yrs of age; **AND**
- Diagnosis of Prader-Willi Syndrome; **AND**
- Submission of medical records (e.g. chart notes) documenting BOTH of the following:
  - Methylation analysis genetic testing used to detect lack of active genetic material in particular region of Chromosome 15
  - Patient has signs and symptoms of hyperphagia (e.g. intense persistent sensation of hunger, food preoccupations that cause distress or disrupt daily activities, lack of normal satiety, extreme drive to consume food); **AND**
- Prescribed by, or in consultation with an endocrinologist, geneticists, neurologists, or specialist trained in the syndrome

Vykat XR® Renewal Criteria:

- Submission of medical records (e.g. chart notes) documenting positive clinical response (i.e. decrease in sensation of hunger, less consumption of food, feeling satiated, decrease in food preoccupations, decrease in BMI, improvement in metabolic markers)

Pyruvate Kinase Activators

Pyrukynd®

PDL Status	NP
Quantity Limit	2 tabs/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Pyrukynd® Initial PA Criteria:

- Patient has diagnosis of hemolytic anemia with pyruvate kinase (PK) deficiency***; AND***
- Patient has at least 2 variant alleles in the PK liver and red blood cell gene of which at least 1 was a missense variant; **AND**
- Hemoglobin is <10 g/dL; **AND**
- One of the following:
  - Patient has symptomatic anemia
  - Patient is transfusion dependent; **AND**
- Prescribed by or in consultation with a hematologist
- **Pyrukynd® Renewal PA Criteria:**
- Documentation of positive clinical response to therapy as evidenced by one of the following:
  - Hemoglobin increase ≥ 1.5 g/dL from baseline
  - Reduction in the number of red blood cell units transfused from baseline

Rett Syndrome

Daybue® oral solution

PDL Status	NP
Quantity Limit	120 mL/day
PA Criteria	See below
PA Form	General PA Form

Daybue® oral solution Initial PA Criteria:

- Patient is > 2 years old; **AND**
- Diagnosis of Rett Syndrome; **AND**
- Prescribed by, or in consultation with, a neurologist, clinical geneticist, or developmental pediatrician

Daybue® oral solution Renewal PA Criteria:

- Documentation of positive clinical response to Daybue® (e.g. improvement or stabilization in purposeful hand skills, spoken language, repetitive hand movements, and gait abnormalities)

Daybue STIX®

PDL Status	NP
Quantity Limit	5,000 mg and 6,000 mg packet: 4/day; 8,000 mg packet: 2/day
PA Criteria	See below
PA Form	General PA Form

Daybue STIX® Initial PA Criteria:

See Daybue® oral solution prior authorization criteria; **AND**

- Submission of medical records (e.g., chart notes) documenting ONE of the following:
  - Allergy or intolerance to inactive ingredient in Daybue oral solution (e.g., dye, filler other excipients); **OR**
  - Clinically valid reason why the patient cannot use the solution (e.g., volume restriction, ketogenic diet)

Sickle Cell Disease

Endari®

PDL Status	NP
Quantity Limit	6 packs/day
PA Criteria	See below
PA Form	General PA Form

Endari® Initial PA Criteria:

- Diagnosis of sickle cell disease; **AND**
- Patient meets ONE of the following:
  - Endari will be used in combination with hydroxyurea; **OR**
  - Trial and failure, contraindications, or intolerance to hydroxyurea; **AND**
- Dosed according to weight-based dosing found in package insert:
- < 30 kg, up to 2 packets per day
  - 30-65 kg, up to 4 packets per day
  - > 65 kg, up to 6 packets per day

Endari® Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., decrease in number of days in crisis, number of days in hospital, occurrence of Acute Chest Syndrome)

L-glutamine pack

PDL Status	NP
Quantity Limit	6 packs/day
PA Criteria	See below
PA Form	General PA Form

L-glutamine pack PA Criteria:

See Endari® prior authorization criteria; **AND**

- Trial and failure, contraindication, or intolerance to Endari®

Siklos®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Siklos® PA Criteria:

- Patient has a diagnosis of sickle cell anemia with recurrent moderate to severe painful crisis; **AND**
- ONE of the following:
  - Documentation of need for dosing that will not allow the use of a preferred hydroxyurea agent
  - Patient unable to swallow hydroxyurea capsules

Xromi®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Xromi® PA Criteria:

- Diagnosis of sickle cell anemia with recurrent moderate to severe painful crises; **AND**
- One of the following:
  - Documentation of need for dosing that will not allow the use of a preferred hydroxyurea agent; **OR**
  - Patient is unable to swallow solid oral dosage forms of hydroxyurea

Somatostatins and Related Agents

Korlym® tablets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Korlym® tablets PA Criteria:

- Diagnosis of Cushing's Syndrome; **AND**
- Patient has type 2 diabetes mellitus or glucose intolerance; **AND**
- Patient has failed surgical treatment **OR** is not candidate for surgery; **AND**
- Will NOT be approved for use during pregnancy

Octreotide injection

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Octreotide injection PA Criteria:

- Diagnosis of acromegaly; **OR**
- Treatment is for severe diarrhea/flushing episodes associated with metastatic carcinoid tumors; **OR**
- Treatment is for profuse watery diarrhea associated with VIP-secreting tumors

Isturisa® tablets

PDL Status	NP
Quantity Limit	1 mg: 4/day; 5 mg: 2/day; 10 mg: 6/day
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Isturisa® tablets Initial PA Criteria:

- Diagnosis of endogenous hypercortisolemia associated with Cushing's syndrome; **AND**
- Patient has failed surgical treatment **OR** is not candidate for surgery; **AND**
- Trial and failure (trial duration ≥ 90 days) or intolerance to oral ketoconazole; **AND**
- Patient is 18 years of age or older; **AND**
- Prescribed by, or in consultation with, an endocrinologist

Isturisa® tablets Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., normalization or reduction of urinary free cortisol, improvement in signs or symptoms of the disease)

Mifepristone 300 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Mifepristone 300 mg tablets PA Criteria:

See Korlym® prior authorization criteria; **AND**

- Clinically valid reason why the preferred Korlym® cannot be used

Mycapssa® DR capsules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Mycapssa® DR capsules PA Criteria:

- Diagnosis of acromegaly; **AND**
- Patient has previously taken, responded to, and tolerated treatment with octreotide or lanreotide; **AND**
- Clinically valid reason why the patient is unable to be maintained on current octreotide or lanreotide therapy

Palsonify® tablets

PDL Status	NP
Quantity Limit	2 tabs/day
PA Criteria	See below
PA Form	General PA Form

Palsonify® tablets Initial PA Criteria:

- Patient is at least 18 years of age or older; **AND**
- Diagnosis of acromegaly; **AND**
- Lab documentation confirming ONE of the following:
  - Patient has confirmed higher than normal insulin-like growth factor (IGF-1) levels (above age and gender adjusted normal range) at the time of diagnosis; **OR**
  - Patient has elevated serum growth hormone (GH) > 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at the time of diagnosis: **OR**
  - Patient has an MRI of pituitary gland confirming acromegaly diagnosis; **AND**
- One of the following:
  - Patient has an inadequate response to surgery; **OR**
  - Patient is not a candidate for surgery; **AND**
- Trial and failure, contraindication or intolerance to octreotide; **AND**
- Prescribed by, or in consultation with, an endocrinologist

Palsonify® tablets Renewal criteria:

- Patient demonstrates positive clinical response to therapy (e.g., clinically significant reduction in IGF-1/GH levels, reduction in Total Acromegaly Symptoms Diary (ASD) score)

Recorlev® tablets

PDL Status	NP
Quantity Limit	8/day
PA Criteria	See below
PA Form	General PA Form

Recorlev® tablets Initial PA Criteria:

- Diagnosis of Cushing's Syndrome; **AND**
- Patient is being treated for endogenous hypercortisolemia (e.g., pituitary adenoma, ectopic tumor, adrenal
- adenoma); **AND**
- Surgery is not an option or has not been curative; **AND**
- Trial and failure (trial duration > 90 days) or intolerance to oral ketoconazole; **AND**
- Patient is 18 years of age or older; **AND**
- Prescribed by or in consultation with an endocrinologist; **AND**
- Patient has had baseline liver enzymes and an electrocardiogram prior to initiating therapy, and prescriber attests to monitor regularly thereafter; **AND**
- Patient does not have hypokalemia and hypomagnesemia, or has been corrected prior to therapy

Recorlev® tablets Renewal PA Criteria:

- Documentation of positive clinical response to therapy (e.g., normalization or reduction of urinary free cortisol, improvement in signs or symptoms of the disease)

Sandostatin® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sandostatin® injection PA Criteria:

See prior authorization criteria for octreotide; **AND**

- Clinically valid reason why preferred octreotide cannot be used

Signifor® injection

PDL Status	NP
Quantity Limit	2 injections/day
PA Criteria	See below
PA Form	General PA Form

Signifor® injection PA Criteria:

- Diagnosis of Cushing's Disease; **AND**
- Surgery is not an option or has not been curative; **AND**
- Prescribed by, or in consultation with, an endocrinologist

Somavert® injection

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Somavert® injection PA Criteria:

- Diagnosis of acromegaly; **AND**
- Tral and failure, intolerance, or contraindication to octreotide

Spinal Muscular Atrophy (SMA)

Evrysdi® solution, tablets

PDL Status	NP
Quantity Limit	Solution: 3 bottles/28 days Tabs: 1/day
PA Criteria	See below
PA Form	General PA Form

Evrysdi® solution, tablets Initial PA Criteria:

- Diagnosis of Spinal Muscular Atrophy (SMA); **AND**
- Submission of medical records (e.g., chart notes, laboratory values) confirming the mutation or deletion of genes in chromosome 5q confirming in one of the following:
  - Homozygous gene deletion or mutation of SMN1 gene; **OR**
  - Compound heterozygous mutation of SMN1 gene; **AND**
- Prescribed by, or in consultation with, a neurologist with expertise in the diagnosis and treatment of SMA; **AND**
- Patient will not receive concomitant survival motor neuron (SMN) modifying therapy (e.g., Spinraza); **AND**
- One of the following:
  - Patient has not previously received gene replacement therapy for the treatment of SMA (e.g., Zolgensma); **OR**
  - Both of the following:
    - Patient has previously received gene therapy for the treatment of SMA (e.g., Zolgensma); **AND**
    - Provider attests that there has been an inadequate response to gene therapy (e.g., sustained decrease in $\geq$ 1 motor test score over a period of 6-months); **AND**
- Advise female patients of reproductive potential to use effective contraception during treatment and for at least 1 month after the last dose

Evrysdi® solution, tablets Renewal PA Criteria

- Patient continues to meet initial criteria; **AND**
- Patient has clinically significant improvement in SMA associated signs and symptoms (progression, stabilization, or decreased decline in motor function)

Thymidine Kinase Deficiency

Kygevvi® packet

PDL Status	NP
Quantity Limit	27 capsules/28 days
PA Criteria	See below
PA Form	General PA Form
PA Approval Duration	Initial: 6 months; Renewal: 1 year

Kygevvi® packet Initial PA Criteria:

- Diagnosis of thymidine kinase 2 deficiency (TK2d) as evidenced by a mutation in the thymidine kinase 2 (TK2) gene; **AND**
- Patient has an age of symptom onset on or before 12 years; **AND**
- Prescriber attests that baseline liver transaminase (e.g., alanine aminotransferase [ALT] and aspartate aminotransferase [AST]) and total bilirubin levels were obtained prior to treatment initiation; **AND**
- Agent is prescribed by either a neurologist, metabolic disease/geneticist specialist, pulmonologist, or a physician specializing in the condition being treated

Kygevvi® packet Renewal PA Criteria

- Patient has disease stabilization or clinical response to therapy; **AND**
- Patient does not have unacceptable toxicity (e.g., elevated ALT, AST, total bilirubin, GI adverse reactions)

Transthyretin Amyloidosis Agents

Attruby®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Attruby® Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Prescribed by or in consultation with a cardiologist, neurologist, or specialist with knowledge of ATTRv; **AND**
- Diagnosis of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) with cardiomyopathy; **AND**
- Patient has New York Heart Association Class I, II or III heart failure; **AND**
- Patient does not meet have any of the following:
  - Impaired renal function (glomerular filtration rate < 15 mL/min/1.73 m2)
  - History of heart transplantation
  - New York Heart Association Class IV; **AND**
- Patient will not use in combination with another medication indicated for the management cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Vyndamax)

Attruby® Renewal PA Criteria

- Patient meets initial criteria; **AND**
- Patient has demonstrated a positive benefit from therapy (e.g., improved clinical symptoms of heart failure)

Vyndamax®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See below
PA Form	General PA Form

Vyndamax® Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Must be prescribed in consultation with a cardiologist neurologist, or specialist with knowledge of ATTRv; **AND**
- Diagnosis of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) with cardiomyopathy; **AND**
- Patient has New York Heart Association Class I, II or III heart failure; **AND**
- Patient does not have any of the following:
  - Impaired renal function (glomerular filtration rate < 25 mL/min/1.73 m2)
  - History of liver or heart transplantation
  - Implanted left ventricular assist device (LVAD) [pacemaker or cardiac defibrillator allowed]
  - Patient is pregnant or breastfeeding
  - New York Heart Association Class IV; **AND**
- Patient will not use in combination with another medication indicated for the management cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Attruby)

Vyndamax® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient has demonstrated a positive benefit from therapy (e.g., improved clinical symptoms of heart failure)

Wainua®

PDL Status	NP
Quantity Limit	1 injector/28 days
PA Criteria	See below
PA Form	General PA Form

Wainua® Initial PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of hereditary transthyretin-mediated amyloidosis (ATTRv amyloidosis) with polyneuropathy; **AND**
- Documentation that patient has a transthyretin (TTR) mutation (e.g., V30 m); **AND**
- Prescribed by, or in consultation with, a neurologist, cardiologist, or specialist with knowledge of ATTRv; **AND**
- Documentation of ONE of the following:
  - Patient has a baseline polyneuropathy disability (PND) score ≤ IIIb
  - Patient has a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2
  - Patient has a baseline neuropathy impairment score (NIS) between 10 and 130; **AND**
- Patient has not had a liver transplant; **AND**
- Will not be used in combination with another medication indicated for the management cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Attruby, Vyndaqel, Vyndamax)

Wainua® Renewal PA Criteria:

- Patient has demonstrated a benefit from therapy (e.g., improved neurologic impairment, motor function, slowing of disease progression, quality of life assessment); **AND**
- Will not be used in combination with another medication indicated for the management cardiomyopathy or neuropathy of transthyretin-mediated amyloidosis (e.g., Amvuttra, Onpattro, Attruby, Vyndaqel, Vyndamax)

Tyrosinemia Type 1

Orfadin® suspension, nitisinone capsules, Nityr® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Orfadin® suspension, nitisinone capsules, Nityr® tablets PA Criteria:

- Diagnosis of hereditary tyrosinemia type 1; **AND**
- Agent is prescribed by a physician specializing in the condition being treated; **AND**
- Patient has a clinically valid reason as to why the Orfadin® capsules cannot be utilized

Urea Cycle Disorders

Carbaglu® tablets, Pheburane® pellets

PDL Status	P
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carbaglu® tablets, Pheburane® pellets PA Criteria:

- Diagnosis of urea cycle disorders

Carglumic acid tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Carglumic acid tablets PA Criteria:

- Diagnosis of urea cycle disorders; **AND**
- Trial and failure, contraindication, or intolerance of Carbaglu®

Glycerol phenylbutyrate liquid

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Glycerol phenylbutyrate liquid PA Criteria:

- Diagnosis of urea cycle disorders; **AND**
- Trial and failure, contraindication, or intolerance of Pheburane®

Olpruva® packets, Ravicti® liquid

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Olpruva® packets, Ravicti® liquid PA Criteria:

- Diagnosis of urea cycle disorders; **AND**
- Trial and failure, contraindication, or intolerance of Pheburane®

Sodium phenylbutyrate tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Sodium phenylbutyrate tablets PA Criteria:

- Diagnosis of urea cycle disorders; **AND**
- Trial and failure, contraindication, or intolerance of Buphenyl®

Wilson Disease

Galzin®

PDL Status	NP
Quantity Limit	None
PA Criteria	See below
PA Form	General PA Form

Galzin® PA Criteria:

- Diagnosis of Wilson's disease; **AND**
- Intolerance of zinc sulfate

Trientine

PDL Status	NP
Quantity Limit	250 mg: 8/day; 500 mg: 4/day
PA Criteria	See below
PA Form	General PA Form

Trientine PA Criteria:
Standard NP Criteria

WHIM Syndrome

Zolremdi®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See below
PA Form	General PA Form

Zolremdi® PA Criteria:
Standard NP Criteria

XV. RENAL AND GENITOURINARY

Agents for BPH

Tadalafil 5 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tadalafil PA Criteria:

- Diagnosis of Benign Prostatic Hyperplasia (BPH); **AND**
- Patient is NOT concurrently receiving nitrates or guanylate cyclase stimulators; **AND**
- Trial and failure, contraindication, or intolerance to at least ONE agent from each of the following classes:
  - Alpha blockers for BPH
  - Androgen Hormone Inhibitors

Cialis® 5 mg tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Cialis® PA Criteria:

- Diagnosis of Benign Prostatic Hyperplasia (BPH); **AND**
- Patient is NOT concurrently receiving nitrates or guanylate cyclase stimulators; **AND**
- Trial and failure, contraindication, or intolerance to at least ONE agent from each of the following classes:
  - Alpha blockers for BPH
  - Androgen Hormone Inhibitors; **AND**
- Clinically valid reason why generic tadalafil tablets cannot be used

Dutasteride/tamsulosin capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Dutasteride/Tamsulosin capsules PA Criteria:

- Patient has a diagnosis of benign prostatic hyperplasia (BPH) with an enlarged prostate; **AND**
- Trial and failure, contraindication, or intolerance to finasteride; **AND**
- Patient is unable to use the individual components

Entadfi® capsules

PDL Status	NP
Quantity Limit	1/day; 182/year
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Entadfi® PA Criteria:

- Diagnosis of Benign Prostatic Hyperplasia (BPH) with an enlarged prostate; **AND**
- Total length of therapy has not exceeded 26 weeks; **AND**
- Trial and failure, contraindication, or intolerance to combination therapy with alpha blocker and androgen
- hormone inhibitor; **AND**
- Clinically valid reason why the individual components of Entadfi® cannot be used (finasteride and tadalafil); **AND**
- Patient is NOT concurrently receiving nitrates or guanylate cyclase stimulators

Alpha Blockers for BPH

Alfuzosin tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Tamsulosin capsules

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Cardura XL® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Cardura XL® tablets PA Criteria:

Standard NP Criteria

Tezruly® solution

PDL Status	NP
Quantity Limit	20 mL/day
PA Criteria	See Below
PA Form	General PA Form

Tezruly® solution PA Criteria:

- Patient is unable to swallow solid dosage forms

Androgen Hormone Inhibitors

Covered for males only.

Dutasteride capsules, finasteride tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Proscar® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Proscar® PA Criteria:
Standard NP Criteria

Kidney Stone Agents

Thiola EC® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Thiola EC® tablets PA Criteria:

- Patient has tried/failed an adequate trial of or is intolerant to two preferred agents; **AND**
- Clinically valid reason why preferred Thiola cannot be used

Tiopronin DR tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tiopronin DR tablets PA Criteria:

- See Thiola EC® PA criteria; **AND**
- Clinically valid reason why brand Thiola EC® cannot be used

Phosphate Binders and Related Agents

Sevelamer carbonate tablets

PDL Status	P
Quantity Limit	9/day
PA Criteria	None

Sevelamer carbonate packs

PDL Status	P
Quantity Limit	0.8 g packets: 6/day; 2.4 g packets: 5/day
PA Criteria	None

Sevelamer carbonate packs PA Criteria:

- Patient is unable to swallow solid dosage forms

Velphoro® chewable tablets

PDL Status	P
Quantity Limit	6/day
PA Criteria	See Below
PA Form	General PA Form

Velphoro® chewable tablets PA Criteria:

- Patient is ≥ 9 years old; **AND**
- Patient has chronic kidney disease (CKD) and is on dialysis; **AND**
- Trial and failure, contraindication, or intolerance of TWO preferred phosphorus binders; **OR**
- Patient is unable to swallow solid dosage forms AND trial and failure, contraindication, or intolerance to sevelamer powder

Auryxia® tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Auryxia® tablets PA Criteria:

- Diagnosis of hyperphosphatemia in chronic kidney disease on dialysis; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents; **OR**
- Diagnosis of iron deficiency anemia in chronic kidney disease NOT on dialysis; **AND**
  - Trial and failure, contraindication, or intolerance to TWO oral iron products (e.g., ferrous sulfate, ferrous gluconate)

Ferric citrate tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Ferric citrate tablets PA Criteria:

- Diagnosis of hyperphosphatemia in chronic kidney disease on dialysis; **AND**
  - Trial and failure, contraindication, or intolerance to TWO preferred agents; **OR**
- Diagnosis of iron deficiency anemia in chronic kidney disease NOT on dialysis; **AND**
- Trial and failure, contraindication, or intolerance to TWO oral iron products (e.g., ferrous sulfate, ferrous gluconate); **AND**
- Clinically valid reason why brand Auryxia® cannot be used.

Fosrenol® packs

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Fosrenol® PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred phosphorus depletors; **AND**
- Contraindication to sevelamer powder for suspension; **AND**
- Patient is unable to swallow solid dosage forms

Renvela® packs

PDL Status	NP
Quantity Limit	0.8 g packets: 6/day 2.4 g packets: 5/day
PA Criteria	See Below
PA Form	General PA Form

Renvela® packs PA Criteria:

- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why preferred generic agent cannot be used

Renvela® tablets

PDL Status	NP
Quantity Limit	9/day
PA Criteria	See Below
PA Form	General PA Form

Renvela® tablets PA Criteria:

Standard NP Criteria

Xphozah®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Xphozah® PA Criteria:

- Patient is 18 years of age or older; **AND**
- Diagnosis of chronic kidney disease (CKD); **AND**
- Patient is currently on dialysis; **AND**
- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Agent will be used as adjunctive therapy to reduce serum phosphorus; **AND**
- Patient does not have known or suspected mechanical gastrointestinal obstruction

Urinary Analgesics

Belladonna/Opium® suppositories

PDL Status	NP
Quantity Limit	12/30 days
PA Criteria	See Below
PA Form	General PA Form

Belladona/Opium® suppositories PA Criteria:

- Patient is 12 years of age or older; **AND**
- Medication is being used for the management of ureteral spasm and pain; **AND**
- Dose must not be greater than 4 suppositories a day; **AND**
- Trial and failure of the preferred product phenazopyridine

Urinary Tract Antispasmodics

Fesoterodine ER tablets, Myrbetriq® tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Oxybutynin ER tablets

PDL Status	P
Quantity Limit	5 mg: 1/day 10, 15 mg: 2/day
PA Criteria	None

Oxytrol® patches

PDL Status	P
Quantity Limit	8 patches/28 days
PA Criteria	None

Solifenacin tablets, tolterodine ER capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Tolterodine tablets

PDL Status	P
Quantity Limit	2/day
PA Criteria	None

Darifenacin tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Darifenacin PA Criteria:

Standard NP Criteria

Flavoxate tablets

PDL Status	NP
Quantity Limit	2 fills/60 days
PA Criteria	See Below
PA Form	General PA Form

Flavoxate tablets PA Criteria:
Standard NP Criteria

Gelnique® gel

PDL Status	NP
Quantity Limit	1 pack (1 gr)/day
PA Criteria	See Below
PA Form	General PA Form

Gelnique® gel PA Criteria:
Standard NP Criteria

Gemtesa® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Gemtesa® tablets PA Criteria:

- Patient is 18 years of age or older: **AND**
- Diagnosis of overactive bladder (OAB); **AND**
- Trial and failure of one preferred anticholinergic agent (e.g., fesoterodine, oxybutynin, solifenacin, tolterodine); **AND**
- Trial and failure, or contraindication, or intolerance to Myrbetriq

Mirabegron suspension

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Mirabegron suspension PA Criteria:

- Diagnosis of neurogenic detrusor overactivity (NDO); **AND**
- Trial and failure, contraindication, or intolerance to oxybutynin solution; **AND**
- Patient is unable to swallow solid dosage forms; **AND**
- Clinically valid reason why brand Myrbetriq suspension cannot be used

Mirabegron tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Mirabegron tablets PA Criteria:

- Clinically valid reason why preferred Myrbetriq® cannot be used

Myrbetriq® suspension

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Myrbetriq® suspension PA Criteria:

- Diagnosis of neurogenic detrusor overactivity (NDO); **AND**
- Trial and failure, contraindication, or intolerance to oxybutynin solution; **AND**
- Patient is unable to swallow solid dosage forms;

Oxybutynin 2.5 mg IR tablets

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Oxybutynin 2.5 mg IR tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product; **AND**
- Strength of requested product is not attainable with preferred PDL agents (e.g., by taking two capsules or half tablet (as approved per FDA and/or manufacturer)

Toviaz® ER tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Toviaz® ER tablets PA Criteria:
Standard NP Criteria

Trospium tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Trospium tablets PA Criteria:
Standard NP Criteria

Trospium XR capsules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Trospium XR capsules PA Criteria:
Standard NP Criteria

VESIcare® suspension

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

VESIcare® suspension PA Criteria:
Standard NP Criteria

VESIcare® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

VESIcare® tablets PA Criteria:
Standard NP Criteria

XVI. RESPIRATORY

Anaphylaxis Therapy Agents

Epinephrine auto injector

PDL Status	P
Quantity Limit	2/Rx
PA Criteria	None

Auvi-Q®, EpiPen®, EpiPen Jr®

PDL Status	NP
Quantity Limit	2/Rx
PA Criteria	See Below
PA Form	General PA Form

Auvi-Q®, EpiPen®, EpiPen Jr® PA Criteria:
Standard NP Criteria

Neffy®

PDL Status	NP
Quantity Limit	2/Rx
PA Criteria	See Below
PA Form	General PA Form

Neffy® PA Criteria:

- Clinically valid reason why the preferred epinephrine auto-injector cannot be used

Anticholinergics, Nasal

Ipratropium 0.3% solution

PDL Status	P
Quantity Limit	2 boxes/30 days
PA Criteria	None

Ipratropium 0.6% solution

PDL Status	P
Quantity Limit	3 boxes/30 days
PA Criteria	None

Antihistamines, Nasal

Azelastine nasal spray 0.1%, 0.15% (Rx), Dymista® nasal spray

PDL Status	P
Quantity Limit	2 bottles/30 days
PA Criteria	None

Olopatadine nasal spray

PDL Status	P
Quantity Limit	1 bottle/30 days
PA Criteria	None

Azelastine/fluticasone nasal spray

PDL Status	NP
Quantity Limit	1 bottle/30 days
PA Criteria	See Below
PA Form	General PA Form

Azelastine/fluticasone PA Criteria:
Standard NP Criteria

Ryaltris® nasal spray

PDL Status	NP
Quantity Limit	1 bottle/30 days
PA Criteria	See Below
PA Form	General PA Form

Ryaltris® nasal spray PA Criteria:

- Diagnosis of Seasonal Allergic Rhinitis; **AND**
- Patient is 6 years of age or older; **AND**
- Trial and failure, contraindication, or intolerance to preferred azelastine/fluticasone combination product; **AND**
- Clinically valid reason as to why the patient is unable to take components of Ryaltris individually (Note: Patient convenience is not an approvable reason)

Antihistamines: First Generation

These agents are covered for recipients < 21 years old only unless otherwise noted.
Please refer to OTC Drug Formulary for coverage of OTC products.

Carbinoxamine maleate 4 mg/5 mL suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Carbinoxamine maleate 4 mg/5 mL suspension PA Criteria:
Standard NP Criteria

Carbinoxamine maleate 6 mg tablets

PDL Status	NP
Quantity Limit	6 tabs/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Carbinoxamine maleate 6 mg tablets PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Carbzah® solution; Clemza® tablets; Corphena® solution

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Carbzah® solution; Clemza® tablets; Corphena® solution PA Criteria:

Standard NP Criteria

Diphenhydramine elixir 12.5 mg/5 mL

PDL Status	NP
Quantity Limit	80 mL/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Diphenhydramine elixir 12.5 mg/5 mL PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Karbinal ER® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Karbinal ER® suspension PA Criteria:

Standard NP Criteria

Ryclora® solution; Ryvent® solution

PDL Status	NP
Quantity Limit	Ryclora: 30 mL/day; Ryvent: 5 tabs/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Ryclora® solution, Ryvent® solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Antihistamines: Non-Sedating, Oral

These agents are covered for recipients < 21 years old only unless otherwise noted.
Please refer to OTC Drug Formulary for coverage of OTC products.

Levocetirizine tablets (Rx only)

PDL Status	P
Quantity Limit	1/day
PA Criteria	None
Notes	Covered for members less than 21 years old only

Clarinex® tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Clarinex® tablets PA Criteria:
Standard NP Criteria

Clarinex D® tablets

PDL Status	NP
Quantity Limit	12 hour: 2/day 24 hour: 1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Clarinex D® tablets PA Criteria:
Standard NP Criteria

Desloratadine solution

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Desloratadine solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Desloratadine tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Desloratadine tablets PA Criteria:

Standard NP Criteria

Desloratadine ODT

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Desloratadine ODT PA Criteria:

- Patient is unable to swallow solid dosage forms

Levocetirizine solution (Rx only)

PDL Status	NP
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form
Notes	Covered for members less than 21 years old only

Levocetirizine solution PA Criteria:

Standard NP Criteria

Antitussives, Non-Narcotic

Benzonatate capsules (100 and 200 mg)

PDL Status	P
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Benzonatate capsules PA Criteria:

- Patient is ≥ 10 years of age; **OR**
- Patient is < 10 years of age and prescriber is aware that, if chewed, benzonatate may cause numbness of the mouth, tongue, throat, and esophagus, increasing the risk of choking

Benzonatate 150 mg capsules

PDL Status	P
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Benzonatate capsules PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Cystic Fibrosis Agents, Inhaled/Injectable

Bethkis® nebulizer solution

PDL Status	P
Quantity Limit	224 mL/56 days
PA Criteria	See Below
PA Form	General PA Form

Bethkis® nebulizer solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection

Pulmozyme® inhalation solution

PDL Status	P
Quantity Limit	5 mL/day
PA Criteria	See Below
PA Form	General PA Form

Pulmozyme® inhalation solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection

Tobramycin solution (300 mg/5 mL)

PDL Status	P
Quantity Limit	10 mL/day
PA Criteria	See Below
PA Form	General PA Form

Tobramycin solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection

Tobramycin vial (excluding 1.2-gram vials)

PDL Status	P
Quantity Limit	None
PA Criteria	See Below
PA Form	General PA Form

Tobramycin vial PA Criteria:

- Claims exceeding $200 will only be approved for diagnosis of Cystic Fibrosis or *Pseudomonas* infection

Bronchitol oral inhalation capsule

PDL Status	NP
Quantity Limit	20/day
PA Criteria	See Below
PA Form	General PA Form

Bronchitol oral inhalation capsule PA Criteria:

- Diagnosis of Cystic Fibrosis
- Patient must not have an episode of hemoptysis (>60 mL) in the last 3 months; **AND**
- Must be 18 years of age or older; **AND**
- Patient must have baseline FEV1 >40% to <90%; **AND**
- Patient has passed the Bronchitol Tolerance Test; **AND**
- Must be used concomitantly with a short-acting bronchodilator; **AND**
- Prescriber attests that the patient has been instructed to administer the agent 5-15 minutes after a short-acting bronchodilator

Cayston® inhalation solution

PDL Status	NP
Quantity Limit	3/day
PA Criteria	See Below
PA Form	General PA Form

Cayston® inhalation solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection; **AND**
- Trial and failure, contraindication, intolerance, or resistance to preferred inhaled tobramycin product

Kitabis Pak® nebulizer solution

PDL Status	NP
Quantity Limit	280 mL/56 days
PA Criteria	See Below
PA Form	General PA Form

Kitabis Pak® nebulizer solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection; **AND**
- Clinically valid reason why the generic tobramycin vials cannot be used

Tobramycin solution (300 mg/4 mL)

PDL Status	NP
Quantity Limit	8 mL/day
PA Criteria	See Below
PA Form	General PA Form

Tobramycin solution PA Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection; **AND**
- Clinically valid reason why preferred Bethkis® cannot be used

TOBI® Podhaler and inhalation solution

PDL Status	NP
Quantity Limit	Podhaler: 224 /28 days Solution: 280 mL/56 days
PA Criteria	See Below
PA Form	General PA Form

TOBI® Podhaler and inhalation solution Criteria:

- Diagnosis of Cystic Fibrosis or *Pseudomonas* infection; **AND**
- Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice versus the preferred agents

Cystic Fibrosis Agents, Oral

Alyftrek® tablets

PDL Status	NP
Quantity Limit	4-20-50 mg: 3/day; 10-50-125 mg: 2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Alyftrek® tablets Initial PA Criteria:

- Diagnosis of cystic fibrosis (CF); **AND**
- Must be prescribed by, or in consultation with, a provider at a CF Center of Excellence or pulmonologist; **AND**
- Age ≥ 6 years old; **AND**
- Lab documentation confirming patient has at least one variant in the CFTR gene that is responsive based on clinical data, in vitro data, or results in production of CFTR protein

Alyftrek® tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decreased pulmonary exacerbations, improvement, or stabilization of lung function)

Kalydeco® granules, tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months
Notes	Will NOT be approved for homozygous F508del mutation in the CFTR gene

Kalydeco® granules, tablets Initial PA Criteria:

- Diagnosis of cystic fibrosis (CF); **AND**
- Must be prescribed by, or in consultation with, a provider at a CF Center of Excellence or pulmonologist**; AND**
- Lab documentation confirming patient has one mutation in the CFTR gene that is responsive to Kalydeco®

Kalydeco® granules, tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decreased pulmonary exacerbations, improvement, or stabilization of lung function)

Orkambi® tablets and granules

PDL Status	NP
Quantity Limit	Tablets: 4/day; Granules: 2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Orkambi® granules, tablets Initial PA Criteria:

- Diagnosis of cystic fibrosis (CF); **AND**
- Must be prescribed by, or in consultation with, a provider at a CF Center of Excellence or pulmonologist**; AND**
- Age ≥ 1 years old; **AND**
- Lab documentation confirming patient has homozygous F508del mutation in the CFTR gene

Orkambi® granules, tablets Renewal PA Criteria:

- Disease response to therapy and tolerability compared to baseline (e.g., decreased pulmonary exacerbations, improvement, or stabilization of lung function)

Symdeko® tablets

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Symdeko® tablets Initial PA Criteria:

- Diagnosis of cystic fibrosis (CF); **AND**
- Must be prescribed by, or in consultation with, a provider at a CF Center of Excellence or pulmonologist**; AND**
- Age ≥ 6 years old; **AND**
- Lab documentation confirming ONE of the following:
  - Patient is homozygous for the F508del mutation in the CFTR gene
  - Patient has ≥ 1 mutation in the CFTR gene that is responsive based on in vitro data

Symdeko® tablets Renewal PA Criteria:

- Patient had not received a lung transplant; **AND**
  - Disease response to therapy and tolerability compared to baseline (e.g., decreased pulmonary exacerbations, improvement or stabilization of lung function); **OR**
- Patient has received a lung transplant; **AND**
- Prescriber attests that the patient continues to experience non-pulmonary CF related symptoms (e.g., sinus, gastrointestinal, diabetes, pancreatic)

Trikafta® granules, tablets

PDL Status	NP
Quantity Limit	Granules: 2/day; Tablets: 3/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Trikafta® granules, tablets Initial PA Criteria:

- Diagnosis of cystic fibrosis (CF); **AND**
- Must be prescribed by, or in consultation with, a provider at a CF Center of Excellence or pulmonologist**; AND**
- Age ≥ 2 years old; **AND**
- Lab documentation confirming patient has at least one variant in the CFTR gene that is responsive based on clinical data, in vitro data, or results in production of CFTR protein; **AND**
- If the request is for Trikafta granules, patient is unable to swallow solid dosage forms

Trikafta® granules, tablets Renewal PA Criteria:

- Patient had not received a lung transplant; **AND**
  - Disease response to therapy and tolerability compared to baseline (e.g., decreased pulmonary exacerbations, improvement or stabilization of lung function); **OR**
- Patient has received a lung transplant; **AND**
- Prescriber attests that the patient continues to experience non-pulmonary CF related symptoms (e.g., sinus, gastrointestinal, diabetes, pancreatic)

DPP-1 Inhibitors

Brinsupri® tablets

PDL Status	NP
Quantity Limit	1 tablet/day
PA Criteria	See below
PA Form	General PA Form

Brinsupri® tablets Initial PA Criteria:

- Diagnosis of non-cystic fibrosis bronchiectasis (NCFB); **AND**
- Diagnosis has been confirmed by a documented computed tomography (CT) scan; **AND**
- Prescribed by, or in consultation with, a pulmonologist or infectious disease specialist; **AND**
- One of the following:
  - Patient is 12 to 17 years of age with at least one documented pulmonary exacerbation in the past 12 months requiring systemic antibiotic therapy; **OR**
  - Patient is 18 years of age or older with at least two documented pulmonary exacerbations in the past 12 months requiring systemic antibiotic therapy;

Brinsupri® tablets Renewal PA Criteria:

- Patient demonstrates positive clinical response to therapy (e.g., stabilization or reduction in number of pulmonary exacerbations, reduced cough or sputum)

Inhaled: Anticholinergics and Anticholinergic Combinations

Anoro Ellipta®

PDL Status	P
Quantity Limit	2 blisters/day
PA Criteria	None

Albuterol/ipratropium nebulizer solution

PDL Status	P
Quantity Limit	18 mL /day
PA Criteria	None

Atrovent HFA®

PDL Status	P
Quantity Limit	2 inhalers/month
PA Criteria	None

Breztri Aerosphere®

PDL Status	P
Quantity Limit	1 inhaler/ month
PA Criteria	See Below
PA Form	General PA Form

Breztri Aerosphere® Initial PA Criteria:

- Patient is the labeled age minimum; **AND**
- One of the following:
  - Diagnosis of chronic obstructive pulmonary disease (COPD); **AND**
    - Trial and failure (as defined by continued symptoms, including exacerbations) of adequate treatment with a LABA/LAMA; **OR**
    - Eosinophil count ≥ 300 cells/ µL as confirmed by submission of medical records; **OR**
  - Diagnosis of asthma; **AND**
    - Trial and failure (as defined by continued symptoms, including exacerbations) of adequate treatment with TWO dual combination ICS/LABA therapies; **AND**
- Must be used as maintenance therapy only

Breztri Aerosphere® Renewal PA Criteria:

- Documentation of continued efficacy via prescriber's medical opinion on patient evaluation; **AND**
- Patient has not experienced any intolerable adverse effects (e.g., hypersensitivity, bronchospasm, worsening of intraocular pressure, increased severe infections)

Ipratropium solution

PDL Status	P
Quantity Limit	10 mL/day
PA Criteria	None

Spiriva Handihaler®

PDL Status	P
Quantity Limit	1 capsule/day
PA Criteria	None

Trelegy Ellipta®

PDL Status	P
Quantity Limit	2 blisters/ day
PA Criteria	See Below
PA Form	General PA Form

Trelegy Ellipta® Initial PA Criteria:

- Diagnosis of chronic obstructive pulmonary disease (COPD); **AND**
  - Trial and failure (as defined by continued symptoms, including exacerbations) of adequate treatment with a LABA/LAMA; **OR**
  - Eosinophil count ≥ 300 cells/ µL as confirmed by submission of medical records; **AND**
- Must be used as maintenance therapy only; **OR**
- Diagnosis of asthma in patients 12 years of age or older; **AND**
  - Trial and failure (as defined by continued symptoms, including exacerbations) of adequate treatment with TWO dual combination ICS/LABA therapies; **AND**
- Must be used as maintenance therapy only; **AND**
- Patient does not have known hypersensitivity to milk proteins

Trelegy Elllipta® Renewal PA Criteria:

- Documentation of continued efficacy via prescriber's medical opinion on patient evaluation; **AND**
- Patient has not experienced any intolerable adverse effects (e.g., hypersensitivity, bronchospasm, worsening of intraocular pressure, increased severe infections)

Bevespi Aerosphere®

PDL Status	NP
Quantity Limit	1 inhaler/ month
PA Criteria	See Below
PA Form	General PA Form

Bevespi Aerosphere® PA Criteria:

Standard NP Criteria

Combivent Respimat®

PDL Status	NP
Quantity Limit	2 inhalers/ month
PA Criteria	See Below
PA Form	General PA Form

Combivent Respimat® PA Criteria:

Standard NP Criteria

Duaklir Pressair®

PDL Status	NP
Quantity Limit	1 inhaler/ month
PA Criteria	See Below
PA Form	General PA Form

Duaklir Pressair® PA Criteria:

Standard NP Criteria

Incruse Ellipta®

PDL Status	NP
Quantity Limit	1 blister/day
PA Criteria	See Below
PA Form	General PA Form

Incruse Ellipta® PA Criteria:

- Diagnosis of Chronic Obstructive Pulmonary Disorder (COPD); **AND**
- Must be used as maintenance therapy only; **AND**
- Trial and failure, contraindication, or intolerance to TWO preferred inhaled anticholinergic/anticholinergic combination agents; **AND**
- Patient must not have severe hypersensitivity to milk proteins

Ipratropium HFA

PDL Status	NP
Quantity Limit	2 inhaler/ month
PA Criteria	See Below
PA Form	General PA Form

Ipratropium HFA PA Criteria:

- Clinically valid reason why brand Atrovent HFA® cannot be used

Spiriva Respimat®

PDL Status	NP
Quantity Limit	1 inhaler/month
PA Criteria	See Below
PA Form	General PA Form

Spiriva Respimat® PA Criteria:

- One of the following:
  - Diagnosis of Asthma; **AND**
    - Patient age ≥ 6 years; **AND**
    - Optimal doses of inhaled steroids and long-acting beta-agonists are being used, and breakthrough symptoms require frequent use of inhaled short-acting bronchodilators; **OR**
  - Diagnosis of Chronic Obstructive Pulmonary Disorder (COPD); **AND**
    - Must be used as maintenance therapy only; **AND**
    - Trial and failure, contraindication, or intolerance to Spiriva Handihaler®

Stiolto Respimat®

PDL Status	NP
Quantity Limit	1 inhaler/month
PA Criteria	See Below
PA Form	General PA Form

Stiolto Respimat® PA Criteria:

Standard NP Criteria

Tiotropium inhalation capsules

PDL Status	NP
Quantity Limit	1 capsule/day
PA Criteria	See Below
PA Form	General PA Form

Tiotropium inhalation PA Criteria:

- Clinically valid reason why the patient cannot use the preferred brand Spiriva Handihaler®

Tudorza®

PDL Status	NP
Quantity Limit	1 inhaler/month
PA Criteria	See Below
PA Form	General PA Form

Tudorza® PA Criteria:
See Incruse Ellipta® prior authorization criteria

Umeclidinium Ellipta

PDL Status	NP
Quantity Limit	1 blister/day
PA Criteria	See Below
PA Form	General PA Form

Umeclidinium Ellipta PA Criteria:
See Incruse Ellipta® prior authorization criteria

- Clinically valid reason why brand Incruse Ellipta® cannot be used

Umeclidinium/vilanterol Ellipta

PDL Status	NP
Quantity Limit	2 blisters/day
PA Criteria	See Below
PA Form	General PA Form

Umeclidinium/vilanterol Ellipta PA Criteria:

- Clinically valid reason why Anora Ellipta® cannot be used

Yupelri®

PDL Status	NP
Quantity Limit	3 mL/day
PA Criteria	See Below
PA Form	General PA Form

Yupelri® Initial PA Criteria:

- Diagnosis Patient must be ≥ 18 years of age; **AND**
- Diagnosis of Chronic Obstructive Pulmonary Disorder (COPD); **AND**
- Trial and failure, contraindication, or intolerance to TWO preferred inhaled anticholinergic/anticholinergic combination agents; **AND**
- Must be used as maintenance therapy only**; AND**
- Patient is unable to master proper inhaler technique, as attested by prescriber; **AND**
- Patient is not prescribed other inhaled long-acting anticholinergic agents.

Yupelri® Renewal PA Criteria:

- Patient continues to meet initial criteria; **AND**
- Patient symptoms are clinically improving, as documented by provider; **AND**
- Patient demonstrates continued compliance, based on fill history (not using PRN); **AND**
- Prescriber documents that nebulized therapy continues to be required

Inhaled: Beta Agonist-Corticosteroid Combination Products

For agents requiring prior authorization in this class, please submit faxed PA requests using the **Beta Agonist-Corticosteroid Combination PA Form**.

Advair HFA®

PDL Status	P
Quantity Limit	1 inhaler/month
PA Criteria	None

Advair Diskus®

PDL Status	P
Quantity Limit	2 blisters/day
PA Criteria	None

Dulera®

PDL Status	P
Quantity Limit	2 inhalers/month
PA Criteria	None

Symbicort®

PDL Status	P
Quantity Limit	2 inhalers/month
PA Criteria	None

Airsupra®

PDL Status	NP
Quantity Limit	2 inhalers/month
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Airsupra® PA Criteria:

- Agent will be used for the treatment of asthma in patients 18 years of age and older; **AND**
- Trial and failure, contraindication, or intolerance to preferred agents Symbicort® and Dulera®

Breo Ellipta®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Breo Ellipta® PA Criteria:

- One of the following:
  - Treatment of asthma in patients 5 years of age or older; **OR**
  - Treatment of COPD where optimal doses of a long-acting beta agonist and/or long-acting muscarinic antagonists are being used and symptoms are still uncontrolled (100/25 mcg strength only); **AND**
- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Patient must not have severe hypersensitivity to milk proteins

Breyna®

PDL Status	NP
Quantity Limit	2 inhalers/month
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Breyna® PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Clinically valid reason why the patient cannot use the preferred brand Symbicort®

Budesonide/formoterol

PDL Status	NP
Quantity Limit	2 inhalers/month
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Budesonide/formoterol PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Clinically valid reason why the patient cannot use the preferred brand Symbicort®

Fluticasone/salmeterol diskus

PDL Status	NP
Quantity Limit	55, 113, 232, -14 mcg: 1 inhaler/month 100, 250, 500, -50 mcg: 2 blisters/month
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Fluticasone/salmeterol diskus PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Clinically valid reason why the patient cannot use the preferred brand Advair HFA® or Advair Diskus®

Fluticasone/salmeterol HFA

PDL Status	NP
Quantity Limit	1 inhaler/ month
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Fluticasone/salmeterol HFA PA Criteria:

Standard NP Criteria

Fluticasone/vilanterol

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Fluticasone/vilanterol PA Criteria:

- See Breo Ellipta® prior authorization criteria; **AND**
- Clinically valid reason why the patient cannot use the preferred brand Breo Ellipta®

Wixela®

PDL Status	NP
Quantity Limit	2 blisters/day
PA Criteria	See Below
PA Form	Beta Agonist-Corticosteroid Combination PA Form

Wixela® PA Criteria:

- Trial and failure, contraindication, or intolerance of TWO preferred agents; **AND**
- Clinically valid reason why the patient cannot use the preferred brand Advair HFA® or Advair Diskus®

Inhaled: Beta Agonists, Long Acting

Serevent Diskus®

PDL Status	P
Quantity Limit	2 blisters/day
PA Criteria	None

Striverdi Respimat®

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Striverdi Respimat® PA Criteria:

- Diagnosis of COPD; **AND**
- Trial and failure, contraindication, or intolerance of the preferred agent (Serevent Diskus®)

Inhaled: Beta Agonists, Short Acting

Albuterol HFA

PDL Status	P
Quantity Limit	2 inhalers/month
PA Criteria	None

Ventolin® HFA

PDL Status	P
Quantity Limit	2 inhalers/month
PA Criteria	None

Xopenex® HFA

PDL Status	P
Quantity Limit	2 canisters/month
PA Criteria	See Below
PA Form	General PA Form

Xopenex® HFA PA Criteria:

- Patients has experienced intolerable side effects to albuterol (e.g., tachycardia, etc.)

Levalbuterol HFA

PDL Status	NP
Quantity Limit	2 canisters/month
PA Criteria	See Below
PA Form	General PA Form

Levalbuterol HFA PA Criteria:

- Patients has experienced intolerable side effects to albuterol (e.g., tachycardia, etc.); **AND**
- Clinically valid rationale for why patient cannot use brand Xopenex HFA®

ProAir RespiClick®

PDL Status	NP
Quantity Limit	2 inhalers/month
PA Criteria	See Below
PA Form	General PA Form

ProAir RespiClick® PA Criteria:
Standard NP Criteria

Inhaled: Nebulizers, Beta Agonists

Albuterol nebulizer solution

PDL Status	P
Quantity Limit	125 nebs/month 3 bottles/month
PA Criteria	None

Arformoterol

PDL Status	P
Quantity Limit	60 nebs/month
PA Criteria	None

Brovana®

PDL Status	NP
Quantity Limit	60 nebs/month (120 mL/month)
PA Criteria	See Below
PA Form	General PA Form

Brovana® PA Criteria:

- Diagnosis of COPD; **AND**
- Difficulty using a dry powder inhaler (DPI); **AND**
- Trial and failure, contraindication, or intolerance of the preferred agent (arformoterol nebulizer)

Formoterol

PDL Status	NP
Quantity Limit	60 nebs/month
PA Criteria	See Below
PA Form	General PA Form

Formoterol PA Criteria:
See Brovana® prior authorization criteria

Levalbuterol

PDL Status	NP
Quantity Limit	96 nebs/month
PA Criteria	See Below
PA Form	General PA Form

Levalbuterol PA Criteria:

- Patients has experienced intolerable side effects to albuterol (e.g., tachycardia)

Perforomist

PDL Status	NP
Quantity Limit	60 nebs/month
PA Criteria	See Below
PA Form	General PA Form

Perforomist PA Criteria:

- See Brovana® prior authorization criteria

Inhaled: Nebulizers, Mast Cell Stabilizers

Cromolyn solution

PDL Status	P
Quantity Limit	120 vials/month
PA Criteria	See Below
PA Form	General PA Form

Cromolyn solution PA Criteria:

- Diagnosis of asthma

Inhaled: Steroids

Alvesco®

PDL Status	P
Quantity Limit	2/30 days
PA Criteria	See Below
PA Form	General PA Form

Alvesco® PA Criteria:

- Diagnosis of asthma; **AND**
- Patient is 12 years of age or older

Arnuity Ellipta®

PDL Status	P
Quantity Limit	1 blister/day
PA Criteria	None

Asmanex HFA®

PDL Status	P
Quantity Limit	1/30 days
PA Criteria	None

Asmanex Twisthaler®

PDL Status	P
Quantity Limit	1/30 days
PA Criteria	None

Budesonide suspension

PDL Status	P
Quantity Limit	0.25, 0.5 mg: 2 vials/day 1 mg: 1 vial/day
PA Criteria	See Below
PA Form	General PA Form
Notes	PA not required for patients ≤ 8 years of age

Budesonide suspension PA Criteria:

- ONE of the following:
  - Diagnosis of asthma; **AND**
    - Patient is ≤ 8 years old; **OR**
  - Diagnosis of Eosinophilic esophagitis (EoE); **AND**
    - Prescribed by, or in consultation with, a gastroenterologist, allergist, or immunologist; **AND**
    - Prescriber attests to both of the following:
      - Esophageal biopsy consists of ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf) following treatment course of a proton pump inhibitor; **AND**
      - Patient is experiencing symptoms of esophageal dysfunction (e.g., feeding difficulties, vomiting, dysphagia)

Fluticasone Diskus

PDL Status	P
Quantity Limit	50 mcg: 2/day 100 mcg: 4/day 250 mcg: 8/day
PA Criteria	None

Fluticasone HFA

PDL Status	P
Quantity Limit	2/30 days
PA Criteria	None

Pulmicort Respules®

PDL Status	P
Quantity Limit	0.25, 0.5 mg: 2 vials/day 1 mg: 1 vial/day
PA Criteria	See Below
PA Form	General PA Form

Pulmicort Respules® PA Criteria:

- Diagnosis of asthma; **AND**
- Patient is ≤ 8 years old

QVAR RediHaler®

PDL Status	P
Quantity Limit	2/30 days
PA Criteria	None

Fluticasone Furoate Ellipta

PDL Status	NP
PA Criteria	See Below
PA Form	General PA Form

Fluticasone furoate Ellipta PA Criteria:

- Clinically valid reason why Arnuity Ellipta cannot be used

Beclomethasone HFA

PDL Status	NP
Quantity Limit	2/30 days
PA Criteria	See Below
PA Form	General PA Form

Beclomethasone HFA PA Criteria:

Standard NP Criteria

Intranasal: Steroids

Please refer to OTC Drug Formulary for coverage of OTC products.

Fluticasone propionate

PDL Status	P
Quantity Limit	1/30 days
PA Criteria	None

Omnaris®

PDL Status	P
Quantity Limit	1/30 days
PA Criteria	None

Flunisolide

PDL Status	NP
Quantity Limit	2/30 days
PA Criteria	See Below
PA Form	General PA Form
Notes	Please refer to the OTC Drug Formulary for coverage of OTC products.

Flunisolide PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred Rx or covered OTC intranasal steroids.

Mometasone furoate (Rx)

PDL Status	NP
Quantity Limit	1/30 days
PA Criteria	See Below
PA Form	General PA Form

Mometasone furoate PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred Rx or covered OTC intranasal steroids

Qnasl®

PDL Status	NP
Quantity Limit	1/30 days
PA Criteria	See Below
PA Form	General PA Form

Qnasl® PA Criteria:

- Trial and failure, contraindication, or intolerance to 2 preferred Rx or covered OTC intranasal steroids

Xhance®

PDL Status	NP
Quantity Limit	2/30 days
PA Criteria	See Below
PA Form	General PA Form

Xhance® PA Criteria:

- Patient has a trial/failure, contraindication, or intolerance to at least 2 preferred nasal corticosteroid agents; **AND**
- Patient has a clinically valid reason as to why preferred fluticasone propionate products cannot be used

Leukotriene Modifiers

Montelukast tablets and chewables

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Accolate®

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Accolate® PA Criteria:

- Patient is 5 years of age or older and has a diagnosis of asthma; **AND**
- Trial and failure, contraindication, or intolerance of a preferred agent (montelukast tablets or chewables)

Montelukast granules

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form
Notes	For patients less than 3 years of age, no prior authorization is required

Montelukast granules PA Criteria:

- Clinically valid reason why the preferred agents cannot be used

Singulair® chewables, granules, tablets

PDL Status	NP
Quantity Limit	1/day
PA Criteria	See Below
PA Form	General PA Form

Singulair® chewables, granules, tablets PA Criteria:

- Clinically valid reason why the preferred agents cannot be used

Zafirlukast

PDL Status	NP
Quantity Limit	2/day
PA Criteria	See Below
PA Form	General PA Form

Zafirlukast PA Criteria:

- See Accolate® prior authorization criteria

Zileuton CR granules

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Zileuton CR PA Criteria:

- Clinically valid reason why the preferred agents cannot be used

Zyflo®

PDL Status	NP
Quantity Limit	4/day
PA Criteria	See Below
PA Form	General PA Form

Zyflo® PA Criteria:

- Clinically valid reason why the preferred agents cannot be used

Phosphodiesterase (PDE) Inhibitors

Roflumilast

PDL Status	P
Quantity Limit	250 mcg: 28/year 500 mcg: 1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Roflumilast Initial PA Criteria:

- Diagnosis of COPD associated with chronic bronchitis; **AND**
- Patient has forced expiratory volume in 1 second [FEV1] < 50%; **AND**
- Patient is currently receiving standard of care COPD treatments, unless contraindicated (short acting β agonists**OR** short acting anticholinergics PLUS long acting β agonists **OR** long-acting anticholinergics); **AND**
- Patient has a history of continued COPD exacerbations on their current COPD treatment regimen

Roflumilast Renewal PA Criteria:

- Positive clinical response to treatment (e.g., improvement in FEV1 from baseline, reduction in COPD exacerbations); **AND**
- Patient is currently receiving standard of care COPD treatments, unless contraindicated (short acting β agonists**OR** short acting anticholinergics PLUS long acting β agonists **OR** long-acting anticholinergics)

Daliresp®

PDL Status	NP
Quantity Limit	250 mcg: 28/year 500 mcg: 1/day
PA Criteria	See Below
PA Form	General PA Form
PA Approval Duration	6 months

Daliresp® PA Criteria:

- See roflumilast prior authorization criteria; **AND**
- Clinically valid reason why the patient cannot use the preferred generic roflumilast

Ohtuvayre®

PDL Status	NP
Quantity Limit	2 ampules/day
PA Criteria	See Below
PA Form	General PA Form

Ohtuvayre® PA Criteria:

- Patient is ≥ 18 years of age; **AND**
- Diagnosis of Chronic Obstructive Pulmonary Disorder (COPD); **AND**
- Submission of medical records (e.g. chart notes) that patient meets ALL the following:
  - Bronchodilator FEV1/FVC ratio of <0.7
  - FEV1 % predicted of ≤ 79%
  - Modified medical research council (mMRC) dyspnea scale score of ≥ 2; **AND**
- Trial and failure (as defined by continued symptoms, including exacerbations) of adequate treatment concomitantly with ONE of the following:
  - A long-acting beta-agonist (LABA) + long-acting antimuscarinic (LAMA) + inhaled corticosteroid; **OR**
  - A long-acting beta-agonist (LABA) and long-acting antimuscarinic (LAMA); **AND**
- Medication must be used as maintenance therapy only

XVII. SMOKING CESSATION

Smoking Cessation Agents

Please refer to OTC Drug Formulary for coverage of OTC products.

*For children, larger quantities may be approved as medically necessary.

Bupropion Sustained Release tablets (generic for Zyban®)

PDL Status	P
Quantity Limit	2/day; 24 weeks/yr*
PA Criteria	None
Notes	I/DD members not required to meet additional I/DD criteria for this agent.

Chantix® tablets

PDL Status	P
Quantity Limit	2/day; 24 weeks/yr*
PA Criteria	None

Varenicline tablets

PDL Status	P
Quantity Limit	2/day; 24 weeks/yr*
PA Criteria	None

Nicotrol® nasal spray

PDL Status	NP
Quantity Limit	24 weeks/yr*
PA Criteria	See Below
PA Form	General PA Form

Nicotrol® nasal spray PA Criteria:
Standard NP Criteria

XVIII. VITAMINS AND ELECTROLYTES

Fluoride Products

Please refer to OTC Drug Formulary for coverage of OTC products.

Denta 5000® Plus cream, Denta 5000® Gel Plus Sensitive gel, Dentagel® 1.1% gel, NA FL/ K nitro gel

PDL Status	P
Quantity Limit	1 package/Rx
PA Criteria	None

Floriva drops (all formulations except 0.25 mg drops)

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below

Floriva PA Criteria:
Standard NP Criteria

Fraiche (all formulations)

PDL Status	NP
Quantity Limit	1 package/Rx
PA Criteria	See Below
PA Form	See General PA Form

Fraiche PA Criteria:
Standard NP Criteria

Pediatric multivitamins with fluoride tablets

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	See General PA Form

Pediatric multivitamins with fluoride tablets PA Criteria:
Standard NP Criteria

Folic Acid Preparations

Please refer to OTC Drug Formulary for coverage of OTC products.

Folic acid solution 0.2 mg/mL

PDL Status	NP
Quantity Limit	5 mL/day
PA Criteria	See Below
PA Form	See General PA Form

Folic acid solution 0.2 mg/mL PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

L-methylfolate

PDL Status	NP
PA Criteria	See Below
PA Form	See General PA Form

L-methylfolate PA Criteria:

- Patient has documented methylenetetrahydrofolate reductase (MTHFR) mutation/deficiency

Quifolic® solution

PDL Status	NP
Quantity Limit	5 mL/day
PA Criteria	See Below
PA Form	See General PA Form

Quifolic® solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Potassium Depletors

Lokelma® Pak

PDL Status	P
Quantity Limit	1/day
PA Criteria	See Below
PA Form	See General PA Form

Lokelma® Pak Initial PA Criteria:

- Patient must be ≥ 18 years of age; **AND**
- Patient has a diagnosis of chronic hyperkalemia; **AND**
- One of the following:
  - Trial and failure, contraindication, or intolerance to a loop or thiazide diuretic; **OR**
  - Trial and failure, contraindication, or intolerance to a preferred potassium deplete agent

Lokelma® Pak Renewal PA Criteria:

- Patient has a positive clinical response to therapy [e.g., decreased serum potassium levels, levels within normal limits)

Veltassa® powder

PDL Status	P
Quantity Limit	1 packet/day
PA Criteria	See Below
PA Form	See General PA Form

Veltassa® powder Initial PA Criteria:

- Patient must be ≥ 12 years of age; **AND**
- Patient has a diagnosis of chronic hyperkalemia; **AND**
- One of the following:
  - Trial and failure, contraindication, or intolerance to a loop or thiazide diuretic; **OR**
  - Trial and failure, contraindication, or intolerance to a preferred potassium deplete agent

Veltassa® powder Renewal PA Criteria:

- Patient has a positive clinical response to therapy [e.g., decreased serum potassium levels, levels within normal limits)

Kionex® suspension

PDL Status	NP
Quantity Limit	None
PA Criteria	See Below
PA Form	See General PA Form

Kionex® suspension PA Criteria:

Standard NP Criteria

Potassium Supplements

Pokonza® powder, solution

PDL Status	NP
Quantity Limit	10 mEq packets: 20/day; 15 mEq packets: 13/day; 5% soln: 300 mL/day
PA Criteria	See Below
PA Form	See General PA Form

Pokonza® powder, solution PA Criteria:

- The requested agent must be used for a medically necessary diagnosis that is FDA-approved, supported by national compendia, or has medical literature demonstrating its use is clinically safe and effective; **AND**
- Patient has an allergy, contraindication, drug-to-drug interaction, or a history of unacceptable/toxic side effects with ALL preferred medications that is not expected with the requested product

Vitamin B Products

Please refer to OTC Drug Formulary for coverage of OTC products.

Cyanocobalamin injection

PDL Status	P
PA Criteria	See Below
PA Form	See General PA Form
Notes	If the medication is being administered in the prescriber's office **OR** by a Home Health Nurse, coverage must be obtained through the patient's MCO.

Cyanocobalamin injection PA Criteria:

- Diagnosis of Pernicious Anemia; **OR**
- Diagnosis of B12 deficiency; **AND**
  - Provider must submit lab documentation confirming deficiency; **AND**
- Product is being administered by the patient, patient's caregiver, or in a long-term care facility

Cyanocobalamin Nasal Spray

PDL Status	P
PA Criteria	See Below
PA Form	See General PA Form

Cyanocobalamin nasal spray PA Criteria:

- Diagnosis of one of the following:
  - Pernicious Anemia
  - B12 deficiency; **AND**
- Provider must submit lab documentation confirming deficiency

Hydroxocobalamin injection, powder

PDL Status	P
PA Criteria	See Below
PA Form	See General PA Form
Notes	If the medication is being administered in the prescriber's office OR by a Home Health Nurse, coverage must be obtained through the patient's MCO.

Hydroxocobalamin injection, powder PA Criteria:

- Product is being administered by the patient, patient's caregiver, or in a long-term care facility; **AND**
- One of the following:
  - Diagnosis of Pernicious Anemia
  - Dietary deficiency of Vitamin B12 occurring in strict vegetarians and their breastfed infants
  - Malabsorption of Vitamin B12
  - Inadequate secretion of intrinsic factor to facilitate Vitamin B12 absorption
  - Competition for Vitamin B by intestinal parasite or bacteria;
  - Provider must provide peer-reviewed medical literature documenting why the requested drug for the requested indication is the only appropriate choice

Nascobal® nasal spray

PDL Status	NP
PA Criteria	See Below
PA Form	See General PA Form

Nascobal nasal spray PA Criteria:

Standard NP Criteria

Vitamin K Products

Phytonadione

PDL Status	P
Quantity Limit	5/Rx
PA Criteria	None

XIX. WEIGHT MANAGEMENT AGENTS

Appetite and Absorption Agents

Please refer to the OTC Supplement Formulary for coverage of OTC products.

Benzphetamine tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Diethylpropion tablets

PDL Status	P
Quantity Limit	4/day
PA Criteria	None

Diethylpropion ER tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Lomaira® tablets, Orlistat (Rx) capsules, Phendimetrazine IR tablets

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

Phendimetrazine ER capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Phentermine capsules, tablets

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Phentermine/topiramate capsules

PDL Status	P
Quantity Limit	1/day
PA Criteria	None

Xenical® capsules

PDL Status	P
Quantity Limit	3/day
PA Criteria	None

GLP-1 Weight Management Agents

For agents in this class, submit faxed PA requests using the **Weight Management Agents PA Form**.

Wegovy® injection

PDL Status	P
Quantity Limit	4 injections/month
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
Notes	Patients not meeting the below renewal criteria may be approved for up to one month to allow for titration off medication

Wegovy® Injection Initial PA Criteria:

- Patient must be the labeled age minimum (Wegovy ≥ 12); **AND**
- Treatment is for the management of obesity; **AND**
- One of the following:
  - For patients < 18 years of age, submission of medical records (e.g. chart notes) documenting the BMI is ≥ 95th percentile standardized for age and sex; **OR**
  - For patients ≥ 18 years of age, one of the following:
    - Submission of medical records (e.g. chart notes) documenting a body mass index (BMI) ≥ 30 kg/m^2; **OR**
    - Submission of medical records (e.g. chart notes) documenting a BMI of greater than 27 kg/m^2 with a weight related comorbidity (e.g. hypertension, dyslipidemia, diabetes, coronary heart disease, MASH/NASH, obstructive sleep apnea); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications FDA approved for obesity/weight loss; **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)

Wegovy® Injection Renewal PA Criteria:

- Patient has continued participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications for obesity/weight loss; **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- One of the following:
  - For patients ≥ 18 years of age, submission of medical records (e.g., chart notes) documenting a weight loss of ≥ 5% of baseline body weight; **OR**
  - For patients < 18 years of age, submission of medical records (e.g., chart notes) documenting a weight loss of ≥ 5% of baseline BMI-for-age

Zepbound® single dose prefilled pen

PDL Status	P
Quantity Limit	4 injections/month
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
Notes	Patients not meeting the below renewal criteria may be approved for up to one month to allow for titration off medication

Zepbound® single dose prefilled pen Initial PA Criteria:

- Patient must be the labeled age minimum (Zepbound ≥ 18); **AND**
- Treatment is for the management of obesity; **AND**
- One of the following:
  - Submission of medical records (e.g. chart notes) documenting a body mass index (BMI) ≥ 30 kg/m^2; **OR**
  - Submission of medical records (e.g. chart notes) documenting a BMI of greater than 27 kg/m^2 with a weight related comorbidity (e.g. hypertension, dyslipidemia, diabetes, coronary heart disease, MASH/NASH, obstructive sleep apnea); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications FDA approved for obesity/weight loss; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)

Zepbound® single dose prefilled pen Renewal PA Criteria:

- Patient has continued participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications for obesity/weight loss; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Submission of medical records (e.g., chart notes) documenting a weight loss of ≥ 5% of baseline body weight

Liraglutide (weight management) injection

PDL Status	NP
Quantity Limit	3 mg/day
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
Notes	Patients not meeting the below renewal criteria may be approved for up to one month to allow for titration off medication

Liraglutide Injection Initial PA Criteria:

- Patient must be the labeled age minimum (liraglutide ≥ 12); **AND**
- Treatment is for the management of obesity; **AND**
- One of the following:
  - For patients < 18 years of age, submission of medical records (e.g. chart notes) documenting the BMI is ≥ 95th percentile standardized for age and sex; **OR**
  - For patients ≥ 18 years of age, one of the following:
    - Submission of medical records (e.g. chart notes) documenting a body mass index (BMI) ≥ 30 kg/m^2; **OR**
    - Submission of medical records (e.g. chart notes) documenting a BMI of greater than 27 kg/m^2 with a weight related comorbidity (e.g. hypertension, dyslipidemia, diabetes, coronary heart disease, MASH/NASH, obstructive sleep apnea); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications FDA approved for obesity/weight loss; **AND**
- Will not be co-administered another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)
- If request is for liraglutide, trial and failure of two preferred agents; **AND**
- Clinically valid reason why brand Saxenda cannot be used.

Liraglutide Injection Renewal PA Criteria:

- Patient has continued participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)
- Medication will not be used in combination with other medications for obesity/weight loss; **AND**
- Submission of medical records (e.g., chart notes) documenting a weight loss of ≥ 5% of baseline body weight

Saxenda® injection

PDL Status	NP
Quantity Limit	3 mg/day
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
Notes	Patients not meeting the below renewal criteria may be approved for up to one month to allow for titration off medication

Saxenda® Injection Initial PA Criteria:

- Patient must be the labeled age minimum (Saxenda® ≥ 12); **AND**
- Treatment is for the management of obesity; **AND**
- One of the following:
  - For patients < 18 years of age, submission of medical records (e.g. chart notes) documenting the BMI is ≥ 95th percentile standardized for age and sex; **OR**
  - For patients ≥ 18 years of age, one of the following:
    - Submission of medical records (e.g. chart notes) documenting a body mass index (BMI) ≥ 30 kg/m^2; **OR**
    - Submission of medical records (e.g. chart notes) documenting a BMI of greater than 27 kg/m^2 with a weight related comorbidity (e.g. hypertension, dyslipidemia, diabetes, coronary heart disease, MASH/NASH, obstructive sleep apnea); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications FDA approved for obesity/weight loss; **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Patient does not have personal or immediate family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 (MEN2)
- If request is for Saxenda®, trial and failure of two preferred agents

Saxenda® Injection Renewal PA Criteria:

- Patient has continued participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Will not be co-administered with another injectable or oral GLP-1 receptor agonists (i.e., exenatide, liraglutide, Ozempic® inj, tabs [formerly Rybelsus®], Zepbound®); **AND**
- Will not be used concomitantly with DPP-4 inhibitors**; AND**
- Medication will not be used in combination with other medications for obesity/weight loss; **AND**
- Submission of medical records (e.g., chart notes) documenting a weight loss of ≥ 5% of baseline body weight

Neurobehavioral Appetite Modulators

For agents in this class, submit faxed PA requests using the Weight Management Agents PA Form.

Amphetamine sulfate tablets

PDL Status	P
Quantity Limit	5 mg: 3/day 10 mg: 6/day
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
PA Approval Duration	3 months
Notes	Patients may be approved for one additional month to allow for titration off medication. PA not required for members < 21 years old.

Initial Amphetamine sulfate PA Criteria:

- Patient ≥ 21 years old; **AND**
- Agent must not be prescribed by a pain clinic; **AND**
- Treatment is for the management of obesity; **AND**
- One of the following:
  - For patients < 18 years of age, submission of medical records (e.g. chart notes) documenting the BMI is ≥ 95th percentile standardized for age and sex; **OR**
  - For patients ≥ 18 years of age, one of the following:
    - Submission of medical records (e.g. chart notes) documenting a body mass index (BMI) ≥ 30 kg/m^2; **OR**
    - Submission of medical records (e.g. chart notes) documenting a BMI of greater than 27 kg/m^2 with a weight related comorbidity (e.g. hypertension, dyslipidemia, diabetes, coronary heart disease, MASH/NASH, obstructive sleep apnea); **AND**
- Prescriber attests patient is participating in complementary nutritional and lifestyle changes (e.g., dietary modification, increased physical activity as medically able, structured behavioral intervention, comprehensive weight management program); **AND**
- Medication will not be used in combination with other medications FDA approved for obesity/weight loss; **AND**
- Trial and failure, contraindication, or intolerance of 2 preferred agents

Evekeo® tablets

PDL Status	NP
Quantity Limit	5 mg: 3/day 10 mg: 6/day
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
PA Approval Duration	3 months
Notes	Patients may be approved for one additional month to allow for titration off medication. PA not required for members < 21 years old.

Evekeo® PA Criteria:

See amphetamine sulfate criteria; **AND**

- Clinically valid reason why generic amphetamine sulfate tablets cannot be used.

Obesity Agents for Rare Disorders

For agents in this class, submit faxed PA requests using the Weight Management Agents PA Form.

Imcivree® injection

PDL Status	NP
Quantity Limit	90 mg/month
PA Criteria	See Below
PA Form	Weight Management Agents PA Form
PA Approval Duration	6 months

Obesity due to POMC, PSCK1, or LEPR gene deficiency or Bardet-Biedl Syndrome

Imcivree® Initial PA Criteria:

- Diagnosis of ONE of the following:
  - Obesity due to POMC, PCSK1, or LEPR gene deficiency confirmed with genetic testing interpreted as pathogenic, likely pathogenic, or of uncertain significance; **OR**
  - Bardet-Biedl syndrome; **AND**
- ONE of the following:
  - Patient < 18 years of age with weight ≥ 95th percentile for age; **OR**
  - Patient ≥ 18 years BMI ≥ 30 kg/m^2; **AND**
- Prescribed by or in consultation with an endocrinologist, a geneticist, or a physician who specializes in metabolic disorders

Imcivree® Renewal PA Criteria:

- Positive clinical response to therapy as defined by one of the following:
  - For patients ≥ 18 years of age, submission of medical records documenting weight loss from baseline weight ≥ 5% if on therapy < 12 months or ≥ 10% if on therapy > 12 months
  - For patients < 18 years of age, submission of medical records documenting weight loss from baseline BMI-for-age of ≥ 5% if on therapy < 12 months or ≥ 10% if on therapy > 12 months

Acquired Hypothalamic Obesity (HO)

Imcivree® injection Initial PA Criteria:

- Diagnosis of Acquired Hypothalamic Obesity (HO) as confirmed by submission of medical records (i.e., chart notes) documenting ONE of the following:
  - Brain tumors (e.g., craniopharyngiomas, gliomas, pituitary adenomas, hamartomas)
  - Neurosurgical intervention
  - Cranial radiation therapy
  - Traumatic brain injury
  - Stroke
  - Non-tumor structural hypothalamic insult/injury due to inflammation, infection, or bleeding; **AND**
- ONE of the following:
  - Patient < 18 years of age with weight ≥ 95th percentile for age; **OR**
  - Patient ≥ 18 years of age with BMI ≥ 30 kg/m^2; **AND**
- Prescribed by, or in consultation with, an endocrinologist, a geneticist, or a physician who specializes in metabolic disorders

Imcivree® injection Renewal PA Criteria:

- Positive clinical response to therapy as defined by one of the following:
  - For patients ≥ 18 years of age, submission of medical records documenting weight loss from baseline weight ≥ 5% if on therapy < 12 months or ≥ 10% if on therapy > 12 months
  - For patients < 18 years of age, submission of medical records documenting weight loss from baseline BMI-for-age ≥ 5% if on therapy < 12 months or ≥ 10% if on therapy > 12 months