

Response to Reviewers - Duraisamy et al., 2025, IRCOB

Reviewer 1

This is a study of low-moment range of motion in male and female osteoligamentous cervical spine segments based on previous testing in the NHTSA database performed. In general, both the intrinsic and size differences between men and women are of substantial interest, but here, no attempt is made to differentiate size related differences through biomechanical scaling, and certain of the underlying assumptions are questionable and/or unspecified.

Methods

1. At some point, the anatomical differences between female and male segments must be mentioned. Anatomically, the females are likely to be smaller AND the female FSUs are higher in the c-cpine in this dataset. Both are going to tend to exacerbate size-related differences in response to low moments and to injury response.

- Although we had already addressed these in the subsections “Limitations” and “Sex-dependent cervical spine biomechanics” in the Discussion, we understand it better to elaborate further on this to give a clearer picture to the reader. We have now expanded more on this with additional references in the section “Sex-dependent cervical spine biomechanics” and added a comment on the cervical spine segments available in the dataset in the “Limitations”.

2. “The hyperprior is given by ...”. It’s unclear why the prior for a male/female comparison comes from a male study as opposed to just using an uninformed prior. Or something else. Please specify what part of [14] was used for the priors since it is experimental and computational. If the experiments from [14] are used, are these some of the same experiments reported in [12]? Please clarify.

- Even though the priors come from a male study, the parameters were defined such that prior is a weakly informative prior. The purpose of a weakly informative prior was to center the prior distributions. We have now added an additional figure in the appendix to show this (Fig A3). If anything, a weakly informative prior in this case will lead to conservative results. With the lack of additional experimental sources and with the lack information in this source as indicated in our responses to #1 [TODO], we can safely assume that specification of the current prior does not affect the conclusion of the study.
- Hyperprior specifies distribution for flexion and extension (not sex-specific priors)
- experimental data from appendix of [14] is used in this paper. This study consisted of only male samples. [12] contains only female samples and hence we can conclude that they are not the same experiments.
- We also checked if [13] and [14] contain the same experiments. The cervical spine levels used in the two studies are different and hence can be concluded that it is not the same samples. Also, the studies acknowledge different funders.

3. Some of the distribution assumptions seem unnecessary if Bayesian nonparametric modelling is used.

- We are using parametric linear regression (shown in equation 1)

4. Generally, some scaling for size must be included in such a study since this is likely a strong predictor of low moment motion.

- The initial plan for our study was to include anthropometric and morphological variables in the analysis. However, these are not recorded in the NHTSA database for these experiments. Height and weight data was found in pdf report for only one PMHS B03 (female).
- All other papers published by Nightingale and Myers were checked to see if any related information could be extracted. Such information is available only for PMHS used for tension tests. PMHS id used in bending tests were different from tension tests,

Results

5. The Bayesian model specification should be in the methods.

- Thank you this is moved the methods

6. “The posterior distribution was verified using various checks.” Please specify what this means. What checks, and what constituted verification in this case.

- Already described in the Appendix. Add pointers in the main text

7. “for the group level-effect, extension and flexion responses for females are seen to be greater than males.” This is unclear. If it means that the extension and flexion angles for females to a given moment level are greater than males, with some anthropometric specifications), then this is likely generally true. However, some discussion of this would be nice here, including the likely observation that size differences play a role here. - Explain better and link with point #1 - (PD) As i mentioned in point 4, we lack the PMHS-specific anthropometric data from the chosen literature/database. - (PD) May be just for the sake of discussion we could use the winkelstein paper (Very less data though), Height shows some parabolic trend with rotation angle and weight have increasing slope. - (PD) When i chose BMI for generalisation, the slope for the flexion is changing wrt the cervical level, but not for extension. But we basically we couldnt conclude as the data chosen were very small (only 5)

9. **One additional limitation is repeated testing on these specimens. At 3 Nm, this is likely not a primary effect, but should be mentioned.**

- Add in Discussion/Results/Methods

Discussion

9. “... male ... has overlapping distributions ... female ... did not show ... overlap”. Likely a part of this is the size differences and the moment angle relative to the overall range of motion in an osteoligamentous c-spine in flexion. The increase in extension moment happens nearer in angle to the nominal position than the flexion. One thing more difficult to understand is why the distributions do not widen in the flexion cases and seem approximately the same as in the extension case.

- Distributions of the rotations are at constant moment of 3Nm.
- The distribution represents the uncertainty in data an variations in the responses from the PMHS samples and doesn’t necessarily have to do with flexion or extension loading modes.

10. “... could perhaps...” This is well known. In fact, it is quite difficult to injure the neck in pure flexion moment before facial structures contact the torso. In contrast, extension moments load the c-spine to failure at relatively small angles.

- The comment here was about the sex-differences in physiological range-of-motion and not about the responses to injurious loading

11. “... may exhibit sex-dependent variations.” From a structural anthropometric point of view, males and females have size-dependent variations. Also, this is true within males and females. So, for such a low-moment study, it is essential to separate sex intrinsic variations and geometric variations. Specimen anthropometry is available for these tests. Using the gross anthropometry and segmental correlations of anthropometry and segmental length (and circumferences), some estimate of the size effect can be obtained.

- Check for info in the data and if available include in the regression.
- (PD) One paper (<https://www.sciencedirect.com/science/article/pii/S0021929006003289>) contains the cervical morphological data for the same PMHS, but only for upper cervical spine, none of those FSUs are used in this study as we are focusing only on lower cervical spine.

Appendix

12. "... there were 48 female and 29 male FSUs...". Please clarify these are tests, using 16 female and 16 male spine as in the studies. They are not independent spines.

- Good point. Update this one

Reviewer 2

This study quantifies sex differences in cervical neck kinematics using retrospective data from previously published experimental work.

13. Introduction: what is the status of cervical neck biofidelity in current HBMs using the morphing techniques and/or limited experimental validation data? Are their biofidelity scores low with these approaches? Please expand on the practical effects for HBMs when sex is not considered in calibration/validation.

- Add comments about current HBMs
- (PD) <https://www.ircobi.org/wordpress/downloads/irc16/pdf-files/116.pdf> THUMS, they did C4-C5 validationn (with Nightingale male paper) using male specific material properties. Shown that 'THUMS v4.02 AM50 (VPS)' C45 FSU performs poorly in 3.5Nm moment loading. The main idea is to ensure gender specific stiffness response.
- (PD) <https://www.sciencedirect.com/science/article/pii/S0001457523002294> Though they scaled GHBM C F50 from GHBM C F05 (which is infact a scaled down male model, Disc 3rd para) Why they scale the mass to an average male (as they say in conclusion 1st para) ?

14. Methods: It would be prudent to include some additional details for the retrospective data so that the reader does not have to search through refs 12 and 13 to understand if the boundary conditions were the same across the published data that were used:

- did they change boundary conditions between tests/sex. It appears that different segmentation of the cervical vertebrae was used at the least.
 - mentioned in appendix AII. Include a statement in methods
- Were the ages of the PMHS similar across sex?
 - check in the data

- (PD) For male age varies from 47 to 74 (61.13 avg), For female age varies from 33 to 58 (48.16 avg)
 - Were there any inclusion/exclusion criteria that may influence the findings of this current paper (especially any cervical degeneration that may influence response)?
 - Check
 - (PD) Nightingale didnt mention any of such findings in PMHS. But most of the COD were related to cardio, so there wont be any stuffs that would potentially influence the Spinal rotations i guess
 - Were data from all cervical segments referenced (C3-C4 and C5-C6 for females; C4-C5 and C6-C7 for males) combined in the Bayesian model? Are the degrees reported in the results an average across all segments? Your abstract states “segmental rotation” but you’re reporting only singular degree values when comparing males and females. Please clarify.
 - Check and Align wordings
15. Please include the criteria or justification or rationale for your selection of “wider distributions” when defining the priors. How could this selection decision affect your results?
- write more on prior selection
 - (PD) Winkelstein used similar test apparatus and methodology like nightingale but tested for only Male PMHS at 2.5 Nm, as the bayesian model is for 3 Nm and also for comparing female, we chose wider prior.
16. Results: The terms “rotation” and “flexion/extension” seem to be used interchangeably? I am confused by the use of segmental rotation in flexion and extension. What anatomical plane is segmental rotation being measured in? This again is likely discussed in the referenced papers but brief clarifications here and/or a figure demonstrating the results would help to interpret your analysis.
- Add/align/review to make it easy for someone not familiar with cervical spine
17. Discussion: It is important, in your discussion of IV disc and ligaments exhibiting sex-specific variation to include some discussion/interpretation based on age (and the age distributions of the males vs. females included).
- Check if available in literature
 - (PD) Most of the papers i found were like relating the age/sex with level of disc degeneration but not with rotation angle.
 - (PD) <https://doi.org/10.1016/j.math.2009.08.006> Check discussion 5.1. Its lumbar, also data was collected from multiple studies.
 - (PD) <https://doi.org/10.3171/2015.1.SPINE14489> This paper speaks about age and sex differences but they just mentioned like they measured ROM, i am not clear about their methodology. It might be based upon just scan image data on the extreme postures in flexion and extension i guess.

- (PD) May be at some stage we have to admit that age might influence the rotation angle, but it depends on the PMHS majorly, can even be attributed to degeneration due to age. Still based upon our bayesian model, we could still argue that the effect of sex and loading mode were more dominant than age.
 - (PD) Also those are live subjects, they might have asked to bend till its painful, Not like a standard testing apparatus used on a cadaver
18. Your findings are asymmetry in flexion/extension response but also between sexes. Are current HBMs allowing for cervical flexion and extension kinematics to be equal? Please elaborate and discuss further the implications of your work presented here.
- Check validation reports from current HBMs. Connect to 13.
 - (PD) We didn't have any literature showing HBMs having same response on Extension and Flexion, But one thing we could say is THUMS didnt validate its cervical for this quasi static moment loading.
 - (PD) moreover in 13, 1st paper they spoke only about Flexion. The author mentioned that the FSU is very stiff (approx. 2 deg. at 3.5 Nm). As it cannot hit the spinous process at that angle, we could expect the same in Extension too. I couldn't find any other literatures from the authors.
19. Again, please clarify your use of the term "spinal rotations." Are you using this interchangeably with flexion/extension movements?
- Align/review terminology
20. In the limitations, you mention that females may have smaller cervical vertebrae and intervertebral dimensions than males that could influence rotations; however, the anatomical features are conserved across sexes such that size would not influence relative movement of the cervical segments. Suggest re-thinking this assertion with an anatomical perspective.
- Anatomical features are similar - that is why we see similar response in extension. But in flexion, same load of 3 Nm is taken by a smaller spinal segment in the females and hence the difference in flexion.