

Mapping to Our Problem

$$\bullet \Theta = \{(\mathcal{U}, \mathcal{V}) : \mathcal{U}, \mathcal{V} \in \mathbb{R}^n\}$$

- w is the Gaussian distribution over Θ

• $\mathcal{P}_{(u,v)}$ is the distribution of $(a/\sqrt{n}) \cdot Q^{(1)}(u \otimes v) + Q^{(1)} \cdot \text{vec}(G)$

- By rotational invariance: $Q^{(1)} \cdot \text{vec}(G) \sim N(0, I_t)$

$$\bullet \mathcal{P}_{(u,v)} = N(\mu = (a/\sqrt{n}) \cdot Q^{(1)}(u \otimes v), \Sigma = I_t)$$

36

Mapping to Our Problem

- $\Theta = \{(u, v) : u, v \in \mathbb{R}^n\}$
- w is the Gaussian distribution over Θ
- $\mathcal{P}_{(u,v)}$ is the distribution of $(\alpha/\sqrt{n}) \cdot Q^{(1)}(u \otimes v) + Q^{(1)} \cdot \text{vec}(G)$
 - By rotational invariance : $Q^{(1)} \cdot \text{vec}(G) \sim N(0, I_t)$
 - $\mathcal{P}_{(u,v)} = N(\mu = (\alpha/\sqrt{n}) \cdot Q^{(1)}(u \otimes v), \Sigma = I_t)$

Main idea