



Mapping to Our Problem

$$\bullet \Theta \equiv \{(\mathcal{U}, \mathcal{V}) : \mathcal{U}, \mathcal{V} \in \mathbb{R}^n\}$$

- $w$  is the Gaussian distribution over  $\Theta$

•  $\mathcal{P}_{(u,v)}$  is the distribution of  $(a/\sqrt{n}) \cdot Q^{(1)}(u \otimes v) + Q^{(1)} \cdot \text{vec}(G)$

- By rotational invariance:  $Q^{(1)} \cdot \text{vec}(G) \sim N(0, I_t)$

$$\bullet \mathcal{P}_{(u,v)} = N(\mu = (\alpha/\sqrt{n}) \cdot Q^{(1)}(u \otimes v), \Sigma = I_t)$$

# Mapping to Our Problem

- $\Theta = \{(u, v) : u, v \in \mathbb{R}^n\}$
- $w$  is the Gaussian distribution over  $\Theta$
- $\mathcal{P}_{(u,v)}$  is the distribution of  $(\alpha/\sqrt{n}) \cdot Q^{(1)}(u \otimes v) + Q^{(1)} \cdot \text{vec}(G)$ 
  - By rotational invariance :  $Q^{(1)} \cdot \text{vec}(G) \sim N(0, I_t)$
  - $\mathcal{P}_{(u,v)} = N(\mu = (\alpha/\sqrt{n}) \cdot Q^{(1)}(u \otimes v), \Sigma = I_t)$



# Main idea