



Pranesh S A 251801234-2025-AIDS ▾

P2

Started on	Monday, 5 October 2026, 9:55 PM
State	Finished
Completed on	Monday, 5 October 2026, 9:56 PM
Time taken	1 min 9 secs
Marks	1.00/1.00
Grade	10.00 out of 10.00 (100%)

Question 1 | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2
3  int main() {
4      int n, k;
5      if (scanf("%d", &n) != 1) return 0;
6
7      int arr[n];
8      for (int i = 0; i < n; i++) {
9          scanf("%d", &arr[i]);
10     }
11
12     scanf("%d", &k);
13
14     int i = 0, j = 1, ans = 0;
15     while (i < n && j < n) {
16         if (i != j && arr[j] - arr[i] == k) {
17             ans = 1;
18             break;
19         } else if (arr[j] - arr[i] < k) {
20             j++;
21         } else {
22             i++;
23         }
24     }
25
26     printf("%d\n", ans);
27     return 0;
28 }
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)