

Hermitian Forms and Zeros of a Polynomial

PRANSHU GABA *

Indian Institute of Science, Bangalore
gabapranshu@ug.iisc.in

November 12, 2017

Abstract

We looked at the general properties of Hermitian (self-adjoint) matrices, and used the Schur-Cohn theorem to find the number of roots of a polynomial lying within and without the unit circle.

1 Introduction

In this paper we see the properties of Hermitian matrices, which are very interesting, as well as useful. We also see and prove the Schur-Cohn theorem to find the number of roots of a polynomial lying within the unit circle.

There are many ways to locate the roots of a polynomial. The Schur-Cohn theorem shows a surprising connection between linear algebra and roots of a polynomial. It will be used to find out how many roots of the polynomial lie inside and outside the unit circle.

2 Definitions

2.1 Adjoint

The adjoint of a matrix $A \in \mathbb{C}_n$, denoted by A^* , is the matrix obtained by taking its transpose, followed by taking the complex conjugate of every element. If the ij^{th} of A is a_{ij} , then the ij^{th} entry of A^* is $\overline{a_{ji}}$. Note that A^* is a linear transformation. The adjoint satisfies $\langle Ax, y \rangle = \langle x, A^*y \rangle$.

2.2 Positive Definite

Definition. Any matrix $B \in \mathbb{M}_n$ that satisfies $\langle Bx, x \rangle \geq 0$ for all $x \in \mathbb{C}^n$ is called a positive semidefinite matrix.

Corollary. *All the eigenvalues of positive semidefinite matrix are non-negative.*

*:)

2.3 Hermitian Matrices

Hermitian matrices (also known as self-adjoint matrices) are matrices that satisfy $A = A^*$.

Theorem. *All the eigenvalues of a Hermitian matrix are real.*

Proof. Left as an exercise. □

A^*A is always positive semidefinite.

2.4 Unitary Matrices

2.5 Matrix S

Let S be the $n \times n$ square matrix
$$\begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix}$$
. Note that S is a nilpotent matrix of order n , i.e. S^n is a zero matrix.

2.6 Norm of a matrix

2.6.1 Operator

2.6.2 Hilbert norm

2.7 Diagonalization

Hermitian matrices can be diagonalized. For every Hermitian matrix A , there exists a diagonal matrix Λ such that $A = U^* \Lambda U$. Here U is some unitary matrix.

Lemma. *If $A \in \mathbb{M}_n(\mathbb{C})$ and $\langle Ax, x \rangle \in \mathbb{R}$ for every x , then $A = A^*$.*

Proof. Let $\alpha \in \mathbb{C}$ and $h, g \in \mathbb{C}^n$. Then □

Corollary. *Every positive semidefinite matrix is Hermitian.*

2.8 Projection

3 Schur-Cohn Theorem

Given any polynomial $p(z) = a_0 z^n + a_1 z^{n-1} + \dots + a_n$ with complex coefficients, we are interested in finding how many of its roots lie within the unit circle and how many roots lie outside. Without loss of generality, let $a_0 = 1$ as it does not change the roots of the polynomial.

Suppose p has roots α_i . Then $p(z) = (z - \alpha_1)(z - \alpha_2) \dots (z - \alpha_n)$.

Then $p(S)$ is
$$\begin{bmatrix} a_n & a_{n-1} & \ddots & \ddots & a_1 \\ 0 & a_n & a_{n-1} & \ddots & \ddots \\ 0 & 0 & a_n & \ddots & \ddots \\ 0 & 0 & 0 & \ddots & a_{n-1} \\ 0 & 0 & 0 & 0 & a_n \end{bmatrix}.$$

This can be factorized as $p(S) = (S - \alpha_1 I)(S - \alpha_2 I) \cdots (S - \alpha_n I)$. Let $B_j = S - \alpha_j I$.

Next, define q as the polynomial $\overline{a_n} z^n + \overline{a_{n-1}} z^{n-1} + \cdots + \overline{a_0}$. Note that its roots are $\frac{1}{\overline{\alpha_i}}$. We get $q(z) = (1 - \overline{\alpha_1} z)(1 - \overline{\alpha_2} z) \cdots (1 - \overline{\alpha_n} z)$. Also, $q(S) = (I - \overline{\alpha_1} S)(I - \overline{\alpha_2} S) \cdots (I - \overline{\alpha_n} S)$. Let $C_j = I - \overline{\alpha_j} S$.

Let H be equal to $\|q(S)x\|^2 - \|p(S)x\|^2$

H can also be written as $\langle (q(S)^* q(S) - p(S)^* p(S))x, x \rangle$.

We can now state the Schur-Cohn theorem:

Theorem. *The polynomial p , it will have k roots inside the circle, and $n - k$ roots outside the circle iff k eigenvalues of H are positive and $n - k$ are negative.*

4 Proof

4.1 Linear Polynomial

Let's write $q(S)$ and $p(S)$ as a product of the linear terms. $q(S)^* q(S) - p(S)^* p(S)$
 $= (C_1 C_2 C_3 \cdots C_n)^* (C_1 C_2 C_3 \cdots C_n) - (B_1 B_2 B_3 \cdots B_n)^* (B_1 B_2 B_3 \cdots B_n)$

Let's look at $C_1^* C_1 - B_1^* B_1$ first. Substituting the values of C_1 and B_1 , we get

$$\begin{aligned} & C_1^* C_1 - B_1^* B_1 \\ &= (I - \overline{\alpha_1} S)^* (I - \overline{\alpha_1} S) - (S - \alpha_1 I)^* (S - \alpha_1 I) \\ &= (I - \alpha_1 S^*) (I - \overline{\alpha_1} S) - (S^* - \overline{\alpha_1} I) (S - \alpha_1 I) \\ &= (I - \alpha_1 S^* - \overline{\alpha_1} S + |\alpha_1|^2 S^* S) - (S^* S - \alpha_1 S^* - \overline{\alpha_1} S + |\alpha_1|^2 I) \\ &= I - |\alpha_1|^2 I - S^* S + |\alpha_1|^2 S^* S \\ &= (1 - |\alpha_1|^2) (I - S^* S) \end{aligned}$$

Note that $I - S^* S$ is a positive definite matrix. If $|\alpha| < 1$, then the root of the linear polynomial lies within the unit circle. Also note that H has one negative eigenvalue. Similarly, if $|\alpha| > 1$, then the root of the linear polynomial lies outside the unit circle, and the eigenvalue of H is positive. This shows that the Schur-Cohn theorem is true for $n = 1$. We will now extend the proof for all n .

4.2 Spectral theorem and proof

4.3 CF theorem and proof

5 Extensions

5.1 Arbitrary radius

5.2 Limitations

6 Conclusion

Thank You!

7 Acknowledgements