

Olympiad problem collection

Compiled by Prasanna Ramakrishnan

Updated: August 27, 2021

Introduction. The best way to get better at olympiad mathematics is to do lots of problems, but sometimes it can be hard to find suitable problems from the vast array of contests available online. This is a collection of nice problems from international olympiads suitable for students aiming to get an Honorable mention or Bronze medal at the IMO. Problems in each section are arranged roughly in order of difficulty (from my perspective). Do not take the arrangement too seriously – they are all around the same level.

Algebra

Easier than IMO

(Centro SL 2008) Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 = 12$. Show that

$$\sqrt{1+a^3} + \sqrt{1+b^3} + \sqrt{1+c^3} \leq 9.$$

(Centro SL 2008) Prove that if x and y are real numbers such that $(x + \sqrt{x^2 + 1})(y + \sqrt{y^2 + 1}) = 1$, then $x + y = 0$.

(Centro 2020/5) Let $P(x)$ be a polynomial with real non-negative coefficients. Let k be a positive integer and x_1, x_2, \dots, x_k positive real numbers such that $x_1 x_2 \cdots x_k = 1$. Prove that

$$P(x_1) + P(x_2) + \cdots + P(x_k) \geq kP(1).$$

(Centro 2020/3) Find all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ satisfying the following property: if a, b and c are integers such that $a + b + c = 0$, then

$$f(a) + f(b) + f(c) = a^2 + b^2 + c^2.$$

(EGMO 2019/1) Find all triples (a, b, c) of real numbers such that $ab + bc + ca = 1$ and

$$a^2b + c = b^2c + a = c^2a + b.$$

IMO level

(IMO 2010/1) Find all function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for all $x, y \in \mathbb{R}$ the following equality holds

$$f([x]y) = f(x)[f(y)]$$

where $[a]$ is greatest integer not greater than a .

(IMO 1990/4) Let \mathbb{Q}^+ be the set of positive rational numbers. Construct a function $f : \mathbb{Q}^+ \rightarrow \mathbb{Q}^+$ such that

$$f(xf(y)) = \frac{f(x)}{y}$$

for all x, y in \mathbb{Q}^+ .

(IMO 2014/1) Let $a_0 < a_1 < a_2 \dots$ be an infinite sequence of positive integers. Prove that there exists a unique integer $n \geq 1$ such that

$$a_n < \frac{a_0 + a_1 + a_2 + \cdots + a_n}{n} \leq a_{n+1}.$$

(IMO 2008/4) Find all functions $f : (0, \infty) \mapsto (0, \infty)$ (so f is a function from the positive real numbers) such that

$$\frac{(f(w))^2 + (f(x))^2}{f(y^2) + f(z^2)} = \frac{w^2 + x^2}{y^2 + z^2}$$

for all positive real numbers w, x, y, z , satisfying $wx = yz$.

(IMO 2007/1) Real numbers a_1, a_2, \dots, a_n are given. For each i , ($1 \leq i \leq n$), define

$$d_i = \max\{a_j \mid 1 \leq j \leq i\} - \min\{a_j \mid i \leq j \leq n\}$$

and let $d = \max\{d_i \mid 1 \leq i \leq n\}$.

(a) Prove that, for any real numbers $x_1 \leq x_2 \leq \dots \leq x_n$,

$$\max\{|x_i - a_i| \mid 1 \leq i \leq n\} \geq \frac{d}{2}. \quad (*)$$

(b) Show that there are real numbers $x_1 \leq x_2 \leq \dots \leq x_n$ such that the equality holds in (*).

(IMO 2019/1) Let \mathbb{Z} be the set of integers. Determine all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that, for all integers a and b ,

$$f(2a) + 2f(b) = f(f(a + b)).$$

(IMO 2004/4) Let $n \geq 3$ be an integer. Let t_1, t_2, \dots, t_n be positive real numbers such that

$$n^2 + 1 > (t_1 + t_2 + \dots + t_n) \left(\frac{1}{t_1} + \frac{1}{t_2} + \dots + \frac{1}{t_n} \right).$$

Show that t_i, t_j, t_k are side lengths of a triangle for all i, j, k with $1 \leq i < j < k \leq n$.

(IMO 2012/4) Find all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that, for all integers a, b, c that satisfy $a + b + c = 0$, the following equality holds:

$$f(a)^2 + f(b)^2 + f(c)^2 = 2f(a)f(b) + 2f(b)f(c) + 2f(c)f(a).$$

(Here \mathbb{Z} denotes the set of integers.)

Combinatorics

(IMO 2018/4) A site is any point (x, y) in the plane such that x and y are both positive integers less than or equal to 20.

Initially, each of the 400 sites is unoccupied. Amy and Ben take turns placing stones with Amy going first. On her turn, Amy places a new red stone on an unoccupied site such that the distance between any two sites occupied by red stones is not equal to $\sqrt{5}$. On his turn, Ben places a new blue stone on any unoccupied site. (A site occupied by a blue stone is allowed to be at any distance from any other occupied site.) They stop as soon as a player cannot place a stone.

Find the greatest K such that Amy can ensure that she places at least K red stones, no matter how Ben places his blue stones.

(IMO 2015/1) We say that a finite set \mathcal{S} of points in the plane is balanced if, for any two different points A and B in \mathcal{S} , there is a point C in \mathcal{S} such that $AC = BC$. We say that \mathcal{S} is centre-free if for any three different points A , B and C in \mathcal{S} , there is no points P in \mathcal{S} such that $PA = PB = PC$.

- (a) Show that for all integers $n \geq 3$, there exists a balanced set consisting of n points.
- (b) Determine all integers $n \geq 3$ for which there exists a balanced centre-free set consisting of n points.

(IMO 2011/4) Let $n > 0$ be an integer. We are given a balance and n weights of weight $2^0, 2^1, \dots, 2^{n-1}$. We are to place each of the n weights on the balance, one after another, in such a way that the right pan is never heavier than the left pan. At each step we choose one of the weights that has not yet been placed on the balance, and place it on either the left pan or the right pan, until all of the weights have been placed.

Determine the number of ways in which this can be done.

(IMO 2003/1) Let A be a 101-element subset of the set $S = \{1, 2, \dots, 1000000\}$. Prove that there exist numbers t_1, t_2, \dots, t_{100} in S such that the sets

$$A_j = \{x + t_j \mid x \in A\}, \quad j = 1, 2, \dots, 100$$

are pairwise disjoint.

(IMO 2002/1) Let n be a positive integer. Each point (x, y) in the plane, where x and y are non-negative integers with $x + y < n$, is coloured red or blue, subject to the following condition: if a point (x, y) is red, then so are all points (x', y') with $x' \leq x$ and $y' \leq y$. Let A be the number of ways to choose n blue points with distinct x -coordinates, and let B be the number of ways to choose n blue points with distinct y -coordinates. Prove that $A = B$.

(IMO 2001/4) Let n be an odd integer greater than 1 and let c_1, c_2, \dots, c_n be integers. For each permutation $a = (a_1, a_2, \dots, a_n)$ of $\{1, 2, \dots, n\}$, define $S(a) = \sum_{i=1}^n c_i a_i$. Prove that there exist permutations $a \neq b$ of $\{1, 2, \dots, n\}$ such that $n!$ is a divisor of $S(a) - S(b)$.

(IMO 2000/4) A magician has one hundred cards numbered 1 to 100. He puts them into three boxes, a red one, a white one and a blue one, so that each box contains at least one card. A member of the audience draws two cards from two different boxes and announces the sum of numbers on those cards. Given this information, the magician locates the box from which no card has been drawn.

How many ways are there to put the cards in the three boxes so that the trick works?

Geometry

- (IMO 2018/1) Let Γ be the circumcircle of acute triangle ABC . Points D and E are on segments AB and AC respectively such that $AD = AE$. The perpendicular bisectors of BD and CE intersect minor arcs AB and AC of Γ at points F and G respectively. Prove that lines DE and FG are either parallel or they are the same line.
- (IMO 2017/4) Let R and S be different points on a circle Ω such that RS is not a diameter. Let ℓ be the tangent line to Ω at R . Point T is such that S is the midpoint of the line segment RT . Point J is chosen on the shorter arc RS of Ω so that the circumcircle Γ of triangle JST intersects ℓ at two distinct points. Let A be the common point of Γ and ℓ that is closer to R . Line AJ meets Ω again at K . Prove that the line KT is tangent to Γ .
- (IMO 2016/1) Triangle BCF has a right angle at B . Let A be the point on line CF such that $FA = FB$ and F lies between A and C . Point D is chosen so that $DA = DC$ and AC is the bisector of $\angle DAB$. Point E is chosen so that $EA = ED$ and AD is the bisector of $\angle EAC$. Let M be the midpoint of CF . Let X be the point such that $AMXE$ is a parallelogram. Prove that BD , FX and ME are concurrent.
- (IMO 2015/4) Triangle ABC has circumcircle Ω and circumcenter O . A circle Γ with center A intersects the segment BC at points D and E , such that B , D , E , and C are all different and lie on line BC in this order. Let F and G be the points of intersection of Γ and Ω , such that A , F , B , C , and G lie on Ω in this order. Let K be the second point of intersection of the circumcircle of triangle BDF and the segment AB . Let L be the second point of intersection of the circumcircle of triangle CGE and the segment CA .
- Suppose that the lines FK and GL are different and intersect at the point X . Prove that X lies on the line AO .
- (IMO 2014/4) Let P and Q be on segment BC of an acute triangle ABC such that $\angle PAB = \angle BCA$ and $\angle CAQ = \angle ABC$. Let M and N be the points on AP and AQ , respectively, such that P is the midpoint of AM and Q is the midpoint of AN . Prove that the intersection of BM and CN is on the circumference of triangle ABC .
- (IMO 2013/4) Let ABC be an acute triangle with orthocenter H , and let W be a point on the side BC , lying strictly between B and C . The points M and N are the feet of the altitudes from B and C , respectively. Denote by ω_1 is the circumcircle of BWN , and let X be the point on ω_1 such that WX is a diameter of ω_1 . Analogously, denote by ω_2 the circumcircle of triangle CWM , and let Y be the point such that WY is a diameter of ω_2 . Prove that X, Y and H are collinear.
- (IMO 2012/1) Given triangle ABC the point J is the centre of the excircle opposite the vertex A . This excircle is tangent to the side BC at M , and to the lines AB and AC at K and L , respectively. The lines LM and BJ meet at F , and the lines KM and CJ meet at G . Let S be the point of intersection of the lines AF and BC , and let T be the point of intersection of the lines AG and BC . Prove that M is the midpoint of ST .
- (The excircle of ABC opposite the vertex A is the circle that is tangent to the line segment BC , to the ray AB beyond B , and to the ray AC beyond C .)

(IMO 2010/4) Let P be a point interior to triangle ABC (with $CA \neq CB$). The lines AP , BP and CP meet again its circumcircle Γ at K , L , respectively M . The tangent line at C to Γ meets the line AB at S . Show that from $SC = SP$ follows $MK = ML$.

(IMO 2009/4) Let ABC be a triangle with $AB = AC$. The angle bisectors of $\angle CAB$ and $\angle ABC$ meet the sides BC and CA at D and E , respectively. Let K be the incentre of triangle ADC . Suppose that $\angle BEK = 45^\circ$. Find all possible values of $\angle CAB$.

(IMO 2009/2) Let ABC be a triangle with circumcentre O . The points P and Q are interior points of the sides CA and AB respectively. Let K, L and M be the midpoints of the segments BP, CQ and PQ , respectively, and let Γ be the circle passing through K, L and M . Suppose that the line PQ is tangent to the circle Γ . Prove that $OP = OQ$.

(IMO 2008/1) Let H be the orthocenter of an acute-angled triangle ABC . The circle Γ_A centered at the midpoint of BC and passing through H intersects the sideline BC at points A_1 and A_2 . Similarly, define the points B_1, B_2, C_1 and C_2 .

Prove that the six points A_1, A_2, B_1, B_2, C_1 and C_2 are concyclic.

(IMO 2007/4) In triangle ABC the bisector of angle BCA intersects the circumcircle again at R , the perpendicular bisector of BC at P , and the perpendicular bisector of AC at Q . The midpoint of BC is K and the midpoint of AC is L . Prove that the triangles RPK and RQL have the same area.

(IMO 2006/1) Let ABC be triangle with incenter I . A point P in the interior of the triangle satisfies

$$\angle PBA + \angle PCA = \angle PBC + \angle PCB.$$

Show that $AP \geq AI$, and that equality holds if and only if $P = I$.

(IMO 2005/1) Six points are chosen on the sides of an equilateral triangle ABC : A_1, A_2 on BC , B_1, B_2 on CA and C_1, C_2 on AB , such that they are the vertices of a convex hexagon $A_1A_2B_1B_2C_1C_2$ with equal side lengths.

Prove that the lines A_1B_2, B_1C_2 and C_1A_2 are concurrent.

(IMO 2004/1) Let ABC be an acute-angled triangle with $AB \neq AC$. The circle with diameter BC intersects the sides AB and AC at M and N respectively. Denote by O the midpoint of the side BC . The bisectors of the angles $\angle BAC$ and $\angle MON$ intersect at R . Prove that the circumcircles of the triangles BMR and CNR have a common point lying on the side BC .

(IMO 2003/4) Let $ABCD$ be a cyclic quadrilateral. Let P, Q, R be the feet of the perpendiculars from D to the lines BC, CA, AB , respectively. Show that $PQ = QR$ if and only if the bisectors of $\angle ABC$ and $\angle ADC$ are concurrent with AC .

(IMO 2001/1) Consider an acute-angled triangle ABC . Let P be the foot of the altitude of triangle ABC issuing from the vertex A , and let O be the circumcenter of triangle ABC . Assume that $\angle C \geq \angle B + 30^\circ$. Prove that $\angle A + \angle COP < 90^\circ$.

(IMO 2000/1) Two circles G_1 and G_2 intersect at two points M and N . Let AB be the line tangent to these circles at A and B , respectively, so that M lies closer to AB than N . Let CD be the line parallel to AB and passing through the point M , with C on G_1 and D on G_2 . Lines AC and BD meet at E ; lines AN and CD meet at P ; lines BN and CD meet at Q . Show that $EP = EQ$.

Number Theory

(IMO 2019/4) Find all pairs (k, n) of positive integers such that

$$k! = (2^n - 1)(2^n - 2)(2^n - 4) \cdots (2^n - 2^{n-1}).$$

(IMO 2017/1) For each integer $a_0 > 1$, define the sequence a_0, a_1, a_2, \dots for $n \geq 0$ as

$$a_{n+1} = \begin{cases} \sqrt{a_n} & \text{if } \sqrt{a_n} \text{ is an integer,} \\ a_n + 3 & \text{otherwise.} \end{cases}$$

Determine all values of a_0 such that there exists a number A such that $a_n = A$ for infinitely many values of n .

(IMO 2016/4) A set of positive integers is called fragrant if it contains at least two elements and each of its elements has a prime factor in common with at least one of the other elements. Let $P(n) = n^2 + n + 1$. What is the least possible positive integer value of b such that there exists a non-negative integer a for which the set

$$\{P(a+1), P(a+2), \dots, P(a+b)\}$$

is fragrant?

(IMO 2013/1) Assume that k and n are two positive integers. Prove that there exist positive integers m_1, \dots, m_k such that

$$1 + \frac{2^k - 1}{n} = \left(1 + \frac{1}{m_1}\right) \cdots \left(1 + \frac{1}{m_k}\right).$$

(IMO 2011/1) Given any set $A = \{a_1, a_2, a_3, a_4\}$ of four distinct positive integers, we denote the sum $a_1 + a_2 + a_3 + a_4$ by s_A . Let n_A denote the number of pairs (i, j) with $1 \leq i < j \leq 4$ for which $a_i + a_j$ divides s_A . Find all sets A of four distinct positive integers which achieve the largest possible value of n_A .

(IMO 2009/1) Let n be a positive integer and let $a_1, a_2, a_3, \dots, a_k$ ($k \geq 2$) be distinct integers in the set $1, 2, \dots, n$ such that n divides $a_i(a_{i+1} - 1)$ for $i = 1, 2, \dots, k-1$. Prove that n does not divide $a_k(a_1 - 1)$.

(IMO 2006/4) Determine all pairs (x, y) of integers such that

$$1 + 2^x + 2^{2x+1} = y^2.$$

(IMO 2005/4) Determine all positive integers relatively prime to all the terms of the infinite sequence

$$a_n = 2^n + 3^n + 6^n - 1, \quad n \geq 1.$$

(IMO 2002/4) Let $n \geq 2$ be a positive integer, with divisors $1 = d_1 < d_2 < \dots < d_k = n$. Prove that $d_1 d_2 + d_2 d_3 + \dots + d_{k-1} d_k$ is always less than n^2 , and determine when it is a divisor of n^2 .