## CHAPTER-9 CIRCLES

## **1 EXERCISE-10.5**

1. A chord of a circle is equal to the radius of the circle. Find the angle subtended by the chord at a point on the minor arc and also at a point on the major arc.

## 2 SOLUTION

The input parameters are

Symbol	Value	Description
r	1	Radius
О	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	circle point
P	$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$	standard basis $vectore_1$
θ	60°	$\angle QOP$
α	130°	$\angle QRP$
β	-40°	$\angle QSP$

Table 1:

Take three points Q,R and P on a unit circle at angles  $\theta, \alpha$ , and  $\beta$ . Then

$$\mathbf{Q} = \begin{pmatrix} \cos\theta \\ \sin\theta \end{pmatrix}, \mathbf{R} = \begin{pmatrix} \cos\alpha \\ \sin\alpha \end{pmatrix}, \mathbf{S} = \begin{pmatrix} \cos\beta \\ \sin\beta \end{pmatrix}$$
(1)

$$\cos \angle QRP = \frac{(\mathbf{Q} - \mathbf{R})(\mathbf{P} - \mathbf{R})}{\|\mathbf{Q} - \mathbf{R}\|\|\mathbf{P} - \mathbf{R}\|}$$
(2)

Where

$$(\mathbf{Q} - \mathbf{R})(\mathbf{P} - \mathbf{R}) = (\cos \theta - \cos \alpha \sin \theta - \sin \alpha)(1 - \cos \alpha \quad o - \sin \alpha)$$
(3)

$$= (\cos \theta - \cos \alpha) \cos \alpha + (\sin \theta - \sin \alpha) \tag{4}$$

$$= 2\sin\frac{\theta - \alpha}{2}\sin\frac{\theta + \alpha}{2}\cos\alpha + 2\cos\frac{\theta + \alpha}{2}\sin\frac{\theta - \alpha}{2}$$
 (5)

$$= (\cos \alpha - \cos \theta) \cos \alpha + (\sin \theta - \sin \alpha) \tag{6}$$

$$\|\mathbf{Q} - \mathbf{R}\|^2 \|\mathbf{P} - \mathbf{R}\|^2 = (\cos \theta - \cos \alpha)^2 + (\sin \theta - \sin \alpha)^2 (1 - \cos \alpha)^2 + (0 - \sin \alpha)^2$$
(7)

$$= (2 - 2\cos\theta\cos\alpha - 2\sin\theta\sin\alpha)(2 - \cos\alpha) \tag{8}$$

substituing the (6) and (8) in (2)

$$\cos\angle QRP = \frac{2.079}{4.323} \tag{9}$$

$$\angle QRP = \cos^{-1} 0.480 \tag{10}$$

$$\angle QRP = 66^{\circ} \tag{11}$$

$$\cos \angle QSP = \frac{(\mathbf{Q} - \mathbf{S})(\mathbf{P} - \mathbf{S})}{\|\mathbf{Q} - \mathbf{S}\|\|\mathbf{P} - \mathbf{S}\|}$$
(12)

$$(\mathbf{Q} - \mathbf{S})(\mathbf{P} - \mathbf{S}) = (\cos \theta - \cos \beta \sin \theta - \sin \beta) (1 - \cos \beta \quad o - \sin \beta)$$
 (13)

$$= (\cos \theta - \cos \beta) \cos \beta + (\sin \theta - \sin \beta) \tag{14}$$

$$= 2\sin\frac{\theta - \beta}{2}\sin\frac{\theta + \beta}{2}\cos\beta + 2\cos\frac{\theta + \beta}{2}\sin\frac{\theta - \beta}{2}$$
 (15)

$$= (\cos \beta - \cos \theta) \cos \beta + (\sin \theta - \sin \beta) \tag{16}$$

$$\|\mathbf{Q} - \mathbf{S}\|^2 \|\mathbf{P} - \mathbf{S}\|^2 = (\cos \theta - \cos \beta)^2 + (\sin \theta - \sin \beta)^2 (1 - \cos \beta)^2 + (0 - \sin \beta)^2)$$
(17)

$$= (2 - 2\cos\theta\cos\beta - 2\sin\theta\sin\beta)(2 - \cos\beta) \tag{18}$$

substituing the (16) and (18) in (12)

$$\cos \angle QSP = \frac{1.048}{1.098} \tag{19}$$

$$\angle QSP = \cos^{-1} 0.954 \tag{20}$$

$$\angle QSP = 17^{\circ} \tag{21}$$

## 3 FIGURE




Figure 1: