


Circle Assignment

Ginna Shreyani- FWC22006

September 2022

1 Problem:


A triangle ABC is drawn to circumscribe a circle of radius 4cm such that the segments BD and DC into which BC is divided by the point of contact D are of lengths 8cm and 6cm respectively. Find the sides AB and AC.



3 Solution:

3.1 Theory:

3.2 Mathematical Calculation:



Let us assume that the center of the circle is origin and point D lies on Y-axis. Therefore, we can get the points B and C.

2 Construction:

Variable	Point/Length	Description
r	4cm	Radius of the given circle
C	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	Origin
c	6cm	Distance from Vertex C to point D
D	$\begin{pmatrix} c \\ 0 \end{pmatrix}$	Point on side BC
O	$\begin{pmatrix} c \\ r \end{pmatrix}$	Center of the circle
CB	$\begin{pmatrix} c+b \\ 0 \end{pmatrix}$	Vertex B

$$D = \begin{pmatrix} 0 \\ -4 \end{pmatrix}$$

$$B = \begin{pmatrix} 8 \\ -4 \end{pmatrix}$$

$$C = \begin{pmatrix} -6 \\ -4 \end{pmatrix}$$

Here point C acts as external point from which tangents can be drawn to the circle. The direction vector \mathbf{m} satisfy the equation

$$\mathbf{m}^T \Sigma \mathbf{m} = 0$$

Assuming the external point as \mathbf{h} , Σ is given as

$$\Sigma = ((\mathbf{V}\mathbf{h} + \mathbf{u})((\mathbf{V}\mathbf{h} + \mathbf{u})^T - ((\mathbf{h}^T \mathbf{V}\mathbf{h} + 2\mathbf{u}^T \mathbf{h} + f) \mathbf{V}$$

Σ can be orthogonally diagonalized as

$$\Sigma = \Gamma^T \mathbf{D} \Gamma \quad (1)$$

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \quad (2)$$

$$\Gamma = \begin{pmatrix} \mathbf{y}_1 & \mathbf{y}_2 \end{pmatrix}, \quad \Gamma^T = \Gamma^{-1} \quad (3)$$

Using (2) and (3) and substituting \mathbf{h} as \mathbf{C} , the normal vector \mathbf{n} of the tangent drawn from \mathbf{C} can be written as

$$\mathbf{n} = \Gamma \begin{pmatrix} \sqrt{|\lambda_1|} \\ \pm \sqrt{|\lambda_2|} \end{pmatrix} \quad (4)$$

Using the vectors \mathbf{n} in (4), the direction vectors \mathbf{m} can be found in 2-dimensional space since they are orthogonal. The points of contact of the tangents are given by

$$\mathbf{T}_i = \mathbf{C} - \frac{\mathbf{m}^T ((\mathbf{V}\mathbf{C} + \mathbf{u}))}{\mathbf{m}^T \mathbf{V} \mathbf{m}}$$

Similarly we can find the contact point from vertex B.

Since, we have all the contact points we can get the line equations of $\mathbf{A} - \mathbf{C}$ and $\mathbf{A} - \mathbf{B}$

$$\mathbf{n}^T \mathbf{X} - \mathbf{T}_i = 0$$

The intersection of these two lines is vertex A. Since, we know all the vertices of the given triangle the length of AB and AC can be calculated. Therefore, length of AB is $\|\mathbf{A} - \mathbf{B}\|$ and length of AC is $\|\mathbf{A} - \mathbf{C}\|$