

Introduction to Symplectic Geometry

Sachchidanand Prasad

Contents

1	Introduction	2
1.1	An overview of geometry	2
2	Symplectic algebra	3
2.1	Some basic definitions	3
2.2	Symplectic vector space	4
2.3	Symplectomorphism	8
2.4	Subspaces of a symplectic vector space	9
3	Compatible Complex Structures	13

Course information:

- **Course name** : Introduction to Symplectic Geometry
- **Instructor** : Sachchidanand Prasad
- **Time** : Wednesday 17:30 - 19:00
- **Course webpage** : [Link to the course website](#)
- **References** :

- [1] *Lectures on Symplectic Geometry*, by Ana Cannas da Silva.
- [2] *Introduction to Symplectic Topology*, by Dusa McDuff and Dietmar Salamon.
- [3] *Lectures on Symplectic Manifolds*, by Alan Weinstein.
- [4] *Symplectic Techniques in Physics*, by Victor Guillemin and Shlomo Sternberg.

1 Introduction

The word *symplectic* was invented by Hermann Weyl in 1939. He replaced the Latin roots in the word *complex*, *com-plexus*, by the corresponding Greek roots *sym-plektikos*.

1.1 An overview of geometry

- **Geometry:** Background Space (smooth manifold) + extra structure (tensor)
 - Riemannian geometry: smooth manifold + **metric structure**
 - **metric structure** = positive-definite symmetric 2-tensor
 - Complex geometry: smooth manifold + **complex structure**
 - **complex structure** = involutive endomorphism $((1,1)$ -tensor)
 - Symplectic geometry: smooth manifold + **symplectic structure**
 - **symplectic structure** = closed non-degenerate 2-form
 - Contact geometry: smooth manifold + **contact structure**
 - **contact structure** = “local contact 1-form”

In both symplectic and Riemannian geometry the main object of study is a smooth manifold equipped with a bilinear form on each tangent space. In the Riemannian manifold, this form is a symmetric, nondegenerate, positive definite form, turning each tangent space into normed vector space. On the other hand, in symplectic geometry, we instead require a skew-symmetric bilinear form on each tangent space, again varying smoothly. We still require that at each point p in our manifold M , a skew-symmetric 2-form ω_p should be nondegenerate, that is,

$$\omega_p(X, Y) = 0 \quad \forall Y \in T_p M, \text{ then } X \equiv 0.$$

Finally, note that because ω is a skew-symmetric 2-form, it must be closed, that is, $d\omega = 0$. We will now compare both geometry and from next lecture onwards we will discuss in more details. We will use the following notations:

- M : real finite dimensional smooth manifold without boundary.
- $C^\infty(M) = \{f : M \rightarrow \mathbb{R} : f \text{ is smooth}\}.$
- $\chi(M) = \{X : M \rightarrow TM : X \text{ is a vector field}\}.$
- $\Omega^k(M) = \{\omega : TM \times TM \times \cdots \times TM \rightarrow \mathbb{R}\}.$

We now start some comparison between Riemannian geometry and symplectic geometry:

(1a) **Riemannian manifold is a pair $(M, \langle \cdot, \cdot \rangle)$, where**

- $\langle \cdot, \cdot \rangle : \chi(M) \times \chi(M) \rightarrow C^\infty(M)$ satisfies $\langle X, Y \rangle = \langle Y, X \rangle$ and $\langle fX + gY, Z \rangle = f \langle X, Z \rangle + g \langle Y, Z \rangle.$
- $\langle \cdot, \cdot \rangle$ is positive definite.

(2a) **Symplectic manifold is a pair (M, ω) where**

- $\omega \in \Omega^2(M)$ is bilinear.
 - ω is nondegenerate.
 - ω is closed, that is $d\omega = 0$.
-

(1b) Every smooth manifold is a Riemannian manifold.

(2b) Not all manifolds are Symplectic. The necessary conditions are:

- $\dim M = \text{even}$.
 - M is oriented.
 - If M is compact, then $H_{dR}^2(M, \mathbb{R}) \neq 0$.
-

(1c) **Isometry**: Two Riemannian manifolds $(M_1, \langle \cdot, \cdot \rangle_1)$ and $(M_2, \langle \cdot, \cdot \rangle_2)$ are isometric if there exists a C^1 map $\varphi : M_1 \rightarrow M_2$ such that

$$\left\langle d\varphi_p(X), d\varphi_p(Y) \right\rangle_{\varphi(p)} = \langle X, Y \rangle_p$$

(2c) Similarly, we have **symplectomorphism** between two symplectic manifolds.

(1d) **Curvature** is a local invariant in Riemannian manifolds.

(2d) There are no local invariants (apart from dimension) in symplectic manifolds. According to the **Darboux-Weinstein theorem**, given any two symplectic manifolds of the same finite dimension, they look alike locally.

2 Symplectic algebra

In this lecture, we will mostly recall the linear algebra preliminaries for our course. More precisely, we will deal with linear symplectic algebra which we will be using through out the course.

2.1 Some basic definitions

Definition 2.1 (Vector sapce). A set $(V, +, \cdot)$ is said to be a **vector space** over a field \mathbb{F} if the operations

$$+ : V \times V \rightarrow V \text{ and } \cdot : \mathbb{F} \times V \rightarrow V,$$

satisfies the following properties. For any v, v_1, v_2, v_3 and $\alpha, \beta \in \mathbb{F}$ we have the following.

1. (Commutativity) $v_1 + v_2 = v_2 + v_1$.
2. (Associativity) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$.
3. (Existence of additive identity) There exists $0 \in V$ such that for any $v \in V$ $0 + v = v = v + 0$.

4. (Existence of additive inverse) For any $v \in V$, there exists w such that $v + w = 0 = w + v$. We will denote $w = -v$.
5. (Multiplicative identity) For any $v \in V$, $1 \cdot v = v$.
6. (Multiplication associativity) $(\alpha\beta) \cdot v = \alpha \cdot (\beta \cdot v)$.
7. (Distribution law)
 - $(\alpha + \beta) \cdot v = \alpha \cdot v + \beta \cdot v$.
 - $\alpha(v_1 + v_2) = \alpha \cdot v_1 + \alpha \cdot v_2$.

Our field will always be either \mathbb{R} or \mathbb{C} .

Definition 2.2 (Linear map). Let $T : V \rightarrow W$ be a map between two vector spaces V and W . Then T is said to be **linear** if,

$$T(\alpha v_1 + \beta v_2) = \alpha T(v_1) + \beta T(v_2),$$

for $v_1, v_2 \in V$ and $\alpha, \beta \in \mathbb{F}$.

Definition 2.3 (Dual space). If V is a vector space over a field \mathbb{F} . Then the **dual space** of V , denoted by V^* , is defined by

$$V^* := \{\varphi : V \rightarrow \mathbb{F} : \varphi \text{ is linear}\}.$$

Definition 2.4 (Bilinear map). Let V, W, S be vector spaces over a field \mathbb{F} . The a **bilinear map** B is a map

$$B : V \times W \rightarrow S$$

such that B is linear in each argument. That is, $B(\cdot, w) : V \rightarrow S$ and $B(v, \cdot) : W \rightarrow S$ is linear for any $v \in V$ and $w \in W$.

Definition 2.5. A **bilinear form** ω on a vector space V is a bilinear map $B : V \times V \rightarrow \mathbb{F}$. The bilinear form ω is said to be **nondegenerate** if the kernel

$$\ker \omega := \{v \in V : \omega(v, w) = 0 \text{ for all } w \in V\}$$

is trivial.

We identify a bilinear form ω on E with the linear mapping $u \mapsto (v \mapsto \omega(u, v))$ for $u, v \in E$.

Definition 2.6. Let ω be a bilinear form on a vector space V .

1. ω is said to be **symmetric** if $\omega(v, w) = \omega(w, v)$ for any $v, w \in V$.
2. ω is said to be **skew-symmetric** if $\omega(v, w) = -\omega(w, v)$ for any $v, w \in V$.

2.2 Symplectic vector space

The first important notions that we introduce are the symplectic form and the symplectic vector space. We also define the concept of canonical form of a symplectic form and the symplectic basis of a symplectic vector space. Throughout this notes, we will assume V to be a vector space of finite dimension.

Definition 2.7. The pair (V, ω) is said to be **symplectic vector space** if $\omega : V \times V \rightarrow \mathbb{R}$ is skew-symmetric, nondegenerate bilinear form. We call ω a **symplectic form** on E .

Remark. It follows from the definition that $\omega(v, v) = 0$ for any $v \in V$.

Example 2.8. On $\mathbb{R}^{2n} = \mathbb{R}^n \times \mathbb{R}^n$ we define ω by

$$\omega((\mathbf{x}, \mathbf{y}), (\mathbf{x}', \mathbf{y}')) := \sum_{i=1}^n (x_i y'_i - x'_i y_i) = \langle \mathbf{x}, \mathbf{y}' \rangle - \langle \mathbf{x}', \mathbf{y} \rangle.$$

We claim that ω is a symplectic form on \mathbb{R}^{2n} . It is clear that ω is a bilinear form. Further, we need to check two things:

(i) ω is skew-symmetric.

For any $(\mathbf{a}, \mathbf{b}), (\mathbf{c}, \mathbf{d}) \in \mathbb{R}^n \times \mathbb{R}^n$, we have

$$\begin{aligned}\omega((\mathbf{c}, \mathbf{d}), (\mathbf{a}, \mathbf{b})) &= \langle \mathbf{b}, \mathbf{c} \rangle - \langle \mathbf{a}, \mathbf{d} \rangle, \text{ and} \\ \omega((\mathbf{a}, \mathbf{b}), (\mathbf{c}, \mathbf{d})) &= \langle \mathbf{a}, \mathbf{d} \rangle - \langle \mathbf{b}, \mathbf{c} \rangle = -\omega((\mathbf{c}, \mathbf{d}), (\mathbf{a}, \mathbf{b})).\end{aligned}$$

(ii) ω is nondegenerate.

Let $\omega((\mathbf{x}, \mathbf{y}), (\mathbf{a}, \mathbf{b})) = 0$ for any $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^n \times \mathbb{R}^n$. We need to show that $(\mathbf{x}, \mathbf{y}) = (\mathbf{0}, \mathbf{0})$. Take $\mathbf{a} = \mathbf{0}$ and $\mathbf{b} = \mathbf{x}$, then

$$\omega((\mathbf{x}, \mathbf{y}), (\mathbf{0}, \mathbf{x})) = 0 \implies \langle \mathbf{x}, \mathbf{x} \rangle - \langle \mathbf{y}, \mathbf{0} \rangle = 0 \implies \mathbf{x} = \mathbf{0}.$$

Similarly, one can show that $\mathbf{y} = \mathbf{0}$ and hence, ω is nondegenerate.

This is called standard symplectic form on $\mathbb{R}^n \times \mathbb{R}^n$.

The above example can also be written in the following form:

Example 2.9. Let $V = \mathbb{R}^{2n}$ with a basis $\{e_1, e_2, \dots, e_n, f_1, f_2, \dots, f_n\}$ and define ω as

$$\omega(e_i, e_j) = 0, \quad \omega(f_i, f_j) = 0 \quad \text{and} \quad \omega(e_i, f_j) = \delta_{ij}.$$

Then ω is standard symplectic form on V .

Example 2.10. Let V be any vector space of dimension n and V^* denotes its dual. If $E = V \oplus V^*$ and define

$$\omega : E \times E \rightarrow \mathbb{R}, \quad \omega((v, \alpha), (v', \alpha')) = \alpha'(v) - \alpha(v'),$$

then (E, ω) is a symplectic vector space.

Since α and α' are linear maps, it is clear that ω is a bilinear form. Let us show it is skew-symmetric and nondegenerate.

(i) ω is skew-symmetric.

For any $v, v' \in V$ and $\alpha, \alpha' \in V^*$, we have

$$\begin{aligned}\omega((v, \alpha), (v', \alpha')) &= \alpha'(v) - \alpha(v') \\ &= -(\alpha(v') - \alpha'(v)) \\ &= \omega((v', \alpha'), (v, \alpha)).\end{aligned}$$

(ii) ω is nondegenerate.

Let $\omega((v, \alpha), (w, \beta)) = 0$ for any $(w, \beta) \in E$. We need to show that $v = 0$ and $\alpha \equiv 0$. Observe that for any $\beta \in V^*$

$$\omega((v, \alpha), (0, \beta)) = \beta(v) = 0 \implies v = 0.$$

Similarly, for any $w \in V$,

$$\omega((v, \alpha), (w, 0)) = \alpha(w) = 0 \implies \alpha = 0.$$

Thus (E, ω) is a symplectic vector space.

Definition 2.11. Let (V, ω) is a symplectic vector space, then for any subspace $W \subseteq V$, we define the ω -orthogonal space

$$W^\omega := \{v \in V : \omega(v, w) = 0, \forall w \in W\}.$$

Proposition 2.12. Let V be a k -dimensional vector space over \mathbb{R} and ω be a bilinear form.

1. If ω is symmetric with rank r , then there exists a basis \mathcal{B} of V such that with respect to \mathcal{B} ,

$$[\omega]_{\mathcal{B}} = \begin{bmatrix} \epsilon_1 & & & & \\ & \ddots & & & \\ & & \epsilon_r & & \\ & & & 0 & \\ & & & & \ddots \\ & & & & & 0 \end{bmatrix}, \text{ where } \epsilon_i = \pm 1, \quad i = 1, 2, \dots, r.$$

2. If ω is skew-symmetric with rank r , then $r = 2n$ and there is a basis \mathcal{B} of V relative to which

$$[\omega]_{\mathcal{B}} = \begin{bmatrix} 0 & I_n & 0 \\ -I_n & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{ where } I_n \text{ is the identity matrix of size } n.$$

Proof. 1. Proof is left.

2. Since $\omega \neq 0$, we can choose $e_1, f_1 \in V$ such that $\omega(e_1, f_1) \neq 0$ (this must implies that both the vectors are linearly independent). By rescaling e_1 , we can further assume that $\omega(e_1, f_1) = 1$. Define $W_1 := \text{span}\{e_1, f_1\}$. Since, ω is skew-symmetric, we have $\omega(e_1, e_1) = 0 = \omega(f_1, f_1)$. Thus, the restriction of ω on W_1 is

$$[\omega]_{\{e_1, f_1\}} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Let W_2 be the ω -orthogonal complement of W_1 , that is, $W_2 = W_1^\omega$. It is clear that $W_1 \cap W_2 = \{0\}$. We claim that $V = W_1 \oplus W_2$. Note that for any $v \in V$, we have

$$\begin{aligned} \omega(e_1, v - \omega(v, f_1)e_1 + \omega(v, e_1)f_1) &= 0 \text{ and} \\ \omega(f_1, v - \omega(v, f_1)e_1 + \omega(v, e_1)f_1) &= 0. \end{aligned}$$

Thus, $v - \omega(v, f_1)e_1 + \omega(v, e_1)f_1 \in W_2^\omega$ and hence $V = W_1 \oplus W_2$. We can repeat the process on W_2 and find e_2 and f_2 such that $\omega(e_2, f_2) = 1$. Now the matrix will be

$$[\omega]_{\{e_1, e_2, f_1, f_2\}} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}$$

Inductively, we get a basis

$$\mathcal{B} = \{e_1, e_2, \dots, e_n, f_1, f_2, \dots, f_n\}$$

such that $[\omega]_{\mathcal{B}}$ will be in the given form.

□

Remark. Since we focus on non-degenerate skew-symmetric bilinear form, that is, $\text{rank} = 2n$, we may consider only the case with matrix representation $\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$ and V must be of even dimension.

We just showed the following corollary.

Corollary 2.13. *Every finite dimensional symplectic vector space (V, ω) has even dimension.*

Exercise 2.14. Show that the space of skew-symmetric bilinear form is isomorphic to the space $\wedge^2 V^*$ of the second exterior product of V^* .

So if $\mathcal{B} = \{e_1, \dots, e_{2n}\}$ is a basis for V , and \mathcal{B}^* is its dual, then for any $\omega \in \wedge^2 V$ with the matrix (ω_{ij}) , relative to \mathcal{B} , can also be written as

$$[\omega]_{\mathcal{B}} = \sum_{i < j} \omega_{ij} e_i^* \wedge e_j^*.$$

Remark. Since elements of $\wedge^2 V^*$ are represented by anti-symmetric matrices and with all the entries of the main diagonal equal to 0, for a vector space V of dimension $2n$ we have $\dim \wedge^2 V^* = \frac{2n(2n-1)}{2} = n(2n-1)$.

Corollary 2.15 (Canonical form of ω). *For every skew-symmetric bilinear form ω , there exists a basis $\mathcal{B} = \{e_1, \dots, e_{2n}\}$ of V such that*

$$[\omega]_{\mathcal{B}} = \sum_{i < j} e_i^* \wedge e_j^*.$$

This representation is called a canonical form of ω and we call \mathcal{B} a symplectic basis of V .

2.3 Symplectomorphism

Definition 2.16. Two symplectic vector spaces (V_1, ω_1) and (V_2, ω_2) are called **symplectomorphic** if there exists an isomorphism $\varphi : V_1 \rightarrow V_2$ of vector space such that $\omega_2(\varphi(x), \varphi(y)) = \omega_1(x, y)$. In other words, $\varphi^* \omega_2 = \omega_1$. We call φ a **symplectomorphism**. We will write $V_1 \cong V_2$.

Exercise 2.17. What can you conclude if $\dim V_1 = \dim V_2$ and $\varphi : V_1 \rightarrow V_2$ satisfies $\varphi^* \omega_2 = \omega_1$?

We claim that φ is injective. If $v \in \ker \varphi$, then for any $v' \in V_1$, we have

$$\omega_1(v, v') = \omega_2(\varphi(v), \varphi(v')) = 0.$$

Since, ω_1 is nondegenerate, $v = 0$. Since the dimension matches, $V_1 \cong V_2$.

Exercise 2.18. Show that the set of all symplectomorphisms of a symplectic vector space (V, ω) forms a group under the composition.

Definition 2.19. The group of symplectomorphism of a symplectic vector space (V, ω) is called **symplectic group** and we will denote this by $\text{Sp}(V)$.

Example 2.20. Some examples on symplectomorphism:

1. $V = \mathbb{R}^{2n}$ and $\omega((\mathbf{a}, \mathbf{b}), (\mathbf{c}, \mathbf{d})) = \langle \mathbf{a}, \mathbf{d} \rangle - \langle \mathbf{c}, \mathbf{b} \rangle$.

- $\varphi(e_i) = f_i$ and $\varphi(f_i) = -e_i$. If we change $\varphi(f_i) = e_i$, will it work?

It is clear that φ is an isomorphism. We just need to show that $\varphi^* \omega = \omega$. Note that

$$\omega(e_i, e_j) = 0 = \omega(f_i, f_j).$$

$$\delta_{ij} = \omega(\varphi(e_i), \varphi(f_j)) = \omega(f_i, -e_j) = -\omega(f_i, e_j) = \omega(e_j, f_i) = \delta_{ij}.$$

- $\varphi(e_i) = e_i + f_i$ and $\varphi(f_i) = f_i$.
- For any invertible matrix X ,

$$\varphi(e_i) = \sum_j X_{ij} e_j \text{ and } \varphi(f_i) = \sum_j (X^{-1})_{ji} f_j.$$

2. We show in **Example 2.10** $E = V \oplus V^*$ is a symplectic vector space. We can give a symplectomorphism on E as follows. Let $T : V \rightarrow V$ be an isomorphism and $T^* : V^* \rightarrow V^*$ be the dual map. Then

$$T \oplus T^* : E \rightarrow E,$$

is a symplectomorphism.

3. Let V be a complex vector space of complex dimension n , with complex, positive definite inner product (=Hermitian metric) $h : V \times V \rightarrow \mathbb{C}$. Then V , viewed as a real vector space, with bilinear form the imaginary part $\omega = \text{Im}(h)$ is a symplectic vector space. Every unitary map $V \rightarrow V$ preserves h , hence also ω and is therefore symplectic.

Exercise 2.21. Show that \mathbb{R}^{2n}, E and the third example are symplectomorphic.

Proposition 2.22. Every symplectic vector space (V, ω) of dimension $2n$ is symplectomorphic to \mathbb{R}^{2n} with the canonical symplectic form.

As a consequence of the above proposition, we have the following theorem, which we call *Linear Darboux theorem*.

Theorem 2.23 (Linear Darboux Theorem). *For any symplectic vector space (V, ω) there exists a basis $\mathcal{B} = \{e_i, f_i\}_{i=1}^n$ of V such that*

$$\omega(e_i, e_j) = 0 = \omega(f_i, f_j) \text{ and } \omega(e_i, f_j) = \delta_{ij} \quad \forall i, j.$$

This basis is called a Darboux basis of V .

The above theorem is equivalent to following statements:

- (i) Any symplectic vector space is even-dimensional.
- (ii) Any even dimensional vector space admits a linear symplectic form.
- (iii) Up to linear symplectomorphisms, there is a unique linear symplectic form on each even dimensional vector space.

2.4 Subspaces of a symplectic vector space

Recall the **Definition 2.11** of ω -perpendicular space. Note that with our assumption that V is finite dimensional, ω is nondegenerate if and only if the map

$$\omega^b : V \rightarrow V^*, \quad \omega^b(v)(w) = \omega(v, w) \quad \forall v, w \in V$$

is an isomorphism.

Note. For any subspace $W \subset V$, we have

$$W^\omega = \left(\omega^b\right)^{-1}(\text{ann}(W)),$$

where $\text{ann}(W)$ is the annihilator of W , that is, the set of all $f \in V^*$ such that $f(w) = 0$ for $w \in W$.

We have

$$v \in \text{ann}(W) \iff \text{for any } w \in W, \left(\omega^b(v)\right)(w) = 0 \iff \omega(v, w) = 0 \iff v \in W^\omega.$$

Definition 2.24. *A subspace $W \subseteq V$ of a symplectic vector space is called*

- (i) **isotropic** if $W \subseteq W^\omega$, that is, $\omega|_{W \times W} = 0$;
- (ii) **co-isotropic** if $W^\omega \subseteq W$, that is, W^ω is isotropic;
- (iii) **Lagrangian** if $W^\omega = W$, that is, W is isotropic and co-isotropic;
- (iv) **symplectic** if $\omega_{W \times W}$ is nondegenerate.

The set of Lagrangian subspaces of V is called the **Lagrangian Grassmannian** and denoted $\text{Lag}(V)$.

Exercise 2.25. Show that W is symplectic if and only if $W \cap W^\omega = \{0\}$.

Exercise 2.26. Let (V, ω) is a symplectic vector space and W be any subspace of V . Consider the map $\varphi : V \rightarrow W^*$, defined by $\varphi(v) = \omega(v)|_W$ for any $v \in V$ and $w \in W$. Show that φ is surjective. Deduce that $\dim W^\omega = \dim V - \dim W$. Also, show that $(W^\omega)^\omega = W$.

Remark. 1. From **Exercise 2.26**, we conclude that if $\dim V = 2n$, then all the isotropic subspaces have dimension smaller or equal n , all the co-isotropic have dimension bigger or equal n and all the Lagrangian subspace have dimension n .

- 2. If $W \subseteq V$ is symplectic subspace, then it follows from the definition that $W \cap W^\omega = \{0\}$ and therefore, from the dimension sum restriction (**Exercise 2.26**), we must have $V = W \oplus W^\omega$.

Example 2.27. Every 1-dimensional subspace of V is isotropic and every subspace with codimension 1 is co-isotropic.

Example 2.28. Consider $V = \mathbb{R}^{2n}$ with canonical symplectic form ω . Define

$$\begin{aligned} W_1 &= \text{span}\{e_1, e_2\}. && \text{Isotropic} \\ W_2 &= \text{span}\{e_1, e_2, \dots, e_n, f_3, f_4, \dots, f_n\}. && \text{Co-isotropic} \\ W_3 &= \text{span}\{e_1, e_2, \dots, e_n\}. && \text{Lagrangian} \\ W_4 &= \text{span}\{e_1, f_1\}. && \text{Symplectic} \end{aligned}$$

Exercise 2.29. Let (V, ω) be a symplectic vector space and W be any subspace of V .

1. Show that if W is isotropic, then $\dim W \leq \frac{1}{2} \dim V$.
2. Show that if W is Lagrangian, then $\dim W = \frac{1}{2} \dim V$.
3. Show that if W is Lagrangian, then any basis $\mathcal{B}_W = \{e_1, e_2, \dots, e_n\}$ of W can be extended to a symplectic basis $\{e_1, \dots, e_n, f_1, \dots, f_n\}$ of V .

Proposition 2.30. *For any symplectic vector space (V, ω) , there exists a Lagrangian subspace L .*

Proof. Since for every $v \in V$ we have $\omega(v, v) = 0$, V has an isotropic subspace. Let $L \subseteq V$ be a maximal isotropic subspace of V , that is, it is not contained in any isotropic subspace of strictly larger dimension. Then we claim that L is Lagrangian, that is, $L^\omega = L$. We only need to show that L is co-isotropic. Suppose not, take $v \in L^\omega \setminus L$, then $L' = L \oplus \text{span}\{v\}$ is isotropic and larger than L . \square

An immediate consequence is that any symplectic vector space V has even dimension: For if L is a Lagrangian subspace

$$\dim V = \dim L + \dim L^\omega = 2 \dim L.$$

From this proof we also conclude that a maximal isotropic subspace is a Lagrangian subspace. Therefore we have the following corollary.

Corollary 2.31. *Every isotropic subspace is contained in a Lagrangian subspace.*

Some properties: Let W, W_1, W_2 be subspaces of a symplectic vector space (V, ω) .

(1) $\dim W + \dim W^\omega = \dim V$.

As per the hint given in [Exercise 2.26](#), consider the map

$$\varphi : V \rightarrow W^*, \quad \varphi(v) = \omega(v, \cdot).$$

Note that

$$\ker \varphi = \{v \in V : \varphi(v) = 0\} = \{v \in V : \omega(v, w) = 0 \forall w \in W\} = W^\omega.$$

Now, we claim that φ is surjective. Let $f \in W^*$, that is, $f : W \rightarrow \mathbb{R}$ is linear. As $W \subseteq V$, we can extend the map f to V , say \tilde{f} . Since $\tilde{\varphi} : V \rightarrow V^*, v \mapsto \omega(v, \cdot)$ is an isomorphism, there exists $v \in V$ such that $\tilde{\varphi}(v) = \tilde{f}$. Thus, $\varphi(v) = f$. So, $\text{Image}(\varphi) = W^*$. Thus, using rank-nullity theorem, we have

$$\dim V = \dim W^* + \dim W^\omega = \dim W + \dim W^\omega.$$

(2) $(W^\omega)^\omega = W$.

Note that for any $w \in W$,

$$\omega(w, v) = 0, \forall v \in W^\omega \implies w \in (W^\omega)^\omega \implies W \subseteq (W^\omega)^\omega.$$

Using the dimension formula, we have

$$\begin{aligned} \dim W^\omega + \dim (W^\omega)^\omega &= \dim V = \dim W + \dim W^\omega \\ \implies \dim W &= \dim (W^\omega)^\omega \implies W = (W^\omega)^\omega. \end{aligned}$$

(3) If $W_1 \subseteq W_2$, then $W_2^\omega \subseteq W_1^\omega$.

Let $w'_2 \in W_2^\omega$. For any $w_2 \in W_2$, $\omega(w_2, w'_2) = 0$. For any $w_1 \in W_1 \subseteq W_2$, we must have

$$\omega(w'_2, w_1) = 0 \implies w_2 \in W_1^\omega.$$

(4) If W is symplectic, then $W \oplus W^\omega = V$.

We have

$$\dim(W + W^\omega) = \dim W + \dim W^\omega - \dim(W \cap W^\omega) \implies W + W^\omega = V.$$

(5) Every 1-dimensional subspace is isotropic.

(6) Every codimensional 1 subspace is co-isotropic.

(7) If W is Lagrangian, then $\dim W = \frac{1}{2} \dim V$.

(8) If W is Lagrangian, then any basis $\{e_1, e_2, \dots, e_n\}$ of W can be extended to a symplectic basis $\{e_1, \dots, e_n, f_1, \dots, f_n\}$.

(9) If W is Lagrangian, then (V, ω) is symplectomorphic to the space $(W \oplus W^*, \Omega)$, where $\Omega(x \oplus \alpha, y \oplus \beta) = \beta(x) - \alpha(y)$.

Example 2.32 (Quotient Space). Let (V, ω) is a symplectic vector space and W be any isotropic subspace of V . We define $V/W := \{[v] = v + W : v \in V\}$. Then there is a natural symplectic form on W^ω/W .

Define

$$\Omega : W^\omega / W \times W^\omega / W \rightarrow \mathbb{R}, \quad \Omega([v], [v']) := \omega(v, v').$$

We need to verify that Ω is well-defined and is a symplectic form on W^ω / W .

(i) **Well-defined:**

For any $v, v' \in W^\omega$ and $w, w' \in W$ we have

$$\omega(v + w, v' + w') = \omega(v, v') + \underbrace{\omega(v, w')}_0 + \underbrace{\omega(w, v')}_0 + \underbrace{\omega(w, w')}_0 = \omega(v, v').$$

Here the middle terms vanish because of orthogonality and last term vanishes because W is isotropic.

(ii) **Bilinear:** Exercise.

(iii) **Skew-symmetric:** Exercise.

(iv) **Non-degenerate**

For any $v, v' \in V$, let

$$\Omega([v], [v']) = 0 \implies \omega(v, v') = 0 \implies v = 0.$$

Proposition 2.33. Given any finite collection of Lagrangian subspaces L_1, L_2, \dots, L_k , of a symplectic there exists a Lagrangian subspace L with $L \cap L_i = \{0\}$ for all $i = 1, 2, \dots, k$.

Proof. We will write the proofs in steps:

1. **Step 1 :** Choose a maximal isotropic subspace L of V such that $L \cap L_i = \{0\}$.

We can choose such an isotropic subspace because **finite union of proper subspace can not be the full space**. Since L_i 's are Lagrangian, so they are proper subspace of V and hence $\bigcup_i L_i \subsetneq V$. Choose a $v \in V$ which is not in any L_i . Then $\text{span}\{v\}$ is an isotropic subspace of V such that intersection with any L_i is trivial.

2. **Step 2 :** We will show that L is Lagrangian. Suppose it is false, that is, $L^\omega \subsetneq L$. From **Example 2.32**, we know that L^ω / L has a symplectic form. Let $\pi : L^\omega \rightarrow L^\omega / L$ be the quotient map. Then we have the following claims.

(a) For each i , the space $\pi(L_i \cap L^\omega)$ is isotropic.

Let Ω as defined in **Example 2.32**. Then we need to show that $\Omega|_{\pi(L_i \cap L^\omega)} = 0$. This is easy as for any $[v], [v'] \in \pi(L_i \cap L^\omega)$

$$\Omega([v], [v']) = \omega(v, v') = 0.$$

The last equality is because $v, v' \in L_i$ which is Lagrangian, in particular it is isotropic.

(b) There exists a one dimensional space $F \subseteq L^\omega / L$ such that F is transversal to each of $\pi(L_i \cap L^\omega)$.

Similar to the step 1, we can choose an element $[v] \in L^\omega / L$ away from each of $\pi(L_i \cap L^\omega)$. Define $F = \text{span}\{[v]\}$. It is clear that $F \cap (\pi(L_i \cap L^\omega)) = \{0\}$, and hence they are transversal.

Now we note that

- $L' := \pi^{-1}(F)$ is isotropic subspace of V ;
- $L \subsetneq L'$ and
- $L' \cap L_i = \{0\}$ for each i .

Combing all this, we get a contradiction to the choice of L . Thus, L is Lagrangian. □

Remark. As a consequence of [Proposition 2.33](#), one can give an alternative proof of the Linear Darboux Theorem ([Theorem 2.23](#)).

Theorem 2.34. *Every symplectic vector space (V, ω) of dimension $2n$ is symplectomorphic to \mathbb{R}^{2n} with the canonical symplectic form.*

Proof. Use [Proposition 2.30](#), let L_1 be a Lagrangian subspace of V . Now, using [Proposition 2.33](#), choose a Lagrangian subspace L_2 which is transversal to L_1 . Then, the map

$$L_1 \times L_2 \rightarrow \mathbb{R}, \quad (l_1, l_2) \mapsto \omega(l_1, l_2).$$

is nondegenerate. This gives an isomorphism between L_1 and L_2^* as shown by the composition

$$\psi : L_1 \xhookrightarrow{i} V \xrightarrow{\omega^b} V^* \xrightarrow{i^*} L_2^*.$$

Note that

$$\begin{aligned} \ker(\psi) &= \{l_1 \in L_1 : \psi(l_1) = i^*(\omega^b(i(l_1))) = 0\} \\ &= \{l_1 \in L_1 : i^*(\omega(l_1, v)) = 0, \quad v \in V\} \\ &= \{0\}. \end{aligned}$$

Now, let $\{e_1, e_2, \dots, e_n\}$ be a basis for L_1 and f_1, f_2, \dots, f_n be the dual basis for L_2^* . Since L_1 and L_2 are transversal to each other, $\{e_1, e_2, \dots, e_n, f_1, f_2, \dots, f_n\}$ is a symplectic basis for V . □

Corollary 2.35. *Let (V_1, ω_1) and (V_2, ω_2) be two symplectic vector spaces of same dimension. Let $L_1, L'_1 \subseteq V_1, L_2, L'_2 \subseteq V_2$ be Lagrangian subspace such that $L_1 \cap L'_1$ and $L_2 \cap L'_2$ are trivial. Then there exists a symplectomorphism $\varphi : V_1 \rightarrow V_2$ such that $\varphi(L_1) = L_2$ and $\varphi(L'_1) = L'_2$.*

3 Compatible Complex Structures

Definition 3.1. *Let V be a real vector space. Then an automorphism $J : V \rightarrow V$ is said to be a **complex structure** on V if $J^2 = -Id$, that is, $J(J(v)) = -v$ for $v \in V$. We will write this as (V, \mathbb{R}, J) .*

Example 3.2. Take $V = \mathbb{R}$. If V has a complex structure J , then $J^2 = -Id$. Let $J(1) = \alpha$.

$$J(J(1)) = -1 \implies \alpha^2 = -1,$$

a contradiction as $\alpha \in \mathbb{R}$. So, on \mathbb{R} , there is no complex structure.

Example 3.3. Tak $V = \mathbb{R}^2$. Define J as

$$J(e_1) = e_2 \text{ and } J(e_2) = -e_1.$$

Then it is easy to see that J is a complex structure on \mathbb{R}^2 .

Example 3.4. Note that if n is odd, then \mathbb{R}^n does not admit a complex structure. As if there is a complex structure J , then we must have

$$J^2 = -Id \implies \det(J^2) = (-1)^n = -1 \implies (\det J)^2 = -1,$$

not possible.

The real vector space \mathbb{R}^n admits a complex structure if and only if n is even. Let $n = 2k$. If $\{e_1, \dots, e_k, f_1, \dots, f_k\}$ is a basis of \mathbb{R}^n , then define $J : \mathbb{R}^{2k} \rightarrow \mathbb{R}^{2k}$, as

$$J(e_i) = f_i \text{ and } J(f_i) = -e_i.$$

Then J is a complex structure on \mathbb{R}^n . On the other hand, if J exists, then

$$\det(J^2) = (-1)^n \implies (\det J)^2 = (-1)^n \implies n \text{ is even.}$$

Theorem 3.5. Every real vector space (V, \mathbb{R}, J) with a complex structure J is even dimensional.

Corollary 3.6. Every symplectic vector space admits a complex structure.

Definition 3.7. A complex structure J on a symplectic vector space (V, ω) is said to be ω -compatible, if

$$G_J(v, w) = \omega(v, J(w))$$

is an inner product.

Remark. We note that J is ω -compatible implies

$$\omega(Jv_1, Jv_2) = \omega(v_1, v_2).$$

By using the properties of g , we have

$$\omega(Jv_1, Jv_2) = g(Jv_1, v_2) = g(v_2, Jv_1) = \omega(v_2, J^2v_1) = \omega(v_2, -v_1) = \omega(v_1, v_2).$$

Indeed,

$$J \text{ is } \omega\text{-compatible} \iff \begin{cases} \omega(Jv_1, Jv_2) = \omega(v_1, v_2) \\ \omega(v, Jv) > 0, \quad \forall v \neq 0. \end{cases}$$

Thus, following the above remark, we get if J is ω -compatible, then $J \in \text{Sp}(V)$.

The next result says that compatible complex structures always exist on symplectic vector space.

Theorem 3.8. Let (V, ω) be a symplectic vector space. Then there is a compatible complex structure J on V .

Proof. Since V is a vector space, we choose an inner product G on V . Since ω and G are nondegenerate, the maps

$$\left. \begin{array}{l} u \in V \xrightarrow{\omega^*(u)} \omega(u, \cdot) \in V^* \\ v \in V \xrightarrow{G^*(v)} G(v, \cdot) \in V^* \end{array} \right\} \text{ are isomorphisms between } V \text{ and } V^*.$$

Hence, we can find a linear map $A : V \rightarrow V$ such that $\omega(u, v) = G(u, Av)$.

Note that for any $u, v \in V$,

$$\begin{aligned}\omega^*(u)(v) &= \omega(u, v) = G(u, Av) = G^*(u)(Av) \\ \implies G^*(u) \circ A &= \omega^*(u) \quad \forall u \in V \implies G^* \circ A = \omega^* \\ \implies A &= G^{*-1} \circ \omega^*.\end{aligned}$$

If $A^2 = -Id$, then A is a compatible complex structure on V . Let us suppose that $A^2 \neq -Id$. Note that,

$$G(A^T u, v) = G(u, Av) = \omega(u, v) = -\omega(v, u) = -G(v, Au) = G(-Au, v),$$

which implies $A^T = -A$, that is, A is skew-symmetric. We also note that

- $(AA^T)^T = AA^T$, that is, AA^T is symmetric and
- for any $u \neq 0$, $G(AA^T u, u) = G(A^T u, A^T u) > 0$, that is, AA^T is positive definite.

This implies that AA^T is diagonalizable with positive eigenvalues λ_i . So there exists $P \in GL(n, \mathbb{R})$ such that

$$AA^T = P \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_{2n}).$$

So, we may take any real power of AA^T . In particular,

$$\sqrt{AA^T} := P \text{diag}(\sqrt{\lambda_1}, \sqrt{\lambda_2}, \dots, \sqrt{\lambda_{2n}}) P.$$

Then $\sqrt{AA^T}$ is symmetric and positive-definite. Let

$$J = (\sqrt{AA^T})^{-1} A.$$

This factorization $A = \sqrt{AA^T} J$ is called *polar decomposition* of A . Since A commutes with $\sqrt{AA^T}$, J must commute with $\sqrt{AA^T}$. Now we have

- $J^T = A^T \sqrt{AA^T}^{-1} = \sqrt{AA^T}(-A) = -J$ and
- Since $A^T = -A$, we have $JJ^T = Id$. Thus, $J^2 = -Id$.

So, J is a complex structure on V . Now it remains to show that it is compatible with ω , that is $G_j(u, v) = \omega(u, Jv)$ is an inner product. Equivalently,

(i) $\omega(Ju, Jv) = \omega(u, v)$.

$$\omega(Ju, Jv) = G(Ju, AJv) = G(u, J^t AJv) = G(u, AJ^t Jv) = G(u, Av) = \omega(u, v).$$

(ii) For any $v \neq 0$, $\omega(v, Jv) > 0$.

$$\omega(v, Jv) = G(v, AJv) = G(v, \sqrt{AA^T} v) > 0.$$

□