

# Derivative

## Engineering Mathematics-I

Dr. (PhD) Sachchidanand Prasad

SPNREC, Araria

October 7, 2024

# Today's Goal

- Understand the importance of the **customer** in business
- Understand the importance of the **employee** in business
- Understand the importance of the **owner** in business
- Understand the importance of the **community** in business
- Understand the importance of the **environment** in business
- Understand the importance of the **technology** in business
- Understand the importance of the **innovation** in business
- Understand the importance of the **competition** in business
- Understand the importance of the **market** in business
- Understand the importance of the **product** in business
- Understand the importance of the **service** in business
- Understand the importance of the **brand** in business
- Understand the importance of the **reputation** in business
- Understand the importance of the **trust** in business
- Understand the importance of the **loyalty** in business
- Understand the importance of the **customer satisfaction** in business
- Understand the importance of the **employee satisfaction** in business
- Understand the importance of the **owner satisfaction** in business
- Understand the importance of the **community satisfaction** in business
- Understand the importance of the **environmental satisfaction** in business
- Understand the importance of the **technological satisfaction** in business
- Understand the importance of the **innovative satisfaction** in business
- Understand the importance of the **competitive satisfaction** in business
- Understand the importance of the **market satisfaction** in business
- Understand the importance of the **product satisfaction** in business
- Understand the importance of the **service satisfaction** in business
- Understand the importance of the **brand satisfaction** in business
- Understand the importance of the **reputation satisfaction** in business
- Understand the importance of the **trust satisfaction** in business
- Understand the importance of the **loyalty satisfaction** in business
- Understand the importance of the **customer satisfaction** in business
- Understand the importance of the **employee satisfaction** in business
- Understand the importance of the **owner satisfaction** in business
- Understand the importance of the **community satisfaction** in business
- Understand the importance of the **environmental satisfaction** in business
- Understand the importance of the **technological satisfaction** in business
- Understand the importance of the **innovative satisfaction** in business
- Understand the importance of the **competitive satisfaction** in business
- Understand the importance of the **market satisfaction** in business
- Understand the importance of the **product satisfaction** in business
- Understand the importance of the **service satisfaction** in business
- Understand the importance of the **brand satisfaction** in business
- Understand the importance of the **reputation satisfaction** in business
- Understand the importance of the **trust satisfaction** in business
- Understand the importance of the **loyalty satisfaction** in business

# Today's Goal

- ▶ What is derivative?

# Today's Goal

- ▶ What is derivative?
- ▶ Geometric meaning of derivative

# Today's Goal

- ▶ What is derivative?
- ▶ Geometric meaning of derivative
- ▶ Some examples

# Today's Goal

- ▶ What is derivative?
- ▶ Geometric meaning of derivative
- ▶ Some examples
- ▶ Rolle's theorem

# Today's Goal

- ▶ What is derivative?
- ▶ Geometric meaning of derivative
- ▶ Some examples
- ▶ Rolle's theorem
- ▶ Lagrange's mean value theorem

# What is derivative?



# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by

# What is derivative?


Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by

$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$

# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by


$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$



# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by


$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$



# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by


$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$



# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by


$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$



# What is derivative?

Suppose we are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the line. Then the slope of the line is calculated by

$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$



# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line




# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).


# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).




# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).




# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).




# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).




# Average Change

If we are given with a curve with two points, then the average rate of change is calculated by the slope of the secant line (*the line which passes through the two points on the curve*).




## Average Change Cont...

## Average Change Cont...







# Average Change Cont...




# Average Change Cont...




# Average Change Cont...



# Average Change Cont...



# Average Change Cont...



## Average Change Cont...

## Average Change Cont...

- ▶ The slope of the secant line is also referred to as the *average rate of change* of  $f$  over the interval  $[x, x + h]$ .

## Average Change Cont...

- ▶ The slope of the secant line is also referred to as the *average rate of change* of  $f$  over the interval  $[x, x + h]$ .
- ▶ Thus, average rate of change of  $f$  over the interval  $[x, x + h]$  will be



## Average Change Cont...

- ▶ The slope of the secant line is also referred to as the *average rate of change* of  $f$  over the interval  $[x, x + h]$ .
- ▶ Thus, average rate of change of  $f$  over the interval  $[x, x + h]$  will be

$$\frac{f(x + h) - f(x)}{h}.$$

## Average Change Cont...

- ▶ The slope of the secant line is also referred to as the *average rate of change* of  $f$  over the interval  $[x, x + h]$ .
- ▶ Thus, average rate of change of  $f$  over the interval  $[x, x + h]$  will be

$$\frac{f(x + h) - f(x)}{h}.$$

- ▶ What will happen if  $h$  “tends” to 0?

# Tangent Line

# Tangent Line

As  $h$  tends to 0 the points  $x$  and  $x + h$  coincide

# Tangent Line

As  $h$  tends to 0 the points  $x$  and  $x + h$  coincide and hence the secant line tends to the tangent line at  $x$ .

# Tangent Line

As  $h$  tends to 0 the points  $x$  and  $x + h$  coincide and hence the secant line tends to the tangent line at  $x$ . See the animation.

# Tangent Line

As  $h$  tends to 0 the points  $x$  and  $x + h$  coincide and hence the secant line tends to the tangent line at  $x$ . See the animation. Thus,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

Slope of the tangent line is known as *Instantaneous Rate of Change*

# Examples



# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \end{aligned}$$

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{x^2 + h^2 + 2xh - x^2}{h}\end{aligned}$$

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{x^2 + h^2 + 2xh - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{h^2 + 2hx}{h}\end{aligned}$$

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{x^2 + h^2 + 2xh - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{h^2 + 2hx}{h} \\&= \lim_{h \rightarrow 0} (h + 2x)\end{aligned}$$

# Examples

Let us check the derivative of  $f(x) = x^2$  is  $2x$ .

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{x^2 + h^2 + 2xh - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{h^2 + 2hx}{h} \\&= \lim_{h \rightarrow 0} (h + 2x) \\&= 2x.\end{aligned}$$

## Example Cont...



## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists.

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

The left hand limit is  $-1$

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

The left hand limit is  $-1$  and the right hand limit is  $1$ ,

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

The left hand limit is  $-1$  and the right hand limit is  $1$ , hence the limit does not exist.

## Example Cont...

Check the differentiability of the function  $f(x) = |x|$ .

Note that for any  $x \in \mathbb{R} \setminus \{0\}$ , the limit

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

exists. For  $x = 0$ , we have

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

The left hand limit is  $-1$  and the right hand limit is  $1$ , hence the limit does not exist. Thus, the function is not differentiable at  $x = 0$ .



# Application of derivative

# Application of derivative

We will discuss the following applications:

# Application of derivative

We will discuss the following applications:

- ▶ Rolle's theorem

# Application of derivative

We will discuss the following applications:

- ▶ Rolle's theorem
- ▶ Mean value theorem

# Rolle's Theorem

# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$

# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$



# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$
- ▶  $f(a) = f(b)$

# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$
- ▶  $f(a) = f(b)$

Then, there exists  $c \in (a, b)$  such that

$$f'(c) = 0.$$

# Rolle's Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$
- ▶  $f(a) = f(b)$


Then, there exists  $c \in (a, b)$  such that

$$f'(c) = 0.$$


# A pictorial view of Rolle's theorem




# A pictorial view of Rolle's theorem




# A pictorial view of Rolle's theorem



# A pictorial view of Rolle's theorem




# A pictorial view of Rolle's theorem





# A pictorial view of Rolle's theorem



# A pictorial view of Rolle's theorem

Show the animation

# Problems on Rolle's Theorem

## Problem

*For the function  $f(x) = x(x^2 - 1)$  test for the applicability of Rolle's theorem in the interval  $[-1, 1]$  and hence find  $c$  such that  $-1 < c < 1$ .*

## Problem cont...

### Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

## Problem cont...

### Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

*We have*

1.  *$f$  is continuous on  $[-1, 1]$ ,*

## Problem cont...

### Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

*We have*

1.  *$f$  is continuous on  $[-1, 1]$ ,*
2.  *$f$  is differentiable on  $(-1, 1)$*

## Problem cont...

### Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

*We have*

1.  *$f$  is continuous on  $[-1, 1]$ ,*
2.  *$f$  is differentiable on  $(-1, 1)$  and*
3.  *$f(-1) = 0 = f(1)$ .*

# Problem cont...

## Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

*We have*

1.  *$f$  is continuous on  $[-1, 1]$ ,*
2.  *$f$  is differentiable on  $(-1, 1)$  and*
3.  *$f(-1) = 0 = f(1)$ .*

*Since  $f$  satisfies the hypothesis of Rolle's theorem,*



# Problem cont...

## Solution

*Given that the function is*

$$f(x) = x(x^2 - 1).$$

*We have*

1.  $f$  is continuous on  $[-1, 1]$ ,
2.  $f$  is differentiable on  $(-1, 1)$  and
3.  $f(-1) = 0 = f(1)$ .

*Since  $f$  satisfies the hypothesis of Rolle's theorem, there exists  $c \in (-1, 1)$  such that  $f'(c) = 0$ . That is,*

## Solution cont...

$$f'(c) = 0$$

## Solution cont...


$$f'(c) = 0 \implies 3c^2 - 1 = 0$$

## Solution cont...

$$f'(c) = 0 \implies 3c^2 - 1 = 0 \implies c = \pm\sqrt{\frac{1}{3}}.$$


## Solution cont...

$$f'(c) = 0 \implies 3c^2 - 1 = 0 \implies c = \pm\sqrt{\frac{1}{3}}.$$



## Solution cont...

$$f'(c) = 0 \implies 3c^2 - 1 = 0 \implies c = \pm\sqrt{\frac{1}{3}}.$$



# Problem 2

## Problem

*Verify the Rolle's theorem for*

$$f(x) = \frac{\sin x}{e^x}, \quad \text{in } (0, \pi).$$

# Problem 3

## Problem

*It is given that the Rolle's theorem holds for the function*

$$f(x) = x^3 + bx^2 + cx, \quad 1 \leq x \leq 2$$

*at the point  $x = \frac{4}{3}$ . Find the value of  $b$  and  $c$ .*



# Lagrange's Mean Value Theorem

# Lagrange's Mean Value Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

# Lagrange's Mean Value Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$

# Lagrange's Mean Value Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$

# Lagrange's Mean Value Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$

Then, there exists  $c \in (a, b)$  such that

$$\frac{f(b) - f(a)}{b - a} = f'(c).$$

# Lagrange's Mean Value Theorem

Suppose  $f$  is defined on  $[a, b]$  such that

- ▶  $f$  is continuous on  $[a, b]$
- ▶  $f$  is differentiable on  $(a, b)$

Then, there exists  $c \in (a, b)$  such that

$$\frac{f(b) - f(a)}{b - a} = f'(c).$$

Proof.


Take

$$g(x) = f(x) - f(a) - \frac{f(b) - f(a)}{b - a}(x - a).$$




A picture

# A picture







# A picture




# A picture



# A picture



# A picture



# Problem-1

## Problem

*Verify the Lagrange's mean value theorem for the function*

$$f(x) = x(x-1)(x-2), \quad a = 0 \text{ and } b = \frac{1}{2}.$$

*Also find  $c$ .*

# Problem-2

## Problem

*Verify the Lagrange's mean value theorem for the function given below and find  $c$*

$$f(x) = \log x, \quad a = 1 \text{ and } b = e.$$

# Cauchy's Mean Value Theorem

Let  $f$  and  $g$  be two functions defined on  $[a, b]$  such that

# Cauchy's Mean Value Theorem

Let  $f$  and  $g$  be two functions defined on  $[a, b]$  such that

- ▶  $f$  and  $g$  are continuous on  $[a, b]$



# Cauchy's Mean Value Theorem

Let  $f$  and  $g$  be two functions defined on  $[a, b]$  such that

- ▶  $f$  and  $g$  are continuous on  $[a, b]$
- ▶  $f$  and  $g$  are differentiable on  $(a, b)$

# Cauchy's Mean Value Theorem

Let  $f$  and  $g$  be two functions defined on  $[a, b]$  such that

- ▶  $f$  and  $g$  are continuous on  $[a, b]$
- ▶  $f$  and  $g$  are differentiable on  $(a, b)$  and
- ▶  $g'(x) \neq 0$  for any  $x \in (a, b)$ .

# Cauchy's Mean Value Theorem

Let  $f$  and  $g$  be two functions defined on  $[a, b]$  such that

- ▶  $f$  and  $g$  are continuous on  $[a, b]$
- ▶  $f$  and  $g$  are differentiable on  $(a, b)$  and
- ▶  $g'(x) \neq 0$  for any  $x \in (a, b)$ .

Then there exists  $c \in (a, b)$  such that

$$\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}.$$