


Validity of epidemiological studies


Dr. Tarun Bhatnagar MD, PhD


nie.gov.in











Threats to validity in epidemiologic studies

Selection bias

Do study participants accurately represent target population?

Bias results from procedures used to select individuals for inclusion in the study and/or analyses

nie.gov.in

HEALTH RESEARCH FUNDAMENTALS



Selection bias in epidemiological studies

- Surveillance Systematic notification of exposed cases
- Screening / diagnosis Systematic case search among exposed
- Admission to health care ICAL RESEA facilities - Systematic admission of:
 - Case-patients exposed / unexposed
 - Control-subjects exposed / unexposed

- Selective survival Systematic inclusion of cases who survived and who may be more or less exposed
- Non response / loss to follow up - Systematic inclusion of subjects more likely to participate who may be:
 - More or less exposed
 - More or less at risk



cma NIE

Dealing with selection bias

Designing stage of a study

- Use incident cases, not prevalent cases
- Case control studies

 - Apply same eligibility criteria for selecting cases and controls
 - Both cases and controls undergo the same diagnostic procedures and intensity of disease surveillance

Dealing with selection bias

Data collection stage of the study

- Minimize nonresponse, nonparticipation and loss to follow-up (Cohort studies)
- Keep a record of all losses and collect baseline data on them
- Make sure that diagnosis of disease is not affected by exposure status (blinding)



Dealing with selection bias

Analysis stage of study

- Compare non-responders/dropouts with responders/non-dropouts with respect to baseline variables
 - Large differences strongly suggest selection bias IOLOGY
 - Small differences do not rule out selection bias
- Use study results and external information to deduce the direction of biases and assess magnitude of biases
 - · Do sensitivity analysis



Threats to validity in epidemiologic studies

Information bias

Do measurements accurately represent phenomena of interest?

Bias results from procedures used to measure "exposures", "outcomes" and "other variables"



icma NIE


Information bias in epidemiological studies

- Case control study
 - Collection of information leaning towards specific exposure status
 - Recall Cases may recall exposure more than controls
 - Better exposure data available on cases compared to controls
- Cohort study
 - Collection of information leaning towards specific outcome status
 - Better outcome data available on exposed compared to unexposed
- Investigator Systematic collection of information supporting expected conclusions (Unconsciously or Consciously) INSTITUTE OF
 - May be examined in the analysis
- Prevarication -Systematic distortion of the truth by subjects



Dealing with information bias

Precise operational definitions of variables Detailed measurement protocols Repeated measurements on key variables Training, certification and re-certification Data audits (of interviewers, of data centers) Data cleaning - visual, computer Re-running all analyses prior to publication





Exposure Outcome

- CONFOUNDER
- Design stage
- Analysis stage
- Restriction
- NDIAN COUNC® Stratification STITUTE OF
- Matching

- Multivariate analysis
- Randomization
 - Experimental studies



How to evaluate associations? Crude Chance Selection bias Information bias Confounding Causal Association Confounding Causal Association Integral Property of the P

Does coffee increase risk of heart attack?

- Truth in Universe
- Population: All Adults
- Actual Coffee intake
- Actual heart attack (MI)





- Truth in Study
- Sample:

Consenting adults Low participation rate

Hospital patients or

- Reported Coffee intake
- Reported/misdiagnosed heart attack (MI)



Does coffee increase risk of heart attack?

- Was the association between coffee and MI due to **CONFOUNDING** by smoking?
 - "A confounder is associated with both the exposure (coffee) and the outcome (MI)."
 - Smoking in
 - 86%

80%

40%

- coffee (+) coffee (-) 27%
- MI (+)
- MI (-)

