



JEE ADVANCED



Sri Chaitanya IIT Academy., India.

★ A.P ★ T.S ★ KARNATAKA ★ TAMILNADU ★ MAHARASTRA ★ DELHI ★ RANCHI

A right Choice for the Real Aspirant

ICON Central Office, Madhapur – Hyderabad

Sec: Sr.Super60(Incoming)_STERLING BT

2022_P1

Date: 18-05-2025

Time: 09.00Am to 12.00Pm

WTA-32

Max. Marks:180

MATHEMATICS

: Areas: Drawing of different graphs, Area under plane curve as a definite integral, Finding out the functions from area functions.

PHYSICS

: MOTIONAL EMF, Motion of a straight conductor in a magnetic field(Theoretical explanation of electromagnetic induction motional emf), Motion of a loop in a magnetic field, Emf induced across a bar rotating in a magnetic field, AC generator.

CHEMISTRY

: GROUP-17: Introduction;Occurrence;atomic and physical properties ;oxidation states;anomalous properties of fluorine;Reactivity towards metals.(General Discussion),preparations,properties and uses (uses as per NCERT) of Cl₂, preparations,properties and uses (uses as per NCERT) of HCl, Structures, preparations, properties and uses of Oxides of chlorine, NOTE:[preparations,properties and uses of F₂,Br₂,I₂(not to be tested); NOTE:[preparations,properties and uses of HF,HBr,HI(not to be tested},, Structures, preparations, properties and uses (uses as per NCERT)of oxoacids of chlorine, preparations,properties and uses of Bleaching powder ; Structures, preparations,properties and uses (uses as per NCERT)of interhalogencompounds. Structures of pseudo halides & Polyhalides NOTE ; GROUP-17: General Introduction: Electronic configuration and general trends in physical and chemical properties of elements across the periods and down the groups; unique behaviour of the first element in each group.(ONLY FOR JEE MAINS), Group-18:Introduction;Occurrence;atomic and physical properties; chemical properties ; compounds of xenon (xenon -fluorine compounds , xenon - oxygen compounds); uses (as per NCERT) of noble gases NOTE: clathrates(not to be tested) NOTE ; GROUP-18: General Introduction: Electronic configuration and general trends in physical and chemical properties of elements across the periods and down the groups; unique behaviour of the first element in each group.(ONLY FOR JEE MAINS)

Name of the Student: _____

H.T. NO:

--	--	--	--	--	--

**JEE-ADVANCE-2022-P1-Model**

Time:3Hr's

IMPORTANT INSTRUCTIONS

Max Marks: 180

MATHEMATICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 1 – 8)	Questions with Numerical Value Answer Type%	+3	0	8	24
Sec – II(Q.N : 9 – 14)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – III(Q.N : 15 – 18)	Matching Type	+3	-1	4	12
Total				18	60

PHYSICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 19 – 26)	Questions with Numerical Value Answer Type	+3	0	8	24
Sec – II(Q.N : 27 – 32)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – III(Q.N : 33 – 36)	Matching Type	+3	-1	4	12
Total				18	60

CHEMISTRY:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 37 – 44)	Questions with Numerical Value Answer Type	+3	0	8	24
Sec – II(Q.N : 45 – 50)	Questions with Multiple Correct Choice with partial mark	+4	-2	6	24
Sec – III(Q.N : 51 – 54)	Matching Type	+3	-1	4	12
Total				18	60





MATHEMATICS

Max Marks: 60

SECTION – I
(NUMERICAL VALUE TYPE)

This section contains **EIGHT (08)** questions.

- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks:+3 **ONLY** if the correct numerical value is entered ;

Partial Mark: 0 In all other cases.

- Let $f(x)$ be a function which satisfy the equation $f(xy) = f(x) + f(y)$ for all $x > 0, y > 0$ such that $f'(1) = 2$. Let A be the area of the region bounded by the curves

$$y = f(x), y = \left| x^3 - 6x^2 + 11x - 6 \right| \text{ and } x = 0, \text{ then find value of } \frac{28}{17} A.$$

- Area of the region bounded by $[x]^2 = [y]^2$, if $x \in [1, 5]$, where $[]$ denotes the greatest integer function, is :

- Let f be a differentiable function satisfying the condition

$$f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)} \left(y \neq 0, f(y) \neq 0 \right) \forall x, y \in R \text{ and } f'(1) = 2. \text{ If the smaller area}$$

enclosed by $y = f(x), x^2 + y^2 = 2$ is A , then find $[A]$, where $[.]$ represents the greatest integer function.

- Consider $y = x^2$ and $f(x)$ where $f(x)$, is a differentiable function satisfying

$$f(x+1) + f(z-1) = f(x+z) \forall x, z \in R \text{ and } f(0) = 0; f'(0) = 4. \text{ If area bounded}$$

by curve $y = x^2$ and $y = f(x)$ is Δ , find the value of $\left(\frac{3}{16}\Delta\right)$.

- The least integer which is greater than or equal to the area of region in $x - y$ plane satisfying $x^6 - x^2 + y^2 \leq 0$ is :

- The set of points (x, y) in the plane satisfying $x^{2/5} + |y| = 1$ form a curve enclosing a region of area $\frac{p}{q}$ square units, where p and q are relatively prime positive integers.

Find $p - q$.

**JEE MAIN
2023**

SINGARAJU
VENKAT KOURDINYA
Sri Chaitanya
6th-12th Class
**300
300
MARKS**



RANK

**JEE Advanced
2023**

VAVILALA
CHIDVILAS REDDY
Sri Chaitanya
6th-12th Class
**341
360
MARKS**



RANK

**NEET
2023**

BORA VARUN
CHAKRAVANTHI
Sri Chaitanya
6th-12th Class
**720
720
MARKS**



RANK



7. Let the function $f : [-4, 4] \rightarrow [-1, 1]$ be defined implicitly by the equation $x + 5y - y^5 = 0$. If the area of triangle formed by tangent and normal to $f(x)$ at $x = 0$ and the line $y = 5$ is A , find $\frac{A}{13}$.
8. If the area bounded by circle $x^2 + y^2 = 4$, the parabola $y = x^2 + x + 1$ and the curve $y = \left[\sin^2 \frac{x}{4} + \cos \frac{x}{4} \right]$, (where $[]$ denotes the greatest integer function) and x -axis is $\left(\sqrt{3} + \frac{2\pi}{3} - \frac{1}{k} \right)$, then the numerical quantity k should be :

SECTION – II

(ONE OR MORE CORRECT ANSWER TYPE)

This section contains **SIX (06)** questions.

- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks :+4 ONLY if (all) the correct option(s) is(are) chosen;

Partial Marks: +3 If all the four options are correct but **ONLY** three options are chosen;


Partial Marks: +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct ;

Partial Marks: +1 If two or more options are correct but **ONLY** two options are chosen, and it is a correct option ;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : -2 In all other cases.

9. Let $f(x)$ be a polynomial function of degree 3 where $a < b < c$ and $f(a) = f(b) = f(c)$. If the graph of $f(x)$ is as shown, which of the following statements are INCORRECT? (Where $c > |a|$)



- A) $\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$
- B) $\int_a^c f(x) dx < 0$
- C) $\int_a^b f(x) dx < \int_c^b f(x) dx$
- D) $\frac{1}{b-a} \int_a^b f(x) dx > \frac{1}{c-b} \int_b^c f(x) dx$

THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN 2023

SINGARAJU VENKAT KOUNDRINA
from 12th Class
300
300 MARKS



RANK

JEE Advanced 2023

VAVILALA CHIDVILAS REDDY
from 12th Class
341
360 MARKS



RANK

NEET 2023

BORA VARUN CHAKRAVARTHI
from 12th Class
720
720 MARKS



RANK



10. $T_n = \sum_{r=2n}^{3n-1} \frac{r}{r^2 + n^2}, S_n = \sum_{r=2n+1}^{3n} \frac{r}{r^2 + n^2}$, then $\forall n \in \{1, 2, 3, \dots\}$:
 A) $T_n > \frac{1}{2} \ln 2$ B) $S_n < \frac{1}{2} \ln 2$ C) $T_n < \frac{1}{2} \ln 2$ D) $S_n > \frac{1}{2} \ln 2$
11. If a curve $y = a\sqrt{x} + bx$ passes through point $(1, 2)$ and the area bounded by curve, line $x = 4$ and x -axis is 8, then :
 A) $a = 3$ B) $b = 3$ C) $a = -1$ D) $b = -1$
12. Area enclosed by the curves $y = x^2 + 1$ and a normal drawn to it with gradient -1; is equal to :
 A) $\frac{2}{3}$ B) $\frac{1}{3}$ C) $\frac{3}{4}$ D) $\frac{4}{3}$
13. The triangle formed by the normal to the curve $f(x) = x^2 - ax + 2a$ at the point $(2, 4)$ and the coordinate axes lies in second quadrant, if its area is $2sq$ units, then α can be
 A) 2 B) $\frac{17}{4}$ C) 5 D) $\frac{19}{4}$
14. Let f and g be continuous function on $a \leq x \leq b$ and set $p(x) = \max\{f(x), g(x)\}$ and $q(x) = \min\{f(x), g(x)\}$, then the area bounded by the curves $y = p(x)$, $y = q(x)$ and the ordinates $x = a$ and $x = b$ is given by
 A) $\int_a^b |f(x) - g(x)| dx$ B) $\int_a^b |p(x) - q(x)| dx$
 C) $\int_a^b \{f(x) - g(x)\} dx$ D) $\int_a^b \{p(x) - q(x)\} dx$

THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN 2023 SINGARAJU VENKAT KOURINDYA Sri Chaitanya 6th-12th Class 300 300 MARKS	RANK 1 JEE Advanced 2023 VAVILALA CHIDVILAS REDDY Sri Chaitanya 6th-12th Class 341 360 MARKS	RANK 1 NEET 2023 BORA VARUN CHAKRAVARTHI Sri Chaitanya 6th-12th Class 720 720 MARKS	RANK 1
---	--	---	---------------



SECTION – III
(MATCHING TYPE)

This section contains **FOUR (04)** Matching List Sets.

- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists : **List-I** and **List-II**.
- **List-I** has **Four** entries (I), (II), (III) and (IV) and **List-II** has **Five** entries (P), (Q), (R), (S) and (T).
- **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks: +3 ONLY if the option corresponding to the correct combination is chosen;

Zero Marks: 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks: -1 In all other cases.

15. Match the statements of Column I with values of Column II

Column – I		Column - II	
(A)	The area bounded by the curve $y = x + \sin x$ and its inverse function between the ordinates $x = 0$ to $x = 2\pi$ is $4s$. Then the value of s is	(p)	2
(B)	The area bounded by $y = xe^{ x }$ and lies $ x =1, y=0$ is	(q)	1
(C)	The area bounded by the curves $y^2 = x^3$ and $ y =2x$ is	(r)	$\frac{32}{5}$
(D)	The smaller are included between the curves $\sqrt{ x } + \sqrt{ y } = 1$ and $ x + y = 1$ is	(s)	$\frac{4}{3}$
		(t)	1 / 3

A) A – p, B – p, C – r, D – s B) A – p, B – s, C – p, D – r

C) A – p, B – p, C – r, D – t D) A – s, B – q, C – r, D – t

16. Match the statements of Column I with values of Column II

Column - I		Column - II	
(A)	Area enclosed by $y = x , x = 2$ and $y = 0$ is	(p)	2
(B)	Area enclosed by the curve $y = \sin x, x = 0, x = \pi$ and $y = 0$ is	(q)	4
(C)	If the area of the region bounded by $x^2 \leq y$ and $y \leq x + 2$ is $\frac{k}{4}$, then k is equal to	(r)	27
(D)	Area of the quadrilateral formed by tangents at the ends of latusrectum of ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$ is	(s)	18

A) A – r, B – p, C – r, D – s B) A – p, B – p, C – s, D – r

C) A – p, B – r, C – s, D – q D) A – p, B – p, C – r, D – r

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN
2023**

SINGARAJU
VENKAT KOURINDYA
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
6th-12th Class
300
300
MARKS



RANK

**JEE Advanced
2023**

VAVILALA
CHIDVILAS REDDY
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
6th-12th Class
341
360
MARKS



RANK

**NEET
2023**

BORA VARUN
CHAKRAVARTHI
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
6th-12th Class
720
720
MARKS



RANK



17. Match the statements of Column I with values of Column II

	Column - I	Column - II
(A)	Area of region formed by points (x, y) satisfying $[x]^2 = [y]^2$ for $0 \leq x \leq 4$ is equal to (where $[]$ denotes greatest integer function)	(p) 48
(B)	The area of region formed by points (x, y) satisfying $x + y \leq 6, x^2 + y^2 \leq 6y$ and $y^2 \leq 8x$ is $\frac{k\pi - 2}{12}$, then $k =$	(q) 27
(C)	The area in the first quadrant bounded by the curve $y = \sin x$ and the line $\frac{2y-1}{\sqrt{2}-1} = \frac{2}{\pi}(6x-\pi)$ is $\left[\frac{\sqrt{3}-\sqrt{2}}{2} - \frac{(\sqrt{2}+1)\pi}{k} \right]$, then $k =$	(r) 7
(D)	If the area bounded by the graph of $y = xe^{-ax}$ ($a > 0$) and the abscissa axis is $\frac{1}{9}$ then the value of ' a ' is equal to	(s) 4
		(t) 3

- A) $A - r, B - q, C - p, D - p$ B) $A - r, B - q, C - p, D - t$
 C) $A - s, B - q, C - p, D - r$ D) $A - s, B - q, C - s, D - t$

18.

	Column - I	Column - II
I)	Area enclosed by $[x]^2 = [y]^2$ for $1 \leq x \leq 4$	(p) 8 sq.units
II)	Area enclosed by $[\lceil x \rceil] + [\lceil y \rceil] = 2$	(q) 6 sq.units
III)	Area enclosed by $[\lceil x \rceil][\lceil y \rceil] = 2$	(r) 4 sq.units
IV)	Area enclosed by $\frac{[\lceil x \rceil]}{[\lceil y \rceil]} = 2, -5 \leq x \leq 5$	(s) 12 sq.units

(Note :- $[\cdot]$ is G.I.F)

- A) $I - q, II - s, III - p, IV - p$ B) $I - r, II - q, III - p, IV - s$
 C) $I - p, II - q, III - s, IV - r$ D) $I - r, II - p, III - q, IV - r$



SECTION – I
(NUMERICAL VALUE TYPE)


This section contains **EIGHT (08)** questions.

- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks: +3 **ONLY** if the correct numerical value is entered ;


Partial Mark: 0 In all other cases.

19. A square metal wire loop of side 10 cm and resistance 1Ω is moved with a constant velocity v_0 in uniform magnetic field of induction $B = 2 \text{ weber}/m^2$ as shown in the figure. The magnetic field lines are perpendicular to the plane of the loop (directed into the paper). The loop is connected to a network of resistors each of value 3Ω . The resistance of the lead wires OS and PQ are negligible. The speed of the loop so as to have a steady current of 1 mA in the loop is $n \times 10^{-2} \text{ m/s}$ then n is




20. A wire is bent as a parabola $y = cx^2$ is located in a uniform magnetic field of magnitude B perpendicular to the xy plane as shown. At the instant $t = 0$ a long metal rod starts translating from the vertex of the parabola with a constant acceleration along positive y-axis. The induced emf across the points of contact of the rod with the wire is given by

$$\varepsilon = 2By\sqrt{\frac{ka}{c}}; \text{ where } k =$$






21. Two conducting rings P and Q of radii r and $2r$ rotate uniformly in opposite directions with centre of mass velocities $2v$ and v respectively on a conducting surface S. There is a uniform magnetic field of magnitude B perpendicular to the plane of the rings. The potential difference between the highest points of the two rings is $nBrv$. Then $n =$
22. In ABCD is a fixed smooth conducting frame in horizontal plane. T is a bulb of power 100W, P is a smooth pulley and OQ is a conducting rod. Neglect the self-inductance of the loop and resistance of any part other than the bulb. The mass M is moving down with constant velocity 10ms^{-1} . Bulb lights at its rated power due to induced emf in the loop due to earth's magnetic field. Find the mass M (in kg) of the block ($g = 10\text{ms}^{-2}$)




23. A conducting rod of length l , resistance R and mass m moving vertically downward due to gravity. Other parts are kept fixed. Magnetic field is B . MN and PQ are vertical, smooth conducting rails. The capacitance of the capacitor is C. The rod is released from rest. The maximum current induced in the circuit is $i_{\max} = \frac{mgBlc}{m + B^n l^2 c}$ then n






24. A thin semicircular conducting ring of radius R is falling with its plane vertical in horizontal magnetic induction \vec{B} . At the position MNQ, the speed of the ring is V, and the potential difference developed across the ring is $\frac{BnR^2}{4}$ then n is




25. A rectangular loop with a sliding connector of length $l = 1.0\text{m}$ is situated in a uniform magnetic field $B = 2\text{T}$ perpendicular to the plane of loop. Resistance of connector is $r = 2\Omega$. Two resistance of 6Ω and 3Ω are connected. The external force required to keep the connector moving with a constant velocity $v = 2\text{ms}^{-1}$ is in _____ N



26. A long straight wire carries a current I_o at distances a and b from it there are two other wires, parallel to the former one, which are interconnected by a resistance R. A connector slides without friction along the wires with a constant velocity. Assuming the resistance of the wires, the connector, the sliding contacts, and the self-inductance of the frame to be negligible, the force required to maintain the connector's velocity

$$\text{constant is } F = \frac{\theta}{R} \left(\frac{\mu_0 i_o}{2\pi} \ln\left(\frac{b}{a}\right) \right)^n \text{ then n is}$$





SECTION – II

(ONE OR MORE CORRECT ANSWER TYPE)

This section contains **SIX (06)** questions.

- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks :+4 ONLY if (all) the correct option(s) is(are) chosen;

Partial Marks: +3 If all the four options are correct but ONLY three options are chosen;


Partial Marks: +2 If three or more options are correct but ONLY two options are chosen, both of which are correct ;

Partial Marks: +1 If two or more options are correct but ONLY two options are chosen, and it is a correct option ;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);


Negative Marks : -2 In all other cases.

27. A conducting rod of length l is moved at constant velocity v_0 on two parallel conducting, smooth fixed rails, that are placed in a uniform constant magnetic field B perpendicular to the plane of the rails as shown in figure. A resistance R is connected between the two ends of the rail. Then which of the following is/are correct?




- A) The thermal power dissipated in the resistor is equal to rate of work done by external person pulling the rod.
- B) If applied external force is doubled then a part of external power increases the velocity of rod
- C) Lenz's law is not satisfied for direction of current in Loop
- D) If resistance R is doubled then power required to maintain the constant velocity V_0 become half
28. In a region there exist a magnetic field B_0 along positive x-axis. A metallic wire of length $2a$ and one side along x-axis and one side parallel of y-axis is rotating about y-axis with an angular velocity w . Then at the instant shown.





- A) Potential difference across PQ is zero
 B) Potential difference across PQ is $\frac{1}{2}B_o\omega a^2$
 C) Potential difference across QR is $\frac{1}{2}B_o\omega a^2$
 D) Potential difference across QR is $B_o\omega a^2$
29. A flexible wire loop in the shape of a circle has a radius that grows linearly with time, as $r = r_o t$. There exists a time varying magnetic field in the region as $B = B_o t$, where B_o and r_o are positive constants with appropriate dimensions
 A) The rate of change of induced emf varies linearly with time
 B) The induced emf varies parabolically with time
 C) The dimensional formula of $\frac{B_o}{r_o}$ is $M^1 L^{-1} T^{-2} A^{-1}$
 D) The dimensions of $\frac{B_o}{r_o}$ is independent of the physical quantity length
30. A conducting rod of length l is hinged at point O. It is free to rotate in a vertical plane. There exists a uniform magnetic field \bar{B} in horizontal direction. The rod is released from the position as shown in fig. Potential difference between the two ends of the rod after rotating an angle θ is proportional to




- A) $l^{3/2}$ B) l^2 C) B D) $(\sin \theta)^{1/2}$






31. A disc of radius r is rolling without sliding on a horizontal surface with a velocity of center of mass v and angular velocity ω in a uniform magnetic field B which is perpendicular to the plane of the disc as shown. O is the center of the disc and P,Q,R and S are the four points on the disc. Which of the following statements is true?



- A) Due to translation, induced emf across PS = Bvr
 B) Due to rotation, induced emf across QS=0
 C) Due to translation, induced emf across RO=0
 D) Due to rotation, induced emf across OQ = Bvr
32. A conducting wire of length l and mass m can slide without friction on two parallel rails and is connected to capacitance C . The whole system lies in a magnetic field B and a constant force F is applied to the rod. Then



- A) The rod moves with constant velocity
 B) The rod moves with an acceleration of $\frac{(F)}{(m + B^2 l^2 C)}$
 C) There is constant charge on the capacitor
 D) Charge on the capacitor increases with time

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN
2023**
SINGARAJU
VENKAT KOURDINYA
Sri Chaitanya
6th-12th Class
**300
300
MARKS**



RANK

**JEE Advanced
2023**
VAVILALA
CHIDVILAS REDDY
Sri Chaitanya
6th-12th Class
**341
360
MARKS**



RANK

**NEET
2023**
BORA VARUN
CHAKRAVANTHI
Sri Chaitanya
6th-12th Class
**720
720
MARKS**



RANK



SECTION – III
(MATCHING TYPE)

This section contains **FOUR (04)** Matching List Sets.



- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists : **List-I** and **List-II**.
- **List-I** has **Four** entries (I), (II), (III) and (IV) and **List-II** has **Five** entries (P), (Q), (R), (S) and (T).
- **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks: +3 ONLY if the option corresponding to the correct combination is chosen;

Zero Marks: 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks: -1 In all other cases.

33. A conducting loop is held in a magnetic field such that the field is oriented perpendicular to the area of the loop as shown. At any instant, magnetic flux density over the entire area has the same value but it varies with time



Column I		Column II	
i.	Induced current in the coil is in the clockwise sense	a.	For $t_2 < t < t_3$
ii	Induced current in the coil is in the anticlockwise sense	b.	For $t_3 < t < t_4$
iii	Induced current is zero	c.	For $t_5 < t < t_6$
iv	Induced current is maximum	d.	For $t_4 < t < t_5$

- A) i – a, ii – b, iii – d, iv – c B) i – b, ii – a, iii – c, iv – c
 C) i – c, ii – d, iii – b, iv – a D) i – c, ii – a, b, iii – d, iv – c

34. Match the dimensional formulae with the corresponding units


Column I		Column II	
A.	$ML^2 T^{-2} A^{-1}$	p.	Farad
B.	$M^{-1} L^{-2} T^4 A^2$	q.	Weber
C.	$ML^2 T^{-3} A^{-1}$	r.	Ohm
D	$ML^2 T^{-3} A^{-2}$	s	emf (volt)

- A) A → q; B → P; C → s; D → r B) A → c; B → q; C → r; D → s
 C) A → q; B → s; C → p; D → r D) A → r; B → s; C → p; D → q





35. The variation of magnetic flux associated with a coil of resistance 2Ω varies with time t as shown match the quantities in column I with their respective values in SI units, in column II



Column I		Column II	
A.	Induced emf produced	p.	4
B.	Induced current	q.	1
C.	Charge flows in 2 sec	r.	8
D.	Heat generated in 2sec	s	2

- A) $A \rightarrow s; B \rightarrow q; C \rightarrow s; D \rightarrow p$ B) $A \rightarrow s; B \rightarrow q; C \rightarrow p; D \rightarrow s$
 C) $A \rightarrow s; B \rightarrow p; C \rightarrow q; D \rightarrow s$ D) $A \rightarrow q; B \rightarrow p; C \rightarrow s; D \rightarrow p$
36. A uniform circular disc of mass M , radius R , having uniformly distributed charge Q is placed on a horizontal rough floor of coefficient of friction μ . A uniform vertically downward magnetic field, varying with time t as $B = B_0 t^2$ is confined in a cylindrical region of radius $r (> R_0)$ coaxial with the disc assuming B_0 to be a positive constant, match quantities in column-I (given as a multiple of k) with their respective k values in Column-II

Column I			Column II	
A.	Rotation of disc begins at time $t_o = k \left(\frac{\mu Mg}{QB_0 R} \right)$		p.	$\frac{2}{3}$
B.	Torque due to friction at $t = 2s$ is $\tau_f = k(\mu MgR)$		q.	$\frac{8}{9}$
C.	Torque due to magnetic field at time values $t = 0, t = t_o$ and $t = 3t_o$ are $k(\mu MgR)$, with multiple respective values of k to be selected from Column-II		r.	Zero
D.	Angular velocity of disc at $t = 2t_0$ is $\omega = k \left(\frac{M\mu^2 g^2}{QB_0 R^2} \right)$		s	$\frac{4}{3}$
			t	2

- A) $A \rightarrow p; B \rightarrow q; C \rightarrow t; D \rightarrow r$ B) $A \rightarrow s; B \rightarrow p; C \rightarrow p, r, t; D \rightarrow q$
 C) $A \rightarrow s; B \rightarrow p; C \rightarrow r, p, t; D \rightarrow q$ D) $A \rightarrow r, t; B \rightarrow s; C \rightarrow p; D \rightarrow q$

THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN

2023

SINGARAJU
VENKAT KOUNDRINA
Sri Chaitanya
6th-12th Class
300
300 MARKS

RANK

JEE Advanced

2023

VAVILALA
CHIDVILAS REDDY
Sri Chaitanya
6th-12th Class
341
360 MARKS

RANK

NEET

2023

BORA VARUN
CHAKRAVANTHI
Sri Chaitanya
6th-12th Class
720
720 MARKS

RANK



SECTION – I
(NUMERICAL VALUE TYPE)


This section contains **EIGHT (08)** questions.

- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks: +3 **ONLY** if the correct numerical value is entered ;

Partial Mark: 0 In all other cases.

37. In the following reaction,



X has 'P' σ -bonds, 'Q' π -bonds and 'R' lone pairs on Xe atom, then the value

$$\text{of } \left(\frac{P+Q+R}{4} \right) \text{ is } \underline{\hspace{2cm}}$$

38. In case of oxides of chlorine

X = oxidation no. of 'Cl' in Cl_2O_6 compound and

Y = the number of oxygen atoms surrounding each 'Cl' atom in Cl_2O_7 compound, then

$$\text{the value of } \left(\frac{x+y}{3} \right) \text{ is } \underline{\hspace{2cm}}$$

39. For oxyacid $HClO_x$, if $x = y = z$ (x, y, z are natural numbers) where

x = Total no. of lone pairs at central atom

y = No. of oxygen atoms

$$z = \text{Total no. of } \pi \text{ electrons in the oxyacid, the value of } \left(\frac{2x+y+z}{5} \right) \text{ is } \underline{\hspace{2cm}}$$

40. Consider the following compounds A to E

- A) XeF_n B) $XeF_{(n+1)}^{\oplus}$ C) XeF_{n+1}^{\ominus} D) XeF_{n+2} E) XeF_{n+4}^{2-}

If $n = 4$, P = total n. of bond pairs and

Q = Total no. of lone pairs on central atom of compounds A to E

$$\text{Then, the value of } \left(\frac{P+Q}{4} \right) \text{ is } \underline{\hspace{2cm}}$$

41. Xenon(Xe) on reaction with O_2F_2 gives is Xe compound 'P'. Three moles of Compound 'P' on complete hydrolysis gives 'x' moles of XeO_3 and 'y' moles of HF. Then the value

$$\text{of } \left(\frac{x+y}{2} \right) \text{ is } \underline{\hspace{2cm}}$$

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN
2023**

SINGARAJU
VENKAT KOURINDYA
Sri Chaitanya
6th-12th Class
300
300
MARKS



RANK

**JEE Advanced
2023**

VAVILALA
CHIDVILAS REDDY
Sri Chaitanya
6th-12th Class
341
360
MARKS



RANK

**NEET
2023**

BORA VARUN
CHAKRAVANTHI
Sri Chaitanya
6th-12th Class
720
720
MARKS



RANK



42. Ozonolysis of ClO_2 produces an oxide of chlorine then if A = No.of π -bonds in (P)
 B) = Average oxidation state of chlorine in (P), then value of $\left(\frac{2B - A}{3}\right)$ is _____
43. When Cl_2 reacts with hot conc. NaOH solution gives corresponding products, then
 If x = the difference in oxidation state of 'Cl' in the products
 Y = no.of moles of 'NaCl' produced during reaction.
 Then the value of (x / y) is _____
44. When 3.55g of bleaching powder is treated with acetic acid and excess of KI liberated iodine, which required 5ml of 0.25N hypo solution. What is the percentage of available Cl_2 ?

SECTION – II

(ONE OR MORE CORRECT ANSWER TYPE)

This section contains **SIX (06)** questions.

- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks :+4 ONLY if (all) the correct option(s) is(are) chosen;

Partial Marks: +3 If all the four options are correct but **ONLY** three options are chosen;


Partial Marks: +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct ;

Partial Marks: +1 If two or more options are correct but **ONLY** two options are chosen, and it is a correct option ;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : -2 In all other cases.

45. The correct order of the following is / are;
 A) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$: Acidic strength
 B) $\text{HF} > \text{HCl} > \text{HBr} > \text{HI}$: thermal stability
 C) $\text{HF} > \text{HCl} > \text{HBr} > \text{HI}$: Boiling point
 D) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$: Bond dissociation enthalpy
46. Which of the following is/are correct statement regarding the following reaction is



- A) XeF_6 can undergo partial hydrolysis
 B) 'Y' is an oxy-acid of xenon
 C) 'X' is an explosive
 D) Both are examples of non-redox reactions

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

JEE MAIN 2023 SINGARAJU VENKAT KOUDINYA 1st Position Sri Chaitanya 6th-12th Class 300 300 MARKS	RANK 1 JEE Advanced 2023 VAVALALA CHIDVILAS REDDY 1st Position Sri Chaitanya 6th-12th Class 341 360 MARKS	RANK 1 NEET 2023 BORA VARUN CHAKRAVARTHI 1st Position Sri Chaitanya 6th-12th Class 720 720 MARKS
--	--	---



47. Which of the following reactions of Xe compounds are feasible?

- A) $XeF_2 + HF \longrightarrow H(XeF_3)$ B) $XeF_4 + PF_5 \longrightarrow [XeF_3]^+ [PF_6]^-$
 C) $XeF_6 + RbF \longrightarrow [XeF_5]^+ [RbF_2]^-$ D) $3XeF_4 + 6H_2O \longrightarrow 2Xe + XeO_3 + 12HF + \frac{3}{2}O_2$

48. The species which are not formed on fluorination of phosphorous pentachloride in polar organic solvent are

- A) PF_3 and PCl_3 B) PF_5 and PCl_3
 C) $[PCl_4]^+ [PCl_4F_2]^\ominus$ and $[PCl_4]^+ [PF_6]^\ominus$ D) $[PF_4]^\oplus [PF_6]^\ominus$ and $[PCl_4]^\oplus [PF_6]^\ominus$

49. With respect to hypochlorite, chlorate and perchlorate ions choose the correct statements

- A) The molecular shape of only chlorate ions are influenced by the lone pair of electrons of ' Cl '
 B) The hypochlorite is the weakest conjugate base
 C) The hypochlorite ion oxidizes the sulphite ion
 D) The hypochlorite and chlorate ions disproportionate to give rise to identical set of ions

50. Partial hydrolysis XeF_6 gives

- A) XeO_2F_2 B) $XeOF_2$ C) $XeOF_4$ D) XeO_3

SECTION – III (MATCHING TYPE)

This section contains **FOUR (04)** Matching List Sets.

- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists : **List-I** and **List-II**.
- **List-I** has **Four** entries (I), (II), (III) and (IV) and **List-II** has **Five** entries (P), (Q), (R), (S) and (T).
- **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.

• Answer to each question will be evaluated according to the following marking scheme :

Full Marks:+3 ONLY if the option corresponding to the correct combination is chosen;

Zero Marks: 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks: -1 In all other cases.

51. Match the xenon compounds given in List-I with corresponding geometries and number of lone pairs on Xe given in list-II based on VSEPR model and choose the correct option

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN
2023**

SINGARAJU
VENKAT KOUNDRINA
B.Tech
Sri Chaitanya
6th-12th Class
**300
300
MARKS**



RANK

**JEE Advanced
2023**

VAVILALA
CHIDVILAS REDDY
B.Tech
Sri Chaitanya
6th-12th Class
**341
360
MARKS**



RANK

**NEET
2023**

BORA VARUN
CHAKRAVANTHI
B.Tech
Sri Chaitanya
6th-12th Class
**720
720
MARKS**



RANK



List-I		List-II	
P)	XeF_4	1)	Tetrahedral and one lone pair of electrons
Q)	XeO_3F_2	2)	Trigonal bipyramidal and 2 lone pair of electrons
R)	XeF_2	3)	Octahedral and 2 lone pair electrons
S)	XeO_3	4)	Trigonal bipyramidal and no lone pair of electrons
		5)	Trigonal bipyramidal and 3 lone pair of electrons

A) $P - 4, Q - 3, R - 5, S - 1$ B) $P - 4, Q - 2, R - 1, S - 3$

C) $P - 3, Q - 4, R - 5, S - 1$ D) $P - 3, Q - 4, R - 2, S - 1$

52. Match the following sets related to compounds and number of lone pairs at central atoms

List-I		List-II	
A)	$HClO_4$	p)	1
B)	$HClO_3$	q)	2
C)	$HClO_2$	r)	0
D)	$HClO$	s)	3

A) $A - s, B - r, C - q, D - p$ B) $A - s, B - q, C - p, D - r$

C) $A - r, B - p, C - s, D - q$ D) $A - r, B - p, C - q, D - s$

53. Match the following

List-I		List-II	
A)	XeF_5^{\ominus}	p)	Octahedral
B)	$HXeO_4^{\ominus}$	q)	Pentagonal planar
C)	XeO_2F_4	r)	See-saw
D)	XeF_5^{\oplus}	s)	Square pyramidal

A) $A - q, B - p, C - r, D - s$ B) $A - q, B - r, C - p, D - s$

C) $A - r, B - p, C - s, D - q$ D) $A - q, B - r, C - s, D - p$

54. Match the following halogens with their properties

List-I		List-II	
A)	F_2	p)	Maximum enthalpy of dissociation
B)	Cl_2	q)	Corrosive liquid
C)	Br_2	r)	Maximum solubility in water
D)	I_2	s)	Minimum standard reduction potential

A) $A - s, B - p, C - q, D - r$ B) $A - r, B - p, C - s, D - q$

C) $A - r, B - p, C - q, D - s$ D) $A - s, B - q, C - p, D - r$

**THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

JEE MAIN
2023

SINGARAJU
VENKAT KOURINDYA
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
B.Sc.-12th Class
300
300
MARKS



RANK

JEE Advanced
2023

VAVILALA
CHIDOVAS REDDY
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
B.Sc.-12th Class
341
360
MARKS



RANK

NEET
2023

BORA VARUN
CHAKRAVANTHI
SRI CHAITANYA
EDUCATIONAL INSTITUTIONS
B.Sc.-12th Class
720
720
MARKS



RANK



Sri Chaitanya
Junior Colleges & Coaching Centres

Infinity Learn
BY SRI CHAITANYA



WITH ALL INDIA RANK 1 IN JEE ADVANCED 2023
SRI CHAITANYA
STANDS AT THE TOP

SEIZES 5 RANKS IN TOP 10 IN ALL-INDIA OPEN CATEGORY

ANDHRA PRADESH STATE TOPPER

VAVILALA CHIDVILAS REDDY
HT.No. 236165088
Sri Chaitanya
6th - 12th Class

RANK



RAMESH SURYA THEJA
HT.No. 23601571



KALRA RISHI
HT.No. 237047209



RAGHAV GOYAL
HT.No. 237020329



BIKKINA ABHINAV CHOWDARY
HT.No. 236103128



RANK

32 TOP RANKS BELOW 100 IN ALL-INDIA OPEN CATEGORY

RANK 12 DASARI SAKETH NAIDU HT.No. 236060148	RANK 14 GUTHIKONDA ABHIRAM HT.No. 23612921	RANK 17 P GNANA KOUSIK REDDY HT.No. 238133520	RANK 18 D VENKATA YUGEESH HT.No. 238049359	RANK 19 MOULIK JINDAL HT.No. 237021004	RANK 22 D PRATAP SINGH HT.No. 232006139	RANK 23 SHAIK SAHIL CHANDA HT.No. 236132870	RANK 26 SONI MAYANK HT.No. 232097025	RANK 33 GOVAL NAMAN HT.No. 232029345
RANK 37 SUMEDH SS HT.No. 231029088	RANK 38 V DEERAJ SATVIK REDDY HT.No. 236057151	RANK 44 V VERGIYA KAUSHAL HT.No. 231004058	RANK 46 SUTHAR HARSHAL HT.No. 231005060	RANK 58 ABHI JAIN HT.No. 232091036	RANK 61 RIDAM JAIN HT.No. 232056802	RANK 62 SUTHAR HARSH HT.No. 231010028	RANK 63 SOJITRA DIPEN HT.No. 231017001	RANK 65 D PANENDRANADHA REDDY HT.No. 238062088
RANK 67 GUNDA SUSHANTH HT.No. 236128732	RANK 68 HARSH TAYA HT.No. 237021776	RANK 69 KRISH GUPTA HT.No. 232068189	RANK 73 KUNCHAKURTHY VIVEK HT.No. 236134093	RANK 81 N DHARMA TEJA REDDY HT.No. 236050138	RANK 84 S VENKAT KOUNDINYA HT.No. 236130678	RANK 87 MRUNAL SHRIKANT HT.No. 23104138	RANK 92 SAMOTA APURVA HT.No. 232099270	RANK 97 N PRINCE BRANHAM REDDY HT.No. 236064112

BELOW
20 »
All India Open
Category Ranks

10
RANKS (50%)

BELOW
100 »
All India Open
Category Ranks

32
All India Open
Category Ranks

BELOW
1000 »
All India Open
Category Ranks

181
All India
Category Ranks
Count

BELOW
100 »
All India
Category Ranks
Count

89
All India
Category Ranks
Count

BELOW
1000 »
All India
Category Ranks
Count

699

NUMBER OF RANKS QUALIFIED
3,621*

ADMISSIONS OPEN » JEE ADVANCED LONG-TERM 2024

040 66 06 06 06

www.srichaitanya.net



Scan QR Code for
JEE Advanced Offers