

3. Let

$$A = \{(\alpha, \beta) \in \mathbf{R} \times \mathbf{R} : |\alpha - 1| \leq 4 \text{ and } |\beta - 5| \leq 6\}$$

and

$$B = \{(\alpha, \beta) \in \mathbf{R} \times \mathbf{R} : 16(\alpha - 2)^2 + 9(\beta - 6)^2 \leq 144\}.$$

Then

$$(1) B \subset A$$

$$(2) A \cup B = \{(x, y) : -4 \leq x \leq 4, -1 \leq y \leq 11\}$$

$$(3) \text{neither } A \subset B \text{ nor } B \subset A$$

$$(4) A \subset B$$

Ans. (1)


Sol. $A : |x-1| \leq 4 \text{ and } |y-5| \leq 6$

$$\Rightarrow -4 \leq x - 1 \leq 4 \Rightarrow -3 \leq x \leq 5$$

$$\Rightarrow -1 \leq y \leq 11$$

$$B : 16(x - 2)^2 + 9(y - 6)^2 \leq 144$$

$$B : \frac{(x-2)^2}{9} + \frac{(y-6)^2}{16} \leq 1$$



From Diagram $B \subset A$

4. If the range of the function $f(x) = \frac{5-x}{x^2-3x+2}$,

$x \neq 1, 2$, is $(-\infty, \alpha] \cup [\beta, \infty)$,

then $\alpha^2 + \beta^2$ is equal to :

$$(1) 190 \quad (2) 192$$

$$(3) 188 \quad (4) 194$$

Ans. (4)

Sol. $y = \frac{5-x}{x^2-3x+2}$
 $yx^2 - 3xy + 2y + x - 5 = 0$
 $yz^2 + (-3y + 1)x + (2y - 5) = 0$

Case I : If $y = 0$ (Accepted)

$$\Rightarrow x = 5$$

Case II : If $y \neq 0$

$$D \geq 0$$

$$(-3y + 1)^2 - 4(y)(2y - 5) \geq 0$$

$$9y^2 + 1 - 6y - 8y^2 + 20y \geq 0$$

$$y^2 + 14y + 1 \geq 0$$

$$(y + 7)^2 - 48 \geq 0$$

$$|y + 7| \geq 4\sqrt{3}$$

$$\Rightarrow y + 7 \geq 4\sqrt{3} \text{ or } y + 7 \leq -4\sqrt{3}$$

$$\Rightarrow y \geq 4\sqrt{3} - 7 \text{ or } y \leq -4\sqrt{3} - 7$$

From Case I and Case II

$$y \in (-\infty, -4\sqrt{3} - 7] \cup [4\sqrt{3} - 7, \infty)$$

$$\text{So } \alpha = -4\sqrt{3} - 7$$

$$\beta = 4\sqrt{3} - 7$$

$$\Rightarrow a^2 + b^2 = (-4\sqrt{3} - 7)^2 + (4\sqrt{3} - 7)^2 \\ = 2(48 + 49) \\ = 194$$

5. A bag contains 19 unbiased coins and one coin with head on both sides. One coin drawn at random is tossed and head turns up. If the probability that the drawn coin was unbiased, is $\frac{m}{n}$, $\text{gcd}(m, n) = 1$, then $n^2 - m^2$ is equal to :

$$(1) 80 \quad (2) 60$$

$$(3) 72 \quad (4) 64$$

Ans. (1)

Sol. $P(H) = \frac{19}{20} \times \frac{1}{2} + \frac{1}{20} \times 1$ Head occurs
Selection of unbiased Selection of biased coin

$$\text{Required probability} = \frac{\frac{19}{20} \times \frac{1}{2}}{\frac{19}{20} \times \frac{1}{2} + \frac{1}{20} \times 1} = \frac{19}{21}$$

$$\therefore \frac{m}{n} = \frac{19}{21}$$

$$\Rightarrow n^2 - m^2 = 441 - 361 = 80$$



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now

Ans. (2)

Sol. Length of LR = $\frac{2b^2}{a} = 10 \Rightarrow [5a = b^2] \dots\dots(1)$

$$f(t) = t^2 + t + \frac{11}{12}$$

$$\frac{df(t)}{dt} = 2t + 1 = 0 \Rightarrow t = -\frac{1}{2}$$

$$\text{Min value of } f(t) = \left(\frac{-1}{2}\right)^2 + \left(\frac{-1}{2}\right) + \frac{11}{12}$$

$$= \frac{1}{4} - \frac{1}{2} + \frac{11}{12} = \frac{3-6+11}{12} = \frac{8}{12} = \frac{2}{3} = e$$

$$e^2 = \frac{1-b^2}{a^2} \Rightarrow \frac{4}{9} = \frac{1-b^2}{a^2}$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{1-4}{a} = \frac{5}{a} \Rightarrow [b^2 = \frac{5a^2}{a}] \dots\dots(2)$$

From (1) & (2)

$$5a = \frac{5a^2}{a} \Rightarrow a = 9, b = \sqrt{45} = 3\sqrt{5}$$

$$\therefore a^2 + b^2 = 81 + 45 = 126$$

13. Let $y = y(x)$ be the solution of the differential equation $(x^2+1)y' - 2xy = (x^4 + 2x^2 + 1) \cos x$,

$y(0) = 1$. Then $\int_{-3}^3 y(x) dx$ is :

(1) 24

(3) 30

(2) 36

(4) 18

Ans. (1)

Sol. $(x^2 + 1)\frac{dy}{dx} - 2xy = (x^4 + 2x^2 + 1)\cos x$

$$\frac{dy}{dx} - \left(\frac{2x}{x^2 + 1}\right)y = \frac{(x^2 + 1)^2 \cos x}{cx^2 + 1} = (x^2 + 1)\cos x$$

(Linear D.E.)

$$P = \frac{-2x}{x^2 + 1}, Q = (x^2 + 1)\cos x$$

$$I.F = e^{\int P dx} = e^{\int \frac{-2x}{x^2 + 1} dx} = \frac{1}{x^2 + 1}$$

$$y \cdot \frac{1}{x^2 + 1} = \int (x^2 + 1)\cos x \cdot \frac{1}{x^2 + 1} dx$$

$$\frac{y}{x^2 + 1} = \sin x + c \Rightarrow y \cos = 1 \Rightarrow c = 1$$

$$y = (x^2 + 1)(\sin x + 1)$$

$$\int_{-3}^3 y dx = \int_{-3}^3 (x^2 + 1)(\sin x + 1) dx$$

$$dx = \int_{-3}^3 x^2 \sin x + x^2 \cos x + 1 dx$$

$$\Rightarrow \int_{-3}^3 x^2 \sin x dx + \int_{-3}^3 x^2 \cos x dx + \int_{-3}^3 1 dx \\ = 0 + 18 + 0 + 6 = 24$$

14. If the equation of the line passing through the point

$$\left(0, -\frac{1}{2}, 0\right) \text{ and perpendicular to the lines} \\ \vec{r} = \lambda(\hat{i} + \hat{a}j + \hat{b}k) \text{ and}$$

$$\vec{r} = (\hat{i} - \hat{j} - 6\hat{k}) + \mu(-\hat{b}i + \hat{a}j + 5\hat{k})$$

is $\frac{x-1}{-2} = \frac{y+4}{d} = \frac{z-c}{-4}$, then $a + b + c + d$ is equal to :

(1) 10 (2) 14

(3) 13 (4) 12

Ans. (2)

Sol. Line is \perp^r to 2 line \Rightarrow line will be parallel to

$$(\hat{i} + \hat{a}j + \hat{b}k) \times (-\hat{b}i + \hat{a}j + 5\hat{k})$$

Parallel vector along the required line is

$$\hat{i}(5a - ab) - \hat{j}(b^2 + 5) + \hat{k}(a + ab)$$

Dr's of required line $\alpha (5a - ab), -(b^2 + 5), (a + ab)$

Also Dr's of required line $\alpha -2, d, -4$

$$\therefore \frac{5a - ab}{-2} = \frac{-(b^2 + 5)}{d} = \frac{a + ab}{-4} \dots\dots(1)$$

Also point $\left(0, -\frac{1}{2}, 0\right)$ will lie on $\frac{x-1}{-2} = \frac{y+4}{d} = \frac{z-c}{-4}$

$$\frac{0-1}{-2} = \frac{\frac{-1}{2} + 4}{d} = \frac{0-c}{-4} \Rightarrow d = 7, c = 2$$

$$\text{From (1)} \frac{5a - ab}{-2} = \frac{-b^2 - 5}{7} = \frac{a + ab}{-4}$$

$$\frac{5a - ab}{-2} = \frac{a + ab}{-4}; \frac{-b^2 - 5}{7} = \frac{a + ab}{-4}$$

$$-20a + 4ab = -2a - 2ab \quad | 4b^2 + 20 = 70 + 7ab$$

$$18a = 6ab$$

$$36 + 20 = 70 + 21a$$

$$b = 3$$

$$56 = 28a \Rightarrow a = 2$$

$$a + b + c + d = 2 + 3 + 2 + 7 = 14$$



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now



Level up your prep for JEE with our **LIVE JEE Courses!**



LIVE classes with top Kota faculty



ALLEN's **study material**



Tests with **national benchmarking**



ALLEN App Advantage: **24/7 doubt support,**
Custom Practice & more

Enrol Now



Secure up to

90% scholarship*

on our Online Courses!

*based on your
JEE Main 2025 scores!*

Enrol Now

