

JEE-MAIN EXAMINATION – JANUARY 2025(HELD ON THURSDAY 23rd JANUARY 2025)

TIME : 3 : 00 PM TO 6 : 00 PM

CHEMISTRY**TEST PAPER WITH SOLUTION****SECTION-A**

51. The effect of temperature on spontaneity of reactions are represented as:

	ΔH	ΔS	Temperature	Spontaneity
(A)	+	-	any T	Non spontaneous
(B)	+	+	low T	spontaneous
(C)	-	-	low T	Non spontaneous
(D)	-	+	any T	spontaneous

(1) (B) and (D) only
 (2) (A) and (D) only
 (3) (B) and (C) only
 (4) (A) and (C) only

Ans. (3)**Sol.** $\because \Delta G = \Delta H - T\Delta S$ For spontaneity of reaction : $\Delta G = -ve$

52. Standard electrode potentials for a few half cells are mentioned below:

$$E_{\text{Cu}^{2+}/\text{Cu}}^{\circ} = 0.34 \text{ V}, E_{\text{Zn}^{2+}/\text{Zn}}^{\circ} = -0.76 \text{ V}$$

$$E_{\text{Ag}^{+}/\text{Ag}}^{\circ} = 0.80 \text{ V}, E_{\text{Mg}^{2+}/\text{Mg}}^{\circ} = -2.37 \text{ V}$$

Which one of the following cells gives the most negative value of ΔG° ?

- (1) $\text{Zn}|\text{Zn}^{2+}(1\text{M})||\text{Ag}^{+}(1\text{M})|\text{Ag}$
 (2) $\text{Zn}|\text{Zn}^{2+}(1\text{M})||\text{Mg}^{2+}(1\text{M})|\text{Mg}$
 (3) $\text{Ag}|\text{Ag}^{+}(1\text{M})||\text{Mg}^{2+}(1\text{M})|\text{Mg}$
 (4) $\text{Cu}|\text{Cu}^{2+}(1\text{M})||\text{Ag}^{+}(1\text{M})|\text{Ag}$

Ans. (1)**Sol.** $\because \Delta G^{\circ} = -nFE^{\circ}$

$$\begin{aligned} \text{Option (1)} \quad E^{\circ} &= 0.8 + 0.76 \\ &= 1.56 \text{ V} \end{aligned}$$

$$\begin{aligned} \therefore \Delta G^{\circ} &= -2 \times F \times 1.56 \\ &= -3.12 \text{ V} \end{aligned}$$

$$\begin{aligned} \text{Option (2)} \quad E^{\circ} &= -2.37 + 0.76 \\ &= -1.61 \text{ V} \end{aligned}$$

$$\begin{aligned} \therefore \Delta G^{\circ} &= -2 \times F \times (-1.61) \\ &= +3.22 \text{ V} \end{aligned}$$

$$\begin{aligned} \text{Option (3)} \quad E^{\circ} &= -2.37 - 0.8 \\ &= -3.17 \text{ V} \end{aligned}$$

$$\begin{aligned} \therefore \Delta G^{\circ} &= -2 \times F \times (-3.17) \\ &= +6.34 \text{ V} \end{aligned}$$

$$\begin{aligned} \text{Option (4)} \quad E^{\circ} &= 0.8 - 0.34 \\ &= 0.46 \text{ V} \end{aligned}$$

$$\begin{aligned} \Delta G^{\circ} &= -2 \times F \times 0.46 \\ &= -0.92 \text{ V} \end{aligned}$$

53. The α - Helix and β - Pleated sheet structures of protein are associated with its:

- (1) quaternary structure
 (2) primary structure
 (3) secondary structure
 (4) tertiary structure


Ans. (3)**Sol.** α -helix and β -pleated sheet belongs to secondary structure of protein, which have hydrogen bonds.

Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!


Enrol Now

54. Given below are two statements:

Consider the following reaction




Statement (I) : In the case of formaldehyde



(H \rightarrow H), K is about 2280, due to small substituents, hydration is faster.

Statement (II) : In the case of trichloro



due to $-I$ effect of $-Cl$.

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) **Statement I** true but **Statement II** is false
- (2) Both **Statement I** and **Statement II** are true
- (3) **Statement I** is false but **Statement II** is true
- (4) Both **Statement I** and **Statement II** are false

Ans. (2)


Sol. $k_{eq} = 2280$ is for HCHO

$k_{eq} = 2000$ is for chloral

Both data is given in clayden and warren book.

$k_{eq} > 1$ because HCHO and chloral are more electrophilic.

55. Consider the reaction



The equation representing correct relationship between the degree of dissociation (x) of $\text{X}_2\text{Y(g)}$ with its equilibrium constant K_p is _____.

Assume x to be very very small.

$$(1) x = \sqrt[3]{\frac{2K_p}{P}}$$


$$(2) x = \sqrt[3]{\frac{2K_p^2}{P}}$$

$$(3) x = \sqrt[3]{\frac{K_p}{2P}}$$

$$(4) x = \sqrt[3]{\frac{K_p}{P}}$$

Ans. (2)

Sol. 1 mole



$$1-x \text{ mole} \quad x \text{ mole} \quad \frac{x}{2} \text{ mole}$$

$$\therefore P_{X_2Y} = \frac{1-x}{1+\frac{x}{2}} \times P$$

$$P_{X_2} = \frac{x}{1+\frac{x}{2}} \times P$$

$$P_{Y_2} = \frac{x/2}{1+\frac{x}{2}} \times P$$

$$\therefore K_p = \left(\frac{x}{1+\frac{x}{2}} \times P \right) \left(\frac{x}{2\left(1+\frac{x}{2}\right)} \times P \right)^{\frac{1}{2}} \left(\frac{1-x}{1+\frac{x}{2}} \times P \right)$$

$$\therefore K_p = \left(\frac{x}{1-x} \right) \left(\frac{x}{2\left(1+\frac{x}{2}\right)} \right)^{\frac{1}{2}} \times P^{\frac{1}{2}}$$

$\because x$ to be very very small

$$\therefore K_p = \frac{x^{3/2}}{(2)^2} \times P^{\frac{1}{2}}$$

$$\therefore x^{\frac{3}{2}} = \frac{K_p \times 2^{\frac{1}{2}}}{P^{\frac{1}{2}}}$$

$$\therefore x^3 = \frac{K_p^2 \times 2}{P}$$


$$x = \left(\frac{K_p^2 \times 2}{P} \right)^{\frac{1}{3}}$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!


Enrol Now

56. Identify A, B and C in the given below reaction sequence



- (1) PbCl_2 , PbSO_4 , PbCrO_4
- (2) PbS , PbSO_4 , PbCrO_4
- (3) PbS , PbSO_4 , $\text{Pb}(\text{CH}_3\text{COO})_2$
- (4) PbCl_2 , $\text{Pb}(\text{SO}_4)_2$, PbCrO_4

Ans. (2)



57. Given below are two statements:

Statement (I): The boiling points of alcohols and phenols increase with increase in the number of C-atoms.

Statement (II): The boiling points of alcohols and phenols are higher in comparison to other class of compounds such as ethers, haloalkanes.

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) Both **Statement I** and **Statement II** are false
- (2) **Statement I** is false but **Statement II** is true
- (3) **Statement I** is true but **Statement II** is false
- (4) Both **Statement I** and **Statement II** are true

Ans. (4)

Sol. $B.P. \propto M.W.$

$B.P. \propto$ Inter molecular hydrogen bonding

Alcohol & Phenol have intermolecular H-bonding

58. When a non-volatile solute is added to the solvent, the vapour pressure of the solvent decreases by 10 mm of Hg. The mole fraction of the solute in the solution is 0.2. What would be the mole fraction of the solvent if decrease in vapour pressure is 20 mm of Hg ?

- (1) 0.6
- (2) 0.4
- (3) 0.2
- (4) 0.8

Ans. (1)

Sol. $\because P^\circ - P \propto X_{\text{solute}}$

and $\therefore 10 \propto 0.2$

$$\therefore 20 \propto 0.4$$

$$\therefore X_{\text{solvent}} = 1 - X_{\text{solute}}$$

$$= 1 - 0.4$$

$$= 0.6$$

59. Given below are two statements:

Statement (I) : For a given shell, the total number of allowed orbitals is given by n^2 .

Statement (II) : For any subshell, the spatial orientation of the orbitals is given by $-l$ to $+l$ values including zero.

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) **Statement I** is true but **Statement II** is false
- (2) **Statement I** is false but **Statement II** is true
- (3) Both **Statement I** and **Statement II** are true
- (4) Both **Statement I** and **Statement II** are false

Ans. (3)

Sol. For a shell total number of orbitals = n^2


Magnetic quantum number have values $(-\ell$ to $+\ell$) including 0.



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now


60. The ascending order of relative rate of solvolysis of following compounds is



- (1) (D) < (A) < (B) < (C)
- (2) (C) < (B) < (A) < (D)
- (3) (D) < (B) < (A) < (C)
- (4) (C) < (D) < (B) < (A)

Ans. (1)

Sol. Solvolysis or $S_N1 \propto$ stability of carbocation
Stability order



61. Match List - I with List - II.





List - I (Isomers of $C_{10}H_{14}$)	List - II (Ozonolysis product)		
(A)	(I)		
(B)	(II)		
(C)	(III)		
(D)	(IV)		

Choose the **correct** answer from the options given below :


- (1) (A)-(II), (B)-(III), (C)-(I), (D)-(IV)
- (2) (A)-(III), (B)-(IV), (C)-(I), (D)-(II)
- (3) (A)-(III), (B)-(II), (C)-(I), (D)-(IV)
- (4) (A)-(I), (B)-(IV), (C)-(III), (D)-(II)

Ans. (2)

Sol. Ozonolysis product



62. Which of the following graphs most appropriately represents a zero order reaction ?



Ans. (2)




Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

Sol. For zero order reaction : $A \rightarrow P$

$$\text{Rate} = k$$

$$[A]_t = a_0 - kt$$



63. Match **List - I** with **List - II**.

List - I		List - II	
(A)	Bronze	(I)	Cu, Ni
(B)	Brass	(II)	Fe, Cr, Ni, C
(C)	UK silver coin	(III)	Cu, Zn
(D)	Stainless Steel	(IV)	Cu, Sn

Choose the **correct** answer from the options given below :

- (1) (A)-(IV), (B)-(II), (C)-(III), (D)-(I)
- (2) (A)-(IV), (B)-(III), (C)-(I), (D)-(II)
- (3) (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
- (4) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)

Ans. (2)

Sol. Bronze \rightarrow Cu, Sn

Brass \rightarrow Cu, Zn

UK silver coin \rightarrow Cu, Ni

Stainless steel \rightarrow Fe, Cr, Ni, C

64. Identify the coordination complexes in which the central metal ion has d^4 configuration.

- (A) $[\text{FeO}_4]^{2-}$
- (B) $[\text{Mn}(\text{CN})_6]^{3-}$
- (C) $[\text{Fe}(\text{CN})_6]^{3-}$
- (D) $\text{Cr}_2(\text{O}-\text{C}-\text{Me})_4(\text{H}_2\text{O})_2$
- (E) $[\text{NiF}_6]^{2-}$

Choose the **correct** answer from the options given below :

- (1) (C) and (E) only (2) (B), (C) and (D) only
- (3) (B) and (D) only (4) (A), (B) and (E) only

Ans. (3)

Sol. $\text{Fe}^{+6} = [\text{Ar}]3d^2$

$\text{Mn}^{+3} = [\text{Ar}]3d^4$

$\text{Fe}^{+3} = [\text{Ar}]3d^5$

$\text{Cr}^{+2} = [\text{Ar}]3d^4$

$\text{Ni}^{+4} = [\text{Ar}]3d^6$

65. Given below are the atomic numbers of some group 14 elements. The atomic number of the element with lowest melting point is :

- (1) 14
- (2) 6
- (3) 82
- (4) 50

Ans. (4)

Sol. Order of M.P. of group 14 : C > Si > Ge > Pb > Sn

element	M.P. (°C)
$Z = 6 = \text{C}$	3730
$Z = 14 = \text{Si}$	1410
$Z = 32 = \text{Ge}$	937
$Z = 50 = \text{Sn}$	232
$Z = 82 = \text{Pb}$	327

66. pH of water is 7 at 25°C. If water is heated to 80°C, its pH will :

- (1) Decrease
- (2) Remains the same
- (3) H^+ concentration increases, OH^- concentration decreases
- (4) Increase

Ans. (1)

Sol. With increase in temperature, K_w of water increases

So, degree of dissociation of water increase






\therefore pH as well as pOH of water decrease.



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

67. Identify the products [A] and [B], respectively in the following reaction :



Ans. (3)

Sol. A is phenol and B is para benzoquinone.

68. Consider a binary solution of two volatile liquid components 1 and 2 x_1 and y_1 are the mole fractions of component 1 in liquid and vapour phase, respectively. The slope and intercept of the linear plot of $\frac{1}{x_1}$ vs $\frac{1}{y_1}$ are given respectively as :

(1) $\frac{P_1^0}{P_2^0}, \frac{P_2^0 - P_1^0}{P_2^0}$

(2) $\frac{P_2^0}{P_1^0}, \frac{P_1^0 - P_2^0}{P_2^0}$

(3) $\frac{P_1^0}{P_2^0}, \frac{P_1^0 - P_2^0}{P_1^0}$

(4) $\frac{P_2^0}{P_1^0}, \frac{P_2^0 - P_1^0}{P_1^0}$

Ans. (1)

Sol. ∵ For liquid solution of two liquids '1' and '2'

$$P_1 = P_T y_1 = P_1^0 x_1$$

$$\therefore \frac{P_T}{x_1} = \frac{P_1^0}{y_1}$$

$$\therefore \frac{P_2^0 + x_1(P_1^0 - P_2^0)}{x_1} = \frac{P_1^0}{y_1}$$

$$\therefore \frac{P_2^0}{x_1} + (P_1^0 - P_2^0) = \frac{P_1^0}{y_1}$$

$$\therefore \frac{1}{x_1} = \left(\frac{P_1^0}{P_2^0} \right) \left(\frac{1}{y_1} \right) + \left(\frac{P_2^0 - P_1^0}{P_2^0} \right)$$

$$\therefore \text{Slope} = \left(\frac{P_1^0}{P_2^0} \right)$$

$$\therefore \text{Intercept} = \left(\frac{P_2^0 - P_1^0}{P_2^0} \right)$$

69. Given below are two statements about X-ray spectra of elements :



Statement (I) : A plot of \sqrt{v} (v = frequency of X-rays emitted) vs atomic mass is a straight line.

Statement (II) : A plot of v (v = frequency of X-rays emitted) vs atomic number is a straight line. In the light of the above statements choose the **correct** answer from the options given below :

- (1) **Statement I** is true but **Statement II** is false
- (2) Both **Statement I** and **Statement II** are true
- (3) Both **Statement I** and **Statement II** are false
- (4) **Statement I** is false but **Statement II** is true

Ans. (3)





Sol.



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!


Enrol Now

74. The bond dissociation enthalpy of X_2 , $\Delta H_{\text{bond}}^\circ$ calculated from the given data is _____ kJ mol⁻¹. (Nearest integer)



[Given : M^+X^- is a pure ionic compound and X forms a diatomic molecule X_2 is gaseous state]

Ans. (200)




$$\therefore \Delta H_f(MX) = \Delta H_{\text{sub}}(M) + \text{I.E.}(M) + \frac{1}{2}[\text{B.E.}(X-X)] + \text{EG}(X) + \text{L.E.}(MX)$$

$$-400 = (100) + (500) + \frac{1}{2}(\text{B.E.}) + (-300) + (-800)$$






$$\therefore \text{B.E.} = 200 \text{ kJ mole}^{-1}$$

75. A compound 'X' absorbs 2 moles of hydrogen and 'X' upon oxidation with $\text{KMnO}_4 | \text{H}^+$ gives



The total number of σ bonds present in the compound 'X' is _____.

Ans. (27)



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now



Level up your prep for JEE Adv. 2025 with our **Online Rank Booster Course!**



LIVE classes for JEE Main & Advanced



Soft copies of ALLEN's study material




Covers important questions across **Physics, Chemistry & Maths**



ALLEN App Advantage: **24/7 doubt support,
Custom Practice & more**

Enrol Now



Win up to
90% scholarship*
with the ALLEN Online Scholarship Test

at just **₹49/-**



1-hour online test
Can be taken from anywhere

Register Now

