


Sol.



Let intersection points of these two parabolas are $A(x_1, y_1)$ & $B(x_2, y_2)$

\therefore equation of parabola I and II are given below

$$\therefore (x - 4)^2 + (y - 3)^2 = x^2 \quad \dots\dots(1)$$

$$\& (x - 4)^2 + (y - 3)^2 = y^2 \quad \dots\dots(2)$$

Here $A(x_1, y_1)$ & $B(x_2, y_2)$ will satisfy with equation
Also from equations (1) & (2), we get $x = y \dots(3)$

Put $x = y$ in equation (1)

We get $x^2 - 14x + 25 = 0$

$$x_1 + x_2 = 14$$

$$x_1 x_2 = 25$$

$$\therefore AB^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$$

$$= 2(x_1 - x_2)^2$$

$$= 2[(x_1 + x_2)^2 - 4x_1 x_2]$$

$$= 192$$

4. Let ABC be a triangle formed by the lines $7x - 6y + 3 = 0$, $x + 2y - 31 = 0$ and $9x - 2y - 19 = 0$, Let the point (h, k) be the image of the centroid of ΔABC in the line $3x + 6y - 53 = 0$. Then $h^2 + k^2 + hk$ is equal to

(1) 37


(3) 40

(2) 47


(4) 36

Ans. (1)

Sol.



$$\therefore \text{centroid of } \Delta ABC = \left(\frac{9+3+5}{3}, \frac{11+4+13}{3} \right) \\ = \left(\frac{17}{3}, \frac{28}{3} \right)$$



Let image of centroid with respect to line mirror is (h, k)

$$\therefore \left(\frac{k - \frac{28}{3}}{h - \frac{17}{3}} \right) \left(-\frac{1}{2} \right) = -1$$

$$\& 3 \left(\frac{h + \frac{17}{3}}{2} \right) + 6 \cdot \left(\frac{\frac{k+28}{3}}{2} \right) = 53$$

Solving (1) & (2) we get $h = 3, k = 4$

$$\therefore h^2 + k^2 + hk = 37$$

5. Let $\vec{a} = 2\hat{i} - \hat{j} + 3\hat{k}$, $\vec{b} = 3\hat{i} - 5\hat{j} + \hat{k}$ and \vec{c} be a vector such that $\vec{a} \times \vec{c} = \vec{c} \times \vec{b}$ and $(\vec{a} + \vec{c}) \cdot (\vec{b} + \vec{c}) = 168$. Then the maximum value of $|\vec{c}|^2$ is :

(1) 77

(2) 462

(3) 308

(4) 154

Ans. (3)

Sol. $\vec{a} = 2\hat{i} - \hat{j} + 3\hat{k}$

$\vec{b} = 3\hat{i} - 5\hat{j} + \hat{k}$

$\vec{a} \times \vec{c} = \vec{c} \times \vec{b}$

$\vec{a} \times \vec{c} + \vec{b} \times \vec{c} = 0$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

9. The number of solutions of the equation

$$\left(\frac{9}{x} - \frac{9}{\sqrt{x}} + 2\right) \left(\frac{2}{x} - \frac{7}{\sqrt{x}} + 3\right) = 0$$

- (1) 2 (2) 4
 (3) 1 (4) 3

Ans. (2)

Sol. Consider $\frac{1}{\sqrt{x}} = \alpha$ $x > 0$

$$\{9\alpha^2 - 9\alpha + 2\} \{2\alpha^2 - 7\alpha + 3\} = 0$$

$$(3\alpha - 2)(3\alpha - 1)(\alpha - 3)(2\alpha - 1) = 0$$

$$\alpha = \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, 3$$

$$x = 9, 4, \frac{9}{4}, \frac{1}{9}$$

So, no. of solutions = 4

10. Let $y = y(x)$ be the solution of the differential equation

$$\cos x (\log_e(\cos x))^2 dy + (\sin x - 3y \sin x \ln(\cos x)) dx = 0,$$

$x \in \left(0, \frac{\pi}{2}\right)$. If $y\left(\frac{\pi}{4}\right) = \frac{-1}{\log_e 2}$, then $y\left(\frac{\pi}{6}\right)$ is :

$$(1) \frac{2}{\log_e(3) - \log_e(4)} \quad (2) \frac{1}{\log_e(4) - \log_e(3)}$$

$$(3) -\frac{1}{\log_e(4)} \quad (4) \frac{1}{\log_e(3) - \log_e(4)}$$

Ans. (4)

Sol.

$$\cos x (\ln(\cos x))^2 dy + (\sin x - 3y \sin x \ln(\cos x)) dx = 0$$

$$\cos x (\ln(\cos x))^2 \frac{dy}{dx} - 3 \sin x \ln(\cos x) y = -\sin x$$

$$\frac{dy}{dx} - \frac{3 \tan x}{\ln(\cos x)} y = \frac{-\tan x}{(\ln(\cos x))^2}$$

$$\frac{dy}{dx} + \frac{3 \tan x}{\ln(\sec x)} y = \frac{-\tan x}{(\ln(\sec x))^2}$$

$$\text{I.F.} = e^{\int \frac{3 \tan x}{\ln(\sec x)} dx} = (\ln(\sec x))^3$$

$$y \times (\ln(\sec x))^3 = - \int \frac{\tan x}{(\ln(\sec x))^2} (\ln(\sec x))^3 dx + C$$

$$y \times (\ln(\sec x))^3 = -\frac{1}{2} (\ln(\sec x))^2 + C$$

$$\text{Given : } x = \frac{\pi}{4}, y = -\frac{1}{\ln 2}$$

$$\frac{-1}{\ln 2} \times (\ln \sqrt{2})^3 = -\frac{1}{2} \times (\ln \sqrt{2})^2 + C$$

$$\Rightarrow \frac{-1}{8 \ln 2} \times (\ln 2)^3 = \frac{-1}{2} \times \frac{1}{4} (\ln 2)^2 + C$$

$$-\frac{1}{8} (\ln 2)^2 = \frac{-1}{8} (\ln 2)^2 + C$$

$$\Rightarrow C = 0$$

$$\therefore y (\ln(\sec x))^3 = \frac{-1}{2} (\ln(\sec x))^2 + 0$$

$$y = \frac{-1}{2 \ln(\sec x)}$$

$$y = \frac{1}{2 \ln(\cos x)}$$

$$\therefore y\left(\frac{\pi}{6}\right) = \frac{1}{2 \ln\left(\cos \frac{\pi}{6}\right)}$$

$$= \frac{1}{2 \ln\left(\frac{\sqrt{3}}{2}\right)}$$

$$= \frac{1}{2 \left(\frac{1}{2} \ln 3 - \ln 2 \right)}$$

$$= \frac{1}{\ln 3 - \ln 4}$$

Option (4)



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

14. Let $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} + 7\hat{j} + 3\hat{k}$. Let

$$L_1 : \vec{r} = (-\hat{i} + 2\hat{j} + \hat{k}) + \lambda \vec{a}, \lambda \in \mathbb{R} \text{ and}$$

$L_2 : \vec{r} = (\hat{j} + \hat{k}) + \mu \vec{b}, \mu \in \mathbb{R}$ be two lines. If the line L_3 passes through the point of intersection of L_1 and L_2 , and is parallel to $\vec{a} + \vec{b}$, then L_3 passes through the point:

- (1) (8, 26, 12) (2) (2, 8, 5)
 (3) (-1, -1, 1) (4) (5, 17, 4)

Ans. (1)

$$\text{Sol. } L_1 : \vec{r} = (-\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} + 2\hat{j} + \hat{k})$$

$$\Rightarrow \vec{r} = (\lambda - 1)\hat{i} + 2(\lambda + 1)\hat{j} + (\lambda + 1)\hat{k}$$

$$L_2 : \vec{r} = (\hat{j} + \hat{k}) + \mu(2\hat{i} + 7\hat{j} + 3\hat{k})$$

$$\Rightarrow \vec{r} = 2\mu\hat{i} + (1 + 7\mu)\hat{j} + (1 + 3\mu)\hat{k}$$

For point of intersection equating respective components

$$\Rightarrow \lambda - 1 = 2\mu \quad \dots(1)$$

$$2(\lambda + 1) = 1 + 7\mu \quad \dots(2)$$

$$\lambda + 1 = 1 + 3\mu \quad \dots(3)$$

We get

$$\Rightarrow \lambda = 3 \text{ and } \mu = 1$$

$$\Rightarrow \vec{a} + \vec{b} = 3\hat{i} + 9\hat{j} + 4\hat{k}$$

$$L_3 : \vec{r} = 2\hat{i} + 8\hat{j} + 4\hat{k} + \alpha(3\hat{i} + 9\hat{j} + 4\hat{k})$$

$$\text{For } \alpha = 2, \vec{r} = 8\hat{i} + 26\hat{j} + 12\hat{k}$$

15. The value of $\lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 5}{(k+3)!} \right)$ is :

(1) $\frac{4}{3}$ (2) 2

(3) $\frac{7}{3}$ (4) $\frac{5}{3}$

Ans. (4)

$$\text{Sol. } \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 5}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 6 - 1}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(k+1)(k+2)(k+3) - 1}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(k+1)(k+2)(k+3)}{(k+3)!} - \frac{1}{(k+3)!}$$

$$= \lim_{k=1}^n \left(\frac{1}{k!} - \frac{1}{(k+3)!} \right)$$

$$= \lim_{k=1} \left(\frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!} - \frac{1}{4!} - \frac{1}{5!} - \frac{1}{6!} - \dots - \frac{1}{(n+3)!} \right)$$

$$= \frac{1}{1} + \frac{1}{2} + \frac{1}{6} = \frac{10}{6} = \frac{5}{3}$$

16. The integral $80 \int_0^{\frac{\pi}{4}} \left(\frac{\sin \theta + \cos \theta}{9 + 16 \sin 2\theta} \right) d\theta$ is equal to :

(1) $3 \log_e 4$

(3) $4 \log_e 3$

(2) $6 \log_e 4$

(4) $2 \log_e 3$

Ans. (3)

$$\text{Sol. } I = 80 \int_0^{\frac{\pi}{4}} \left(\frac{\sin \theta + \cos \theta}{9 + 16(2 \sin \theta \cdot \cos \theta)} \right) d\theta$$

$$= 80 \int_0^{\frac{\pi}{4}} \frac{\sin \theta + \cos \theta}{9 - 16(1 - 2 \sin \theta \cdot \cos \theta - 1)} d\theta$$

$$= 80 \int_0^{\frac{\pi}{4}} \frac{\sin \theta + \cos \theta}{9 + 16 - 16(\sin \theta - \cos \theta)^2} d\theta$$

Let $\sin \theta - \cos \theta = t$

$$(\cos \theta + \sin \theta) d\theta = dt$$

$$= 80 \int_{-1}^0 \frac{dt}{25 - 16t^2}$$

$$= \frac{80}{16} \int_{-1}^0 \frac{dt}{\left(\frac{5}{4}\right)^2 - t^2}$$

$$= \frac{5}{2 \left(\frac{5}{4}\right)} \ln \left(\frac{\frac{5}{4} + t}{\frac{5}{4} - t} \right) \Big|_{-1}^0$$

$$= 2 \ln(1) + 4 \ln 3$$

$$= 4 \ln 3$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

Sol. $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$

$$A^2 = \begin{bmatrix} 3 & -2 \\ 2 & -1 \end{bmatrix}, A^3 = \begin{bmatrix} 4 & -3 \\ 3 & -2 \end{bmatrix}, A^4 = \begin{bmatrix} 5 & -4 \\ 4 & -3 \end{bmatrix}$$

and so on

$$A^6 = \begin{bmatrix} 7 & -6 \\ 6 & -5 \end{bmatrix}$$

$$A^m = \begin{bmatrix} m+1 & -m \\ m & -m-1 \end{bmatrix},$$

$$A^{m^2} = \begin{bmatrix} m^2+1 & -m^2 \\ m^2 & -(m^2-1) \end{bmatrix}$$

$$A^{m^2} + A^m = 3I - A^{-6}$$

$$\begin{bmatrix} m+1 & -m^2 \\ m^2 & -(m^2-1) \end{bmatrix} + \begin{bmatrix} m+1 & -m \\ m & -(m-1) \end{bmatrix}$$

$$= 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} -5 & 6 \\ -6 & 7 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & -6 \\ 6 & -4 \end{bmatrix}$$

$$= m^2 + 1 + m + 1 = 8$$

$$= m^2 + m - 6 = 0 \Rightarrow m = -3, 2$$

$$n(s) = 2$$

23. Let $[t]$ be the greatest integer less than or equal to t . Then the least value of $p \in \mathbb{N}$ for which

$$\lim_{x \rightarrow 0^+} \left(x \left(\left[\frac{1}{x} \right] + \left[\frac{2}{x} \right] + \dots + \left[\frac{p}{x} \right] \right) - x^2 \left(\left[\frac{1}{x^2} \right] + \left[\frac{2^2}{x^2} \right] + \dots + \left[\frac{9^2}{x^2} \right] \right) \right) \geq 1$$

is equal to ____.

Ans. (24)

$$\lim_{x \rightarrow 0^+} \left(x \left(\left[\frac{1}{x} \right] + \left[\frac{2}{x} \right] + \dots + \left[\frac{p}{x} \right] \right) - x^2 \left(\left[\frac{1}{x^2} \right] + \left[\frac{2^2}{x^2} \right] + \dots + \left[\frac{9^2}{x^2} \right] \right) \right) \geq 1$$

$$(1 + 2 + \dots + p) - (1^2 + 2^2 + \dots + 9^2) \geq 1$$

$$\frac{p(p+1)}{2} - \frac{9 \cdot 10 \cdot 19}{6} \geq 1$$

$$p(p+1) \geq 572$$

Least natural value of p is 24

24. The number of 6-letter words, with or without meaning, that can be formed using the letters of the word MATHS such that any letter that appears in the word must appear at least twice, is 4 ____.

Ans. (1405)

- Sol. (i) Single letter is used, then no. of words = 5
(ii) Two distinct letters are used, then no. of words

$${}^5C_2 \times \left(\frac{6!}{2!4!} \times 2 + \frac{6!}{3!3!} \right) = 10(30 + 20) = 500$$

- (iii) Three distinct letters are used, then no. of words

$${}^5C_3 \times \frac{6!}{2!2!2!} = 900$$

Total no. of words = 1405

25. Let $S = \{x : \cos^{-1} x = \pi + \sin^{-1} x + \sin^{-1}(2x+1)\}$.

Then $\sum_{x \in S} (2x-1)^2$ is equal to ____.

Ans. (5)

Sol. $\cos^{-1} x = \pi + \sin^{-1} x + \sin^{-1}(2x+1)$

$$2\cos^{-1} x - \sin^{-1}(2x+1) = \frac{3\pi}{2}$$

$$2\alpha - \beta = \frac{3\pi}{2} \text{ where } \cos^{-1} x = \alpha, \sin^{-1}(2x+1) = \beta$$

$$2\alpha = \frac{3\pi}{2} + \beta$$

$$\cos 2\alpha = \sin \beta$$

$$2\cos^2 \alpha - 1 = \sin \beta$$

$$2x^2 - 1 = 2x + 1$$

$$x^2 - x - 1 = 0$$

$$\Rightarrow n = \frac{1 \pm \sqrt{5}}{2} = \begin{cases} n = \frac{1 + \sqrt{5}}{2} \text{ rejected} \\ n = \frac{1 - \sqrt{5}}{2} \end{cases}$$

$$\therefore 4x^2 - 4x = 4$$

$$(2x-1)^2 = 5$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

[Enrol Now](#)



Level up your prep for JEE Adv. 2025 with our **Online Rank Booster Course!**



LIVE classes for JEE Main & Advanced



Soft copies of ALLEN's study material




Covers important questions across **Physics, Chemistry & Maths**



ALLEN App Advantage: **24/7 doubt support,
Custom Practice & more**

Enrol Now



Win up to
90% scholarship*
with the ALLEN Online Scholarship Test

at just **₹49/-**



1-hour online test
Can be taken from anywhere

Register Now

