

Sol. $f_1(x) = \log_5(18x - x^2 - 77)$

$$\therefore 18x - x^2 - 77 > 0$$

$$x^2 - 18x + 77 < 0$$


$$x \in (7, 11) \quad \alpha = 7, \beta = 11$$

$$f_2(x) = \log_{(x-1)}\left(\frac{2x^2+3x-2}{x^2-3x-4}\right)$$

$$\therefore x-1 > 0, x-1 \neq 1, \frac{2x^2+3x-2}{x^2-3x-4} > 0$$

$$x > 1, x \neq 2, \frac{(2x-1)(x+2)}{(x-4)(x+1)} > 0$$

$$x > 1, x \neq 2,$$



$$\therefore x \in (4, \infty)$$

$$\therefore \gamma = 4$$

$$\therefore \alpha^2 + \beta^2 + \gamma^2 = 49 + 121 + 16 = 186$$

5. Let the function $f(x) = (x^2 - 1)|x^2 - ax + 2| + \cos|x|$ be not differentiable at the two points $x = \alpha = 2$ and $x = \beta$. Then the distance of the point (α, β) from the line $12x + 5y + 10 = 0$ is equal to :

$$(1) 3$$

$$(2) 4$$

$$(3) 2$$

$$(4) 5$$

Ans. Allen Ans. (BONUS)

NTA Ans. (1)

Sol. $\cos|x|$ is always differentiable

\therefore we have to check only for $|x^2 - ax + 2|$

\therefore Not differentiable at

$$x^2 - ax + 2 = 0$$

One root is given, $\alpha = 2$

$$\therefore 4 - 2a + 2 = 0$$

$$a = 3$$

$$\therefore$$
 other root $\beta = 1$

but for $x = 1$ $f(x)$ is differentiable

(Drop)

- 6.** Let a straight line L pass through the point $P(2, -1, 3)$ and be perpendicular to the lines $\frac{x-1}{2} = \frac{y+1}{1} = \frac{z-3}{-2}$ and $\frac{x-3}{1} = \frac{y-2}{3} = \frac{z+2}{4}$.

If the line L intersects the yz-plane at the point Q, then the distance between the points P and Q is :

$$(1) 2 \quad (2) \sqrt{10}$$

$$(3) 3 \quad (4) 2\sqrt{3}$$

Ans. (3)

Sol. Vector parallel to 'L'

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 1 & 3 & 4 \end{vmatrix} = 10\hat{i} - 10\hat{j} + 5\hat{k}$$

$$= 5(2\hat{i} - 2\hat{j} + \hat{k})$$

Equation of 'L'

$$\frac{x-2}{2} = \frac{y+1}{-2} = \frac{z-3}{1} = \lambda \text{ (say)}$$

Let $Q(2\lambda + 2, -2\lambda - 1, \lambda + 3)$

$$\Rightarrow 2\lambda + 2 = 0 \Rightarrow \lambda = -1$$

$$\Rightarrow Q(0, 1, 2)$$

$$d(P, Q) = 3$$

- 7.** Let $S = \mathbb{N} \cup \{0\}$. Define a relation **R** from S to R

by :

$$R = \left\{ (x, y) : \log_e y = x \log_e \left(\frac{2}{5}\right), x \in S, y \in R \right\}.$$

Then, the sum of all the elements in the range of **R**

is equal to

$$(1) \frac{3}{2} \quad (2) \frac{5}{3}$$

$$(3) \frac{10}{9} \quad (4) \frac{5}{2}$$

Ans. (2)




Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

Sol. $S = \{0, 1, 2, 3, \dots\}$

$$\log_e y = x \log_e \left(\frac{2}{5}\right)$$

$$\Rightarrow y = \left(\frac{2}{5}\right)^x$$



Required

$$\text{Sum} = 1 + \left(\frac{2}{5}\right)^1 + \left(\frac{2}{5}\right)^2 + \left(\frac{2}{5}\right)^3 + \dots = \frac{1}{1 - \frac{2}{5}} = \frac{5}{3}$$

8. Let the line $x + y = 1$ meet the axes of x and y at A and B , respectively. A right angled triangle AMN is inscribed in the triangle OAB , where O is the origin and the points M and N lie on the lines OB and AB , respectively. If the area of the triangle AMN is $\frac{4}{9}$ of the area of the triangle OAB and

$AN : NB = \lambda : 1$, then the sum of all possible value(s) of λ is :

(1) $\frac{1}{2}$


(2) $\frac{13}{6}$

(3) $\frac{5}{2}$

(4) 2

Ans. (4)

Sol.



$$\text{Area of } \triangle OAB = \frac{1}{2}$$

$$\text{Area of } \triangle AMN = \frac{4}{9} \times \frac{1}{2} = \frac{2}{9}$$

Equation of AB is $x + y = 1$

$OA = 1$, $AM = \sec(45^\circ - \theta)$

$AN = \sec(45^\circ - \theta) \cos \theta$

$MN = \sec(45^\circ - \theta) \sin \theta$

$$\text{Ar}(\triangle AMN) = \frac{1}{2} \times \sec^2(45^\circ - \theta) \sin \theta \cos \theta = \frac{2}{9}$$

$$\Rightarrow \tan \theta = 2, \frac{1}{2}$$

$\tan \theta = 2$ is rejected

$$\frac{AN}{NB} = \frac{\lambda}{1} = \cot \theta = 2$$

9. If $\alpha x + \beta y = 109$ is the equation of the chord of the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$, whose mid point is $\left(\frac{5}{2}, \frac{1}{2}\right)$, then $\alpha + \beta$ is equal to


(1) 37

(2) 46

(3) 58

(4) 72

Ans. (3)



Equation of chord $T = S_1$

$$\frac{5}{2} \left(\frac{x}{9}\right) + \frac{1}{2} \left(\frac{y}{4}\right) = \frac{25}{36} + \frac{1}{16}$$

$$\Rightarrow \frac{5x}{18} + \frac{y}{8} = \frac{100+9}{144} = \frac{109}{144}$$

$$\Rightarrow 40x + 18y = 109$$

$$\Rightarrow \alpha = 40, \beta = 18$$

$$\Rightarrow \alpha + \beta = 58$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

10. If all the words with or without meaning made using all the letters of the word "KANPUR" are arranged as in a dictionary, then the word at 440th position in this arrangement, is :

- (1) PRNAKU (2) PRKANU
 (3) PRKAUN (4) PRNAUK

Ans. (3)

Sol. A, K, N, P, R, U

$$\boxed{A} \dots = |5| = 120$$

$$\boxed{K} \dots = |5| = 120$$

$$\boxed{N} \dots = |5| = 120$$

$$\boxed{P} \boxed{A} \dots = |4| = 24$$

$$\boxed{P} \boxed{K} \dots = |4| = 24$$

$$\boxed{P} \boxed{N} \dots = |4| = 24$$

$$\boxed{P} \boxed{R} \boxed{A} \dots = |3| = 6$$

$$\boxed{P} \boxed{R} \boxed{K} \boxed{A} \boxed{N} \boxed{U} = 1$$

$$\boxed{P} \boxed{R} \boxed{K} \boxed{A} \boxed{U} \boxed{N} = 1$$

$$\text{Total} = 440$$

$$\Rightarrow 440^{\text{th}} \text{ word}$$

11. Let α, β ($\alpha \neq \beta$) be the values of m , for which the equations $x + y + z = 1$; $x + 2y + 4z = m$ and $x + 4y + 10z = m^2$ have infinitely many solutions.

Then the value of $\sum_{n=1}^{10} (n^\alpha + n^\beta)$ is equal to :

- (1) 440 (2) 3080
 (3) 3410 (4) 560

Ans. (1)

$$\text{Sol. } \Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 10 \end{vmatrix} = 1(20 - 16) - 1(10 - 4) + 1(4 - 2) \\ = 4 - 6 + 2 = 0$$

For infinite solutions

$$\Delta_x = \Delta_y = \Delta_z = 0$$

$$m^2 - 3x + 2 = 0$$

$$m = 1, 2$$

$$\alpha = 1, \beta = 2$$

$$\therefore \sum_{n=1}^{10} (n^\alpha + n^\beta) = \sum_{n=1}^{10} n^1 + \sum_{n=1}^{10} n^2$$

$$= \frac{10(11)}{2} + \frac{10(11)(21)}{6}$$

$$= 55 + 385$$

$$= 440$$

12. Let $A = [a_{ij}]$ be a matrix of order 3×3 , with $a_{ij} = (\sqrt{2})^{i+j}$. If the sum of all the elements in the third row of A^2 is $\alpha + \beta\sqrt{2}$, $\alpha, \beta \in \mathbb{Z}$, then $\alpha + \beta$ is equal to

- (1) 280 (2) 168
 (3) 210 (4) 224

Ans. (4)

$$\text{Sol. } A = \begin{bmatrix} (\sqrt{2})^2 & (\sqrt{2})^3 & (\sqrt{2})^4 \\ (\sqrt{2})^3 & (\sqrt{2})^4 & (\sqrt{2})^5 \\ (\sqrt{2})^4 & (\sqrt{2})^5 & (\sqrt{2})^6 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 2\sqrt{2} & 4 \\ 2\sqrt{2} & 4 & 4\sqrt{2} \\ 4 & 4\sqrt{2} & 8 \end{bmatrix}$$

$$A^2 = 2^2 \begin{bmatrix} 1 & \sqrt{2} & 2 \\ \sqrt{2} & 2 & 2\sqrt{2} \\ 2 & 2\sqrt{2} & 4 \end{bmatrix} \begin{bmatrix} 1 & \sqrt{2} & 2 \\ \sqrt{2} & 2 & 2\sqrt{2} \\ 2 & 2\sqrt{2} & 4 \end{bmatrix}$$

$$= 4 \begin{bmatrix} - & - & - \\ - & - & - \\ (2+4+8) & (2\sqrt{2}+4\sqrt{2}+8\sqrt{2}) & (4+8+16) \end{bmatrix}$$

$$\text{Sum of elements of 3rd row} = 4(14 + 14\sqrt{2} + 28)$$

$$= 4(42 + 14\sqrt{2})$$

$$= 168 + 56\sqrt{2}$$

$$\alpha + \beta\sqrt{2}$$

$$\therefore \alpha + \beta = 168 + 56 = 224$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

[Enrol Now](#)

Sol. Let $\vec{v} = \hat{i} + \hat{j} + \hat{k}$

$$\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 3 \\ 2 & 3 & -1 \end{vmatrix}$$

$$= -7\hat{i} + 7\hat{j} + 7\hat{k}$$

$$= -7(\hat{i} - \hat{j} - \hat{k})$$

$$\text{Now } \hat{a} = \frac{\hat{i} - \hat{j} - \hat{k}}{\sqrt{3}} \text{ or } \frac{-\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$$

↓

↓

$$\cos \theta = \frac{\hat{a} \cdot \vec{v}}{|\vec{v}|} = \frac{1-1-1}{\sqrt{3}\sqrt{3}} = \frac{-1}{3} \quad \cos \theta = \frac{\hat{a} \cdot \vec{v}}{|\vec{v}|} = \frac{-1+1+1}{3} = \frac{1}{3}$$

(rejected)

$$\Rightarrow \hat{a} = \frac{\hat{i} - \hat{j} - \hat{k}}{\sqrt{3}}$$

$$\text{Now } \cos \frac{\pi}{3} = \frac{\hat{a} \cdot (\hat{i} + \alpha \hat{j} + \hat{k})}{\sqrt{1+\alpha^2+1}}$$

$$\Rightarrow \frac{1}{2} = \frac{1-\alpha-1}{\sqrt{3}\sqrt{\alpha^2+2}}$$

$$\Rightarrow \frac{\sqrt{3}}{2} \sqrt{\alpha^2+2} = -\alpha \quad (\therefore \alpha < 0)$$

$$3\alpha^2 + 6 = 4\alpha^2$$

$$\Rightarrow \alpha = -\sqrt{6}$$

20. If for the solution curve $y = f(x)$ of the differential

$$\frac{dy}{dx} + (\tan x)y = \frac{2 + \sec x}{(1 + 2 \sec x)^2},$$

$x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, $f\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{10}$, then $f\left(\frac{\pi}{4}\right)$ is equal to:

$$(1) \frac{9\sqrt{3}+3}{10(4+\sqrt{3})}$$

$$(2) \frac{\sqrt{3}+1}{10(4+\sqrt{3})}$$

$$(3) \frac{5-\sqrt{3}}{2\sqrt{2}}$$

$$(4) \frac{4-\sqrt{2}}{14}$$

Ans. (4)

Sol. If $e^{\int \tan x dx} = e^{\ln(\sec x)} = \sec x$

$$\therefore y \cdot \sec x = \int \left\{ \frac{2 + \sec x}{(1 + 2 \sec x)^2} \right\} \sec x dx$$

$$= \int \frac{2 \cos x + 1}{(\cos x + 2)^2} dx \quad \text{Let } \cos x = \frac{1-t^2}{1+t^2}$$

$$= \int \frac{2\left(\frac{1-t^2}{1+t^2}\right)+1}{\left(\frac{1-t^2}{1+t^2}+2\right)^2} 2dt$$

$$= \int \frac{2-2t^2+1+t^2}{(1-t^2+2+2t^2)^2} \times 2dt$$

$$= 2 \int \frac{3-t^2}{(t^2+3)^2} dt$$

$$\text{Let } t + \frac{3}{t} = u$$

$$\left(1 - \frac{3}{t^2}\right) dt = du$$

$$= -2 \int \frac{du}{u^2}$$

$$y \cdot (\sec x) = \frac{2}{u} + c$$

$$\boxed{y \cdot \sec x = \frac{2}{t + \frac{3}{t}} + c} \quad \dots \dots \dots \text{(I)}$$

$$\text{At } x = \frac{\pi}{3}, t = \tan \frac{x}{2} = \frac{1}{\sqrt{3}}$$

$$2 \cdot \frac{\sqrt{3}}{10} = \frac{2}{\frac{1}{\sqrt{3}} + 3\sqrt{3}} + c$$

$$2 \cdot \frac{\sqrt{3}}{10} = \frac{2\sqrt{3}}{10} + c \Rightarrow C = 0$$

$$\text{At } x = \frac{\pi}{4}, t = \tan \frac{x}{2} = \sqrt{2} - 1$$

$$\therefore y \cdot \sqrt{2} = \frac{2}{\sqrt{2}-1+\frac{3}{\sqrt{2}-1}}$$

$$y \cdot \sqrt{2} = \frac{2(\sqrt{2}-1)}{6-2\sqrt{2}}$$

$$y = \frac{\sqrt{2}(\sqrt{2}-1)}{2(3-\sqrt{2})} = \frac{1}{\sqrt{2}} \times \frac{2\sqrt{2}-1}{7} = \frac{4-\sqrt{2}}{14}$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

SECTION-B

21. If $24 \int_0^{\frac{\pi}{4}} \left(\sin \left| 4x - \frac{\pi}{12} \right| + [2 \sin x] \right) dx = 2\pi + \alpha$, where $[\cdot]$ denotes the greatest integer function, then α is equal to _____.

Ans. (12)

Sol. $= 24 \int_0^{\frac{\pi}{4}} -\sin \left(4x - \frac{\pi}{12} \right) + \int_{\pi/6}^{\pi/4} \sin \left(4x - \frac{\pi}{12} \right) + \int_0^{\frac{\pi}{6}} [0] dx + \int_1^{\pi/4} [2 \sin x] dx$

 $= 24 \left[\frac{\left(1 - \cos \frac{\pi}{12} \right)}{4} - \frac{\left(-\cos \frac{\pi}{12} - 1 \right)}{4} \right] + \frac{\pi}{4} - \frac{\pi}{6}$
 $= 24 \left(\frac{1}{2} + \frac{\pi}{12} \right) = 2\pi + 12$

$\alpha = 12$

22. If $\lim_{t \rightarrow 0} \left(\int_0^1 (3x+5)^t dx \right)^{\frac{1}{t}} = \frac{\alpha}{5e} \left(\frac{8}{5} \right)^{\frac{2}{3}}$, then α is equal to _____.

Ans. (64)

Sol. 1^∞ form

$$\text{Now } L = e^{t \rightarrow 0} \frac{1}{t} \left(\left. \frac{(3x+5)^{t+1}}{3(t+1)} \right|_0^1 - 1 \right)$$
 $= e^{t \rightarrow 0} \frac{8^{t+1} - 5^{t+1} - 3t - 3}{3t(t+1)}$
 $= e \frac{8\ell n 8 - 5\ell n 5 - 3}{3}$
 $= \left(\frac{8}{5} \right)^{2/3} \left(\frac{64}{5} \right) = \frac{\alpha}{5e} \left(\frac{8}{5} \right)^{2/3}$

On comparing

$\alpha = 64$

23. Let $a_1, a_2, \dots, a_{2024}$ be an Arithmetic Progression such that $a_1 + (a_5 + a_{10} + a_{15} + \dots + a_{2020}) + a_{2024} = 2233$. Then $a_1 + a_2 + a_3 + \dots + a_{2024}$ is equal to _____.

Ans. (11132)

- Sol.** $a_1 + a_5 + a_{10} + \dots + a_{2020} + a_{2024} = 2233$
In an A.P. the sum of terms equidistant from ends is equal.
 $a_1 + a_{2024} = a_5 + a_{2020} = a_{10} + a_{2015} \dots$
 $\Rightarrow 203$ pairs
 $\Rightarrow 203(a_1 + a_{2024}) = 2233$
Hence,

$$S_{2024} = \frac{2024}{2} (a_1 + a_{2024})$$
 $= 1012 \times 11$
 $= 11132$

24. Let integers $a, b \in [-3, 3]$ be such that $a + b \neq 0$. Then the number of all possible ordered pairs

$$(a, b), \text{ for which } \left| \frac{z-a}{z+b} \right| = 1 \text{ and } \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix}$$

$= 1, z \in C$, where ω and ω^2 are the roots of $x^2 + x + 1 = 0$, is equal to _____.

Ans. (10)

- Sol.** $a, b \in I, -3 \leq a, b \leq 3, a + b \neq 0$
 $|z - a| = |z + b|$

$$\begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow \begin{vmatrix} z & z & z \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow z \begin{vmatrix} 1 & 1 & 1 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow z \begin{vmatrix} 1 & 0 & 0 \\ \omega & z+\omega^2-\omega & 1-\omega \\ \omega^2 & 1-\omega^2 & z+\omega-\omega^2 \end{vmatrix} = 1$$

$$\Rightarrow z^3 = 1$$

$$\Rightarrow z = \omega, \omega^2, 1$$

Now

$$|1-a|=|1+b|$$

$\Rightarrow 10$ pairs



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

25. Let $y^2 = 12x$ the parabola and S be its focus. Let PQ be a focal chord of the parabola such that $(SP) \cdot (SQ) = \frac{147}{4}$. Let C be the circle described taking PQ as a diameter. If the equation of a circle C is $64x^2 + 64y^2 - \alpha x - 64\sqrt{3}y = \beta$, then $\beta - \alpha$ is equal to _____.

Ans. (1328)

Sol. $y^2 = 12x$ $a = 3$ $SP \times SQ = \frac{147}{4}$

Let $P(3t^2, 6t)$ and $t_1 t_2 = -1$
(ends of focal chord)

So, $Q\left(\frac{3}{t^2}, \frac{-6}{t}\right)$

$S(3, 0)$

$SP \times SQ = PM_1 \times QM_2$

(dist. from directrix)

$$= (3 + 3t^2) \left(3 + \frac{3}{t^2} \right) = \frac{147}{4}$$

$$\Rightarrow \frac{(1+t^2)^2}{t^2} = \frac{49}{12}$$

$$t^2 = \frac{3}{4}, \frac{4}{3}$$

$$t = \pm \frac{\sqrt{3}}{2}, \pm \frac{2}{\sqrt{3}}$$

considering $t = \frac{-\sqrt{3}}{2}$

$P\left(\frac{9}{4}, -3\sqrt{3}\right)$ and $Q\left(4, 4\sqrt{3}\right)$

Hence, diametric circle:

$$(x-4)\left(x - \frac{9}{4}\right) + (y+3\sqrt{3})(y-4\sqrt{3}) = 0$$

$$\Rightarrow x^2 + y^2 - \frac{25}{4}x - \sqrt{3}y - 27 = 0$$

$$\Rightarrow \alpha = 400, \beta = 1728$$

$$\beta - \alpha = 1328$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now



Level up your prep for JEE Adv. 2025 with our **Online Rank Booster Course!**



LIVE classes for JEE Main & Advanced



Soft copies of ALLEN's study material




Covers important questions across **Physics, Chemistry & Maths**



ALLEN App Advantage: **24/7 doubt support,
Custom Practice & more**

Enrol Now



Win up to
90% scholarship*

with the ALLEN Online Scholarship Test

at just **₹49/-**



1-hour online test
Can be taken from anywhere

Register Now

