

JEE-MAIN EXAMINATION – JANUARY 2025(HELD ON THURSDAY 23rd JANUARY 2025)

TIME : 9 : 00 AM TO 12 : 00 NOON

CHEMISTRY**TEST PAPER WITH SOLUTIONS****SECTION-A**

- 51.** The element that does not belong to the same period of the remaining elements (modern periodic table) is:
 (1) Palladium
 (2) Iridium
 (3) Osmium
 (4) Platinum

Sol. (1)Palladium \Rightarrow 5th periodIridium, Osmium, Platinum \Rightarrow 6th Period

- 52.** Heat treatment of muscular pain involves radiation of wavelength of about 900 nm. Which spectral line of H atom is suitable for this ?

Given: Rydberg constant

$$R_H = 10^5 \text{ cm}^{-1}, h = 6.6 \times 10^{-34} \text{ J s}, c = 3 \times 10^8 \text{ m/s}$$

- (1) Paschen series, $\infty \rightarrow 3$
 (2) Lyman series, $\infty \rightarrow 1$
 (3) Balmer series, $\infty \rightarrow 2$
 (4) Paschen series, $5 \rightarrow 3$

Sol. (1)

$$\lambda = 900 \text{ nm}$$

$$= 9 \times 10^{-5} \text{ cm}$$

$$R_H = 10^5 \text{ cm}^{-1}$$

$$\text{Rydberg eq.} = \frac{1}{\lambda} = R_H Z^2 \times \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\Rightarrow \frac{1}{\lambda \times R_H} = \frac{1}{n_1^2} - \frac{1}{n_2^2}$$

$$\Rightarrow \frac{1}{9 \times 10^{-5} \text{ cm} \times 10^5 \text{ cm}^{-1}} = \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\Rightarrow \frac{1}{n_1^2} - \frac{1}{n_2^2} = \frac{1}{9}$$

It is possible when $n_1 = 3, n_2 = \infty$ Possible series : $\infty \rightarrow 3$

- 53.** The **incorrect** statements among the following is
 (1) PH₃ shows lower proton affinity than NH₃.
 (2) PF₃ exists but NF₅ does not.
 (3) NO₂ can dimerise easily.
 (4) SO₂ can act as an oxidizing agent, but not as a reducing agent.

Sol. (4)SO₂ can oxidise as well as reduce.

Hence it can act as both oxidising and reducing agent.

- 54.** CrCl₃.xNH₃ can exist as a complex. 0.1 molal aqueous solution of this complex shows a depression in freezing point of 0.558°C. Assuming 100% ionisation of this complex and coordination number of Cr is 6, the complex will be
 (Given K_f = 1.86 K kg mol⁻¹)

- (1) [Cr(NH₃)₆] Cl₃
 (2) [Cr(NH₃)₄Cl₂] Cl
 (3) [Cr(NH₃)₅Cl] Cl₂
 (4) [Cr(NH₃)₃Cl₃]

Sol. (3)Given : $\Delta T_f = 0.558^\circ\text{C}$

$$k_f = 1.86 \frac{\text{K} \times \text{kg}}{\text{mol}}$$

0.1 m aq. sol.

$$\Rightarrow \Delta T_f = i \times k_f \times m$$

$$\Rightarrow 0.558 = i \times 1.86 \times 0.1$$

$$\Rightarrow i = 3$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

59. Ice at -5°C is heated to become vapor with temperature of 110°C at atmospheric pressure. The entropy change associated with this process can be obtained from :

$$(1) \int_{268\text{K}}^{383\text{K}} C_p dT + \frac{\Delta H_{\text{melting}}}{273} + \frac{\Delta H_{\text{boiling}}}{373}$$

$$(2) \int_{268\text{K}}^{273\text{K}} \frac{C_{p,m}}{T} dT + \frac{\Delta H_{m,\text{fusion}}}{T_f} + \frac{\Delta H_{m,\text{vaporisation}}}{T_b}$$


$$+ \int_{273\text{K}}^{373\text{K}} \frac{C_{p,m}}{T} dT + \int_{373\text{K}}^{383\text{K}} \frac{C_{p,m}}{T} dT$$

$$(3) \int_{268\text{K}}^{383\text{K}} C_p dT + \frac{q_{\text{rev}}}{T}$$

$$(4) \int_{268\text{K}}^{273\text{K}} C_{p,m} dT + \frac{\Delta H_{m,\text{fusion}}}{T_f} + \frac{\Delta H_{m,\text{vaporisation}}}{T_b}$$

$$+ \int_{273\text{K}}^{373\text{K}} C_{p,m} dT + \int_{373\text{K}}^{383\text{K}} C_{p,m} dT$$

Sol. (2)



$$\Delta S_{\text{overall}} = \Delta S_1 + \Delta S_2 + \Delta S_3 + \Delta S_4 + \Delta S_5$$

$$\Delta S_2 = \frac{\Delta H_{m,\text{fusion}}}{273} \quad T_f = 273 \text{ 'K'}$$

$$\Delta S_3 = \int_{273}^{373} \frac{C_{p,m}}{T} dT \quad T_b = 373 \text{ 'K'}$$

$$\Delta S_4 = \frac{\Delta H_{m,\text{vaporisation}}}{373}$$

$$\Delta S_5 = \int_{373}^{383} \frac{C_{p,m}}{T} dT$$

Answer = (2)

60. The d-electronic configuration of an octahedral Co(II) complex having magnetic moment of 3.95 BM is :

$$(1) t_{2g}^6 e_g^1 \quad (2) t_{2g}^3 e_g^0$$

$$(3) t_{2g}^5 e_g^2 \quad (4) e^4 t_2^3$$

Sol. (3)

$$\text{Co}^{+2} = (\text{Ar})_{18} 3d^7 4s^0$$

$$\begin{array}{c} 1 & 1 & e_g^2 \\ \Downarrow & \Downarrow & \Downarrow \\ 1 & 1 & t_{2g}^5 \end{array}$$

61. The complex that shows Facial – Meridional isomerism is

$$(1) [\text{Co}(\text{NH}_3)_3\text{Cl}_3] \quad (2) [\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$$

$$(3) [\text{Co}(\text{en})_3]^{3+} \quad (4) [\text{Co}(\text{en})_2\text{Cl}_2]^+$$

Sol. (1)

Ma_3b_3 type complexes show Facial - Meridional isomerism

$$(i) [\text{Co}(\text{NH}_3)_3\text{Cl}_3] \Rightarrow \text{Ma}_3\text{b}_3$$

$$(ii) [\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+ \Rightarrow \text{Ma}_4\text{b}_2$$


$$(iii) [\text{Co}(\text{en})_3]^{3+} \Rightarrow \text{M(AA)}_3$$

$$(iv) [\text{Co}(\text{en})_2\text{Cl}_2]^+ \Rightarrow \text{M(AA)}_2\text{b}_2$$

a, b, = NH_3 , Cl^-

AA = en

62. The major product of the following reaction is :



$$(1) \text{CH}_3-\text{CH}_2-\text{CH}_2-\text{OH}$$


$$(2) \text{CH}_3-\underset{\text{CH}_2-\text{OH}}{\underset{|}{\text{CH}}}-\text{CH=O}$$

$$(3) \text{CH}_3-\underset{\text{CH}_2-\text{OH}}{\underset{|}{\text{C}}}-\text{CH}_2-\text{OH}$$

$$(4) \text{CH}_3-\underset{\text{CH}_2}{\underset{\parallel}{\text{C}}}-\text{CH=O}$$

Sol. (3)

This is an example of Tollen's reaction i.e. multiple cross aldol followed by cross Cannizaro reaction



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

Sol. (1)

Condition for precipitation $Q_{\text{ip}} > K_{\text{sp}}$

For $[\text{A(OH)}_2]$

$$[\text{A}^{2+}] [\text{OH}^-]^2 > 9 \times 10^{-10}$$

$$[\text{A}^{2+}] = 1 \text{ M}$$

$$\Rightarrow [\text{OH}^-] > 3 \times 10^{-5} \text{ M}$$

For $[\text{B(OH)}_3]$

$$[\text{B}^{3+}] [\text{OH}^-]^3 > 27 \times 10^{-18}$$

$$[\text{B}^{3+}] = 1 \text{ M}$$

$$\Rightarrow [\text{OH}^-] > 3 \times 10^{-6} \text{ M}$$

So, B(OH)_3 will precipitate before A(OH)_2

69. Match the LIST-I with LIST-II

LIST-I (Classification of molecules based on octet rule)		LIST-II (Example)	
A.	Molecules obeying octet rule	I.	NO, NO_2
B.	Molecules with incomplete octet	II.	$\text{BCl}_3, \text{AlCl}_3$
C.	Molecules with incomplete octet with odd electron	III.	$\text{H}_2\text{SO}_4, \text{PCl}_5$
D.	Molecules with expanded octet	IV.	$\text{CCl}_4, \text{CO}_2$





Choose the **correct** answer from the options given below :

- (1) A-IV, B-II, C-I, D-III
- (2) A-III, B-II, C-I, D-IV
- (3) A-IV, B-I, C-III, D-II
- (4) A-II, B-IV, C-III, D-I

Sol. (1)

- (A) A \rightarrow IV
- (B) B \rightarrow II
- (C) C \rightarrow I
- (D) D \rightarrow III

70. Which among the following react with Hinsberg's reagent?



Choose the correct answer from the options given below :

- (1) B and D only
- (2) C and D only
- (3) A, B and E only
- (4) A, C and E only

Sol. (4)

B and D are 3° amine which does not have replaceable H on N, So does not react.

SECTION-B

71. If 1 mM solution of ethylamine produces $\text{pH} = 9$, then the ionization constant (K_b) of ethylamine is 10^{-x} . The value of x is _____ (nearest integer). [The degree of ionization of ethylamine can be neglected with respect to unity.]

Sol. (7)



$$\text{C} = 10^{-3} \text{ M}$$

$$\text{C}(1 - \alpha)$$

$$\text{C}\alpha$$

$$\Rightarrow \text{C} = 10^{-3} = 10^{-5} = 10^{-5}$$

$$1 - \alpha \approx 1$$

$$\text{Given, } \text{P}^\text{H} = 9 \Rightarrow \text{P}^\text{OH} = 5 \Rightarrow [\text{OH}^\ominus] = 10^{-5} \text{ M}$$

$$\text{Now, } K_b = \frac{[\text{C}_2\text{H}_2\text{NH}_3^+][\text{OH}^\ominus]}{[\text{C}_2\text{H}_5\text{NH}_2]}$$

$$\Rightarrow K_b = \frac{10^{-5} \times 10^{-5}}{10^{-3}} = 10^{-7}$$



Level up your prep for JEE Adv. 2025 with ALLEN Online's LIVE Rank Booster Course!

[Enrol Now](#)

72. During "S" estimation, 160 mg of an organic compound gives 466 mg of barium sulphate. The percentage of Sulphur in the given compound is _____ %.


(Given molar mass in g mol⁻¹ of Ba : 137, S : 32, O : 16)

Sol. (40)

$$\text{Millimoles of BaSO}_4 = \frac{466}{233} = 2 \text{ mmol}$$

$$\% \text{S} = \frac{\frac{466}{233} \times 32}{160} \times 100 = 40\%$$


73. Consider the following sequence of reactions to produce major product (A)



Molar mass of product (A) is _____ g mol⁻¹.

(Given molar mass in g mol⁻¹ of C : 12, H : 1, O : 16, Br : 80, N : 14, P : 31)

Sol. (171)



Molar mass of product ($\text{C}_7\text{H}_7\text{Br}$) (A) is 171 g mol⁻¹

74. For the thermal decomposition of $\text{N}_2\text{O}_5(\text{g})$ at constant volume, the following table can be formed, for the reaction mentioned below :



S.No.	Time/s	Total pressure / (atm)
1.	0	0.6
2.	100	'x'


$$x = \text{_____} \times 10^{-3} \text{ atm [nearest integer]}$$

Given : Rate constant for the reaction is $4.606 \times 10^{-2} \text{ s}^{-1}$.

Sol. (900)

NTA. (897)

$$K_{\text{N}_2\text{O}_5} = 2 \times 4.606 \times 10^{-2} \text{ s}^{-1}$$



P_i	0.6	0	0
P_f	$0.6 - P$	P	$\frac{P}{2}$

$$2 \times 4.606 \times 10^{-2} = \frac{2.303}{100} \log \frac{0.6}{0.6 - P}$$

$$4 \log_{10} \frac{0.6}{0.6 - P}$$

$$10^4 = \frac{0.6}{0.6 - P}$$

$$\Rightarrow 0.6 \times 10^4 - 10^4 P = 0.6$$

$$\Rightarrow 10^4 P = 0.6(10^4 - 1)$$

$$P = (6000 - 0.6) \times 10^{-4}$$

$$= 5999. \times 10^{-4}$$

$$= 0.59994$$

$$P_{\text{Total}} = 0.6 + \frac{P}{2}$$

$$= 0.6 + 0.29997$$

$$= 0.89997$$

$$= 899.97 \times 10^{-3}$$


Ans. 900



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

Enrol Now

Given by NTA



$$\begin{array}{cccc} t = 0 & 0.6 & 0 & 0 \\ t = 100\text{s} & 0.6 - x & x & x/2 \end{array}$$

$$P_{\text{Total}} = 0.6 + \frac{x}{2}$$

As given in equation

$$K_r = 4.606 \times 10^{-2} \text{ sec}^{-1}$$

(Here language conflict in question)

$$(K_r = \frac{KA}{2} \text{ not considered})$$

$$K_r t = \ln \frac{0.6}{0.6 - x}$$

$$4.606 \times 10^{-2} \times 100 = 2.303 \log \frac{0.6}{0.6 - x}$$

$$P_{\text{Total}} = 0.6 + \frac{0.594}{2} = 0.897 \text{ atm}$$

$$= 897 \times 10^{-3} \text{ atm}$$

75. The standard enthalpy and standard entropy of decomposition of N_2O_4 to NO_2 are 55.0 kJ mol^{-1} and 175.0 J/K/mol respectively. The standard free energy change for this reaction at 25°C in J mol^{-1} is _____ (Nearest integer)

Sol. (2850)

$$\Delta H_{\text{rxn}}^{\circ} = 55 \text{ kJ/mol}, \quad T = 298 \text{ K}$$

$$\Delta S_{\text{rxn}}^{\circ} = 175 \text{ J/mol}$$

$$\Delta G_{\text{rxn}}^{\circ} = \Delta H_{\text{rxn}}^{\circ} - T\Delta S_{\text{rxn}}^{\circ}$$

$$\Rightarrow \Delta G_{\text{rxn}}^{\circ} = 55000 \text{ J/mol} - 298 \times 175 \text{ J/mol}$$

$$\Rightarrow \Delta G_{\text{rxn}}^{\circ} = 55000 - 52150$$

$$\Rightarrow \Delta G_{\text{rxn}}^{\circ} = 2850 \text{ J/mol}$$



Level up your prep for JEE Adv. 2025 with
ALLEN Online's LIVE Rank Booster Course!

[Enrol Now](#)



Level up your prep for JEE Adv. 2025 with our **Online Rank Booster Course!**



LIVE classes for JEE Main & Advanced



Soft copies of ALLEN's study material




Covers important questions across **Physics, Chemistry & Maths**



ALLEN App Advantage: **24/7 doubt support,
Custom Practice & more**

Enrol Now



Win up to
90% scholarship*

with the ALLEN Online Scholarship Test

at just **₹49/-**



1-hour online test
Can be taken from anywhere

Register Now

