

Sol. $2x - y = 0$

$$\{0, 0\} \{-1, -2\} \{1, 2\}$$

$$2x - y = 1$$

$$\{0, -1\} \{1, 1\} \{2, 3\} \{-1, -3\}$$

Total $(0, 0) (-1, -2), (1, 2) (0, -1), (1, 1) (2, 3) (-1, -3)$

Reflexive $m = 5$ & $\ell = 7$

Symm. $n = 5$ $\ell + m + n = 17$

option (2)

- 5.** Let the product of $\omega_1 = (8 + i)\sin\theta + (7 + 4i)\cos\theta$ and $\omega_2 = (1 + 8i)\sin\theta + (4 + 7i)\cos\theta$ be $\alpha + i\beta$, $i = \sqrt{-1}$. Let p and q be the maximum and the minimum values of $\alpha + \beta$ respectively.

(1) 140

(2) 130

(3) 160

(4) 150

Ans. (2)

Sol. $\omega_1 = (8 \sin \theta + 7 \cos \theta) + i(\sin \theta + 4 \cos \theta)$

$\omega_2 = (\sin \theta + 4 \cos \theta) + i(8 \sin \theta + 7 \cos \theta)$

$$\begin{aligned}\omega_1\omega_2 &= 8 \sin^2 \theta + 7 \sin \theta \cos \theta + 32 \sin \theta \cos \theta + \\ &28 \cos^2 \theta - 8 \sin^2 \theta - 32 \sin \theta \cos \theta - 7 \sin \theta \cos \theta \\ &- 28 \cos^2 \theta + i(\sin^2 \theta + 4 \sin \theta \cos \theta + 4 \sin \theta \cos \theta \\ &+ 16 \cos^2 \theta + 64 \sin^2 \theta + 56 \sin \theta \cos \theta + 56 \sin \theta \\ &\cos \theta + 49 \cos^2 \theta)\end{aligned}$$

$$\omega_1\omega_2 = 0 + i(65 \sin^2 \theta + 120 \sin \theta \cos \theta + 65 \cos^2 \theta)$$

$$\alpha + \beta = 65 + 60 \sin 2\theta$$

$$\alpha + \beta|_{\max} = 125$$

$$\alpha + \beta|_{\min} = 5$$

$$\text{Ans. } = 125 + 5 = 130$$

option (2)

- 6.** Let the values of p, for which the shortest distance between the lines $\frac{x+1}{3} = \frac{y}{4} = \frac{z}{5}$ and $\vec{r} = (\hat{p}\hat{i} + 2\hat{j} + \hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 4\hat{k})$ is $\frac{1}{\sqrt{6}}$, be a, b, (a < b). Then the length of the latus rectum of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is :-

(1) 9

(2) $\frac{3}{2}$

(3) $\frac{2}{3}$

(4) 18

Ans. (3)

Sol. shortest distance = $\frac{|(\bar{a} - \bar{b})| \cdot (\bar{p} \times \bar{q})}{|\bar{p} \times \bar{q}|}$

where

$$\bar{a} = -\hat{i} + 0\hat{j} + 0\hat{k}$$

$$> \bar{a} - \bar{b} = (-1 - p)\hat{i} - 2\hat{j} - \hat{k}$$

$$\bar{b} = p\hat{i} + 2\hat{j} + \hat{k}$$

$$\bar{p} = 3\hat{i} + 4\hat{j} + 5\hat{k}$$

$$\bar{q} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\frac{1}{16} = \frac{|-1 - p + 4 - 1|}{\sqrt{6}}$$

$$|-p + 2| = 1$$

$$p = 3 \quad \& \quad q = 1$$

$$\frac{x^2}{1^2} + \frac{y^2}{3^3} = 1$$

$$L.R = \frac{2a^2}{b} = \frac{2 \times 1}{3} = \frac{2}{3}$$

option (3)



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now

Sol. $\sum_{r=1}^{20} \frac{4r}{4+3r^2+r^4}$

$$\sum_{r=1}^{20} \frac{4r}{(r^2+r+2)(r^2-r+2)}$$

$$2 \sum_{r=1}^{20} \left(\frac{1}{r^2-r+2} - \frac{1}{r^2+r+2} \right)$$

$$2 \left(\frac{1}{2} - \frac{1}{4} \right)$$

$$\frac{1}{4} - \frac{1}{8}$$

$$\frac{1}{8} - \frac{1}{14}$$

$$\left(\frac{1}{382} - \frac{1}{422} \right)$$

$$= 2 \left(\frac{1}{2} - \frac{1}{422} \right)$$

$$= \frac{420}{422}$$

$$= \frac{210}{211}$$

option (3)

13. If

$$1^2 \cdot \binom{15}{1} + 2^2 \cdot \binom{15}{2} + 3^2 \cdot \binom{15}{3} + \dots + 15^2 \cdot \binom{15}{15} =$$

$2^m \cdot 3^n \cdot 5^k$, where $m, n, k \in \mathbb{N}$, then $m + n + k$ is equal to :-

(1) 19

(2) 21

(3) 18

(4) 20

Ans. (1)

Sol. $\sum_{r=1}^{15} r^2 \binom{15}{r} \Rightarrow 15 \sum_{r=1}^{15} r^{14} \binom{14}{r-1}$

$$15 \sum_{r=1}^{15} (r-1+1)^{14} \binom{14}{r-1}$$

$$15 \cdot 14 \sum_{r=1}^{15} \binom{13}{r-2} + 15 \sum_{r=1}^{15} \binom{14}{r-1}$$

$$15 \cdot 14 \cdot 2^{13} + 15 \cdot 2^{14}$$

$$3^1 \cdot 2^{13} (70 + 10)$$

$$3^1 \cdot 2^{13} \cdot 80$$

$$3^1 \cdot 5^1 \cdot 2^{17}$$

$$m = 17 \quad n = 1 \quad k = 1$$

option (1)

14. Let for two distinct values of p the lines $y = x + p$ touch the ellipse $E : \frac{x^2}{4^2} + \frac{y^2}{3^2} = 1$ at the points A and B. Let the line $y = x$ intersect E at the points C and D. Then the area of the quadrilateral ABCD is equal to

(1) 36

(2) 24

(3) 48

(4) 20

Ans. (2)

Sol. Point of contact are $\left(\frac{\pm a^2 m}{\sqrt{a^2 m^2 + b^2}}, \frac{\pm b^2}{\sqrt{a^2 m^2 + b^2}} \right)$

$$A \left(\frac{-16}{5}, \frac{9}{5} \right) B \left(\frac{16}{5}, \frac{-9}{5} \right)$$

$$\text{Point D is } \left(\frac{12}{5}, \frac{12}{5} \right)$$

$$\text{Area of ABD} = \frac{1}{2} \begin{vmatrix} -\frac{16}{5} & \frac{9}{5} & 1 \\ \frac{16}{5} & -\frac{9}{5} & 1 \\ \frac{12}{5} & \frac{12}{5} & 1 \end{vmatrix}$$

$$= 12$$

Area of ABCD is = 24

option (2)

15. Consider two sets A and B, each containing three numbers in A.P. Let the sum and the product of the elements of A be 36 and p respectively and the sum and the product of the elements of B be 36 and q respectively. Let d and D be the common differences of AP's in A and B respectively such that $D = d + 3$, $d > 0$. If $\frac{p+q}{p-q} = \frac{19}{5}$, then $p - q$ is

equal to

(1) 600

(2) 450

(3) 630

(4) 540

Ans. (4)

Sol. Let $A(a-d, a, a+d)$ $B(b-D, b, b+D)$

$$a = 12$$

$$b = 12$$

$$p = 12(144 - d^2)$$

$$q = 12(144 - D^2)$$



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now

18. Let $f(x) + 2f\left(\frac{1}{x}\right) = x^2 + 5$ and

$2g(x) - 3g\left(\frac{1}{2}\right) = x$, $x > 0$. If $\alpha = \int_1^2 f(x) dx$, and

$\beta = \int_1^2 g(x) dx$, then the value of $9\alpha + \beta$ is :

(1) 1

(2) 0

(3) 10

(4) 11

Ans. (4)

$$\text{Sol. } f(x) + 2f\left(\frac{1}{x}\right) = x^2 + 5$$

$$f\left(\frac{1}{x}\right) + 2f(x) = \frac{1}{x^2} + 5$$

$$f(x) = \frac{2}{3x^2} - \frac{x^2}{3} + \frac{5}{3}$$

$$\alpha = \int_1^2 \left(\frac{2}{3x^2} - \frac{x^2}{3} + \frac{5}{3} \right) dx$$

$$\left(-\frac{2}{3x} - \frac{x^3}{9} + \frac{5x}{3} \right)_1^2$$

$$-\frac{1}{3} - \frac{8}{9} + \frac{10}{3} + \frac{2}{3} + \frac{1}{9} - \frac{5}{3}$$

$$\alpha = 2 - \frac{7}{9} = \frac{11}{9}$$

$$2g(x) - 3g\left(\frac{1}{2}\right) = x$$

$$g\left(\frac{1}{2}\right) = -\frac{1}{2}$$

$$g(x) = \frac{x}{2} - \frac{3}{4}$$

$$\beta = \int_1^2 \left(\frac{x}{2} - \frac{3}{4} \right) dx$$

$$\left(\frac{x^2}{4} - \frac{3x}{4} \right)_1^2 = 1 - \frac{3}{2} - \frac{1}{4} + \frac{3}{4} = 0$$

$$9\alpha + \beta = 11$$

option (4)

19. Let A be the point of intersection of the lines

$$L_1 : \frac{x-7}{1} = \frac{y-5}{0} = \frac{z-3}{-1} \text{ and}$$

$$L_2 : \frac{x-1}{3} = \frac{y+3}{4} = \frac{z+7}{5}.$$

Let B and C be the points on the lines L_1 and L_2 respectively such that $AB = AC = \sqrt{15}$. Then the square of the area of the triangle ABC is :

(1) 54

(2) 63

(3) 57

(4) 60

Ans. (1)

Sol. Angle between both lines

$$\cos\theta = \left| \frac{3+0-5}{\sqrt{2}\sqrt{50}} \right|$$


$$\sin\theta = \frac{2}{10} = \frac{1}{5}$$

$$\sin\theta = \frac{\sqrt{24}}{5}$$

$$\text{area} = \frac{1}{2} \text{absin}\theta$$

$$\frac{1}{2} \sqrt{15} \sqrt{15} \frac{\sqrt{24}}{5}$$

$$\text{square of area } \frac{15 \cdot 15 \cdot 24}{4 \cdot 25}$$



option (1)



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now

20. Let the mean and the standard deviation of the observation 2, 3, 3, 4, 5, 7, a, b be 4 and $\sqrt{2}$ respectively. Then the mean deviation about the mode of these observations is :

(1) 1

(2) $\frac{3}{4}$

(3) 2

(4) $\frac{1}{2}$

Ans. (1)

Sol. $\frac{24+a+b}{8} = 4$

$a+b=8$

$2 = \frac{4+1+1+0+1+9+(a-4)^2+(b-4)^2}{8}$

$16 = 48 + a^2 + b^2 - 8a - 8b$

$a^2 + b^2 = 32$

$32 = 2ab$

$ab = 16$

$a = 4 \quad b = 4$

$\text{mode} = 4$

$\text{mean deviation} = \frac{2+1+1+0+1+3+0+0}{8} = 1$

option (1)

SECTION-B

21. If α is a root of the equation $x^2 + x + 1 = 0$ and

$\sum_{k=1}^n \left(\alpha^k + \frac{1}{\alpha^k} \right)^2 = 20$, then n is equal to _____

Ans. (11)

Sol. $\alpha = \omega$

$\therefore \left(\omega^k + \frac{1}{\omega^k} \right)^2 = \omega^{2k} + \frac{1}{\omega^{2k}} + 2$

$= \omega^{2k} + \omega^k + 2$

$\because \omega^{3k} = 1$

$\therefore \sum_{k=1}^n (\omega^{2k} + \omega^k + 2) = 20$

$\Rightarrow (\omega^2 + \omega^4 + \omega^6 + \dots + \omega^{2n}) + (\omega + \omega^2 + \omega^3 + \dots + \omega^n) + 2n = 20$

Now if $n = 3m$, $m \in \mathbb{N}$ Then $0 + 0 + 2n = 20 \Rightarrow n = 10$ (not satisfy)if $n = 3m+1$, then

$\omega^2 + \omega + 2n = 20$

$-1 + 2n = 20 \Rightarrow n = \frac{21}{2}$ (not possible)

if $n = 3m+2$,

$(\omega^8 + \omega^{10}) + (\omega^4 + \omega^5) + 2n = 20$

$\Rightarrow (\omega^2 + \omega) + (\omega + \omega^2) + 2n = 20$

$2n = 22$

 $n = 11$ satisfy $n = 3m + 2$

$\therefore n = 11$

22. If $\int \frac{\left(\sqrt{1+x^2} + x \right)^{10}}{\left(\sqrt{1+x^2} - x \right)^9} dx =$

$\frac{1}{m} \left(\left(\sqrt{1+x^2} + x \right)^n \left(n\sqrt{1+x^2} - x \right) \right) + C$ where C is the constant of integration and $m, n \in \mathbb{N}$, then $m+n$ is equal to

Ans. (379)**Sol.** rationalise

$\Rightarrow \int \frac{\left(\sqrt{1+x^2} + x \right)^{10}}{\left(\sqrt{1+x^2} - x \right)^9} \times \frac{\left(\sqrt{1+x^2} + x \right)^9}{\left(\sqrt{1+x^2} + x \right)^9} dx$

$\Rightarrow \int \frac{\left(\sqrt{1+x^2} + x \right)^{19}}{1} dx$

Put $\sqrt{1+x^2} + x = t$

$\left(\frac{x}{\sqrt{1+x^2}} + 1 \right) dx = dt$



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now

$$dx = \frac{dt}{t} \sqrt{1+x^2}$$

Now as $\sqrt{1+x^2} + x = t$

$$\text{so } \sqrt{1+x^2} - x = \frac{1}{t}$$

$$\therefore \sqrt{1+x^2} = \frac{1}{2} \left(t + \frac{1}{t} \right)$$

$$\text{Thus } I = \int t^{19} \cdot \frac{dt}{t} \cdot \frac{1}{2} \left(t + \frac{1}{t} \right)$$

$$\Rightarrow \frac{1}{2} \int (t^{19} + t^{17}) dt$$

$$= \frac{1}{2} \left(\frac{t^{20}}{20} + \frac{t^{18}}{18} \right) + C$$

$$= \frac{t^{19}}{360} \left[9t + \frac{10}{t} \right] + C$$

$$= \frac{t^{19}}{360} \left[9 \left(t + \frac{1}{t} \right) + \frac{1}{t} \right] + C$$

$$\Rightarrow \frac{(\sqrt{1+x^2} + x)^{19}}{360} \left[9(2\sqrt{1+x^2}) + (\sqrt{1+x^2} - x) \right] + C$$

$$\Rightarrow \frac{(\sqrt{1+x^2} + x)^{19}}{360} [19\sqrt{1+x^2} - x] + C$$

$$\therefore m = 360, n = 19$$

$$\therefore m + n = 379$$

23. A card from a pack of 52 cards is lost. From the remaining 51 cards, n cards are drawn and are found to be spades. If the probability of the lost card to be a spade is $\frac{11}{50}$, the n is equal to

Ans. (2)

Sol. n cards are drawn & are found all spade, thus remaining spades = $13 - x$
remaining total cards = $52 - x$

$$\text{Now given that } P(\text{lost card is spade}) = \frac{11}{50}$$

$$\text{i.e. } \frac{\binom{13-n}{1}}{\binom{52-n}{1}} = \frac{11}{50}$$

$$50(13 - n) = 11(52 - n)$$

$$39n = 78$$

$$n = 2$$

24. Let m and n , ($m < n$) be two 2-digit numbers. Then the total numbers of pairs (m, n) , such that $\gcd(m, n) = 6$, is _____

Ans. (64)

Sol. Let $m = 6a, n = 6b$

$$m < n \Rightarrow a < b$$

where a & b are co-prime numbers

also since m & n are 2 digit nos, so

$$10 \leq m \leq 99 \text{ & } 10 \leq n \leq 99$$

$$\text{i.e. } 2 \leq a \leq 16 \text{ & } 2 \leq b \leq 16$$

($\because a$ is integer)

Now

$$2 \leq a < b \leq 16 \text{ & } a \text{ & } b \text{ are co-prime}$$

\therefore if

$$a = 2, b = 3, 5, 7, 9, 11, 13, 15$$

$$a = 3, b = 4, 5, 7, 8, 10, 11, 13, 14, 16$$

$$a = 4, b = 5, 7, 9, 11, 13, 15$$

$$a = 5, b = 6, 7, 8, 9, 11, 12, 13, 14, 16$$

$$a = 6, b = 7, 11, 13$$

$$a = 7, b = 8, 9, 10, 11, 12, 13, 15, 16$$

$$a = 8, b = 9, 11, 13, 15$$

$$a = 9, b = 10, 11, 13, 14, 16$$

$$a = 10, b = 11, 13$$

$$a = 11, b = 12, 13, 14, 15, 16$$

$$a = 12, b = 13$$

$$a = 13, b = 14, 15, 16$$

$$a = 14, b = 15$$

$$a = 15, b = 16$$

64 ordered pairs



Level up your prep for JEE with
ALLEN Online's LIVE JEE course!


Enrol Now

25. Let the three sides of a triangle ABC be given by the vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $3\hat{i} - 4\hat{j} - 4\hat{k}$.

Let G be the centroid of the triangle ABC. Then

$$6(\lvert\overrightarrow{AG}\rvert^2 + \lvert\overrightarrow{BG}\rvert^2 + \lvert\overrightarrow{CG}\rvert^2)$$

Ans. (164)



By given data

$$\overrightarrow{AB} + \overrightarrow{AC} = \overrightarrow{CB}$$

Let pv of \vec{A} are \vec{O} then

$$\overrightarrow{AB} = \vec{B} - \vec{A}$$

i.e. pv of \vec{B} = $2\hat{i} - \hat{j} + \hat{k}$

$$\overrightarrow{CA} = \vec{A} - \vec{C}$$

i.e. pv of \vec{C} = $-(\hat{i} - 3\hat{j} - 5\hat{k})$

Now pv of centroid

$$(\vec{G}) = \frac{\vec{A} + \vec{B} + \vec{C}}{3} = \frac{\vec{O} + (2, -1, 1) + (-1, 3, 5)}{3}$$

$$\vec{G} = \frac{1}{3}(\hat{i} + 2\hat{j} + 6\hat{k})$$

$$\text{Now } \overrightarrow{AG} = \frac{1}{3}(\hat{i} + 2\hat{j} + 6\hat{k})$$

$$\Rightarrow \lvert\overrightarrow{AG}\rvert^2 = \frac{1}{9} \times 41$$

$$\overrightarrow{BG} = \left(\frac{1}{3} - 2\right)\hat{i} + \left(\frac{2}{3} + 1\right)\hat{j} + (2 - 1)\hat{k}$$

$$\Rightarrow \lvert\overrightarrow{BG}\rvert^2 = \frac{59}{9}$$

$$\overrightarrow{CG} = \left(\frac{1}{3} + 1\right)\hat{i} + \left(\frac{2}{3} - 3\right)\hat{j} + (2 - 5)\hat{k}$$

$$\Rightarrow \lvert\overrightarrow{CG}\rvert^2 = \frac{146}{9}$$

Now

$$6(\lvert\overrightarrow{AG}\rvert^2 + \lvert\overrightarrow{BG}\rvert^2 + \lvert\overrightarrow{CG}\rvert^2) = 6 \times \left[\frac{41}{9} + \frac{59}{9} + \frac{146}{9} \right]$$

$$= 6 \times \frac{246}{9} = 164$$

Level up your prep for JEE with
ALLEN Online's LIVE JEE course!

Enrol Now



Level up your prep for JEE with our **LIVE JEE Courses!**



LIVE classes with top Kota faculty



ALLEN's **study material**



Tests with **national benchmarking**



ALLEN App Advantage: **24/7 doubt support,
Custom Practice & more**

Enrol Now



Secure up to

90% scholarship*

on our Online Courses!

*based on your
JEE Main 2025 scores!*

Enrol Now

