

DISCRETE-TO-CONTINUUM LIMITS OF LONG-RANGE ELECTRICAL INTERACTIONS IN NANOSTRUCTURES*

PRASHANT K. JHA[†], TIMOTHY BREITZMAN[‡], AND KAUSHIK DAYAL[§]

Abstract. We consider electrostatic interactions in two classes of nanostructures: (1) helical nanotubes, and (2) thin films with uniform bending (i.e., constant mean curvature). Starting from the atomic scale with a discrete distribution of dipoles, we obtain the continuum limit of the electrostatic energy; the continuum energy depends on the geometric parameters that define the nanostructure, such as the pitch and twist of the helical nanotubes and the curvature of the thin film. We find that the limiting energy is local in nature. This can be rationalized by noticing that the decay of the dipole kernel is sufficiently fast when the lattice sums run over one and two dimensions, and is also consistent with prior work on dimension reduction of continuum micromagnetic bodies to the thin film limit.

1. Introduction. Electrical and magnetic interactions are long-range; that is, a charge or dipole interacts with all the other charges and dipoles in the system, and the interactions cannot be truncated because the decay with distance is slow [Tou56, Bro63, JM94, MD14]. We consider such electrostatic interactions in nanostructures, specifically helical geometries and thin films with uniform bending. These geometries are ubiquitous in nanotechnology; while not periodic, their structure has significant symmetry that we exploit in this paper, using the framework of Objective Structures [Jam06]. We exploit this symmetry to adapt periodic calculations of the continuum energy to the setting of these nanostructures. Specifically, starting from a discrete atomic-scale description of the electrostatic energy, we find the limit energy when the discrete lengthscale of the nanostructures goes to zero.

For simplicity, we assume in this paper that the charge density can be approximated as composed of discrete dipoles. The electrostatic energy of such a system is the sum of all pairwise dipole-dipole interactions. Unlike short-range bonded atomic interactions that typically scale as r^{-6} with distance r , the dipole-dipole interactions decay slowly with distance as r^{-3} . Consequently, we cannot simply truncate after a few neighbors, and naive truncation can lead to qualitatively incorrect results in numerical calculations [MD14, GD20a, GD20b]. While we use the setting of discrete electrical dipoles, the setting of magnetic dipoles has an identical mathematical structure and physical interpretation [Bro63, JM94, MS02, SS09], and we borrow key ideas from that literature. Further, while discrete dipoles provide the simplest setting to illustrate the physics, it can be extended to the more realistic and general setting of a charge density field following ideas from [Xia05]. A key physical distinction between the electrical and magnetic situations is the possibility of electrical monopoles that does not exist for magnetic case, but we examine this elsewhere [SWB⁺21] and assume here that there are no free charges.

We turn to the question of dealing with the non-periodic geometry of the nanostructures. While neither helices nor thin films with curvature are periodic, the framework of Objective Structures (OS) introduced in [Jam06] provides a powerful approach to deal with such geometries. In brief, OS provides a group-theoretic description of these nanostructures that enables a parallel to be made with periodic lattices. This parallel to periodic lattices has enabled the adaptation of various methods developed for lattices to the setting of helices and thin-films, e.g. [DJ07, HTJ12, ADE13a, ADE13b]. Our strategy in this work is to use the OS framework to adapt continuum limit calculations from the

*Submitted to the editors December 16, 2020.

Funding: We acknowledge financial support from NSF (1635407), ARO (W911NF-17-1-0084, MURI W911NF-19-1-0245), ONR (N00014-18-1-2528), BSF (2018183), AFOSR (MURI FA9550-18-1-0095), and the AFRL Visiting Faculty program.

[†]Oden Institute for Computational Engineering and Sciences, The University of Texas at Austin (pjha@utexas.edu).

[‡]Air Force Research Laboratory (timothy.breiztman.1@us.af.mil).

[§]Center for Nonlinear Analysis, Department of Mathematical Sciences, Carnegie Mellon University; Department of Materials Science and Engineering, Carnegie Mellon University; Department of Civil and Environmental Engineering, Carnegie Mellon University (Kaushik.Dayal@cmu.edu).

setting of periodic lattices to the setting of nanostructures.

Our work is focused on obtaining discrete-to-continuum limits of the energy. This multiscale approach has proven very powerful in enabling the systematic reduction of the very large number of degrees of freedom associated to the discrete problem to a much more tractable continuum problem. This overall idea has played an important role in developing models, often in conjunction with variational tools such as Γ -convergence, both for bulk crystals [BLBL02, BLBL07, Sch09] as well as for thin films [Sch06]. Further, these ideas have played a role in the development of numerical multiscale atomistic methods such as the quasicontinuum method [TOP96, MT02, TM11, KO01, LLO12, DLO10].

However, all the work in the previous paragraph is restricted to the setting of short-range bonded atomic interactions. In the context of electrical and magnetic interactions, the calculation of continuum limit energies in the context of discrete-to-continuum has been examined both formally and rigorously using a discrete dipoles on a periodic 3-d lattice [Tou56, Bro63, JM94, MS02, SS09]. Further, this has been examined formally for periodic charge distributions, also in 3-d, [MD14]. All of these works show that the continuum limit energy consists of a local part and a nonlocal part. In contrast, in this work, we consider topologically low-dimensional structures: a 1-d helical nanotube and a 2-d thin film with constant bending curvature. In the limit that the discrete lengthscale characterizing the nanotube and thin film goes to 0, we find that the limit continuum energy is entirely local.

The absence of nonlocality in the limit can be rationalized by observing that the decay of the interactions as r^{-3} is sufficiently fast that we obtain a local limit if summed over a (topologically) 1-d or 2-d object. We highlight a complementary body of work that applies dimension reduction techniques to go from a 3-d continuum to a 2-d or 1-d continuum. In the context of electrical and magnetic interactions, [GJ97] and subsequent works [Car01, KS05, KSZ15] (for thin films) and [GH15, CH15] (for thin wires) find, as we do, that the limit energy is not nonlocal.

The techniques employed in this work are broadly based on the rigorous results provided in [JM94] on the continuum limit of magnetic dipole interactions on a 3-d lattice, with appropriate generalizations and modifications for our setting. The overall strategy of [JM94] is as follows. First, the operator that associates the discrete dipole lattice to the generated electric field is shown to be bounded for smooth test functions; next, the pointwise limit of the action of the operator on smooth test functions is obtained; and, finally, using the boundedness of the operator and the density of the test functions, the limit of the energy density is obtained. For the helical and thin film nanostructures considered in this work, we adapt this strategy to account for the fact that the lattice sites and dipoles are not related by a translation transformation, but by a more general isometric transformation.

The key results of this work are as follows. First, the limit energy is rigorously derived and found to be local. Second, the limiting energy density depends on the macroscopic geometric parameters, such as the pitch, radius and so on for the helical nanotube, and on the stretch and curvature for the thin film. These parameters can be related to macroscopic measures of deformation, and link the macroscopic deformation to the small-scale structure. Third, while the limiting energy is local, there are energetic contributions from both the normal and the tangential components of the dipole. This is in contrast to the result obtained by dimension reduction, and is due to the fact that we start with a single unit cell thickness in the normal direction and take the limit along the length (for the helix) or in the plane (for the thin film). In contrast, in dimension reduction approaches, there is no discreteness at all, and the limit is related to the ratio of the dimensions in and out of plane.

Organization. In section 2, we discuss prior work, primarily on dimension reduction from a 3-d continuum to a 2-d continuum, and highlight the local nature of the limiting energy. We then discuss heuristically the scaling of electrostatic interactions that lead to this locality generically for topologically low-dimensional nanostructures. In section 3, we present the main results for helical nanotubes and thin films with constant bending curvature. We prove various claims in section 4. In section 5, we summarize the results.

90 *Notation.* We denote the real line and set of integers by \mathbb{R} and \mathbb{Z} respectively; $\mathbb{R}^d, \mathbb{Z}^d$ denote
91 these in dimension $d = 1, 2, 3$. For any $c, c_1, c_2 \in \mathbb{R}$, $c\mathbb{Z}^d$ denotes the set $\{cz; z \in \mathbb{Z}^d\}$ and $c_1\mathbb{Z} \times c_2\mathbb{Z}$
92 denotes the set $\{(c_1z_1, c_2z_2); z_1, z_2 \in \mathbb{Z}\}$. \mathcal{L} and U denote the set of lattice sites and the lattice unit
93 cell respectively; \mathcal{L}_λ and U_λ denote these in the lattice scaled by λ , with \mathcal{L}_1, U_1 denoting $\mathcal{L}_\lambda, U_\lambda$
94 for $\lambda = 1$. We use $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$ to denote the point in space with components x_i in the
95 orthonormal basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ for \mathbb{R}^3 . We follow the standard notation wherein scalars are denoted
96 by lowercase letters, vectors by bold lowercase letters, and second order tensors by bold uppercase
97 letters. $|\mathbf{x}| = \sqrt{\sum_{i=1}^n x_i^2}$ denotes the Euclidean norm of vector $\mathbf{x} \in \mathbb{R}^n$; $|\mathbf{A}| = \sqrt{\mathbf{A} : \mathbf{A}}$ denotes the
98 norm of tensor \mathbf{A} ; and $\mathbf{A} : \mathbf{B} = A_{ij}B_{ij}$ denotes the inner product of the two tensors. For any vector
99 $\mathbf{a} \in \mathbb{R}^n$ and tensor \mathbf{A} , we have $|\mathbf{A}\mathbf{a}| \leq |\mathbf{A}||\mathbf{a}|$. We use $|\Omega|$ to denote the Lebesgue measure of the
100 set $\Omega \in \mathbb{R}^n$. We use $\chi_A = \chi_A(\mathbf{x})$ to denote the indicator function of set $A \subset \mathbb{R}^d$. We use $L^2(A, B)$
101 to denote the space of Lebesgue square-integrable functions $u: A \subset \mathbb{R}^n \rightarrow B \subset \mathbb{R}^m$; $(u, v)_{L^2(A, B)}$
102 for the inner product of functions $u, v \in L^2(A, B)$; and $\|u\|_{L^2(A, B)}$ the L^2 norm of $u \in L^2(A, B)$.
103 When there is no ambiguity, we will suppress $L^2(A, B)$ and write (u, v) and $\|u\|$. $C_0^\infty(\mathbb{R}^n, \mathbb{R}^m)$
104 denotes the space of infinitely differentiable test functions $u: \mathbb{R}^n \rightarrow \mathbb{R}^m$ with compact support in
105 \mathbb{R}^n . $\mathcal{L}(V, W)$ is the space of bounded linear maps $T: V \rightarrow W$ where V, W are Hilbert spaces. The
106 norm of the map $T \in \mathcal{L}(V, W)$ is denoted by $\|T\|_{\mathcal{L}(V, W)}$, and is given by the expression:

$$107 \quad (1.1) \quad \|T\|_{\mathcal{L}(V, W)} = \sup_{\|f\|_V \neq 0} \frac{\|Tf\|_W}{\|f\|_V}.$$

108 We use $u_\lambda \xrightarrow[\lambda \rightarrow 0]{} u$ to denote the strong convergence of $u_\lambda \in V$ to $u \in V$ as $\lambda \rightarrow 0$, i.e.,
109 $\|u_\lambda - u\|_V \rightarrow 0$ as $\lambda \rightarrow 0$.

110 **2. Prior Results on Dimension Reduction, and Dipole Interaction Scalings.** We briefly
111 revisit the results of [GJ97] and [CH15]. Respectively, they performed dimension reduction from the
112 3-d continuum to the 2-d thin film and 1-d thin wire limits for the magnetostatic energy.

113 Consider a material domain $\Omega_h = S \times [0, h]$, where $S \subset \mathbb{R}^2$ is a 2-d domain in the plane spanned
114 by $(\mathbf{e}_1, \mathbf{e}_2)$, and $h > 0$ is the material thickness in the normal direction \mathbf{e}_3 . Suppose $\mathbf{d}: \Omega_h \rightarrow \mathbb{R}^3$,
115 with $\mathbf{d} = \mathbf{0}$ on $\mathbb{R}^3 \setminus \Omega_h$, is the dipole field in the material. The electrostatic energy density is given by

$$116 \quad e_h(\mathbf{d}) = \frac{1}{|\Omega_h|} \int_{\Omega_h} \frac{1}{2} \nabla \phi(\mathbf{x}) \cdot \mathbf{d} \, d\mathbf{x},$$

117 where $|\Omega_h|$ is the volume of Ω_h , and ϕ is the electric potential that satisfies the electrostatic equation:

$$118 \quad \operatorname{div}(-\nabla \phi + \mathbf{d}) = 0 \quad \text{on } \mathbb{R}^3, \quad |\nabla \phi(\mathbf{x})| \rightarrow \mathbf{0} \text{ as } |\mathbf{x}| \rightarrow \infty$$

119 with $|\mathbf{d}| = d$. Let $\Omega_1 = S \times [0, 1]$, and $\mathbf{y}(\mathbf{x}) = (x_1, x_2, x_3/h) \in \Omega_1$ for $\mathbf{x} \in \Omega_h$ is the map from
120 Ω_h to Ω_1 . For fixed $h > 0$, consider the dipole field $\mathbf{d}_h: \Omega_h \rightarrow \mathbb{R}^3$ and $\tilde{\mathbf{d}}_h: \Omega_1 \rightarrow \mathbb{R}^3$ such that

$$121 \quad \tilde{\mathbf{d}}_h(\mathbf{y}(\mathbf{x})) = \mathbf{d}_h(\mathbf{x}), \quad \forall \mathbf{x} \in \Omega_h.$$

122 Let \mathbf{d}_h be the sequence of dipole field for $h > 0$, and $\tilde{\mathbf{d}}_h$ is defined as above. Assume that dipole
123 field $\tilde{\mathbf{d}}_h$ is such that, first, $\tilde{\mathbf{d}}_h = \mathbf{0}$ on $\mathbb{R}^3 \setminus \Omega_1$, and second, it converges to $\tilde{\mathbf{d}}_0$ in $L^2(\mathbb{R}^3)$; then the limit
124 of the energy density $e_h = e_h(\mathbf{d}_h)$ is [GJ97]:

$$125 \quad e_h(\mathbf{d}_h) \rightarrow e_0(\tilde{\mathbf{d}}_0) = \frac{1}{2|\Omega_1|} \int_{\Omega_1} |\tilde{\mathbf{d}}_0|^2 \, d\mathbf{x}$$

That is, the limiting energy e_0 is local, and only the normal component of the dipole moment appears in the expression.

Next, consider a thin straight wire with axis along e_1 , denoted by $\Omega_h = (-1, 1) \times B_2(\mathbf{0}, h)$, where $B_2(\mathbf{0}, h)$ is a ball of size h centered at $\mathbf{0}$ in the plane spanned by (e_2, e_3) . The limiting energy density is [CH15]:

$$\frac{1}{2|\Omega_1|} \int_{\Omega_1} (|\tilde{d}_{02}|^2 + |\tilde{d}_{03}|^2) \, d\mathbf{x},$$

where $\Omega_1 = (-1, 1) \times B_2(\mathbf{0}, 1)$ is the rescaled domain of Ω_h , and \tilde{d}_0 is the limiting field. We notice that the limiting energy is again local, and only the components of the dipole moment perpendicular to the wire appear.




Fig. 2.1: A schematic of the unit cell with a dipole (a), and generic 1-d, 2-d, 3-d periodic lattices (b, c, d).

The absence of nonlocality in the limiting energy in the results above, as well as in our results in section 3 below, can be physically understood through the fact that these structures are 1-d or 2-d topologically. To see this, we consider a system of discrete dipoles associated to the uniform 1-d, 2-d, and 3-d periodic lattices with the unit cell of size 1 (Figure 2.1). The energy of a lattice of dipoles is given by [Bro63, JM94]:

$$(2.1) \quad E = -\frac{1}{2} \sum_i \sum_{j, j \neq i} \mathbf{d}_i \cdot \mathbf{K}(\mathbf{x}_j - \mathbf{x}_i) \mathbf{d}_j = \sum_i |U_i| \left[-\frac{1}{|U_i|} \frac{1}{2} \sum_{j, j \neq i} \mathbf{d}_i \cdot \mathbf{K}(\mathbf{x}_j - \mathbf{x}_i) \mathbf{d}_j \right],$$

where the sum is over the cells in the lattice and the term inside square bracket denotes the energy density of a cell i . \mathbf{d}_i denotes the dipole in cell i , \mathbf{x}_i denotes the coordinate of lattice site i , and \mathbf{K} is the dipole field kernel defined as:

$$(2.2) \quad \mathbf{K}(\mathbf{x}) = -\frac{1}{4\pi|\mathbf{x}|^3} \left(\mathbf{I} - 3 \frac{\mathbf{x}}{|\mathbf{x}|} \otimes \frac{\mathbf{x}}{|\mathbf{x}|} \right), \quad \mathbf{x} \neq \mathbf{0}.$$

We use these expressions to heuristically understand the scaling of the energy for systems with different topological dimensions. For simplicity, we assume below that the volume of the unit cell and the magnitude of the dipole are both 1, i.e., $|U_i| = 1$ and $|\mathbf{d}_i| = 1$ for each i , and some constant factors are neglected.

REMARK 2.1 (1-d lattice). We can estimate an upper bound on the energy density e of a typical unit cell as follows:

$$e \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times |\mathbf{d}| \times (\text{number of dipoles at } r) \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times 1 \times 1 = \sum_{r=1}^{\infty} \frac{1}{r^3}.$$

We use that the total dipole moment at a distance r from a given unit cell is, at most, that of another dipole in the unit cell at a distance r . This sum is well-behaved and bounded.

REMARK 2.2 (2-d lattice). As in the 1-d setting, we first bound the net dipole at a distance r from a given unit cell. Since the structure is a 2-d lattice, the number of unit cells at a distance r is of order $2\pi r$. Therefore, an upper bound on the energy density is:

$$e \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times |\mathbf{d}| \times (\text{number of dipoles at } r) \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times 1 \times 2\pi r = 2\pi \sum_{r=1}^{\infty} \frac{1}{r^2}.$$

This sum is also well-behaved and bounded.

REMARK 2.3 (3-d lattice). Following the argument of the 2-d lattice, we now have that the net dipole at a distance r from a given unit cell is, at most, of the order $4\pi r^2$. Therefore, an upper bound on the energy density is:

$$e \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times |\mathbf{d}| \times (\text{number of dipoles at } r) \leq \sum_{r=1}^{\infty} \frac{1}{r^3} \times 1 \times 4\pi r^2 = 4\pi \sum_{r=1}^{\infty} \frac{1}{r}.$$

This sum is divergent. However, through a more careful analysis that accounts for the signs – not just the magnitudes – of the dipole interactions, the energy density can be shown to be conditionally convergent [Tou56, JM94].

When the lattice sum is bounded and converges unconditionally, it is possible to truncate after a finite distance and obtain sufficient numerical accuracy. When the lattice sum is conditionally convergent, that can be physically related to nonlocality; specifically, the slow convergence does not allow for truncation, and the far-field values play an important role. [MD14] discusses this from a physical perspective.

3. Results on Continuum Limits of the Electrostatic Energy. We consider two classes of nanostructures: helical nanotubes and thin films, the latter allowing for a constant bending curvature (i.e., nonzero constant mean curvature and zero Gauss curvature), and obtain the corresponding continuum limit electrostatic energy. In both cases, we start with discrete dipoles, where the discreteness is parametrized by the scale $\lambda > 0$, and examine the limit $\lambda \rightarrow 0$. We show that the dipole-dipole interaction energy density – per unit cross-sectional area in the case of nanotubes, and per unit thickness in the case of films – converges to a local energy density in the limit.

3.1. Helical Nanotube. We consider a discrete helix with axis \mathbf{e}_3 characterized by the angle θ and length δ . Suppose $\mathbf{x}_0 \in \mathbb{R}^3$ is a point on the helix. Then, the other points on the helix are related by an isometric transformation of \mathbf{x}_0 . Let $s \in \mathbb{R}$ be the parametric coordinate of a point on the helix. Then, the map $\bar{\mathbf{x}}: \mathbb{R} \rightarrow \mathbb{R}^3$ that takes a point in the parametric space to a unique point on the helix can be expressed as

$$(3.1) \quad \bar{\mathbf{x}}(s) = \mathbf{Q}(s\theta)\mathbf{x}_0 + s\delta\mathbf{e}_3.$$

Here $\mathbf{Q}(\theta)$ is the rotational tensor represented by the matrix in the orthonormal basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ as:

$$(3.2) \quad \mathbf{Q}(\theta) := \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We notice that these definitions imply that the pitch of the helix is $2\pi\delta/\theta$.

Without loss of generality, we assume $\mathbf{x}_0 = \mathbf{e}_1$. The tangent vector to the helix at s is given by

$$(3.3) \quad \mathbf{t}(s) = \frac{d\bar{\mathbf{x}}(s)}{ds} = \theta \mathbf{Q}'(s\theta) \mathbf{e}_1 + \delta \mathbf{e}_3.$$

Let $\hat{\mathbf{t}}(s) = \mathbf{t}(s)/\sqrt{\theta^2 + \delta^2}$ denote the unit tangent vector. We define the second order projection tensors $\mathbf{P}_{||} = \mathbf{P}_{||}(s)$ and $\mathbf{P}_{\perp} = \mathbf{P}_{\perp}(s)$, for $s \in \mathbb{R}$, as follows

$$(3.4) \quad \mathbf{P}_{||}(s) = \hat{\mathbf{t}}(s) \otimes \hat{\mathbf{t}}(s), \quad \mathbf{P}_{\perp}(s) = \mathbf{I} - \mathbf{P}_{||}(s).$$

For any vector \mathbf{a} and any $s \in \mathbb{R}$, we have

$$(3.5) \quad \mathbf{a} = \mathbf{P}_{||}(s) \mathbf{a} + \mathbf{P}_{\perp}(s) \mathbf{a}, \quad \text{with} \quad \mathbf{P}_{||}(s) \mathbf{a} \cdot \mathbf{P}_{\perp}(s) \mathbf{a} = 0.$$

3.1.1. Lattice Geometry and Dipole Moment. Let $\mathcal{L} = \mathbb{Z}$ denote the set of parametric coordinates of the points on helix. We consider a discrete system of dipole moments $\mathbf{d}: \mathcal{L} \rightarrow \mathbb{R}^3$ associated to the points on the helix given by \mathcal{L} , see Figure 3.1. The magnitudes of the dipoles at the lattice sites are equal, but they are oriented differently; in particular, the orientations of dipoles at lattice sites follow the relation:

$$(3.6) \quad \mathbf{d}(s+1) = \mathbf{Q}(\theta) \mathbf{d}(s), \quad s \in \mathcal{L}$$

We associate a unit cell to each lattice site. Let $U(s) = [s, s+1)$ denote the unit cell in the parametric space at the site s , for $s \in \mathcal{L}$. Let $S(r)$, $r \in \mathbb{R}$, be given by

$$S(r) = \{\mathbf{x}; (\mathbf{x} - \bar{\mathbf{x}}(r)) \cdot \mathbf{t}(r) = 0, |\mathbf{x} - \bar{\mathbf{x}}(r)|^2 < R^2\}.$$

Note that $|S(r)| = |S(0)| = \pi R^2$. The unit cell in real space is defined by $\bar{U}(s) = \{\mathbf{x} \in S(r); r \in U(s)\}$.

We take, without loss of generality, $R^2 = 1/(\pi\sqrt{\theta^2 + \delta^2})$ so that $|\bar{U}(s)| = \text{area}(S) \times \text{length}(\{\bar{\mathbf{x}}(r); r \in U(s)\}) = \pi R^2 \sqrt{\theta^2 + \delta^2} = 1$.

We now consider the setting in which the cells are of size $\lambda > 0$ so that as $\lambda \rightarrow 0$ the density of cells in the helix increases. For $\lambda > 0$, suppose $\mathcal{L}_{\lambda} = \lambda\mathbb{Z}$ denotes the parametric coordinates of the sites in a scaled lattice, and $\mathbf{d}_{\lambda}: \mathcal{L}_{\lambda} \rightarrow \mathbb{R}^3$ denotes the corresponding system of dipole moments. Associated to $s \in \mathcal{L}_{\lambda}$, let $U_{\lambda}(s) = [s, s+\lambda)$ denote the cell in the parametric space. The 3-d cell is given by $\bar{U}_{\lambda}(s) = \{\mathbf{x} \in S_{\lambda}(r); r \in U_{\lambda}(s)\}$, where $S_{\lambda}(r) = \{\mathbf{x}; (\mathbf{x} - \bar{\mathbf{x}}(r)) \cdot \mathbf{t}(r) = 0, |\mathbf{x} - \bar{\mathbf{x}}(r)|^2 < \lambda^2 R^2\}$ is the scaled cross-section. Note that $\text{area}(S_{\lambda}(r)) = \pi \lambda^2 R^2$ and $|\bar{U}_{\lambda}(s)| = \pi \lambda^2 R^2 \times \lambda \sqrt{\theta^2 + \delta^2} = \lambda^3$. Let $\tilde{\mathbf{d}}_{\lambda}: \mathbb{R} \rightarrow \mathbb{R}^3$ be the piecewise constant extension of \mathbf{d}_{λ} given by

$$(3.7) \quad \tilde{\mathbf{d}}_{\lambda}(s) = \frac{\mathbf{d}_{\lambda}(i)}{|\bar{U}_{\lambda}(s)|} = \frac{\mathbf{d}_{\lambda}(i)}{\lambda^3}, \quad \forall s \in U_{\lambda}(i), \quad \forall i \in \mathcal{L}_{\lambda}.$$

To compute the limit of the dipole-dipole interaction energy as $\lambda \rightarrow 0$, we assume that dipole moment density field $\tilde{\mathbf{d}}_{\lambda}$ converges to some field $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$ in the L^2 norm. As in [JM94], instead of

215 working with $\tilde{\mathbf{d}}_\lambda$, as defined above, we could assume that the dipole moment $\mathbf{d}_\lambda(i)$, for $i \in \mathcal{L}_\lambda$, is
 216 due to the background dipole moment density field $\mathbf{f}_\lambda \in L^2(\mathbb{R}, \mathbb{R}^3)$ such that

$$217 \quad (3.8) \quad \mathbf{d}_\lambda(i) = \sqrt{\theta^2 + \delta^2} \int_{U_\lambda(i)} \int_{S_\lambda(r)} \mathbf{f}_\lambda(r) \, dS_\lambda(r) \, dr = \lambda^2 \int_{U_\lambda(i)} \mathbf{f}_\lambda(r) \, dr,$$

218 where $dS_\lambda(r)$ is the area measure for surface S_λ . We assumed that the background field is uniform
 219 in $S_\lambda(r)$ for all $r \in \mathbb{R}$ and used $R^2 = 1/(\pi\sqrt{\theta^2 + \delta^2})$. The existence of one such background field
 220 \mathbf{f}_λ is evident: we can define $\mathbf{f}_\lambda = \tilde{\mathbf{d}}_\lambda$. The physical dimension of \mathbf{f}_λ is dipole moment per unit
 221 volume. We have the following lemma that relates the convergence of the background dipole moment
 222 field and the piecewise constant extension.

223 **LEMMA 3.1.** *Let \mathbf{f}_λ , $\lambda > 0$, be the sequence of $L^2(\mathbb{R}, \mathbb{R}^3)$ functions and let $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$ such
 224 that $\mathbf{f}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathbb{R}, \mathbb{R}^3)$. Let $\mathbf{d}_\lambda: \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ be given by (3.8) and let $\tilde{\mathbf{d}}_\lambda$ be a piecewise constant
 225 L^2 extension of \mathbf{d}_λ given by (3.7). Then, $\tilde{\mathbf{d}}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathbb{R}, \mathbb{R}^3)$.*

226 *On the other hand, if $\mathbf{d}_\lambda: \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ is such that $\tilde{\mathbf{d}}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathbb{R}, \mathbb{R}^3)$, then there exists a
 227 background field $\mathbf{f}_\lambda \in L^2(\mathbb{R}, \mathbb{R}^3)$ such that \mathbf{d}_λ is given by (3.8).*

228 The proof is similar to the proof of Theorem 4.1 from [JM94].

229 **REMARK 3.1.** *Since the discrete dipole field \mathbf{d}_λ has helical symmetry, from (3.8) we can see that
 230 \mathbf{f} will also have helical symmetry.*




Fig. 3.1: Discrete dipole moments (red arrows) lying on the helix. (a) and (b) show the view in (e_1, e_3) and (e_1, e_2) planes respectively. The dipole moments corresponding to different sites are related by (3.6). For the parametric coordinate s , $\bar{\mathbf{x}}(s)$ gives the coordinate of the point on the helix.

231 **3.1.2. Electrostatic Energy.** For $\lambda > 0$, the energy associated to the system of dipole moments
 232 \mathbf{d}_λ can be expressed as [Bro63, JM94]:

$$233 \quad E_\lambda = -\frac{1}{2} \sum_{\substack{s, s' \in \mathcal{L}_\lambda, \\ s \neq s'}} \mathbf{d}_\lambda(s) \cdot \mathbf{K}(\bar{\mathbf{x}}(s') - \bar{\mathbf{x}}(s)) \mathbf{d}_\lambda(s') = |S_\lambda| e_\lambda,$$

234 where e_λ is the energy per unit area given by

$$235 \quad (3.9) \quad e_\lambda = -\frac{1}{2|S_\lambda|} \sum_{\substack{s, s' \in \mathcal{L}_\lambda, \\ s \neq s'}} \mathbf{d}_\lambda(s) \cdot \mathbf{K}(\bar{\mathbf{x}}(s') - \bar{\mathbf{x}}(s)) \mathbf{d}_\lambda(s').$$

236 Substituting (3.8) above and proceeding similar to Section 6 of [JM94], e_λ can be written as

$$237 \quad (3.10) \quad e_\lambda = (\mathbf{f}_\lambda, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)},$$

238 where $T_\lambda: L^2(\mathbb{R}, \mathbb{R}^3) \rightarrow L^2(\mathbb{R}, \mathbb{R}^3)$ is the map given by

$$239 \quad (3.11) \quad (T_\lambda \mathbf{f})(s) = \lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) \mathbf{f}(s') \, ds'$$

240 and $\mathbf{K}_\lambda(s', s)$, for $s, s' \in \mathbb{R}$, is the discrete dipole field kernel given by

$$241 \quad (3.12) \quad \mathbf{K}_\lambda(s', s) = \sum_{\substack{u, v \in \mathcal{L}_\lambda, \\ u \neq v}} \chi_{U_\lambda(v)}(s') \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u)) \chi_{U_\lambda(u)}(s).$$

242 *Scaling Property of \mathbf{K}_λ .* For any $a, b \in \mathbb{R}$, we have

$$243 \quad (3.13) \quad \begin{aligned} \bar{\mathbf{x}}(\lambda a) - \bar{\mathbf{x}}(\lambda b) &= \mathbf{Q}(\lambda a \theta) \mathbf{e}_1 + \delta \lambda a \mathbf{e}_3 - \mathbf{Q}(\lambda b \theta) \mathbf{e}_1 - \delta \lambda b \mathbf{e}_3 \\ &= \lambda \left(\bar{\mathbf{x}}(a) - \bar{\mathbf{x}}(b) + \underbrace{\left[\frac{\mathbf{Q}(\lambda a \theta) - \mathbf{Q}(\lambda b \theta) - (\lambda \mathbf{Q}(a \theta) - \lambda \mathbf{Q}(b \theta))}{\lambda} \right]}_{=: \mathbf{A}_\lambda(a, b)} \mathbf{e}_1 \right). \end{aligned}$$

244 Using the relation above, it is easy to show

$$245 \quad \mathbf{K}_\lambda(s', s) = \frac{1}{\lambda^3} \sum_{\substack{u, v \in \mathcal{L}_1, \\ u \neq v}} \chi_{U_1(v)}(s'/\lambda) \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u) + \mathbf{A}_\lambda(v, u) \mathbf{e}_1) \chi_{U_1(u)}(s/\lambda),$$

246 where we recall that $U_1(u) = [u, u+1)$, $u \in \mathcal{L}_1$, is the lattice cell in the parametric space for $\lambda = 1$.

247 We define a discrete kernel $\mathbf{K}_{1,\lambda}(s, s')$, for $s, s' \in \mathbb{R}$, as follows

$$248 \quad (3.14) \quad \mathbf{K}_{1,\lambda}(s', s) = \sum_{\substack{u, v \in \mathcal{L}_1, \\ u \neq v}} \chi_{U_1(v)}(s') \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u) + \mathbf{A}_\lambda(v, u) \mathbf{e}_1) \chi_{U_1(u)}(s).$$

249 We then have

$$250 \quad \mathbf{K}_\lambda(s', s) = \frac{1}{\lambda^3} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda).$$

251 **3.1.3. Limit of Electrostatics Energy.** In this section, we obtain the limit of the energy per
 252 unit surface area e_λ as $\lambda \rightarrow 0$ assuming that the background dipole field density (or equivalently the
 253 dipole moment density $\tilde{\mathbf{d}}_\lambda$) \mathbf{f}_λ converges to some density field \mathbf{f} in L^2 . The idea is to first show that
 254 the map T_λ in (3.11) is bounded and obtain its limit. With that, the limit of e_λ follows.

255 *Limit of Discrete Electric Field.* Let $T_{1,\lambda}$ be the map with kernel $\mathbf{K}_{1,\lambda}$. For any function
 256 $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$, we have

$$257 \quad (3.15) \quad (T_{1,\lambda}\mathbf{f})(s) = \int_{\mathbb{R}} \mathbf{K}_{1,\lambda}(s', s) \mathbf{f}(s') \, ds'.$$

258 We have the following main result on the map T_λ .

259 **PROPOSITION 3.2.** *The map $T_{1,\lambda}$ and T_λ are bounded in $L^2(\mathbb{R}, \mathbb{R}^3)$ for all $\lambda > 0$ and satisfy*

$$260 \quad (3.16) \quad \|T_\lambda\|_{\mathcal{L}(L^2, L^2)} = \|T_{1,\lambda}\|_{\mathcal{L}(L^2, L^2)}.$$

261 *Further, for $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$,*

$$262 \quad (T_\lambda\mathbf{f})(s) \xrightarrow{\lambda \rightarrow 0} -h_0(\mathbf{I} - 3\mathbf{P}_\parallel(s))\mathbf{f}(s) = -h_0(\mathbf{P}_\perp(s) - 2\mathbf{P}_\parallel(s))\mathbf{f}(s)$$

263 *pointwise, where $\mathbf{P}_\perp(s)$ and $\mathbf{P}_\parallel(s)$ are projection tensors that project onto the normal plane and*
 264 *the tangent line to the helix respectively (see (3.4)). h_0 is a constant given by*

$$265 \quad (3.17) \quad h_0 = \sum_{\substack{v \in \mathbb{Z}, \\ v \neq 0}} \frac{1}{4\pi|v|^3(\theta^2 + \delta^2)^{3/2}}.$$

266 We provide the proof of [Proposition 3.2](#) in [subsection 4.1](#).

267 *Limit of Energy.*

268 **THEOREM 3.3.** *Let $\mathbf{f}_\lambda \in L^2(\mathbb{R}, \mathbb{R}^3)$ be a sequence of functions for $\lambda > 0$ with $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$*
 269 *such that $\mathbf{f}_\lambda \xrightarrow{\lambda \rightarrow 0} \mathbf{f}$ in L^2 . Let the system of dipole moments $\mathbf{d}_\lambda: \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ be given by (3.8). Then*

$$270 \quad e_\lambda \xrightarrow{\lambda \rightarrow 0} \frac{1}{2}h_0 \left[\|\mathbf{P}_\perp \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}^2 - 2\|\mathbf{P}_\parallel \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}^2 \right],$$

272 *where h_0 is a constant defined in (3.17).*

273 *Proof.* Since T_λ is bounded and $\mathbf{f}_\lambda \rightarrow \mathbf{f}$, we have

$$274 \quad \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}_\lambda) = \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}) + \lim_{\lambda \rightarrow 0} (T_\lambda (\mathbf{f}_\lambda - \mathbf{f})) = \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}).$$

275 We only need to analyze $T_\lambda \mathbf{f}$ in the limit for $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$. Let $\mathbf{f}^k \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$ be a sequence
 276 of functions such that $\mathbf{f}^k \rightarrow \mathbf{f}$. Using [Proposition 3.2](#), we have

$$\begin{aligned} 277 \quad \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}) &= \lim_{k \rightarrow \infty} \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}^k) + \lim_{k \rightarrow \infty} \lim_{\lambda \rightarrow 0} (T_\lambda (\mathbf{f} - \mathbf{f}^k)) = \lim_{k \rightarrow \infty} \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f}^k) \\ 278 \quad (3.18) \quad &= \lim_{k \rightarrow \infty} (\mathbf{H}_0 \mathbf{f}^k) = \mathbf{H}_0 \mathbf{f}. \end{aligned}$$

280 Using the expression in (3.10) for e_λ , we have

$$\begin{aligned} 281 \quad e_\lambda &= -\frac{1}{2} (\mathbf{f}_\lambda, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)} = -\frac{1}{2} \left[(\mathbf{f}_\lambda - \mathbf{f}, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)} + (\mathbf{f}, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)} \right] \\ 282 \quad &= -\frac{1}{2} \left[(\mathbf{f}_\lambda - \mathbf{f}, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)} + (\mathbf{f}, T_\lambda \mathbf{f})_{L^2(\mathbb{R}, \mathbb{R}^3)} + (\mathbf{f}, T_\lambda (\mathbf{f}_\lambda - \mathbf{f}))_{L^2(\mathbb{R}, \mathbb{R}^3)} \right]. \end{aligned}$$

284 The first and third terms are zero in the limit. Taking the limit of the remaining term and using (3.18),
 285 we have

$$286 \quad \lim_{\lambda \rightarrow 0} e_\lambda = \lim_{\lambda \rightarrow 0} -\frac{1}{2} (\mathbf{f}_\lambda, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathbb{R}, \mathbb{R}^3)} = \frac{1}{2} h_0 \left[\|\mathbf{P}_\perp \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}^2 - 2\|\mathbf{P}_\parallel \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}^2 \right].$$

287 This completes the proof. □

REMARK 3.2. *The limiting energy only comprises of a local self-field energy. In the limit, any point on the helix sees the uniform 1-d system of dipole moments along the tangent line. Further, we see that both the normal components and the tangential component of the dipole moment contribute to the energy and electric field. This is in contrast to [CH15] where the thin wire limit of the magnetostatic energy has contribution only from the normal components.*

3.2. Nanofilm with Constant Bending Curvature. Let $S = (-\bar{\theta}, \bar{\theta}) \times \mathbb{R}$ be the parametric space for a surface with a constant bending curvature κ . The map that takes a point in the parametric space to a unique point on the film is given by

$$(3.19) \quad \bar{\mathbf{x}}(s_1, s_2) = \mathcal{R}\mathbf{Q}(s_1)\mathbf{e}_1 + s_2\delta\mathbf{e}_3,$$

where $\mathcal{R} = 1/\kappa$ is the inverse of curvature, $\bar{\theta} > 0$ is the angular size of the film, and δ is the spacing in the flat direction. $\kappa, \delta, \bar{\theta}$ are fixed parameters for a given film. Here, $\mathbf{Q}(\theta)$ is the rotational tensor with the axis \mathbf{e}_3 , see definition (3.2). The tangent vectors at $\mathbf{s} := \{s_1, s_2\} \in S$ are

$$(3.20) \quad \mathbf{t}_1(\mathbf{s}) = \frac{d\bar{\mathbf{x}}}{ds_1} = \mathcal{R}\mathbf{Q}'(s_1)\mathbf{e}_1, \quad \mathbf{t}_2(\mathbf{s}) = \frac{d\bar{\mathbf{x}}}{ds_2} = \delta\mathbf{e}_3$$

and the normal vector is

$$(3.21) \quad \mathbf{n}(\mathbf{s}) = \mathbf{Q}(s_1)\mathbf{e}_1.$$

3.2.1. Lattice Geometry and Dipole Moment. We consider a lattice embedded on the film $\bar{\mathbf{x}}$. We assume that the lattice is one lattice cell thick in the direction \mathbf{n} normal to the film. Suppose $\mathcal{L} \subset S$ is the set of parametric coordinates of the discrete lattice sites. Let \mathcal{L} and the lattice cell U (in the parametric space S) be given by

$$(3.22) \quad \begin{aligned} \mathcal{L} &= \{\mathbf{s} = (s_1, s_2) \in S; s_1 = i\theta_l, s_2 = j, i, j \in \mathbb{Z}\} = (-\bar{\theta}, \bar{\theta}) \cap \theta_l\mathbb{Z} \times \mathbb{Z} \\ U(\mathbf{s}) &= [s_1, s_1 + \theta_l] \times [s_2, s_2 + 1), \quad \forall \mathbf{s} \in \mathcal{L}. \end{aligned}$$

Here, θ_l is the angular width of the lattice cell. We assume that θ_l is such that the set of sites in the angular direction, $(-\bar{\theta}, \bar{\theta}) \cap \theta_l\mathbb{Z}$, is not empty, and in fact is sufficiently large so that the continuum limit approximation of the energy density is justified. We assume that the lattice has unit thickness in the normal direction, and suppose that the film given by $\bar{\mathbf{x}}$ passes through the center of the lattice in the normal direction. Then, the unit cell for a given site $\mathbf{s} \in \mathcal{L}$ is $\bar{U}(\mathbf{s}) = \{\mathbf{x}; \mathbf{x} = \bar{\mathbf{x}}(\mathbf{s}) + t\mathbf{n}(\mathbf{s}), \mathbf{s} \in U(\mathbf{s}), t \in (-1/2, 1/2)\}$. On the lattice \mathcal{L} , we define a discrete system of dipole moments $\mathbf{d}: \mathcal{L} \rightarrow \mathbb{R}^3$. Similar to the case of the helical nanotube, the lattice cells in real space are related by an isometric transformation, so the magnitudes of the dipoles at the lattice sites are equal, but they are oriented differently. In particular, we have

$$(3.23) \quad \mathbf{d}(\mathbf{s} + \mathbf{r}) = \mathbf{Q}(r_1)\mathbf{d}(\mathbf{s}), \quad \mathbf{s} \in \mathcal{L},$$

where $\mathbf{r} = (r_1, r_2) \in \mathcal{L}$ such that $\mathbf{r} + \mathbf{s} \in \mathcal{L}$ (i.e. all the translations within \mathcal{L}). We see that the dipole orientation depends only on the angular (first) parameter and is invariant with respect to the second parameter.

We next consider the setting when the lattice is scaled by $\lambda > 0$. The scaled lattice \mathcal{L}_λ and the associated lattice cell U_λ are defined by the natural scaling of \mathcal{L} and U as follows

$$(3.24) \quad \begin{aligned} \mathcal{L}_\lambda &= \{\mathbf{s} \in S; s_1 = i\theta_l\lambda, s_2 = j\lambda, i, j \in \mathbb{Z}\} = (-\bar{\theta}, \bar{\theta}) \cap \lambda\theta_l\mathbb{Z} \times \lambda\mathbb{Z} \\ U_\lambda(\mathbf{s}) &= [s_1, s_1 + \lambda\theta_l] \times [s_2, s_2 + \lambda), \quad \forall \mathbf{s} \in \mathcal{L}_\lambda. \end{aligned}$$

After scaling, the thickness of the lattice cell in the normal direction is λ and the unit cell for $\mathbf{s} \in \mathcal{L}_\lambda$ is $\{\mathbf{x} \in \mathbb{R}^3; \mathbf{x} = \bar{\mathbf{x}}(\mathbf{s}) + t\mathbf{n}(\mathbf{s}), \mathbf{s} \in U_\lambda(\mathbf{s}), t \in (-\lambda/2, \lambda/2)\}$. We can show that the unit cell in




Fig. 3.2: Discrete dipole moments on a nanofilm with uniform bending curvature. (a) and (b) show the view from different perspectives.

the scaled lattice has volume $\lambda^3 \mathcal{R} \theta_l$. Suppose $\tilde{\mathbf{d}}_\lambda : \mathcal{S} \rightarrow \mathbb{R}$ denotes the piecewise constant extension of \mathbf{d}_λ and is given by

$$(3.25) \quad \tilde{\mathbf{d}}_\lambda(\mathbf{s}) = \frac{\mathbf{d}_\lambda(\mathbf{a})}{\lambda^3 \mathcal{R} \theta_l}, \quad \forall \mathbf{s} \in U_\lambda(\mathbf{a}), \forall \mathbf{a} \in \mathcal{L}_\lambda.$$

We are interested in the limit of the energy when $\tilde{\mathbf{d}}_\lambda$ converges to \mathbf{f} in $L^2(\mathcal{S}, \mathbb{R}^3)$. As in the case of the helix and following [JM94], we suppose that there exists a background dipole moment density field $\mathbf{f}_\lambda \in L^2(\mathcal{S}, \mathbb{R}^3)$ such that the dipole moment at site $\mathbf{s} \in \mathcal{L}_\lambda$ is given by

$$(3.26) \quad \mathbf{d}_\lambda(\mathbf{s}) = \int_{-\lambda/2}^{\lambda/2} \left[\int_{U_\lambda(\mathbf{s})} \mathbf{f}_\lambda(\mathbf{t}) \mathcal{R} \, dt_1 \, dt_2 \right] dt_3 = \mathcal{R} \lambda \int_{U_\lambda(\mathbf{s})} \mathbf{f}_\lambda(\mathbf{t}) \, dt,$$

where $d\mathbf{t} = dt_1 \, dt_2$ is the area measure (note that $d\mathbf{t}$ does not include \mathcal{R}). The existence of one such background field \mathbf{f}_λ is evident: We can define $\mathbf{f}_\lambda = \tilde{\mathbf{d}}_\lambda$. Similar to the case of the helix, we have the following lemma that relates the convergence of the background dipole moment field and the piecewise constant extension.

LEMMA 3.4. *Let \mathbf{f}_λ , $\lambda > 0$, be a sequence of $L^2(\mathcal{S}, \mathbb{R}^3)$ functions and let $\mathbf{f} \in L^2(\mathcal{S}, \mathbb{R}^3)$ be such that $\mathbf{f}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathcal{S}, \mathbb{R}^3)$. Let $\mathbf{d}_\lambda : \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ be given by (3.26) and let $\tilde{\mathbf{d}}_\lambda$ be a piecewise constant L^2 extension of \mathbf{d}_λ given by (3.25). Then, $\tilde{\mathbf{d}}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathcal{S}, \mathbb{R}^3)$.*

On the other hand, if $\mathbf{d}_\lambda : \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ is such that $\tilde{\mathbf{d}}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathcal{S}, \mathbb{R}^3)$ then there exists a background field $\mathbf{f}_\lambda \in L^2(\mathcal{S}, \mathbb{R}^3)$ such that \mathbf{d}_λ is given by (3.26).

The proof follows directly from the proof of Theorem 4.1 of [JM94].

REMARK 3.3. *As different unit cells are related by isometric transformations, the dipole moments in different unit cells are related by the rotational part of the isometric transformation.*

3.2.2. Electrostatic Energy. As before, the energy associated to the system of dipole moments \mathbf{d}_λ , for $\lambda > 0$, is given by

$$(3.27) \quad E_\lambda = -\frac{1}{2} \sum_{\substack{\mathbf{s}, \mathbf{s}' \in \mathcal{L}_\lambda, \\ \mathbf{s} \neq \mathbf{s}'}} \mathbf{d}_\lambda(\mathbf{s}) \cdot \mathbf{K}(\bar{\mathbf{x}}(\mathbf{s}') - \bar{\mathbf{x}}(\mathbf{s})) \mathbf{d}_\lambda(\mathbf{s}') = |(-\lambda/2, \lambda/2)| \hat{e}_\lambda,$$

where $|(-\lambda/2, \lambda/2)| = \lambda$ is the thickness of the lattice in normal direction, and \hat{e}_λ is the energy per unit length given by

$$(3.27) \quad \hat{e}_\lambda = -\frac{1}{2\lambda} \sum_{\substack{s, s' \in \mathcal{L}_\lambda, \\ s \neq s'}} \mathbf{d}_\lambda(s) \cdot \mathbf{K}(\bar{\mathbf{x}}(s') - \bar{\mathbf{x}}(s)) \mathbf{d}_\lambda(s').$$

For convenience, we normalize \hat{e}_λ by $\mathcal{R}\theta_l$, where $\mathcal{R}\theta_l$ is independent of λ and gives the size of original lattice in the angular direction. We let

$$(3.28) \quad e_\lambda = \frac{\hat{e}_\lambda}{\mathcal{R}\theta_l} \Rightarrow E_\lambda = \lambda(\mathcal{R}\theta_l)e_\lambda.$$

Substituting (3.26) and proceeding similar to the case of the helix, we can express e_λ as

$$(3.29) \quad e_\lambda = (\mathbf{f}_\lambda, T_\lambda \mathbf{f}_\lambda)_{L^2(\mathcal{S}, \mathbb{R}^3)},$$

where $T_\lambda: L^2(\mathcal{S}, \mathbb{R}^3) \rightarrow L^2(\mathcal{S}, \mathbb{R}^3)$ is the map defined as

$$(3.30) \quad (T_\lambda \mathbf{f})(s) = \frac{\mathcal{R}}{\theta_l} \lambda \int_{\mathcal{S}} \mathbf{K}_\lambda(s', s) \mathbf{f}(s') \, ds'$$

and $\mathbf{K}_\lambda(s', s)$, for $s, s' \in \mathcal{S}$, is the discrete dipole field kernel given by

$$(3.31) \quad \mathbf{K}_\lambda(s', s) = \sum_{\substack{u, v \in \mathcal{L}_\lambda, \\ u \neq v}} \chi_{U_\lambda(v)}(s') \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u)) \chi_{U_\lambda(u)}(s).$$

Scaling Property of \mathbf{K}_λ . As in the case of the helix, it is convenient to first rescale the lattice \mathcal{L}_λ such that the lattice cell size is independent of λ after rescaling, and define a new map on the rescaled lattice. This is considered next.

Let $\mathcal{S}_{1,\lambda} = (-\bar{\theta}/\lambda, \bar{\theta}/\lambda) \times \mathbb{R}$ so that $s \in \mathcal{S}$ implies $s/\lambda \in \mathcal{S}_{1,\lambda}$. We define a rescaled lattice $\mathcal{L}_{1,\lambda}$ such that $s \in \mathcal{L}_\lambda$ implies $s/\lambda \in \mathcal{L}_{1,\lambda}$. It is given by

$$(3.32) \quad \mathcal{L}_{1,\lambda} = \{s \in \mathcal{S}_{1,\lambda}; s_1 = i\theta_l, s_2 = j, i, j \in \mathbb{Z}\} = (-\bar{\theta}/\lambda, \bar{\theta}/\lambda) \cap \theta_l \mathbb{Z} \times \mathbb{Z}.$$

The lattice cell for $s \in \mathcal{L}_{1,\lambda}$ is given by $U_1(s)$, where $U_1(s)$ is defined in (3.22) (using $\lambda = 1$ in U_λ).

For $\mathbf{a}, \mathbf{b} \in \mathcal{S}_{1,\lambda}$, we have

$$(3.33) \quad \bar{\mathbf{x}}(\lambda \mathbf{a}) - \bar{\mathbf{x}}(\lambda \mathbf{b}) = \lambda (\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b}) \mathbf{e}_1),$$

where

$$(3.34) \quad \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b}) = \frac{\mathcal{R}}{\lambda} [\mathbf{Q}(\lambda \mathbf{a}_1) - \mathbf{Q}(\lambda \mathbf{b}_1) - \lambda \mathbf{Q}(\mathbf{a}_1) + \lambda \mathbf{Q}(\mathbf{b}_1)].$$

Keeping in mind these definitions, for $\mathbf{u} \in \mathcal{L}_{1,\lambda}$, we also note

$$(3.35) \quad \chi_{U_\lambda(\lambda \mathbf{u})}(s) = \begin{cases} 1 & \text{if } s \in U_\lambda(\lambda \mathbf{u}), \\ 0 & \text{otherwise} \end{cases} = \begin{cases} 1 & \text{if } s/\lambda \in U_1(\mathbf{u}), \\ 0 & \text{otherwise} \end{cases} = \chi_{U_1(\mathbf{u})}(s/\lambda).$$

Using the above relation and (3.33), we can show, for any $s, s' \in \mathcal{S}$,

$$(3.36) \quad \mathbf{K}_\lambda(s', s) = \frac{1}{\lambda^3} \sum_{\substack{u, v \in \mathcal{L}_{1,\lambda}, \\ u \neq v}} \chi_{U_1(v)}(s'/\lambda) \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u) + \mathbf{A}_\lambda(v, u) \mathbf{e}_1) \chi_{U_1(u)}(s/\lambda).$$

380 If we introduce the discrete dipole field kernel $\mathbf{K}_{1,\lambda}(\mathbf{s}', \mathbf{s})$, for $\mathbf{s}, \mathbf{s}' \in \mathcal{S}_{1,\lambda}$, defined on $\mathcal{L}_{1,\lambda}$ as:

$$381 \quad (3.36) \quad \mathbf{K}_{1,\lambda}(\mathbf{s}', \mathbf{s}) = \sum_{\substack{\mathbf{u}, \mathbf{v} \in \mathcal{L}_{1,\lambda}, \\ \mathbf{u} \neq \mathbf{v}}} \chi_{U_1(\mathbf{v})}(\mathbf{s}') \mathbf{K}(\bar{\mathbf{x}}(\mathbf{v}) - \bar{\mathbf{x}}(\mathbf{u}) + \mathbf{A}_\lambda(\mathbf{v}, \mathbf{u}) \mathbf{e}_1) \chi_{U_1(\mathbf{u})}(\mathbf{s}),$$

382 we have shown that:

$$383 \quad \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) = \frac{1}{\lambda^3} \mathbf{K}_{1,\lambda}(\mathbf{s}'/\lambda, \mathbf{s}/\lambda), \quad \forall \mathbf{s}, \mathbf{s}' \in \mathcal{S}.$$

384 **3.2.3. Limit of Electrostatics Energy.** In this section, we obtain the limit of the energy per
385 unit length e_λ . The broad strategy is similar to the helical nanotube. We first show that the map T_λ
386 is bounded and obtain its limit. The continuum limit of the energy density e_λ then follows easily.

387 *Limit of Discrete Electric Field.* Let $T_{1,\lambda}: L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3) \rightarrow L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$ be the map with kernel
388 $\mathbf{K}_{1,\lambda}$. For any function $\mathbf{f} \in L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$, we have

$$389 \quad (3.37) \quad (T_{1,\lambda} \mathbf{f})(\mathbf{s}) = \frac{\mathcal{R}}{\theta_l} \int_{\mathcal{S}_{1,\lambda}} \mathbf{K}_{1,\lambda}(\mathbf{s}', \mathbf{s}) \mathbf{f}(\mathbf{s}') \, d\mathbf{s}', \quad \forall \mathbf{s} \in \mathcal{S}_{1,\lambda}.$$

390 Let $\mathbf{H}_\lambda = \mathbf{H}_\lambda(\mathbf{s})$ be the zeroth order moment (with respect to the first argument) of kernel \mathbf{K}_λ
391 given by

$$392 \quad (3.38) \quad \mathbf{H}_\lambda(\mathbf{s}) = \frac{\mathcal{R}\lambda}{\theta_l} \int_{\mathbf{s}' \in \mathcal{S}} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) \, d\mathbf{s}', \quad \forall \mathbf{s} \in \mathcal{S}.$$

393 We now state the limit result of T_λ .

394 **PROPOSITION 3.5.** Suppose $0 < \theta < \pi/4$. The maps $T_{1,\lambda}$ and T_λ are bounded in L^2 for all
395 $\lambda > 0$ and satisfy

$$396 \quad \|T_\lambda\|_{\mathcal{L}(L^2(\mathcal{S}, \mathbb{R}^3), L^2(\mathcal{S}, \mathbb{R}^3))} = \|T_{1,\lambda}\|_{\mathcal{L}(L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3), L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3))}.$$

397 Further, for $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$,

$$398 \quad (T_\lambda \mathbf{f})(\mathbf{s}) \xrightarrow{\lambda \rightarrow 0} \mathbf{H}_0(\mathbf{s}) \mathbf{f}(\mathbf{s}),$$

399 pointwise, where $\mathbf{H}_0(\mathbf{s})$, for $\mathbf{s} \in \mathcal{S}$, is given by

$$400 \quad \mathbf{H}_0(\mathbf{s}) = \lim_{\lambda \rightarrow 0} \mathbf{H}_\lambda(\mathbf{s}) = \mathcal{R} \sum_{\substack{\mathbf{u}=(u_1, u_2) \in \theta_l \mathbb{Z} \times \mathbb{Z}, \\ \mathbf{u} \neq \mathbf{0}}} \mathbf{K}(\mathbf{u}_1 \mathbf{t}_1(\mathbf{s}) + \mathbf{u}_2 \mathbf{t}_2(\mathbf{s})).$$

$$401 \quad \mathbf{t}_i(\mathbf{s}) = \frac{d\bar{\mathbf{x}}(\mathbf{s})}{ds_i}, \quad i = 1, 2, \text{ are tangent vectors on the film.}$$

402 We provide the proof of [Proposition 3.5](#) in [subsection 4.2](#). Based on the proposition above, we
403 state the main result for the thin film.

404 *Limit of Energy.*

405 **THEOREM 3.6.** Let $\mathbf{f}_\lambda \in L^2(\mathcal{S}, \mathbb{R}^3)$ be a sequence of functions for $\lambda > 0$ with $\mathbf{f} \in L^2(\mathcal{S}, \mathbb{R}^3)$
406 such that $\mathbf{f}_\lambda \rightarrow \mathbf{f}$ in $L^2(\mathcal{S}, \mathbb{R}^3)$. Let the system of dipole moments $\mathbf{d}_\lambda: \mathcal{L}_\lambda \rightarrow \mathbb{R}^3$ be given by [\(3.26\)](#).
407 Let e_λ , given by [\(3.29\)](#), be the energy per unit length normalized by $\mathcal{R}\theta_l$. Then

$$408 \quad e_\lambda \xrightarrow{\lambda \rightarrow 0} -\frac{1}{2} (\mathbf{f}, \mathbf{H}_0 \mathbf{f})_{L^2(\mathcal{S}, \mathbb{R}^3)}.$$

409 $\mathbf{H}_0 = \mathbf{H}_0(\mathbf{s})$ is defined in [Proposition 3.5](#).

The proof of [Theorem 3.6](#) follows from the proof of [Theorem 3.3](#) and using [Proposition 3.5](#).

REMARK 3.4. Note that, for $s \in \mathcal{S}$,

$$(3.39) \quad \mathbf{H}_0(s) = \mathbf{Q}(s_1)\mathbf{H}_0(0)\mathbf{Q}(-s_1).$$

Thus, if the limiting dipole moment field \mathbf{f} is uniform in the \mathbf{e}_3 direction, the electric field $\mathbf{H}_0(s)\mathbf{f}(s)$ will be independent of the \mathbf{e}_3 -coordinate. It is easy to see from the expression of \mathbf{H}_0 that both the normal component and the tangential components of the dipole field contribute to the electric field and energy. This is in contrast to [\[GJ97\]](#) where the thin film limit of magnetostatic energy has contribution only from the normal component.

4. Proof of Claims.

4.1. Helical Nanotube. In this section, we prove [Proposition 3.2](#). First, we collect some important results, and then show that T_λ is bounded and extends from $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$ to $L^2(\mathbb{R}, \mathbb{R}^3)$. We then obtain the limit of the map T_λ .

LEMMA 4.1. 1. For any $a, b \in \mathbb{R}$,

$$(4.1) \quad \bar{\mathbf{x}}(b) - \bar{\mathbf{x}}(a) = \mathbf{Q}(a\theta)[(\mathbf{Q}((b-a)\theta) - \mathbf{I})\mathbf{e}_1 + \delta(b-a)\mathbf{e}_3],$$

where $\bar{\mathbf{x}}$ is the map [\(3.1\)](#), \mathbf{Q} is the rotational tensor [\(3.2\)](#), θ and δ define the helix.

2. For any $\theta \in (0, \pi)$,

$$(4.2) \quad \delta \leq \min_{a, b \in \mathcal{L}_1, a \neq b} |\bar{\mathbf{x}}(b) - \bar{\mathbf{x}}(a)|,$$

where \mathcal{L}_1 is $\mathcal{L}_\lambda = \lambda\mathbb{Z}$ for $\lambda = 1$.

3. For any $a, b \in \mathcal{L}_1$ and $\lambda > 0$,

$$(4.3) \quad \delta|a - b| \leq |\bar{\mathbf{x}}(a) - \bar{\mathbf{x}}(b) + \mathbf{A}_\lambda(a, b)\mathbf{e}_1|,$$

where $\mathbf{A}_\lambda(a, b)$ is given by

$$\mathbf{A}_\lambda(a, b) = \frac{\mathbf{Q}(\lambda a\theta) - \mathbf{Q}(\lambda b\theta) - (\lambda\mathbf{Q}(a\theta) - \lambda\mathbf{Q}(b\theta))}{\lambda}.$$

4. For any $s, s' \in \mathbb{R}$ such that $|s - s'| \geq 1$, suppose $a, b \in \mathcal{L}_1$ are such that $s \in [a, a+1)$, $s' \in [b, b+1)$, then

$$(4.4) \quad \frac{|s - s'|}{|a - b|} < 3.$$

Proof. 1. For any $\alpha, \beta \in \mathbb{R}$, we have the identities

$$(4.5) \quad \mathbf{Q}^T(\alpha) = \mathbf{Q}(-\alpha), \quad \mathbf{Q}(\alpha)\mathbf{Q}(\beta) = \mathbf{Q}(\alpha + \beta), \quad \mathbf{Q}(\alpha)\mathbf{e}_3 = \mathbf{e}_3,$$

where the last relation shows that \mathbf{e}_3 is the axis of \mathbf{Q} . By noting the definition of $\bar{\mathbf{x}}$ in [\(3.1\)](#) and using the identities above, [\(4.1\)](#) follows.

2. To show [\(4.2\)](#), we use [\(4.1\)](#) to get

$$|\bar{\mathbf{x}}(b) - \bar{\mathbf{x}}(a)|^2 = |(\mathbf{Q}((b-a)\theta) - \mathbf{I})\mathbf{e}_1|^2 + \delta^2|b - a|^2 \geq \delta^2|b - a|^2 \geq \delta^2,$$

where we used the fact that $|b - a| \geq 1$ for $a, b \in \mathcal{L}_1, a \neq b$.

3. To show (4.3), we substitute the definition of \mathbf{A}_λ to get

$$(4.6) \quad \bar{\mathbf{x}}(a) - \bar{\mathbf{x}}(b) + \mathbf{A}_\lambda(a, b)\mathbf{e}_1 = \frac{\mathbf{Q}(\lambda a\theta) - \mathbf{Q}(\lambda b\theta)}{\lambda} \mathbf{e}_1 + (a - b)\delta \mathbf{e}_3.$$

Since $\mathbf{Q}(\alpha)\mathbf{e}_1$ is orthogonal to \mathbf{e}_3 for any α , we have

$$|\bar{\mathbf{x}}(a) - \bar{\mathbf{x}}(b) + \mathbf{A}_\lambda(a, b)\mathbf{e}_1| \geq \delta|a - b|.$$

4. To show (4.4), we note that for $s, s' \in \mathbb{R}$ such that $|s - s'| \geq 1$ with $a, b \in \mathcal{L}_1$ and $s \in [a, a+1), s' \in [b, b+1)$, we can write $s = a + \Delta s$ and $s' = b + \Delta s'$ with $0 \leq \Delta s, \Delta s' < 1$. Thus

$$\frac{|s - s'|}{|a - b|} = \frac{|a - b + (\Delta s - \Delta s')|}{|a - b|} \leq \frac{|a - b| + |\Delta s - \Delta s'|}{|a - b|} < 1 + \frac{2}{|a - b|} \leq 3,$$

where in the last step we used the fact that $|a - b| \geq 1$ for $a, b \in \mathcal{L}_1, a \neq b$ (which is ensured when $|s - s'| \geq 1$).

This completes the proof. \square

4.1.1. Boundedness. We next show that T_λ is a bounded map. Let $S_\lambda: L^2(\mathbb{R}, \mathbb{R}^3) \rightarrow L^2(\mathbb{R}, \mathbb{R}^3)$ be an isometry defined as

$$(S_\lambda \mathbf{f})(s) := \lambda^{1/2} \mathbf{f}(\lambda s).$$

It is easy to see that $\|S_\lambda \mathbf{f}\|_{L^2} = \|\mathbf{f}\|_{L^2}$. The inverse of S_λ is given by

$$(S_\lambda^{-1} \mathbf{f})(s) = \lambda^{-1/2} \mathbf{f}(s/\lambda).$$

Using S_λ , we can show – noting the definition of T_λ in (3.11) – for $\mathbf{f} \in L^2(\mathbb{R}, \mathbb{R}^3)$,

$$\begin{aligned} (T_\lambda \mathbf{f})(s) &= \lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) \mathbf{f}(s') \, ds' = \lambda^2 \int_{\mathbb{R}} \frac{1}{\lambda^3} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \left(\lambda^{-1/2} (S_\lambda \mathbf{f})(s'/\lambda) \right) \, ds' \\ &= \lambda^{-3/2} \int_{\mathbb{R}} \mathbf{K}_{1,\lambda}(s', s/\lambda) (S_\lambda \mathbf{f})(s') \lambda \, ds' = \lambda^{-1/2} (T_{1,\lambda} (S_\lambda \mathbf{f}))(s/\lambda) \\ &= (S_\lambda^{-1} T_{1,\lambda} S_\lambda \mathbf{f})(s), \end{aligned}$$

where we used a change of variable and the fact that $(S_\lambda^{-1} \mathbf{f})(s) = \lambda^{(-1/2)} \mathbf{f}(s/\lambda)$. It follows from the above equation that

$$\begin{aligned} \|T_\lambda\|_{\mathcal{L}(L^2, L^2)} &= \sup_{\|\mathbf{f}\| \neq 0} \frac{\|T_\lambda \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}}{\|\mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}} = \sup_{\|\mathbf{f}\| \neq 0} \frac{\|S_\lambda^{-1} (T_{1,\lambda} S_\lambda \mathbf{f})\|_{L^2(\mathbb{R}, \mathbb{R}^3)}}{\|\mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}} \\ &= \sup_{\|\mathbf{f}\| \neq 0} \frac{\|T_{1,\lambda} (S_\lambda \mathbf{f})\|_{L^2(\mathbb{R}, \mathbb{R}^3)}}{\|\mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}} = \sup_{\|S_\lambda \mathbf{f}\| \neq 0} \frac{\|T_{1,\lambda} (S_\lambda \mathbf{f})\|_{L^2(\mathbb{R}, \mathbb{R}^3)}}{\|S_\lambda \mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}} \\ &= \|T_{1,\lambda}\|_{\mathcal{L}(L^2, L^2)}. \end{aligned}$$

This completes the proof of (3.16) in Proposition 3.2. Next, we show that $T_{1,\lambda}$ is a bounded map to prove the boundedness of T_λ . We first analyze the discrete dipole field kernel $\mathbf{K}_{1,\lambda}$, which is defined as

$$(4.7) \quad \mathbf{K}_{1,\lambda}(s', s) = \sum_{\substack{u, v \in \mathcal{L}_1, \\ u \neq v}} \chi_{U_1(v)}(s') \mathbf{K}(\bar{\mathbf{x}}(v) - \bar{\mathbf{x}}(u) + \mathbf{A}_\lambda(v, u)\mathbf{e}_1) \chi_{U_1(u)}(s),$$

where $U_\lambda(s) = [s, s + \lambda)$ for $s \in \mathcal{L}_\lambda$, and $\mathbf{A}_\lambda(a, b)$ is given by (3.13).

Consider some typical $s, s' \in \mathbb{R}$ and the corresponding $a, b \in \mathcal{L}_1$ such that $s \in [a, a + 1)$, $s' \in [b, b + 1)$. From (4.7), we have, for all $s, s' \in \mathbb{R}$ such that $|s - s'| < 1$,

- If $a = b$, then $\mathbf{K}_{1,\lambda}(s, s') = \mathbf{0}$.
- If $a \neq b$, then from (4.2), we have

$$|\mathbf{K}_{1,\lambda}(s, s')| \leq \sqrt{6}/(4\pi\delta^3)$$

using $|\mathbf{A}a| \leq |\mathbf{A}| |a|$ and $|\mathbf{I} - 3(\mathbf{x}/|\mathbf{x}|) \otimes (\mathbf{x}/|\mathbf{x}|)| \leq \sqrt{6}$, $\forall \mathbf{x} \neq \mathbf{0}$.

Combining the two cases above, $|\mathbf{K}_{1,\lambda}(s, s')| \leq \sqrt{6}/(4\pi\delta^3)$.

We now consider a case when $|s - s'| \geq 1$. Noting that for this case, $a \neq b$. We proceed as follows

$$\begin{aligned} |\mathbf{K}_{1,\lambda}(s, s')| &\leq \frac{\sqrt{6}}{4\pi|s - s'|^3} \frac{|s - s'|^3}{|a - b|^3} \frac{|a - b|^3}{|\bar{\mathbf{x}}(a) - \bar{\mathbf{x}}(b) + \mathbf{A}_\lambda(a, b)\mathbf{e}_1|^3} \\ &\leq \frac{\sqrt{6}}{4\pi|s - s'|^3} 3^3 \frac{|a - b|^3}{\delta^3|a - b|^3} = \frac{3^3\sqrt{6}}{4\pi\delta^3} \frac{1}{|s - s'|^3} \end{aligned}$$

where we used the bounds (4.3) and (4.4). Combining the above bound for $|s - s'| \geq 1$ with the bound for $|s - s'| < 1$, and renaming the constants, we can write

$$(4.8) \quad |\mathbf{K}_{1,\lambda}(s, s')| \leq \frac{C_1}{C_2 + |s - s'|^3}.$$

Since the kernel $\mathbf{K}_{1,\lambda}$ satisfies (4.8), we have

$$(4.9) \quad \int_{\mathbb{R}} |\mathbf{K}_{1,\lambda}(s', s)| \, ds' \leq C_3, \quad \int_{\mathbb{R}} |\mathbf{K}_{1,\lambda}(s', s)| \, ds \leq C_3,$$

for some fixed $C_3 < \infty$ independent of λ . Using Young's inequality (e.g., Theorem 0.3.1 [Sog17]), we have

$$(4.10) \quad \|T_{1,\lambda}\mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)} \leq C_3 \|\mathbf{f}\|_{L^2(\mathbb{R}, \mathbb{R}^3)}$$

showing that $T_{1,\lambda}$ is a bounded linear map for all $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$. Since $C_0^\infty(\mathbb{R}, \mathbb{R}^3)$ is dense in $L^2(\mathbb{R}, \mathbb{R}^3)$, it follows that T_λ is also bounded in $L^2(\mathbb{R}, \mathbb{R}^3)$, and extends as a bounded linear map from $C_0^\infty(\mathbb{R}, \mathbb{R}^3)$ to $L^2(\mathbb{R}, \mathbb{R}^3)$.

4.1.2. Limit of the Map T_λ . Let $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$. Write $T_\lambda\mathbf{f}$ as:

$$\begin{aligned} T_\lambda(\mathbf{f})(s) &= \lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) \mathbf{f}(s') \, ds' \\ &= \underbrace{\left[\lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) \, ds' \right]}_{=: \mathbf{H}_\lambda(s)} \mathbf{f}(s) + \lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) (\mathbf{f}(s') - \mathbf{f}(s)) \, ds', \end{aligned} \quad (4.11)$$

The second term above is zero in the limit $\lambda \rightarrow 0$. To see this, consider any $R > 0$, and then using the bound on $\mathbf{K}_{1,\lambda}$ in (4.8), we have

$$\begin{aligned} \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \, ds' &\leq \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \frac{C_1}{C_2 + |s/\lambda - s'/\lambda|^3} \, ds' \\ &= \lambda^2 \int_{|s-s'| \geq R\lambda} \frac{C_1}{C_2\lambda^3 + |s - s'|^3} \, ds' = \lambda^2 \int_{|t| \geq R} \frac{C_1}{C_2\lambda^3 + |\lambda t|^3} \lambda \, dt \\ &= \int_{|t| \geq R} \frac{C_1}{C_2 + |t|^3} \, dt, \end{aligned} \quad (4.12)$$

In the intermediate step, we changed variables $t = (s' - s)/\lambda$. Thus, the upper bound on

$$\frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \, ds'$$

is independent of λ , and goes to zero as $R \rightarrow \infty$. For the second term in (4.11), using that

$$\mathbf{K}_\lambda(s', s) = \frac{1}{\lambda^3} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda), \text{ we get}$$

$$\begin{aligned} & \left| \lambda^2 \int_{\mathbb{R}} \mathbf{K}_\lambda(s', s) (\mathbf{f}(s') - \mathbf{f}(s)) \, ds' \right| \\ &= \left| \frac{1}{\lambda} \int_{\mathbb{R}} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) (\mathbf{f}(s') - \mathbf{f}(s)) \, ds' \right| \\ &= \left| \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \mathbf{f}(s') \, ds' - \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \, ds' \mathbf{f}(s) \right. \\ &\quad \left. + \frac{1}{\lambda} \int_{|s-s'| \leq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) (\mathbf{f}(s') - \mathbf{f}(s)) \, ds' \right| \\ &= \left| \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \mathbf{f}(s') \, ds' \right| + \left| \frac{1}{\lambda} \int_{|s-s'| \geq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) \, ds' \mathbf{f}(s) \right| \\ &\quad + \left| \frac{1}{\lambda} \int_{|s-s'| \leq R\lambda} \mathbf{K}_{1,\lambda}(s'/\lambda, s/\lambda) (\mathbf{f}(s') - \mathbf{f}(s)) \, ds' \right|. \end{aligned}$$

We let $\lambda \rightarrow 0$ and then $R \rightarrow \infty$. In the limit $\lambda \rightarrow 0$, the third term above will go to zero, as $\mathbf{f} \in C_0^\infty(\mathbb{R}, \mathbb{R}^3)$ and $\mathbf{K}_{1,\lambda}$ is bounded. The first and second terms will go to zero using (4.12). Thus, we have from (4.11) that

$$(4.13) \quad \lim_{\lambda \rightarrow 0} T_\lambda(\mathbf{f})(s) = \left[\lim_{\lambda \rightarrow 0} \mathbf{H}_\lambda(s) \right] \mathbf{f}(s),$$

We next compute the limit of $\mathbf{H}_\lambda(s)$. Fix $s \in \mathbb{R}$ and suppose $a \in \mathcal{L}_\lambda$ such that $s \in U_\lambda(a)$. Using the definition of $\mathbf{K}_\lambda(s', s)$, we have

$$\begin{aligned} \mathbf{H}_\lambda(s) &= \lambda^2 \int_{\mathbb{R}} \mathbf{K}(s', s) \, ds' = \lambda^2 \sum_{\substack{u \in \mathcal{L}_1, \\ u \neq a}} \mathbf{K}(\bar{\mathbf{x}}(u) - \bar{\mathbf{x}}(a)) \int_{U_\lambda(u)} dt \\ &= \lambda^3 \sum_{\substack{u \in \lambda\mathbb{Z}, \\ u \neq a}} \mathbf{K}(\bar{\mathbf{x}}(u) - \bar{\mathbf{x}}(a)). \end{aligned}$$

From (4.5), we have $\bar{\mathbf{x}}(u) - \bar{\mathbf{x}}(a) = \mathbf{Q}(a\theta)((\mathbf{Q}((u-a)\theta) - \mathbf{I})\mathbf{e}_1 + (u-a)\delta\mathbf{e}_3)$. Using the identity $\mathbf{K}(\mathbf{Q}\mathbf{x}) = \mathbf{Q}\mathbf{K}(\mathbf{x})\mathbf{Q}^T$ and $\mathbf{K}(\lambda\mathbf{x}) = \mathbf{K}(\mathbf{x})/\lambda^3$, we get

$$\begin{aligned} \mathbf{H}_\lambda(s) &= \mathbf{Q}(a\theta) \left[\sum_{u \in \lambda\mathbb{Z}, u \neq a} \mathbf{K}((\mathbf{Q}((u-a)\theta) - \mathbf{I})/\lambda\mathbf{e}_1 + (u-a)\delta/\lambda\mathbf{e}_3) \right] \mathbf{Q}(-a\theta) \\ &= \mathbf{Q}(a\theta) \left[\sum_{i \in \mathbb{Z}, i \neq 0} \mathbf{K}((\mathbf{Q}(i\lambda\theta) - \mathbf{I})/\lambda\mathbf{e}_1 + i\delta\mathbf{e}_3) \right] \mathbf{Q}(-a\theta), \end{aligned}$$

528 where we changed variables $i = (u - a)/\lambda$. Note that $a \in \lambda\mathbb{Z}$, and, therefore, $(u - a) \in \lambda\mathbb{Z}$ for
 529 $u \in \lambda\mathbb{Z}$, which implies $i \in \mathbb{Z}$. Since s is related to a by $s \in U_\lambda(a)$, we have $a \rightarrow s$ in the limit
 530 $\lambda \rightarrow 0$. Therefore, we get

$$531 \quad \mathbf{H}_0(s) := \lim_{\lambda \rightarrow 0} \mathbf{H}_\lambda(s) = \mathbf{Q}(s\theta) \left[\lim_{\lambda \rightarrow 0} \sum_{i \in \mathbb{Z} - \{0\}} \mathbf{K}((\mathbf{Q}(i\lambda\theta) - \mathbf{I})/\lambda\mathbf{e}_1 + i\delta\mathbf{e}_3) \right] \mathbf{Q}(-s\theta).$$

532 To take the limit inside the summation, we show that the sum is absolutely convergent for all $\lambda > 0$
 533 as follows:

$$534 \quad a_\lambda := \sum_{i \in \mathbb{Z} - \{0\}} |\mathbf{K}((\mathbf{Q}(i\lambda\theta) - \mathbf{I})/\lambda\mathbf{e}_1 + i\delta\mathbf{e}_3)| \leq \sum_{i \in \mathbb{Z} - \{0\}} \frac{c}{|(\mathbf{Q}(i\lambda\theta) - \mathbf{I})/\lambda\mathbf{e}_1 + i\delta\mathbf{e}_3|^3}$$

$$535 \quad = \sum_{i \in \mathbb{Z} - \{0\}} \frac{c}{(4\sin^2(i\lambda\theta/2)/\lambda^2 + i^2\delta^2)^{3/2}} \leq \sum_{i \in \mathbb{Z} - \{0\}} \frac{c}{|i|^3} < \infty, \quad \forall \lambda > 0.$$

536
 537 Now, we can write

$$538 \quad \mathbf{H}_0(s) = \mathbf{Q}(s\theta) \left[\sum_{i \in \mathbb{Z} - \{0\}} \lim_{\lambda \rightarrow 0} \mathbf{K} \left(\frac{\mathbf{Q}(i\lambda\theta) - \mathbf{I}}{i\lambda\theta} (i\theta\mathbf{e}_1) + i\delta\mathbf{e}_3 \right) \right] \mathbf{Q}(-s\theta).$$

539 Note that for a fixed $i \in \mathbb{Z}$

$$540 \quad \lim_{\lambda \rightarrow 0} \frac{(\mathbf{Q}(i\lambda\theta) - \mathbf{I})}{i\lambda\theta} i\theta\mathbf{e}_1 + i\delta\mathbf{e}_3 = \lim_{h=i\lambda\theta \rightarrow 0} \frac{(\mathbf{Q}(h) - \mathbf{I})}{h} i\theta\mathbf{e}_1 + i\delta\mathbf{e}_3 = i\theta\mathbf{Q}'(0)\mathbf{e}_1 + i\delta\mathbf{e}_3,$$

541 where $\mathbf{Q}'(0) = \text{d}/\text{d}x \mathbf{Q}(x)|_{x=0}$. Now, using the equation above, and the fact that $\mathbf{K}(x)$ is smooth
 542 away from $x = \mathbf{0}$ (which is ensured in the summation), we get

$$543 \quad \mathbf{H}_0(s) = \mathbf{Q}(s\theta) \left[\sum_{i \in \mathbb{Z} - \{0\}} \mathbf{K}(i\theta\mathbf{Q}'(0)\mathbf{e}_1 + i\delta\mathbf{e}_3) \right] \mathbf{Q}(-s\theta) = \mathbf{Q}(s\theta) \mathbf{H}_0(0) \mathbf{Q}(-s\theta).$$

544 Combining this with (4.13), we get

$$545 \quad \lim_{\lambda \rightarrow 0} T_\lambda(\mathbf{f})(s) = \mathbf{H}_0(s) \mathbf{f}(s) = \mathbf{Q}(s\theta) \mathbf{H}_0(0) \mathbf{Q}(-s\theta) \mathbf{f}(s).$$

546 Next, we simplify $\mathbf{H}_0(s)$. Using $\mathbf{Q}\mathbf{K}(x)\mathbf{Q}^T = \mathbf{K}(\mathbf{Q}x)$ and $\mathbf{Q}(s\theta)\mathbf{Q}'(0) = \mathbf{Q}'(s\theta)$, we can show

$$547 \quad \mathbf{H}_0(s) = \sum_{i \in \mathbb{Z} - 0} \mathbf{K}(i\theta\mathbf{Q}'(s\theta)\mathbf{e}_1 + i\delta\mathbf{e}_3) = \sum_{i \in \mathbb{Z} - 0} \mathbf{K}(i|\mathbf{t}(s)|\hat{\mathbf{t}}(s)),$$

548 where $\mathbf{t}(s) = \theta\mathbf{Q}'(s\theta)\mathbf{e}_1 + \delta\mathbf{e}_3$ is the tangent vector, and $\hat{\mathbf{t}}(s) = \mathbf{t}(s)/|\mathbf{t}(s)|$ with $|\mathbf{t}(s)| = \sqrt{\theta^2 + \delta^2}$.
 549 Using above and substituting the form of dipole field kernel \mathbf{K} , it is easy to show that

$$550 \quad \mathbf{H}_0(s) = -h_0 [\mathbf{I} - 3\hat{\mathbf{t}}(s) \otimes \hat{\mathbf{t}}(s)] = -h_0 [\mathbf{P}_\perp \mathbf{f}(s) - 2\mathbf{P}_\parallel(s)],$$

551 with h_0 defined as

$$552 \quad (4.14) \quad h_0 = \sum_{i \in \mathbb{Z} - 0} \frac{1}{4\pi|i|^3(\theta^2 + \delta^2)^{3/2}}$$

553 and projection tensors $\mathbf{P}_\parallel(s) = \hat{\mathbf{t}}(s) \otimes \hat{\mathbf{t}}(s)$ and $\mathbf{P}_\perp(s) = \mathbf{I} - \mathbf{P}_\parallel(s)$.

4.2. Nanofilm with Uniform Bending. In this section, we prove [Proposition 3.5](#). The outline of the proof is similar to the case of the helix in [subsection 4.1](#).

LEMMA 4.2. *1. Suppose $s, s' \in \mathcal{S}_{1,\lambda} = (-\bar{\theta}/\lambda, \bar{\theta}/\lambda) \times \mathbb{R}$ such that $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda} = (-\bar{\theta}/\lambda, \bar{\theta}/\lambda) \cap \theta_l \mathbb{Z} \times \mathbb{Z}$ with $s \in U_1(\mathbf{a}) = [a_1, a_1 + \theta_l) \times [a_2, a_2 + 1)$, $s' \in U_1(\mathbf{b})$. When $|s - s'| \geq \min\{\theta_l, 1\}$, we have $\mathbf{a} \neq \mathbf{b}$ and*

$$(4.15) \quad \frac{|s - s'|}{|\mathbf{a} - \mathbf{b}|} < 1 + \frac{\theta_l + 1}{\min\{\theta_l, 1\}} =: c_L.$$

2. For any $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda}$, we have

$$(4.16) \quad c_A |\mathbf{a} - \mathbf{b}| \leq |\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|,$$

where $\bar{\mathbf{x}}$ is given by [\(3.19\)](#) and $\mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})$ is defined as

$$\mathbf{A}_\lambda(\mathbf{a}, \mathbf{b}) = \frac{\mathcal{R}}{\lambda} [\mathbf{Q}(\lambda a_1) - \mathbf{Q}(\lambda b_1) - \lambda \mathbf{Q}(a_1) + \lambda \mathbf{Q}(b_1)].$$

Here $c_A = \min\{\delta, \mathcal{R}\sqrt{1 - \bar{\theta}^2/3}\}$ is the constant independent of λ . Note that $c_A > 0$ for $0 < \bar{\theta} < \pi/2$.

Proof. To show [\(4.15\)](#), we proceed as follows. For $s, s' \in \mathcal{S}_{1,\lambda}$ and corresponding $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda}$, there exists $\Delta s, \Delta s'$ such that $s = \mathbf{a} + \Delta s, s' = \mathbf{b} + \Delta s'$ with $0 \leq \Delta s_1, \Delta s'_1 < \theta_l, 0 \leq \Delta s_2, \Delta s'_2 < 1$. We have the bound

$$(4.17) \quad \frac{|s - s'|}{|\mathbf{a} - \mathbf{b}|} \leq 1 + \frac{|\Delta s_1 - \Delta s'_1| + |\Delta s_2 - \Delta s'_2|}{|\mathbf{a} - \mathbf{b}|} < 1 + \frac{\theta_l + 1}{|\mathbf{a} - \mathbf{b}|} \leq 1 + \frac{\theta_l + 1}{\min\{\theta_l, 1\}},$$

where in the last step we used the fact that any $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda}$, satisfying $\mathbf{a} \neq \mathbf{b}$, are at least $\min\{\theta_l, 1\}$ distance apart.

We next show [\(4.16\)](#). Using

$$\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1 = \frac{\mathcal{R}}{\lambda} (\mathbf{Q}(\lambda a_1) - \mathbf{Q}(\lambda b_1))\mathbf{e}_1 + \delta(a_2 - b_2)\mathbf{e}_3$$

and

$$|(\mathbf{Q}(\theta_1) - \mathbf{Q}(\theta_2))\mathbf{e}_1|^2 = (\cos \theta_1 - \cos \theta_2)^2 + (\sin \theta_1 - \sin \theta_2)^2 = 2(1 - \cos(\theta_1 - \theta_2)),$$

we have that

$$(4.18) \quad |\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|^2 = \delta^2 |a_2 - b_2|^2 + \frac{2\mathcal{R}^2}{\lambda^2} (1 - \cos(\lambda a_1 - \lambda b_1)).$$

Let $r = a_1 - b_1$. Then, using a Taylor expansion and the mean value theorem, there exists ξ such that

$$1 - \cos(\lambda r) = \frac{1}{2} \lambda^2 r^2 - \frac{1}{24} \lambda^4 r^4 \cos(\xi).$$

Since $-1 \leq \cos(\xi) \leq 1$, it follows

$$1 - \cos(\lambda r \xi) \geq \frac{1}{2} \lambda^2 r^2 - \frac{1}{24} \lambda^4 r^4.$$

Substituting the relation above in [\(4.18\)](#), we get

$$|\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|^2 \geq \delta^2 |a_2 - b_2|^2 + \mathcal{R}^2 r^2 \left(1 - \frac{1}{12} \lambda^2 r^2\right).$$

584 Since $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda}$, we have $-2\bar{\theta} < \lambda r < 2\bar{\theta}$, and

$$585 \quad 1 - \frac{1}{12}\lambda^2 r^2 \geq 1 - \frac{1}{12}\bar{\theta}^2 4 = 1 - \frac{\bar{\theta}^2}{3}.$$

586 Using the two equations above and defining the constant c_A as in [Lemma 4.2\(2\)](#), [\(4.16\)](#) can be easily
587 shown. \square

588 **4.2.1. Boundedness.** Let $S_\lambda: L^2(\mathcal{S}, \mathbb{R}^3) \rightarrow L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$ be a map such that, for any $\mathbf{f} \in$
589 $L^2(\mathcal{S}, \mathbb{R}^3)$,

$$590 \quad (S_\lambda \mathbf{f})(\mathbf{s}) = \lambda \mathbf{f}(\lambda \mathbf{s}), \quad \forall \mathbf{s} \in \mathcal{S}_{1,\lambda}.$$

591 It is easy to see that S_λ is an isometry. The inverse of S_λ is given by

$$592 \quad (S_\lambda^{-1} \mathbf{f})(\mathbf{s}) = \lambda^{-1} \mathbf{f}(\mathbf{s}/\lambda), \quad \forall \mathbf{s} \in \mathcal{S}.$$

593 Following the similar steps in [subsubsection 4.1.1](#), we can show that

$$594 \quad \|T_\lambda\|_{\mathcal{L}(L^2, L^2)} = \|T_{1,\lambda}\|_{\mathcal{L}(L^2, L^2)}.$$

595 Thus, to show that T_λ is a bounded map, it is sufficient to show that $T_{1,\lambda}$ is bounded. Towards that
596 goal, we first establish that

$$597 \quad (4.19) \quad |\mathbf{K}_{1,\lambda}(\mathbf{s}, \mathbf{s}')| \leq \frac{C_1}{C_2 + |\mathbf{s} - \mathbf{s}'|^3}, \quad \forall \mathbf{s}', \mathbf{s} \in \mathcal{S}_{1,\lambda},$$

598 where C_1, C_2 are constants that may depend on the parameters R, θ, δ defining the surface \mathcal{S} and are
599 independent of λ . Similar to the case of the helix ([subsubsection 4.1.1](#)), we apply Theorem 0.3.1
600 of [\[Sog17\]](#) to show that, for any $\lambda > 0$, $T_{1,\lambda}$ is a bounded linear map on $C_0^\infty(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$. Using the
601 density of $C_0^\infty(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$ in $L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$, $T_{1,\lambda}$ extends as a bounded linear map to $L^2(\mathcal{S}_{1,\lambda}, \mathbb{R}^3)$.

602 It remains to show [\(4.19\)](#). We recall that $\bar{\theta}$ is the fixed angular extent of the film and satisfies the
603 bound $0 < \bar{\theta} < \pi/2$ (in fact we restrict it such that $0 < \bar{\theta} < \pi/4$). Let $\mathbf{s}, \mathbf{s}' \in \mathcal{S}_{1,\lambda}$ be any two generic
604 points, and let $\mathbf{a}, \mathbf{b} \in \mathcal{L}_{1,\lambda}$ be such that $\mathbf{s} \in U_1(\mathbf{a})$ and $\mathbf{s}' \in U_1(\mathbf{b})$. We refer to [subsubsection 3.2.1](#)
605 and [subsubsection 3.2.2](#) for the notation appearing in this section.

606 First, consider \mathbf{s}, \mathbf{s}' such that $|\mathbf{s} - \mathbf{s}'| \geq \min\{\theta_l, 1\}$. For this case, we have $\mathbf{a} \neq \mathbf{b}$. Noting that
607 $|\mathbf{I} - 3(\mathbf{x}/|\mathbf{x}|) \otimes (\mathbf{x}/|\mathbf{x}|)| = \sqrt{6}, \forall \mathbf{x} \neq \mathbf{0}$, we have

$$\begin{aligned} 608 \quad |\mathbf{K}_{1,\lambda}(\mathbf{s}, \mathbf{s}')| &\leq \frac{\sqrt{6}}{4\pi|\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|^3} \\ 609 \quad &= \frac{\sqrt{6}}{4\pi|\mathbf{s} - \mathbf{s}'|^3} \frac{|\mathbf{s} - \mathbf{s}'|^3}{|\mathbf{a} - \mathbf{b}|^3} \frac{|\mathbf{a} - \mathbf{b}|^3}{|\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|^3} \\ 610 \quad (4.20) \quad &\leq \frac{\sqrt{6}}{4\pi|\mathbf{s} - \mathbf{s}'|^3} c_L^3 \frac{1}{c_A^3}, \\ 611 \end{aligned}$$

612 where we used the bounds [\(4.15\)](#) and [\(4.16\)](#).

613 Next, we consider the case when $|\mathbf{s} - \mathbf{s}'| < \min\{\theta_l, 1\}$. This can be further divided in two cases:

- 614 • *Case 1:* $\mathbf{a} = \mathbf{b}$ which implies $|\mathbf{K}_{1,\lambda}(\mathbf{s}', \mathbf{s})| = 0$.
- 615 • *Case 2:* $\mathbf{a} \neq \mathbf{b}$. For this case, we have

$$616 \quad (4.21) \quad |\mathbf{K}_{1,\lambda}(\mathbf{s}, \mathbf{s}')| \leq \frac{\sqrt{6}}{4\pi|\bar{\mathbf{x}}(\mathbf{a}) - \bar{\mathbf{x}}(\mathbf{b}) + \mathbf{A}_\lambda(\mathbf{a}, \mathbf{b})\mathbf{e}_1|^3} \leq \frac{\sqrt{6}}{4\pi c_A^3 |\mathbf{a} - \mathbf{b}|^3}.$$

20

Note that when $\mathbf{a} \neq \mathbf{b}$, we can have (A) $a_1 = b_1, a_2 = b_2 \pm 1$, or (B) $a_1 = b_1 \pm \theta_l, a_2 = b_2$, or (C) $a_1 = b_1 \pm \theta_l, a_2 = b_2 \pm 1$. For all the cases mentioned above, the denominator in (4.21) is bounded from below as follows $|\mathbf{a} - \mathbf{b}| \geq \min\{\theta_l, 1\}$. Thus, we have

$$(4.22) \quad |\mathbf{K}_{1,\lambda}(\mathbf{s}, \mathbf{s}')| \leq \frac{\sqrt{6}}{4\pi c_A^3 (\min\{\theta_l, 1\})^3}.$$

In summary (4.22) holds for any \mathbf{s}, \mathbf{s}' such that $|\mathbf{s} - \mathbf{s}'| < \min\{\theta_l, 1\}$.

Combining the bound for the case $|\mathbf{s} - \mathbf{s}'| < \min\{\theta_l, 1\}$ with the bound for the case $|\mathbf{s} - \mathbf{s}'| \geq \min\{\theta_l, 1\}$, we can write

$$(4.23) \quad |\mathbf{K}_{1,\lambda}(\mathbf{s}, \mathbf{s}')| \leq \frac{C_1}{C_2 + |\mathbf{s} - \mathbf{s}'|^3},$$

where we have renamed the constants for convenience. This completes the proof of the boundedness of $T_{1,\lambda}$. We next obtain the limit of the map T_λ .

4.2.2. Limit of Map T_λ . Let $\mathbf{f} \in C_0^\infty(\mathcal{S}, \mathbb{R}^3)$. We write $T_\lambda \mathbf{f}$ as follows

$$(4.24) \quad \begin{aligned} (T_\lambda \mathbf{f})(\mathbf{s}) &= \frac{\mathcal{R}\lambda}{\theta_l} \int_{\mathcal{S}} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) \mathbf{f}(\mathbf{s}') \, d\mathbf{s}' \\ &= \underbrace{\left[\frac{\mathcal{R}\lambda}{\theta_l} \int_{\mathcal{S}} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) \, d\mathbf{s}' \right]}_{=: \mathbf{H}_\lambda(\mathbf{s})} \mathbf{f}(\mathbf{s}) + \frac{\mathcal{R}\lambda}{\theta_l} \int_{\mathcal{S}} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}', \end{aligned}$$

We next show that the second term in (4.24) is zero in the limit $\lambda \rightarrow 0$. Fix $\bar{R} > 0$, then using the relation $\mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) = \mathbf{K}_{1,\lambda}(\mathbf{s}'/\lambda, \mathbf{s}/\lambda)/\lambda^3$ and the bound on $\mathbf{K}_{1,\lambda}$ in (4.23), we have

$$(4.25) \quad \begin{aligned} \left| \lambda \int_{\mathcal{S}} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| &\leq \lambda \int_{\mathbb{R}^2} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') |\mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s}))| \, d\mathbf{s}' \\ &\leq \frac{1}{\lambda^2} \int_{\mathbb{R}^2} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') \frac{C_1}{C_2 + |\mathbf{s}/\lambda - \mathbf{s}'/\lambda|^3} \, d\mathbf{s}'. \end{aligned}$$

Using the change of variable $\mathbf{t}' = \mathbf{s}'/\lambda$ (so $d\mathbf{s}' = \lambda^2 d\mathbf{t}'$) and $\mathbf{t} = \mathbf{s}/\lambda$ to have

$$(4.26) \quad \left| \lambda \int_{\mathcal{S}} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| \leq \int_{\mathbb{R}^2} \chi_{|\mathbf{t}'-\mathbf{t}| \geq \bar{R}}(\mathbf{t}') \frac{C_1}{C_2 + |\mathbf{t}'-\mathbf{t}|^3} \, d\mathbf{t}'.$$

From above, we have that

$$(4.27) \quad \lim_{\bar{R} \rightarrow \infty} \left[\lim_{\lambda \rightarrow 0} \left| \lambda \int_{\mathcal{S}} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| \right] = 0.$$

We bound the second term in (4.24) by splitting it into two parts as follows:

$$\begin{aligned} \left| \lambda \int_{\mathcal{S}} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| &\leq \left| \lambda \int_{\mathcal{S}} \chi_{|\mathbf{s}-\mathbf{s}'| \geq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| \\ &\quad + \left| \lambda \int_{\mathcal{S}} \chi_{|\mathbf{s}-\mathbf{s}'| \leq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right|. \end{aligned}$$

644 Since $|\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})| \leq 2\|\mathbf{f}\|_{L^\infty} < \infty$ and from (4.27), the first term in the equation above is zero
 645 in the limit $\bar{R} \rightarrow \infty$ following the limit $\lambda \rightarrow 0$. For the second term, we proceed as follows

$$\begin{aligned}
 646 \quad & \left| \lambda \int_S \chi_{|\mathbf{s}-\mathbf{s}'| \leq \bar{R}\lambda}(\mathbf{s}') \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| \\
 647 \quad & \leq \frac{1}{\lambda^2} \int_S \chi_{|\mathbf{s}-\mathbf{s}'| \leq \bar{R}\lambda}(\mathbf{s}') |\mathbf{K}_{1,\lambda}(\mathbf{s}'/\lambda, \mathbf{s}/\lambda)| |\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})| \, d\mathbf{s}' \\
 648 \quad (4.28) \quad & \leq \frac{1}{\lambda^2} \int_{\mathbb{R}^2} \chi_{|\mathbf{s}-\mathbf{s}'| \leq \bar{R}\lambda}(\mathbf{s}') |\mathbf{K}_{1,\lambda}(\mathbf{s}'/\lambda, \mathbf{s}/\lambda)| |\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})| \, d\mathbf{s}'.
 \end{aligned}$$

650 Since $\mathbf{K}_{1,\lambda}$ is bounded as $\mathbf{s}' \rightarrow \mathbf{s}$ (see (4.23)), $|\{\mathbf{s}' \in \mathbb{R}^2; |\mathbf{s} - \mathbf{s}'| \leq \bar{R}\lambda\}| = \pi \bar{R}^2 \lambda^2$, and
 651 $\sup_{|\mathbf{s}' - \mathbf{s}| \leq \bar{R}\lambda} |\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})| \rightarrow 0$ as $\lambda \rightarrow 0$, we have

$$652 \quad (4.29) \quad \lim_{\bar{R} \rightarrow \infty} \left[\lim_{\lambda \rightarrow 0} \left| \lambda \int_S \chi_{|\mathbf{s}-\mathbf{s}'| \leq \bar{R}\lambda} \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| \right] = 0.$$

653 We have shown

$$654 \quad (4.30) \quad \lim_{\lambda \rightarrow 0} \left| \lambda \int_S \mathbf{K}_\lambda(\mathbf{s}', \mathbf{s}) (\mathbf{f}(\mathbf{s}') - \mathbf{f}(\mathbf{s})) \, d\mathbf{s}' \right| = 0.$$

655 Thus, from (4.24), we have

$$656 \quad (4.31) \quad \lim_{\lambda \rightarrow 0} (T_\lambda \mathbf{f})(\mathbf{s}) = \left[\lim_{\lambda \rightarrow 0} \mathbf{H}_\lambda(\mathbf{s}) \right] \mathbf{f}(\mathbf{s}).$$

657 *Limit of \mathbf{H}_λ .* Consider a typical $\mathbf{s} \in \mathcal{S}$ such that $\mathbf{s} \in U_\lambda(\mathbf{a})$ where $\mathbf{a} \in \mathcal{L}_\lambda$. Recall that \mathcal{L}_λ is
 658 the lattice for $\lambda > 0$ and $U_\lambda(\mathbf{a}) = [a_1, a_1 + \theta_1 \lambda) \times [a_2, a_2 + \lambda)$ is the lattice cell. In the definition
 659 of \mathbf{H}_λ , we substitute \mathbf{K}_λ , to get

$$660 \quad (4.32) \quad \mathbf{H}_\lambda(\mathbf{s}) = \frac{\mathcal{R}\lambda}{\theta_1} \sum_{\mathbf{u} \in \mathcal{L}_\lambda, \mathbf{u} \neq \mathbf{a}} \mathbf{K}(\bar{\mathbf{x}}(\mathbf{u}) - \bar{\mathbf{x}}(\mathbf{a})) \int_{U_\lambda(\mathbf{u})} d\mathbf{s}' = \mathcal{R}\lambda^3 \sum_{\mathbf{u} \in \mathcal{L}_\lambda, \mathbf{u} \neq \mathbf{a}} \mathbf{K}(\bar{\mathbf{x}}(\mathbf{u}) - \bar{\mathbf{x}}(\mathbf{a})).$$

661 Substituting the definition of transformation $\bar{\mathbf{x}}$ in (3.19), we can show for $\mathbf{a}, \mathbf{u} \in \mathcal{L}_\lambda$ that

$$662 \quad \bar{\mathbf{x}}(\mathbf{u}) - \bar{\mathbf{x}}(\mathbf{a}) = \mathbf{Q}(a_1 \theta) [\mathcal{R}(\mathbf{Q}(u_1 - a_1) - \mathbf{I}) \mathbf{e}_1 + (u_2 - a_2) \delta \mathbf{e}_3].$$

663 Using the identities $\mathbf{K}(\mathbf{Q}(t)\mathbf{x}) = \mathbf{Q}(t)\mathbf{K}(\mathbf{x})\mathbf{Q}^T(t)$ and $\mathbf{K}(\lambda\mathbf{x}) = \mathbf{K}(\mathbf{x})/\lambda^3$, from (4.32), we have
 (4.33)

$$664 \quad \mathbf{H}_\lambda(\mathbf{s}) = \mathcal{R}\mathbf{Q}(a_1) \underbrace{\left[\sum_{\mathbf{u} \in \mathcal{L}_\lambda, \mathbf{u} \neq \mathbf{a}} \mathbf{K}(\mathcal{R}(\mathbf{Q}(u_1 - a_1) - \mathbf{I})/\lambda \mathbf{e}_1 + (u_2 - a_2)\delta/\lambda \mathbf{e}_3) \right]}_{=:\bar{\mathbf{H}}_\lambda(\mathbf{s})} \mathbf{Q}(-a_1),$$

665 We analyze $\bar{\mathbf{H}}_\lambda$ as follows. First, we expand the sum $\mathbf{u} \in \mathcal{L}_\lambda$

$$\begin{aligned}
 667 \quad \bar{\mathbf{H}}_\lambda(\mathbf{s}) &= \sum_{u_2 \in \lambda\mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda\theta_1\mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, u_2) \neq \mathbf{a}}} \mathbf{K} \left(\mathcal{R} \frac{\mathbf{Q}(u_1 - a_1) - \mathbf{I}}{\lambda} \mathbf{e}_1 + \delta \frac{u_2 - a_2}{\lambda} \mathbf{e}_3 \right) \right] \\
 668 \quad (4.34) \quad &= \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda\theta_1\mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R} \frac{\mathbf{Q}(u_1 - a_1) - \mathbf{I}}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right], \\
 669
 \end{aligned}$$

where we introduced the new variable $t'_2 = (u_2 - a_2)/\lambda$. Since $u_2, a_2 \in \lambda\mathbb{Z}$, we have $t'_2 \in \mathbb{Z}$. Using a Taylor expansion and the mean value theorem, we have an identity

$$(4.35) \quad \mathbf{Q}(u_1 - a_1) - \mathbf{I} = \mathbf{Q}'(\xi)(u_1 - a_1),$$

where $\xi = \xi(u_1 - a_1) \in (-\bar{\theta}, \bar{\theta})$ depends on $u_1 - a_1$. Based on the above observation, we decompose (4.34) as follows

$$(4.36) \quad \begin{aligned} & \bar{H}_\lambda(s) \\ &= \underbrace{\sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda\theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R}\mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right]}_{=: \bar{H}_\lambda^{(1)}(s)} \\ &+ \underbrace{\sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda\theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, t'_2) \neq (a_1, 0)}} \left\{ \mathbf{K} \left(\mathcal{R} \frac{\mathbf{Q}(u_1 - a_1) - \mathbf{I}}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) - \mathbf{K} \left(\mathcal{R}\mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right\} \right]}_{=: \bar{H}_\lambda^{(2)}(s)} \end{aligned}$$

where two new terms are defined for convenience.

Step 1: We show $\bar{H}_\lambda^{(2)}$ goes to zero in the limit $\lambda \rightarrow 0$. Let

$$(4.37) \quad \mathbf{x}_1 = \mathcal{R} \frac{\mathbf{Q}(u_1 - a_1) - \mathbf{I}}{\lambda} \mathbf{e}_1, \quad \mathbf{x}_2 = \mathcal{R}\mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1, \quad \mathbf{z} = \delta t'_2 \mathbf{e}_3.$$

Consider a function $\mathbf{y}: [0, 1] \rightarrow \mathbb{R}^3$ defined as

$$(4.38) \quad \mathbf{y}(r) = \mathbf{x}_1 + r(\mathbf{x}_2 - \mathbf{x}_1).$$

Note that, since $(u_1, t'_2) \neq (a_1, 0)$ and $t'_2 \in \mathbb{Z}$,

$$(4.39) \quad |\mathbf{y}(r) + \mathbf{z}| \geq \min\{\delta, \min_{r \in [0, 1], u_1 \in \lambda\theta_l \mathbb{Z} - \{a_1\} \cap (-\bar{\theta}, \bar{\theta})} |\mathbf{y}(r)|\}.$$

We show that $\min_{r \in [0, 1], u_1 \in \lambda\theta_l \mathbb{Z} - \{a_1\} \cap (-\bar{\theta}, \bar{\theta})} |\mathbf{y}(r)| > 0$ and the lower bound is independent of λ .

For convenience, let $t = u_1 - a_1$. Since $u_1, a_1 \in \lambda\theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta})$, and $u_1 \neq a_1$, we have $t \in \lambda\theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})$. The hypothesis of Proposition 3.5 restricts $\bar{\theta}$ such that

$$(4.40) \quad 0 < \bar{\theta} < \pi/4 \Rightarrow 1 > \cos(2\bar{\theta}) > 0.$$

With $t = u_1 - a_1$, writing out the action of $\mathbf{Q}(t)$ and $\mathbf{Q}'(0)$ on \mathbf{e}_1 , we get

$$(4.41) \quad \begin{aligned} \mathbf{x}_1 &= \mathcal{R} \frac{\mathbf{Q}(t) - \mathbf{I}}{\lambda} \mathbf{e}_1 = \frac{\mathcal{R}}{\lambda} [(\cos(t) - 1)\mathbf{e}_1 + \sin(t)\mathbf{e}_2] \\ \mathbf{x}_2 &= \mathcal{R}\mathbf{Q}'(0) \frac{t}{\lambda} \mathbf{e}_1 = \frac{\mathcal{R}t}{\lambda} [-\sin(t)\mathbf{e}_1 + \cos(t)\mathbf{e}_2]. \end{aligned}$$

Through elementary calculations, we can show

$$(4.42) \quad |\mathbf{y}(r)|^2 = |\mathbf{x}_1 + r(\mathbf{x}_2 - \mathbf{x}_1)|^2 = \frac{\mathcal{R}^2}{\lambda^2} [2(1-r)^2(1 - \cos(t)) + r^2 t^2 + 2r(1-r)t \sin(t)].$$

697 Using a Taylor expansion and noting that $t \in \lambda\theta_l\mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})$, there exists $\xi_1, \xi_2 \in (-2\bar{\theta}, 2\bar{\theta})$
 698 with $\xi_1 = \xi_1(t)$, $\xi_2 = \xi_2(t)$ such that

$$699 \quad (4.43) \quad 1 - \cos(t) = \cos(\xi_1)t^2/2, \quad \sin(t) = t \cos(\xi_2).$$

700 Thus

$$\begin{aligned} 701 \quad |\mathbf{y}(r)|^2 &= \frac{\mathcal{R}^2}{\lambda^2} [2(1-r)^2 \cos(\xi_1)t^2/2 + r^2t^2 + 2r(1-r)t \cos(\xi_2)t] \\ 702 &= \frac{\mathcal{R}^2t^2}{\lambda^2} [(1-r)^2 \cos(\xi_1) + r^2 + 2r(1-r) \cos(\xi_2)] \\ 703 &\geq \frac{\mathcal{R}^2t^2}{\lambda^2} \left[(1-r)^2 \min_{\xi \in (-2\bar{\theta}, 2\bar{\theta})} \cos(\xi) + r^2 + 2r(1-r) \min_{\xi \in (-2\bar{\theta}, 2\bar{\theta})} \cos(\xi) \right] \\ 704 &= \frac{\mathcal{R}^2t^2}{\lambda^2} [(1-r)^2 \cos(2\bar{\theta}) + r^2 + 2r(1-r) \cos(2\bar{\theta})] \\ 705 &\geq \frac{\mathcal{R}^2t^2}{\lambda^2} \min_{r \in [0,1]} [(1-r)^2 \cos(2\bar{\theta}) + r^2 + 2r(1-r) \cos(2\bar{\theta})] \\ 706 \quad (4.44) &= \frac{\mathcal{R}^2t^2}{\lambda^2} \cos(2\bar{\theta}), \end{aligned}$$

708 where we used the fact that $\min_{\xi \in (-2\bar{\theta}, 2\bar{\theta})} \cos(\xi) = \cos(2\bar{\theta})$ in the fourth equation, and $\cos(2\bar{\theta})$ is the
 709 minimum with respect to $r \in [0, 1]$ of the function in the square bracket in the fifth equation. Further,
 710 since $t \in \lambda\theta_l\mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})$, we have

$$711 \quad (4.45) \quad 0 < C_y := \frac{\mathcal{R}^2\lambda^2\theta_l^2}{\lambda^2} \cos(2\bar{\theta}) = (\mathcal{R}\theta_l)^2 \cos(2\bar{\theta}) \leq |\mathbf{y}(r)|^2,$$

712 for any $t \in \lambda\theta_l\mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})$ and $r \in [0, 1]$. The lower bound on $|\mathbf{y}(r)|$ is independent of λ
 713 and r . Finally, combining (4.45) with (4.39) to get

$$714 \quad (4.46) \quad 0 < C_{yz} := \min\{\delta, \mathcal{R}\theta_l\sqrt{\cos(2\bar{\theta})}\} \leq |\mathbf{y}(r) + \mathbf{z}|.$$

715 Proceeding further, we have, from the fundamental theorem of calculus,

$$\begin{aligned} 716 \quad \mathbf{K}(\mathbf{x}_1 + \mathbf{z}) - \mathbf{K}(\mathbf{x}_2 + \mathbf{z}) &= \int_0^1 \frac{d}{dr} \mathbf{K}(\mathbf{y}(r) + \mathbf{z}) dr = \int_0^1 \nabla \mathbf{K}(\mathbf{y}(r) + \mathbf{z}) \frac{d}{dr} \mathbf{y}(r) dr \\ 717 \quad (4.47) &= \int_0^1 \nabla \mathbf{K}(\mathbf{y}(r) + \mathbf{z})(\mathbf{x}_2 - \mathbf{x}_1) dr. \end{aligned}$$

719 Note that because of (4.46), $\nabla \mathbf{K}(\mathbf{y}(r) + \mathbf{z})$ exists and is bounded. From the definition of $\bar{\mathbf{H}}_\lambda^{(2)}$ in
 720 (4.36), a change of variable $t = u_1 - a_1$, the definition of $\mathbf{x}_1, \mathbf{x}_2, \mathbf{z}$ in (4.37) and (4.41), and noting

the identity (4.47), we have

$$\begin{aligned}
& |\bar{\mathbf{H}}_\lambda^{(2)}(\mathbf{s})| \\
& \leq \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda \theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, t'_2) \neq (a_1, 0)}} \left| \mathbf{K} \left(\mathcal{R} \frac{\mathbf{Q}(u_1 - a_1) - \mathbf{I}}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) - \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right| \right] \\
& \leq \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t \in \lambda \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})} |\mathbf{K}(\mathbf{x}_1 + \mathbf{z}) - \mathbf{K}(\mathbf{x}_2 + \mathbf{z})| \right] \\
& \leq \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t \in \lambda \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})} \int_0^1 |\nabla \mathbf{K}(\mathbf{y}(r) + \mathbf{z})| |\mathbf{x}_2 - \mathbf{x}_1| \, dr \right] \\
& \leq \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t \in \lambda \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})} \int_0^1 \frac{C}{|\mathbf{y}(r) + \mathbf{z}|^4} |\mathbf{x}_2 - \mathbf{x}_1| \, dr \right] \\
& = \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t \in \lambda \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})} \int_0^1 \frac{C}{(|\mathbf{y}(r)|^2 + |\mathbf{z}|^2)^2} |\mathbf{x}_2 - \mathbf{x}_1| \, dr \right],
\end{aligned}
\tag{4.48}$$

where we utilized the bound on the gradient of \mathbf{K} with constant $C > 0$ fixed.

Next, we get an upper bound on $|\mathbf{x}_1 - \mathbf{x}_2|$ in terms of t . From (4.41), we have

$$\mathbf{x}_2 - \mathbf{x}_1 = \frac{\mathcal{R}}{\lambda} [t \mathbf{Q}'(0) - \mathbf{Q}(t) + \mathbf{I}] \mathbf{e}_1.$$

By a Taylor expansion and the mean value theorem, we have $\mathbf{Q}(t) = \mathbf{I} + t \mathbf{Q}'(0) + (t^2/2) \mathbf{Q}''(\xi)$ where $\xi = \xi(t) \in (-2\bar{\theta}, 2\bar{\theta})$ depends on t . Substituting this and using the bound $|\mathbf{Q}''_{ij}(\xi)| \leq 1$ gives

$$|\mathbf{x}_2 - \mathbf{x}_1| = \frac{\mathcal{R}}{\lambda} \frac{|t|^2}{2} |\mathbf{Q}''(\xi)| \leq \frac{\mathcal{R}}{\lambda} \frac{|t|^2}{2}.$$

Combining the equation above with (4.48), we get

$$\begin{aligned}
& |\bar{\mathbf{H}}_\lambda^{(2)}(\mathbf{s})| \\
& \leq \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t \in \lambda \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}, 2\bar{\theta})} \int_0^1 \frac{C}{(|\mathbf{y}(r)|^2 + |\mathbf{z}|^2)^2} \frac{\mathcal{R}}{\lambda} \frac{|t|^2}{2} \, dr \right] \\
& = \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t' \in \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}/\lambda, 2\bar{\theta}/\lambda)} \int_0^1 \frac{C}{(|\mathbf{y}(r)|^2 + |\mathbf{z}|^2)^2} \frac{\mathcal{R}}{\lambda} \frac{\lambda^2 |t'|^2}{2} \, dr \right] \\
& \leq \lambda \left\{ \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t' \in \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}/\lambda, 2\bar{\theta}/\lambda)} \int_0^1 \frac{C}{(|\mathbf{y}(r)|^2 + |\mathbf{z}|^2)^2} \frac{\mathcal{R} |t'|^2}{2} \, dr \right] \right\},
\end{aligned}$$

where in the third line we introduced the variable $t' = t/\lambda$. We only have to show that the term inside the braces is bounded as $\lambda \rightarrow 0$ to conclude that $|\bar{\mathbf{H}}_\lambda^{(2)}(\mathbf{s})| \rightarrow 0$ as $\lambda \rightarrow 0$. First, note from (4.44),

we have

$$(4.50) \quad |\mathbf{y}(r)|^2 \geq \frac{\mathcal{R}^2}{\lambda^2} |t|^2 \cos(2\bar{\theta}) = \mathcal{R}^2 |t'|^2 \cos(2\bar{\theta}).$$

Therefore,

$$(4.51) \quad \frac{C}{(|\mathbf{y}(r)|^2 + |\mathbf{z}|^2)^2} \leq \frac{C}{(\mathcal{R}^2 |t'|^2 \cos(2\bar{\theta}) + |\mathbf{z}|^2)^2}.$$

Thus

$$(4.52) \quad |\bar{\mathbf{H}}_\lambda^{(2)}(\mathbf{s})| \leq \lambda \left\{ \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{t' \in \theta_l \mathbb{Z} - \{0\} \cap (-2\bar{\theta}/\lambda, 2\bar{\theta}/\lambda)} \int_0^1 \frac{C}{(\mathcal{R}^2 |t'|^2 \cos(2\bar{\theta}) + |\mathbf{z}|^2)^2} \frac{\mathcal{R} |t'|^2}{2} dr \right] \right\}.$$

Note that the integrand is independent of r . Further, the numerator has $|t'|^2$ whereas the denominator has $(|t'|^2 c + |\mathbf{z}|^2)^2$, therefore, the sum inside the braces is absolutely convergent and finite. Hence, due to the factor λ , we have shown $\lim_{\lambda \rightarrow 0} |\bar{\mathbf{H}}_\lambda^{(2)}(\mathbf{s})| = 0$.

This completes the step 1. We next study $\bar{\mathbf{H}}_\lambda^{(1)}$.

Step 2: We have from (4.36)

$$(4.53) \quad \begin{aligned} \bar{\mathbf{H}}_\lambda^{(1)}(\mathbf{s}) &= \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda \theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta}), \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right] \\ &= \underbrace{\sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda \theta_l \mathbb{Z}, \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right]}_{=: I_1} \\ &\quad - \underbrace{\sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in [\lambda \theta_l \mathbb{Z}] - [\lambda \theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta})], \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right]}_{=: I_2}, \end{aligned}$$

where we have used the notation $[\lambda \theta_l \mathbb{Z}] - [\lambda \theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta})]$ to denote the set $\{t \in \lambda \theta_l \mathbb{Z}; t \notin \lambda \theta_l \mathbb{Z} \cap (-\bar{\theta}, \bar{\theta})\}$. Using the decay property of the dipole field kernel \mathbf{K} , we can show that $|I_2| \rightarrow 0$ in the limit $\lambda \rightarrow 0$. Therefore, we have

$$(4.54) \quad \begin{aligned} \lim_{\lambda \rightarrow 0} \bar{\mathbf{H}}_\lambda^{(1)}(\mathbf{s}) &= \lim_{\lambda \rightarrow 0} \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{u_1 \in \lambda \theta_l \mathbb{Z}, \\ (u_1, t'_2) \neq (a_1, 0)}} \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) \frac{u_1 - a_1}{\lambda} \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right] \\ &= \sum_{t'_2 \in \mathbb{Z}} \left[\sum_{\substack{t'_1 \in \theta_l \mathbb{Z}, \\ (t'_1, t'_2) \neq (0, 0)}} \mathbf{K} \left(\mathcal{R} \mathbf{Q}'(0) t'_1 \mathbf{e}_1 + \delta t'_2 \mathbf{e}_3 \right) \right] \end{aligned}$$

where we introduced the new variable $t'_1 = (u_1 - a_1)/\lambda$. Since $u_1 \in \lambda\theta_1\mathbb{Z}$ and $a_1 \in \lambda\theta_1\mathbb{Z} \cap (-\bar{\theta}, \bar{\theta})$, we have $t'_1 \in \theta_1\mathbb{Z}$. This completes step 2. Note that $\lim_{\lambda \rightarrow 0} \bar{H}_\lambda(s)$ is independent of $s \in \mathcal{S}$.

Upon substituting the limit of $\bar{H}_\lambda^{(1)}$ and $\bar{H}_\lambda^{(2)}$ in (4.36), we have shown

$$(4.55) \quad \lim_{\lambda \rightarrow 0} \bar{H}_\lambda(s) = \lim_{\lambda \rightarrow 0} \bar{H}_\lambda(\mathbf{0}) = \sum_{\substack{\mathbf{u}=(u_1, u_2) \in \theta_1\mathbb{Z} \times \mathbb{Z}, \\ \mathbf{u} \neq \mathbf{0}}} \mathbf{K}(\mathcal{R}\mathbf{Q}'(0)u_1\mathbf{e}_1 + \delta u_2\mathbf{e}_3).$$

Recall that $\mathbf{s} \in \mathcal{S}$ was fixed such that $\mathbf{s} \in U_\lambda(\mathbf{a})$, which implies that $\mathbf{a} \rightarrow \mathbf{s}$ as $\lambda \rightarrow 0$. With this observation and (4.55), we have from (4.33),

$$(4.56) \quad \mathbf{H}_0(s) = \lim_{\lambda \rightarrow 0} \mathbf{H}_\lambda(s) = \mathcal{R}\mathbf{Q}(s_1) \left[\sum_{\substack{\mathbf{u}=(u_1, u_2) \in \theta_1\mathbb{Z} \times \mathbb{Z}, \\ \mathbf{u} \neq \mathbf{0}}} \mathbf{K}(\mathcal{R}\mathbf{Q}'(0)u_1\mathbf{e}_1 + \delta u_2\mathbf{e}_3) \right] \mathbf{Q}(-s_1).$$

Next we simplify $\mathbf{H}_0(s)$. Given the parametric map $\bar{\mathbf{x}} = \bar{\mathbf{x}}(s)$, the two tangent vectors at $\mathbf{s} = (s_1, s_2)$ are

$$(4.57) \quad \mathbf{t}_1(s) = \frac{d\bar{\mathbf{x}}}{ds_1} = \mathcal{R}\mathbf{Q}'(s_1)\mathbf{e}_1, \quad \mathbf{t}_2(s) = \frac{d\bar{\mathbf{x}}}{ds_2} = \delta\mathbf{e}_3.$$

Using $\mathbf{Q}\mathbf{K}(\mathbf{x})\mathbf{Q}^T = \mathbf{K}(\mathbf{Q}\mathbf{x})$ and $\mathbf{Q}(r)\mathbf{Q}'(0) = \mathbf{Q}'(r)$, we write,

$$(4.58) \quad \mathbf{H}_0(s) = \sum_{\substack{\mathbf{u}=(u_1, u_2) \in \theta_1\mathbb{Z} \times \mathbb{Z}, \\ \mathbf{u} \neq \mathbf{0}}} \mathbf{K}(u_1\mathbf{t}_1(s) + u_2\mathbf{t}_2(s)).$$

This completes the proof of Proposition 3.5.

5. Summary of Results. We have shown rigorously that certain low-dimensional nanostructures do not have long-range dipole-dipole interaction in the continuum limit. The energy density in the limit is entirely due to Maxwell-self field. In 1-d and 2-d lattices (in a 3-d ambient space), the dipole field kernel decay is sufficiently fast that long-range interactions do not contribute to the limit energy.

While our calculations show that the energy is local in the continuum limit for 1-d and 2-d discrete systems, in agreement with dimension reduction approaches that reduce a 3-d continuum to a 1-d or 2-d continuum (e.g., [GJ97, CH15] and others), we note an interesting difference. As shown in [GJ97] and other work following it, the thin film limit of the continuum electrostatic energy is due to the component of dipole moment field along the normal direction to the film. Similarly, the thin wire limit of the energy is due to components of dipole moment field in the plane normal to the wire, see [CH15]. This is different from the limit energy in the discrete-to-continuum limit: for the case of a helical nanotube, the limiting energy density is given by

$$h_0 \int_{\mathbb{R}} [|\mathbf{P}_\perp \mathbf{f}|^2 - 2|\mathbf{P}_\parallel \mathbf{f}|^2] \, ds,$$

where h_0 is a constant, and $\mathbf{P}_\parallel \mathbf{f}$ and $\mathbf{P}_\perp \mathbf{f}$ are the projections of the dipole moment field \mathbf{f} along the axis of the helix and in the plane normal to the axis of the helix respectively. Therefore, unlike the thin wire limit using dimension reduction, the discrete-to-continuum energy has contributions from both the normal and tangential components of the dipole moment field. For the case of a thin film with curvature, the limiting energy density is given by

$$-\frac{1}{2} \int_{\mathcal{S}} \mathbf{f}(s) \cdot \mathbf{H}_0(s) \mathbf{f}(s) \, ds,$$

where

$$H_0(\mathbf{s}) = \mathcal{R} \sum_{\substack{\mathbf{u}=(u_1, u_2) \in \theta_l \mathbb{Z} \times \mathbb{Z}, \\ \mathbf{u} \neq \mathbf{0}}} K(u_1 \mathbf{t}_1(\mathbf{s}) + u_2 \mathbf{t}_2(\mathbf{s})).$$

Here, \mathcal{S} is the parametric domain of the film, \mathcal{R} is the inverse of the curvature, θ_l is the angular width of the unit cell, and $\mathbf{t}_i(\mathbf{s})$, $i = 1, 2$, are the tangent vectors at coordinate $\mathbf{s} \in \mathcal{S}$. For simplicity, we fix $\mathbf{s} \in \mathcal{S}$ and assume $\mathbf{t}_1 = \mathbf{e}_1$ and $\mathbf{t}_2 = \mathbf{e}_2$; then the lattice sum above is over a 2-d lattice in $(\mathbf{e}_1, \mathbf{e}_2)$ plane. By substituting the form of \mathbf{K} and computing $H_0(\mathbf{s})\mathbf{f}(\mathbf{s})$, we can show that both the normal and the tangential components of \mathbf{f} are present in the final expression for the energy above. The difference between the dimension reduction approach on the one hand, and the discrete-to-continuum limit on the other hand, is due to the fact that we assume that the discrete structures consist of a finite unit cell in the thickness direction and do not take the limit in the thickness direction.

Acknowledgments. This paper draws from the doctoral dissertation of Prashant K. Jha at Carnegie Mellon University [Jha16]. We thank Richard D. James for useful discussions, and AFRL for hosting a visit by Kaushik Dayal.

REFERENCES

- [ADE13a] Amin Aghaei, Kaushik Dayal, and Ryan S Elliott. Anomalous phonon behavior of carbon nanotubes: First-order influence of external load. *Journal of Applied Physics*, 113(2):023503, 2013.
- [ADE13b] Amin Aghaei, Kaushik Dayal, and Ryan S Elliott. Symmetry-adapted phonon analysis of nanotubes. *Journal of the Mechanics and Physics of Solids*, 61(2):557–578, 2013.
- [BLBL02] Xavier Blanc, Claude Le Bris, and P-L Lions. From molecular models to continuum mechanics. *Archive for Rational Mechanics and Analysis*, 164(4):341–381, 2002.
- [BLBL07] Xavier Blanc, Claude Le Bris, and Pierre-Louis Lions. Atomistic to continuum limits for computational materials science. *ESAIM: Mathematical Modelling and Numerical Analysis*, 41(2):391–426, 2007.
- [Bro63] William Fuller Brown. *Micromagnetics*. Number 18. interscience publishers, 1963.
- [Car01] Gilles Carbou. Thin layers in micromagnetism. *Mathematical Models and Methods in Applied Sciences*, 11(09):1529–1546, 2001.
- [CH15] Khaled Chacouche and Rejeb Hadji. Ferromagnetic of nanowires of infinite length and infinite thin films. *Zeitschrift für angewandte Mathematik und Physik*, 66(6):3519–3534, 2015.
- [DJ07] Traian Dumitrică and Richard D James. Objective molecular dynamics. *Journal of the Mechanics and Physics of Solids*, 55(10):2206–2236, 2007.
- [DLO10] Matthew Dobson, Mitchell Luskin, and Christoph Ortner. Sharp stability estimates for the force-based quasicontinuum approximation of homogeneous tensile deformation. *Multiscale Modeling & Simulation*, 8(3):782–802, 2010.
- [GD20a] Matthew Grasinger and Kaushik Dayal. Architected elastomer networks for optimal electromechanical response. *Journal of the Mechanics and Physics of Solids*, page 104171, 2020.
- [GD20b] Matthew Grasinger and Kaushik Dayal. Statistical mechanical analysis of the electromechanical coupling in an electrically-responsive polymer chain. *Soft Matter*, 2020.
- [GH15] Antonio Gaudiello and Kamel Hamdache. A reduced model for the polarization in a ferroelectric thin wire. *Nonlinear Differential Equations and Applications NoDEA*, 22(6):1883–1896, 2015.
- [GJ97] Gustavo Gioia and Richard D James. Micromagnetics of very thin films. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 453(1956):213–223, 1997.
- [HTJ12] Ye Hakobyan, Ellad B Tadmor, and Richard D James. Objective quasicontinuum approach for rod problems. *Physical Review B*, 86(24):245435, 2012.
- [Jam06] Richard D James. Objective structures. *Journal of the Mechanics and Physics of Solids*, 54(11):2354–2390, 2006.
- [Jha16] Prashant Jha. *Coarse Graining of Electric Field Interactions with Materials*. PhD thesis, Carnegie Mellon University, 2016.
- [JM94] Richard D James and Stefan Müller. Internal variables and fine-scale oscillations in micromagnetics. *Continuum Mechanics and Thermodynamics*, 6(4):291–336, 1994.
- [KO01] J Knap and M Ortiz. An analysis of the quasicontinuum method. *Journal of the Mechanics and Physics of Solids*, 49(9):1899–1923, 2001.
- [KS05] Robert V Kohn and Valeriy V Slastikov. Another thin-film limit of micromagnetics. *Archive for rational mechanics and analysis*, 178(2):227–245, 2005.

- [KSZ15] Martin Kružík, Ulisse Stefanelli, and Chiara Zanini. Quasistatic evolution of magnetoelastic plates via dimension reduction. *Discrete & Continuous Dynamical Systems-A*, 35(12):5999, 2015.
- [LLO12] Xingjie Helen Li, Mitchell Luskin, and Christoph Ortner. Positive definiteness of the blended force-based quasicontinuum method. *Multiscale Modeling & Simulation*, 10(3):1023–1045, 2012.
- [MD14] Jason Marshall and Kaushik Dayal. Atomistic-to-continuum multiscale modeling with long-range electrostatic interactions in ionic solids. *Journal of the Mechanics and Physics of Solids*, 62:137–162, 2014.
- [MS02] Stefan Müller and Anja Schlömerkemper. Discrete-to-continuum limit of magnetic forces. *Comptes Rendus Mathématique*, 335(4):393–398, 2002.
- [MT02] Ronald E Miller and Ellad B Tadmor. The quasicontinuum method: Overview, applications and current directions. *Journal of Computer-Aided Materials Design*, 9(3):203–239, 2002.
- [Sch06] Bernd Schmidt. A derivation of continuum nonlinear plate theory from atomistic models. *Multiscale Modeling & Simulation*, 5(2):664–694, 2006.
- [Sch09] Bernd Schmidt. On the derivation of linear elasticity from atomistic models. *Networks & Heterogeneous Media*, 4(4):789, 2009.
- [Sog17] Christopher D Sogge. *Fourier integrals in classical analysis*, volume 210. Cambridge University Press, 2017.
- [SS09] Anja Schlömerkemper and Bernd Schmidt. Discrete-to-continuum limit of magnetic forces: dependence on the distance between bodies. *Archive for rational mechanics and analysis*, 192(3):589–611, 2009.
- [SWB⁺21] Shoham Sen, Yang Wang, Timothy Breitzman, , and Kaushik Dayal. Two-scale analysis of the polarization density in ionic solids. *in preparation*, 2021.
- [TM11] Ellad B Tadmor and Ronald E Miller. *Modeling materials: continuum, atomistic and multiscale techniques*. Cambridge University Press, 2011.
- [TOP96] Ellad B Tadmor, Michael Ortiz, and Rob Phillips. Quasicontinuum analysis of defects in solids. *Philosophical magazine A*, 73(6):1529–1563, 1996.
- [Tou56] Richard A Toupin. The elastic dielectric. *Journal of Rational Mechanics and Analysis*, 5(6):849–915, 1956.
- [Xia05] Yu Xiao. *The influence of oxygen vacancies on domain patterns in ferroelectric perovskites*. PhD thesis, California Institute of Technology, 2005.