

A Multiscale Model of Glioblastoma-Mesenchymal Stromal Cells Tumor Spheroids

Alzbeta Prasilova¹, Jarmila Havelkova¹, Olena Rogulska¹, Eliska Vavrinova¹, Bela Charvatova², Alzbeta Stetkova², Tomas Bily², Yuriy Petrenko¹

¹Institute of Experimental Medicine CAS, Prague; ²Department of Software Engineering, Faculty of Mathematics and Physics, Charles University, Prague




1 EXPERIMENTAL OBSERVATION: MIXED SPHEROIDS GROW NON-LINEARLY

MOTIVATION & CONTEXT


- Tumor growth is shaped not only by cancer cells, but also by interactions with the surrounding microenvironment.
- GBM spheroids are models mimicking tumor-like organization in 3D. The cell-cell contact, nutrient/metabolite gradients, and stressed inner regions in spheroids offer better representation than 2D cultures.
- Mesenchymal stromal cells (MSCs) are context-dependent: they can either suppress or support GBM growth depending on local conditions.
- To understand MSC effects, we need to look beyond overall spheroid size and examine spatial and dynamic processes inside the spheroid.

Observed vs. linearly interpolated growth curves




Mixed GBM-MSC spheroids grew larger than expected from pure GBM and pure MSC controls, indicating a non-additive stromal effect.

Magnitude of non-linearity



METHODS / GROWTH ASSAY

- U87MG glioblastoma + mesenchymal stromal cell (MSC) spheroids
- 10,000 cells seeded per spheroid
- Conditions: GBM only, MSC only, GBM:MSC 3:1, 1:1, 1:3
- 7-day live imaging of spheroid growth
- Spheroid area quantified from brightfield images



What internal mechanisms could produce this non-linear, pro-tumor MSC effect?

2 OUR APPROACH: A HYBRID 3D COMPUTATIONAL MODEL

OVERVIEW

- We developed a hybrid 3D agent-based model of GBM-MSC spheroid growth that couples:
- Cellular Potts Model (CPM) for cell mechanics
 - Reaction-Diffusion fields for nutrients, metabolites, ECM components and signals
 - Probabilistic logic-based rules for cell behavior and fate decisions

CELLS

- Cells are discrete agents
 - Evolves by minimizing an energy function (Hamiltonian)
 - Cells types & states:
- $H = \sum_{(i,j)} J(\tau_i, \tau_j) \cdot (1 - \delta_{\tau_i, \tau_j}) + \lambda_V (V - V_{\text{target}})^2 + \lambda_S (S - S_{\text{target}})^2$
- adhesion volume surface
- GBM: P, Q, N; GSC: P, Q; MSC: anti, pro
- P = proliferative, Q = quiescent, N = necrotic


FIELDS

- Reaction-diffusion equations
 - Actions:
- diffuse, decay, consume, produce, inject

RULES


- Local sensing of environment conditions
- Probabilistic state transitions
- Cell-cycle and division control
- Context-dependent MSC phenotype switching

SIMULATION WORKFLOW




3 THE MODEL RECAPITULATES HALLMARK FEATURES OF GBM SPHEROIDS

Emergent spheroid organization: nutrient gradients restrict proliferation to the outer rim and drive formation of quiescent and necrotic inner regions.



4 PRO-TUMOR MSC PHENOTYPE NEEDED TO REPRODUCE EXPERIMENTAL TRENDS

MODEL-EXPERIMENT AGREEMENT




Qualitative agreement in curve ordering and overall trajectory
Comparison is qualitative (2D projected area vs 3D simulated volume)

PRO-TUMOR MSC RULES

- 1. TRIGGERED BY**
 - high lactate / acidity
 - hypoxia
 - GBM contact
- 2. MODELED ACTIONS**
 - growth-signal secretion
 - metabolic support
 - indirect buffering of stress / field gradients
- 3. RESULT**
 - qualitative match to co-culture growth trends
 - delayed quiescent / necrotic core onset
 - sustained GBM proliferation

5 TOWARD DRUG RESPONSE PREDICTION



SUMMARY

MSC EFFECT

GBM-MSC growth was **non-linear**: mixed spheroids could not be explained by pure GBM + pure MSC behavior alone.

MODEL

3D hybrid model captured **spheroid physiology**: viable rim, nutrient/metabolite gradients, and quiescent/necrotic core formation.

MECHANISTIC INTERPRETATION

Qualitative agreement required **MSCs** to act as context-dependent **pro-tumor** modifiers of the local microenvironment.


DRUG RESPONSE

Model was extended to include **signaling** and **metabolic** pathways. Functionality was compared to **Buparlisib** drug response experimental data.

Any questions? Contact me!

email: alzbeta.prasilova@iem.cas.cz

Acknowledgement: The study was supported by the Czech Science Foundation grants GAČR 26-21666S and ExRegMed project No CZ.02.01.01/00/22_008/0004562, of the Ministry of Education, Youth and Sports, co-funded by the European Union.



More info on my GitHub:

