

myCSG © 2026 | Area: ADAM | Concept: C1003 : Treatment emergence, pre and post treatment occurrences | Lesson: ADAM_C1003_L201F : ADCM - Prior and concomitant medication flagging |

Lesson Description

Spec:

[SAS Program](#)

[R Program](#)

ADVERTISEMENT

Variable	Derivation
PREFL	A medication is flagged as prior to first treatment if the medication start datetime occurs before the treatment start datetime. If either datetime is missing, the comparison is based on the available start dates. If the medication start date is missing but the treatment start date is present, the medication is also classified as prior. The flag is set to "Y" for records meeting any of these conditions and is left null otherwise.
ONTRTFL	A medication is flagged as on-treatment if the medication start datetime occurs during the treatment period or if the medication begins before treatment but continues into the treatment period. The comparison is first performed using start and end datetimes relative to treatment start and end datetimes. If datetime values are missing, the comparison is based on the available start and end dates. Medications that start before treatment and have no end date or end on or after the treatment start date are considered on-treatment. Medications that start during the treatment period are also considered on-treatment. If the medication start date is missing but the end date occurs on or after the treatment start date, or if both start and end dates are missing while the treatment start date is available, the medication is also classified as on-treatment. The flag is set to "Y" for records meeting any of these conditions and is left null otherwise.

ADVERTISEMENT