Background

- In a clinical trial, we collect information related to medical history, prior medication and concomitant medications, adverse events etc.
- For these 'events' or 'interventions', we collect the start and end dates of occurrence
 - Start and end date of a particular medical condition
 - Start and end date of a specific medication used
 - Start and end date of any adverse event experienced by the clinical trial participant (subject)
- Often, subjects do not remember the exact date and could only partially recollect the dates
 - In 2010
 - In 2015 January
 - In between 10 am and 11 am of 15Jan2019 (not sure of the exact time).
 - Do not remember the date at all
 - etc.
- This results in partial dates in the collected data
- For analysis purposes, we may want to convert these partial dates into full dates using a predefined algorithm
- This process of converting partial dates to full dates for analysis purposes based on predetermined rules is called imputation of dates

ADaM standard for date imputation related variables

- ADaM allows for imputation of dates
- We need traceability as per ADaM standard - that is ability to identify the source values and indicate what changed
- In terms of traceability for date imputation, we have variables to indicate the imputation of a date on a record in the ADaM dataset
 - *DTF variable to indicate the imputation of date value
 - *TMF variable to indicate the imputation of time value
 - *DTMF variable indicate the imputation of date and time value
- These three variables are used to hold a value that represents the highest level of imputation done
- The possible values in *DTF variable are
 - Y - when year is imputed
 - M - when month is imputed
 - D- when day of the month is imputed
- The possible values in *TMF variable are
 - H - when hour is imputed
 - M - when minutes are imputed
 - S - when seconds are imputed