

Background

- In some clinical trials, a functional test named six-minute walk test is used to measure the distance the patient can walk in six minutes.
- Let us assume that in a specific clinical trial, to assess the impact of physical exertion of the walk test, Oxygen Saturation (SPO2) is measured before and after a 6-minute walk test.
- And, the results for oxygen saturation are to be reported in terms of a summary of pre-test and post-test oxygen saturation level (decay) as shown below.

Parameter	Timepoint	Statistic	
Oxygen Saturation	Week 1, Pre-test	n	x
		Mean (SD)	xx.x
Oxygen Saturation	Week 1, Post-test	n	x
		Mean (SD)	xx.x
Oxygen Saturation Decay	Week 1	n	x
		Mean (SD)	xx.x

Structuring the ADaM dataset

- For this analysis, we need to derive the decay observed (post-test saturation minus pre-test saturation).
- There are two rules in ADaM implementation guide that govern the cases of analysis data points related to this analysis:
 - Rule 3: A function of one or more rows within the same parameter for the purpose of creating an analysis data point should be added as a new parameter.
 - Rule 4: A function of multiple rows within a parameter should be added as a new parameter.
- Here, as the analysis datapoint involves more than one row within same parameter but not analyzed as a single row, we will follow Rule 4.

What will we learn in this lesson

- In this lesson, we will see how to derive a new parameter that follows rule 4.