

ADVERTISEMENT

TEAE by Preferred Term - with fisher's exact test p-value				
Safety Analysis Set				
Copyright @ www.mycsg.in				
Preferred Term	Placebo (N=15) n (%)	Low Dose (N=15) n (%)	High Dose (N=15) n (%)	Low dose vs Placebo p-value [1]
Overall	11 (73.3%)	13 (86.7%)	15 (100.0%)	0.6513
ACTINIC KERATOSIS	0	0	1 (6.7%)	-
AGITATION	0	0	1 (6.7%)	-
APPLICATION SITE DERMATITIS	0	1 (6.7%)	1 (6.7%)	1.0000
APPLICATION SITE ERYTHEMA	1 (6.7%)	7 (46.7%)	10 (66.7%)	0.0352
APPLICATION SITE IRRITATION	0	5 (33.3%)	4 (26.7%)	0.0421
APPLICATION SITE PAIN	0	0	2 (13.3%)	-
APPLICATION SITE PERSPIRATION	0	0	1 (6.7%)	-
APPLICATION SITE PRURITUS	2 (13.3%)	9 (60.0%)	12 (80.0%)	0.0209
APPLICATION SITE URTICARIA	0	2 (13.3%)	0	0.4828
APPLICATION SITE VESICLES	0	2 (13.3%)	5 (33.3%)	0.4828
ARTHRALGIA	0	1 (6.7%)	0	1.0000
ATRIOVENTRICULAR BLOCK SECOND DEGREE	1 (6.7%)	0	0	1.0000
BACK PAIN	1 (6.7%)	0	0	1.0000
BENIGN PROSTATIC HYPERPLASIA	0	0	1 (6.7%)	-
BIOPSY PROSTATE	0	0	1 (6.7%)	-
BUNDLE BRANCH BLOCK LEFT	1 (6.7%)	0	0	1.0000
CALCULUS URETHRAL	0	0	1 (6.7%)	-
CELLULITIS	0	1 (6.7%)	0	1.0000
CHEST DISCOMFORT	0	0	1 (6.7%)	-
CHILLS	0	1 (6.7%)	0	1.0000
CONFUSIONAL STATE	0	1 (6.7%)	0	1.0000
CONTUSION	0	0	1 (6.7%)	-
COUGH	0	0	2 (13.3%)	-
DEPRESSED MOOD	0	1 (6.7%)	0	1.0000
DIARRHOEA	4 (26.7%)	0	0	0.0996
DIZZINESS	0	0	1 (6.7%)	-
DYSPHONIA	0	1 (6.7%)	0	1.0000
DYSPNOEA	0	1 (6.7%)	0	1.0000
EPISTAXIS	0	0	2 (13.3%)	-
ERYTHEMA	1 (6.7%)	3 (20.0%)	2 (13.3%)	0.5977
EYE ALLERGY	1 (6.7%)	0	0	1.0000

ADVERTISEMENT