

ADVERTISEMENT

ADVERTISEMENT

TEAE by PT - Odds Ratio			
Safety Analysis Set			
Copyright @ www.mycsg.in			
Preferred Term	Placebo (N=15) n (%)	Active (N=15) n (%)	OR (95% CI)
Overall	11 (73.3%)	13 (86.7%)	2.4 (0.4,15.5)
APPLICATION SITE DERMATITIS	0	1 (6.7%)	-
APPLICATION SITE ERYTHEMA	1 (6.7%)	7 (46.7%)	12.3 (1.3,118.4)
APPLICATION SITE IRRITATION	0	5 (33.3%)	-
APPLICATION SITE PRURITUS	2 (13.3%)	9 (60.0%)	9.8 (1.6,59.7)
APPLICATION SITE URTICARIA	0	2 (13.3%)	-
APPLICATION SITE VESICLES	0	2 (13.3%)	-
ARTHRALGIA	0	1 (6.7%)	-
ATRIOVENTRICULAR BLOCK SECOND DEGREE	1 (6.7%)	0	-
BACK PAIN	1 (6.7%)	0	-
BUNDLE BRANCH BLOCK LEFT	1 (6.7%)	0	-
CELLULITIS	0	1 (6.7%)	-
CHILLS	0	1 (6.7%)	-
CONFUSIONAL STATE	0	1 (6.7%)	-
DEPRESSED MOOD	0	1 (6.7%)	-
DIARRHOEA	4 (26.7%)	0	-
DYSPHONIA	0	1 (6.7%)	-
DYSPNOEA	0	1 (6.7%)	-
ERYTHEMA	1 (6.7%)	3 (20.0%)	3.5 (0.3,38.2)
EYE ALLERGY	1 (6.7%)	0	-
EYE LASER SURGERY	1 (6.7%)	0	-
EYE PRURITUS	1 (6.7%)	0	-
EYE SWELLING	1 (6.7%)	0	-
FATIGUE	0	2 (13.3%)	-
HIATUS HERNIA	1 (6.7%)	0	-
HYPERHIDROSIS	1 (6.7%)	0	-
INCONTINENCE	0	1 (6.7%)	-
INCREASED APPETITE	1 (6.7%)	0	-
INSOMNIA	1 (6.7%)	0	-
JOINT DISLOCATION	0	1 (6.7%)	-
MICTURITION URGENCY	1 (6.7%)	1 (6.7%)	1.0 (0.1,17.6)
MUSCULAR WEAKNESS	0	1 (6.7%)	-
NASAL CONGESTION	1 (6.7%)	1 (6.7%)	1.0 (0.1,17.6)