

ADVERTISEMENT

Treatment Disposition Safety Analysis Set Copyright @ www.mycsg.in			
	Placebo (N=57)	Low Dose (N=62)	High Dose (N=59)
Treatment disposition			
Ongoing	19 (33.3%)	16 (25.8%)	16 (27.1%)
Completed	20 (35.1%)	20 (32.3%)	17 (28.8%)
Discontinued	18 (31.6%)	26 (41.9%)	26 (44.1%)
Primary reason for discontinuing treatment			
Adverse Event	1 (1.8%)	4 (6.5%)	2 (3.4%)
Protocol Deviation	1 (1.8%)	1 (1.6%)	3 (5.1%)
Lost to Follow-up	1 (1.8%)	1 (1.6%)	5 (8.5%)
Withdrawal by Subject	0	3 (4.8%)	0
Investigator Discretion	0	1 (1.6%)	2 (3.4%)
Intolerable Toxicity	6 (10.5%)	3 (4.8%)	3 (5.1%)
Progressive Disease per RECIST	5 (8.8%)	4 (6.5%)	6 (10.2%)
Clinical Progression Without Radiological PD by RECIST	0	7 (11.3%)	2 (3.4%)
Other	4 (7.0%)	2 (3.2%)	3 (5.1%)

ADVERTISEMENT