

ADVERTISEMENT

Treatment Disposition - coding approach 2			
Safety Analysis Set			
Copyright @ www.mycsg.in			
	Placebo (N=48)	Low Dose (N=63)	High Dose (N=67)
Treatment disposition			
Ongoing	13 (27.1%)	21 (33.3%)	21 (31.3%)
Completed	14 (29.2%)	25 (39.7%)	23 (34.3%)
Discontinued	21 (43.8%)	17 (27.0%)	23 (34.3%)
Primary reason for discontinuing treatment			
Adverse Event	2 (4.2%)	3 (4.8%)	3 (4.5%)
Protocol Deviation	3 (6.3%)	2 (3.2%)	1 (1.5%)
Lost to Follow-up	2 (4.2%)	1 (1.6%)	2 (3.0%)
Withdrawal by Subject	1 (2.1%)	1 (1.6%)	4 (6.0%)
Investigator Discretion	2 (4.2%)	3 (4.8%)	3 (4.5%)
Intolerable Toxicity	3 (6.3%)	2 (3.2%)	4 (6.0%)
Progressive Disease per RECIST	1 (2.1%)	2 (3.2%)	1 (1.5%)
Clinical Progression Without Radiological PD by RE	3 (6.3%)	2 (3.2%)	5 (7.5%)
Other	4 (8.3%)	1 (1.6%)	0

ADVERTISEMENT