

ADSL

Treatment Disposition
Safety Analysis Set
Copyright @ www.mycsg.in

SAFFL

TRT01A

	Placebo (N=54)	Low Dose (N=61)	High Dose (N=63)
Treatment disposition			
Ongoing	24 (44.4%)	18 (29.5%)	18 (28.6%)
Completed	16 (29.6%)	18 (29.5%)	28 (44.4%)
Discontinued	14 (25.9%)	25 (41.0%)	17 (27.0%)
Primary reason for discontinuing treatment			
Adverse Event	0	7 (11.5%)	1 (1.6%)
Protocol Deviation	4 (7.4%)	1 (1.6%)	4 (6.3%)
Lost to Follow-up	1 (1.9%)	4 (6.6%)	2 (3.2%)
Withdrawal by Subject	0	2 (3.3%)	2 (3.2%)
Investigator Discretion	3 (5.6%)	1 (1.6%)	1 (1.6%)
Intolerable Toxicity	3 (5.6%)	3 (4.9%)	1 (1.6%)
Progressive Disease per RECIST	0	1 (1.6%)	1 (1.6%)
Clinical Progression Without Radiological PD by RECIST	2 (3.7%)	3 (4.9%)	1 (1.6%)
Other	1 (1.9%)	3 (4.9%)	4 (6.3%)