

ADVERTISEMENT

25

▼ entries per page

Search:

ADVERTISEMENT

ROWORDER ↕	VARIABLE_LABEL ↕	METHOD ↕	FORMULA_FOR_EGFR_DERIVATION ↕
001		CKD-EPI	eGFR = 141 * min(Scr/k,1)**a * max(Scr/k,1)**-1.209 * 0.993**Age * [1.018 if female] * [1.159 if black] Where: - Scr = serum creatinine (mg/dL; to convert from umol/L to mg/dL, multiply the umol/L value by 0.0113) - k = 0.7 for females and 0.9 for males - a = -0.329 for females and -0.411 for males - min = the minimum of Scr/k or 1 - max = the maximum of Scr/k or 1
002		MDRD	eGFR mL/min/1.73m2=a * Scr**-1.154 * age**-0.203 * [1.210if black] * [0.742if female] Where: - Scr = serum creatinine (mg/dL; to convert from umol/L to mg/dL, multiply the umol/L value by 0.0113) - a = 186 if scr_unit="mg/dL", 32788 if scr_unit="umol/L"
003		Based on Cystatin C	eGFR =133 * (PCysC/0.8)**a * 0.996**Age * (0.932 if female) Where: - PCysC = Plasma Cystatin C in mg/L - a = -0.499 if PCysC <= 0.8 mg/L, -1.328 if PCysC > 0.8 mg/L

Showing 1 to 3 of 3 entries