
```
%Pratap %luitel
%ENGS 111
%HW 7 B
%code representation of the plane is based on
%examples in http://www2.math.umd.edu/~jmr/241/lines planes.html
filename = 'stack2-1.mat';
imIn = load(filename);
stack = imIn.stack2;
[xMax,yMax,zMax] = size(stack);
%region of interest (ROI)
r1 = [100,0,zMax]-[0,100,0];%vector // to the plane
r2 = [xMax - 100,0,0]-[0,yMax,0];%vector // to the plane
normal = cross(r1, r2);%normal to the palne
P1 = [xMax,yMax-100,0];%point on the plane
syms x y z
P = [x, y, z];
planefunction = dot(normal, P-P1);%ROI
zplane = solve(planefunction,z);
ezmesh(zplane, [0, xMax, 0, yMax]);
% roiPlane = zeros(xMax,yMax);
% for i = 1:xMax
      for j = 1:yMax
응
응
          for k = 1:zMax
응
              if(31744*i + 19344*y-10000*z == 11144128)
응
                  roiPlane(i,j) = stack(i,j,k);
%
              end
응
          end
응
      end
% end
```

I calculated two vectors that are parallel to the plane of interest. Based on the question, p1 = (xMax, yMax-100, 0) is a point on the plane. With two vectors parallel to the plane, I calculated the cross product between these vectors to yield a normal to the plane. The normal dotted with the vector(p-p1) defines the plane of interest.

I was not sure about how to project the points on the stack to the plane of interest. Thus, I have only plotted the plane.



Published with MATLAB® R2014a