

Evidence Summary - Technology #03

Catalytic Contact Time Lock for Amyloid Beta Clearance

External Evidence Sources (5 total):

Source 1: Neprilysin-Mediated Amyloid Beta Clearance and Its Therapeutic

URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC11651204>

Excerpt: Neprilysin (NEP) is a neutral endopeptidase, important for the degradation of amyloid beta (A β) cerebrospinal fluid (CSF) that is aimed at removing excess A β ...

Source 2: [PDF] Blood-brain barrier penetrating neprilysin degrades monomeric

URL: <https://uu.diva-portal.org/smash/get/diva2:1651329/FULLTEXT01.pdf>

Excerpt: Neprilysin (NEP) is a membrane-bound metallopeptidase and one of the major A β -degrading enzymes. position of amyloid plaques. Neprilysin-2 is an important β - ...

Source 3: Brain Shuttle Neprilysin reduces central Amyloid- β levels | PLOS One

URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0229850>

Excerpt: Mar 10, 2020 · As previously shown, NEP was highly effective at degrading A β in blood but not in the CSF compartment after systemic administration. In contrast ...

Source 4: Inhibiting Neprilysin Is Good for the Heart: What About the Brain?

URL: <https://www.alzforum.org/news/research-news/inhibiting-neprilysin-good-heart-what-about-brain>

Excerpt: Sep 12, 2014 · LCZ696 contains an inhibitor of neprilysin, an enzyme that breaks down the A β peptide that forms amyloid plaques in the brain.

Source 5: Advances in Amyloid- β Clearance in the Brain and Periphery

URL: <https://www.en-journal.org/journal/view.html?uid=633>

Excerpt: Aug 31, 2023 · The clearance of amyloid- β can occur through enzymatic or non-enzymatic pathways in the brain, including neuronal and glial cells, blood-brain ...

*Evidence Classification: This dossier contains source-native information, derived engineering analysis, engineering proposals, and unresolved questions.
No evidence has been physically validated.*