



Author's reply to commentary on 'altered in vivo early neurogenesis traits in patients with depression: Evidence from neuron-derived extracellular vesicles and electroconvulsive therapy' by Dr. Yakovlev

Dear Editors

In our previous paper [1], Dr. Yakovlev addresses a critical point [2] about CD81 and CD63 Western blot bands appearing at approximately 50 kDa, instead of at 25 kDa and 30 kDa, respectively. Given that small extracellular vesicles (EVs) may contain IgG, we cannot definitively conclude that there is no heavy chain contamination in the CD81 and CD63 blots from our previous paper [1].

To address this concern, we conducted additional Western blot tests for another three EV markers, CD9 (25 kDa, Abcam, Catalog# ab263019), Alix (96 kDa, Proteintech, Catalog #67715-1-Ig), and HSP70 (70 kDa, Elabscience, Catalog# E-AB-22005), which do not have target molecular weights around 50 kDa, to minimize the risk of potential contamination from heavy chains. In addition, although our earlier paper [1] showed NCAM1 bands at the expected molecular weight (93 kDa), we also tested another neuronal marker, SNAP25 (25 kDa, Invitrogen, Catalog # MA5-17610), which do not have target molecular weights around 50 kDa, neither. The results, presented in the supplementary file, showed bands at the expected molecular weights.

For clarity, it's important to note that in our prior work [1], western blotting was used alongside nanoparticle tracking analysis (NTA) and transmission electron microscopy (TEM) solely to verify the presence of neuron-derived EVs (NDEVs). The primary analysis was relied on quantitative data from enzyme-linked immunosorbent assay kits, not on western blotting images.

Finally, although the NDEVs method [3] were widely used [4–7], we are grateful for Dr. Yakovlev's insightful critique, which has contributed to enhancing the reliability of our research.

CRedit authorship contribution statement

Shu-xian Xu: Investigation, Writing – original draft. **Zhongchun Liu:** Funding acquisition, Supervision, Writing – review & editing.

Declaration of competing interest

The authors declare that there is no conflict of interest in this work.

Acknowledgement

This work was supported by grant from the National Natural Science Foundation of China (grant number: U21A20364). This work has not received funding/assistance from any commercial organizations. The funding source had no roles in the design of this study and will not have any roles during the execution, analyses, interpretation of the data, or decision to submit results.

<https://doi.org/10.1016/j.brs.2024.05.001>

Received 30 April 2024; Accepted 2 May 2024

Available online 10 May 2024

1935-861X/© 2024 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brs.2024.05.001>.

References

- [1] Xie X-H, Xu S-X, Yao L, Chen M-M, Zhang H, Wang C, et al. Altered in vivo early neurogenesis traits in patients with depression: evidence from neuron-derived extracellular vesicles and electroconvulsive therapy. *Brain Stimul* 2023;17:19–28. <https://doi.org/10.1016/j.brs.2023.12.006>.
- [2] Yakovlev A. Commentary on the Article titled "Altered in vivo early neurogenesis traits in patients with depression: evidence from neuron-derived extracellular vesicles and electroconvulsive therapy". Xie et al. *Brain Stimul* 2024;17:419–20. <https://doi.org/10.1016/j.brs.2024.03.014>.
- [3] Mustapic M, Eitan E, Werner JK, Berkowitz ST, Lazaropoulos MP, Tran J, et al. Plasma extracellular vesicles enriched for neuronal origin: a potential window into brain pathologic processes. *Front Neurosci* 2017;11:278. <https://doi.org/10.3389/fnins.2017.00278>.
- [4] Jia L, Qiu Q, Zhang H, Chu L, Du Y, Zhang J, et al. Concordance between the assessment of Aβ42, T-tau, and P-T181-tau in peripheral blood neuronal-derived exosomes and cerebrospinal fluid. *Alzheimer's Dementia* 2019;15:1071–80. <https://doi.org/10.1016/j.jalz.2019.05.002>.
- [5] Athauda D, Gulyani S, Karnati H kumar, Li Y, Tweedie D, Mustapic M, et al. Utility of neuronal-derived exosomes to examine molecular mechanisms that affect motor function in patients with Parkinson disease: a secondary analysis of the exenatide-PD trial. *JAMA Neurol* 2019;76:420. <https://doi.org/10.1001/jamaneurol.2018.4304>.
- [6] Dutta S, Hornung S, Kruayatidee A, Maina KN, Del Rosario I, Paul KC, et al. α-Synuclein in blood exosomes immunoprecipitated using neuronal and oligodendroglial markers distinguishes Parkinson's disease from multiple system atrophy. *Acta Neuropathol* 2021;142:495–511. <https://doi.org/10.1007/s00401-021-02324-0>.
- [7] Nasca C, Dobbin J, Bigio B, Watson K, de Angelis P, Kautz M, et al. Insulin receptor substrate in brain-enriched exosomes in subjects with major depression: on the path of creation of biosignatures of central insulin resistance. *Mol Psychiatry* 2021;26:5140–9. <https://doi.org/10.1038/s41380-020-0804-7>.

Shu-xian Xu

Department of Psychiatry, Renmin Hospital of Wuhan University, Wuhan, Hubei, PR China

Zhongchun Liu*

Department of Psychiatry, Renmin Hospital of Wuhan University, Wuhan, Hubei, PR China

Taikang Center for Life and Medical Sciences, Wuhan University, Wuhan, PR China

* Corresponding author. Department of Psychiatry, Renmin Hospital of Wuhan University, No. 99 Jiefang Road, Wuchang District, Wuhan, Hubei, PR China, 430060.

E-mail address: zcliu6@whu.edu.cn (Z. Liu).

Handling Editor: M. George