

Juvenile proportions GAM by zone

Effects of year




Figure 1: Lemming cycle effect appears dampened in more recent years.


With random effects

```
##
## Family: binomial
## Link function: logit
##
## Formula:
## propjuv ~ s(Breeding_year) + s(Breeding_year, by = zone1) + s(days) +
##       s(days, by = zone1) + s(Food_type, Breeding_year, bs = "re") +
##       s(Observer, Breeding_year, bs = "re") + s(flocksize, Breeding_year,
##       bs = "re")
##
## Parametric coefficients:
##               Estimate Std. Error z value Pr(>|z|)
## (Intercept) -1.54149    0.04969  -31.02   <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Approximate significance of smooth terms:
##                                     edf Ref.df Chi.sq  p-value
## s(Breeding_year)                   8.1077172   8.766  67.784 5.35e-11 ***
## s(Breeding_year):zone1IJsselmeer  1.0000483   1.000   0.021  0.88471
## s(Breeding_year):zone1Rhinelands  1.0000359   1.000   0.831  0.36214
## s(Breeding_year):zone1Southwest   1.0000200   1.000   2.381  0.12282
## s(days)                           1.0000074   1.000   0.851  0.35634
## s(days):zone1IJsselmeer           1.0000622   1.000   0.284  0.59434
## s(days):zone1Rhinelands           1.0001023   1.000   0.327  0.56770
```


Juvenile proportions GAM by zone

Effect of year and days by zone

juvenile proportion



juvenile proportion



Including mixed effects

- Including mixed effects reduces AIC significantly.
- AIC (GAM without mixed effects): 2490.3678094
- AIC (GAM with mixed effects): 2133.6031601